



The Sauzei-/Humphriesianum zonal boundary (Lower Bajocian, Middle Jurassic) in the type area of the Gosheim Formation (W Swabian Alb, SW Germany)

Authors: Dietze, Volker, Schreiber, Dietmar, and Schweigert, Günter

Source: Palaeodiversity, 15(1) : 13-43

Published By: Stuttgart State Museum of Natural History

URL: <https://doi.org/10.18476/pale.v15.a2>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

The Sauzei-/Humphriesianum zonal boundary (Lower Bajocian, Middle Jurassic) in the type area of the Gosheim Formation (W Swabian Alb, SW Germany)

VOLKER DIETZE, DIETMAR SCHREIBER † & GÜNTER SCHWEIGERT

Abstract

The Bajocian ammonite faunas of the ‘Oberer Blaukalk’ Member (Wedelsandstein Formation) and of the lower part of the ‘Humphriesioolith’ Member (Gosheim Formation) in the vicinity of Gosheim (SW Swabian Alb) are studied. At least three biohorizons of the Sauzei Zone can be distinguished: *dilatus*, *pseudocontrahens*, *macrum* and *?carinodiscus* biohorizon, probably followed by the *deltafalcata* biohorizon of the basal Humphriesianum Zone (Pinguis Subzone). This succession is correlated with other sites in SW Germany and abroad.

Key words: Sauzei Zone, Humphriesianum Zone, ammonite biohorizons, SW Germany, Bajocian.

In memoriam NORBERT WANNENMACHER (1961–2020)

1. Introduction

In the Swabian Alb, the middle part of the Braunjura Group is mostly formed by claystones and often covered by thick hillside debris from overlying Upper Jurassic limestones. An exception is the vicinity of the village of Gosheim in the western part of the Swabian Alb. There, the Braunjura forms a steep slope and an adjacent wide plane. In the Bajocian, relatively thick iron oolites occur; the iron content of which, however, is too low for successful industrial exploitation, although some attempts have been undertaken in the 19th century (see WANNENMACHER 2017). FISCHER (1912, 1924), WEISERT (1932) and BERZ (1933) analysed in detail the succession and macrofossil content of the ‘Humphriesioolith’ in this area, which is today a member of the lower Bajocian Gosheim Formation (DIETZE et al. 2015). Modern descriptions of the lithological succession and ammonites of the ‘Humphriesioolith’ and the underlying ‘Oberer Blaukalk’ members can be found in DIETL (1977), DIETL & RIEBER (1980), FRANZ (1988) and OHMERT (1990). DIETL (1978) was the first researcher, who recognized that basal parts of the ‘Humphriesioolith’ Member (Fig. 1) still belong to the Sauzei Zone and not to the Humphriesianum Zone as often stated in generalized mainstream literature (e.g., GEYER & GWINNER 1984, 2011). More recently, DIETZE et al. (2008) introduced the *macrum* Biohorizon (Sauzei Zone), which comprises the beds with *Skirroceras macrum* in the basal part of the ‘Humphriesioolith’ Member at Gosheim. In addition to that preliminary report, we here complete our study of the complex bio- and lithostratigraphy at the Sauzei-/Humphriesianum zonal boundary in the type area of the Gosheim Formation.

2. Material and methods

Ammonites are rare in the basal part of the ‘Humphriesioolith’ Member at Gosheim. Due to this scarcity, a systematic bed-by-bed sampling of specimens is almost impossible. Only by chance single ammonites can be recovered directly from temporary sections exposed in construction pits, sewer trenches or other excavations. Most outcrops were accessible only for a short time, and we were not always informed quick enough about the opening of such outcrops. The excavated material was usually transported to the local earth dump of Gosheim, where it was levelled. Reconstruction of the origination of loose material from there is only possible for material with very characteristic lithologies such as the ‘Spathulatusbank’ or the ‘Oberer Blaukalk’ Member. Specimens that originate from higher parts of the ‘Humphriesioolith’ Member collected at the earth dump can hardly be assigned precisely to the section since the lithology within this part of the member is too uniform. When sometimes the excavated material was deposited in numerous gigantic boulders, it became obvious that ammonites from the basal part of the ‘Humphriesioolith’ Member are even extremely rare. Despite these enormous difficulties, several age-diagnostic ammonites could be sampled from their bed or at least with detailed information over the last 30 years. This material allowed a high-resolution biostratigraphical subdivision of the basal part of the ‘Humphriesioolith’ Member into biohorizons (for methodology, see CALLOMON 1995). Further material originates from the same interval, but could not be assigned precisely to these biohorizons. Deciphering information from specimens and outcrops was necessary to get an idea about the complete biostrati-




Fig. 1. Middle Jurassic beds in SW Germany. The herein studied location Gosheim in the western part of the Swabian Alb is indicated by an arrow.

graphic succession in the area. Additionally, in the basal part of the 'Humphriesioolith' Member the lithological succession and the thicknesses of individual beds change laterally rapidly. In fresh state, the 'Humphriesioolith' Member often appears as a homogenous unit; only after weathering individual beds are distinguishable. This phenomenon hampers both a detailed measuring of the sections and the correlation of individual beds.

For a better comparison among each other we illustrated all ammonites on the plates in half of their size. Only the smaller specimens of Fig. 3 were illustrated in natural size. The studied material is stored in the collection of the Stuttgart Natural History Museum (acronym SMNS).

3. Outcrops and sections

We here describe outcrops at Gosheim which had provided relevant stratigraphic data over the last decades,

especially precisely collected ammonites. Additionally, we briefly describe further sections and outcrops mentioned in the literature, from where biostratigraphically relevant ammonites are stored in various collections. Since the lithological succession at Gosheim is principally similar in all outcrops except that within the industrial area Sturmbühl, we here provide a slightly idealized section (Fig. 2).

3.1. Brücklestraße (extension of Grimm Company)

In this construction pit, the 'Oberer Blaukalk' Member and the lower 4.5 m of the 'Humphriesioolith' Member were exposed. Since this succession was very heterogeneous, we do not provide a detailed description of the section. Two age-diagnostic ammonites were recovered from the bed: *Emileia vagabunda* (Pl. 1, Figs. 5, 6) and *Skirroceras* cf. *nodosum* (Pl. 6, Figs. 1, 2). Although the excavated material was deposited for months on the earth dump




Fig. 3. Ammonites of the Gosheim Formation, ‘Humphriesiolith’ Member, Lower Bajocian, Humphriesianum Zone, Pinguis Subzone, *?deltafalcata* biohorizon, Gosheim, road near ski lift [leg. N. WANNENMACHER †]. (1) *Pelokodites sulcatum* (BUCKMAN, 1893), a: lateral view, b: ventral view, SMNS 70635/1. (2) “*Dorsetensia*” sp., a: lateral view, b: ventral view, SMNS 70635/2. (3, 4) “*Dorsetensia*” *punctatissima* (HAUG, 1893), (3a, b) SMNS 70635/3 (4a, b) SMNS 70635/4. Scale bar equals 1 cm. All ammonites in natural size. Beginning of the body chamber is indicated by an asterisk.

recorded at the basis of the ‘Humphriesiolith’ Member, followed by an alternation of rusty red to bluish, partly iron-oolitic marly limestones, calcareous marls and marls. In this construction pit, we collected two ammonites from a marly, strongly iron-oolitic horizon c. 1.8 m above the basis of the ‘Humphriesiolith’ Member: *Skirroceras macrum* (Pl. 3, Figs. 1, 2) and *Sonninia* cf. *patella* (Pl. 4, Fig. 1).

3.3. Embankment of street between Hermle Company and ski lift

Unfortunately, we noticed this temporary outcrop when the sewer trench was refilled and the street was completed. Therefore, it was only possible to measure the section along the embankment of the street. The ‘Humphriesiolith’ Member was remarkably thick, but its basis was no longer exposed. Most likely, however, the basis of this member was only a little below the basis of our measured section. A *Pelokodites sulcatum* specimen was collected from the screw, but close to the exposed beds assigned to the *?deltafalcata* biohorizon. From the base to the top:

Gosheim Formation

‘Humphriesiolith’ Member

c. 3.5 m alternation of reddish to gray and gray-brown, usually strongly iron-oolitic, more or less lithified calcareous marls and marly limestones and softer iron-oolitic marls.

c. 0.6 m [*?deltafalcata* biohorizon]: soft, iron-oolitic marls with scattered marly limestone nodules. Shell fragments of *Ctenostreon* and other bivalves (*Pinna*, *Oxytoma*) and echinoid spines. Ammonites: “*Dorsetensia*” sp., “*D.*” *punctatissima*.

c. 3.10 m: alternation of iron-oolitic beds of calcareous marl, marly limestone and marl.

0.15 m [*gervillii/cycloides* biohorizon]: gray clayey marl with re-sedimented nodules. Ammonites: *Chondroceras* sp., *Itinsaites* sp.

0.4 m [*umbilicum* biohorizon]: irregularly bedded rusty red to brown iron-oolitic marl, rich in fossils.

Ammonites: *Stephanoceras* sp., *Itinsaites* sp.

0.3–0.4 m: three hard gray-brown to greenish iron-oolitic marly limestone beds; *Actinostreon* and belemnites abundant.

‘Blagdenischichten’ Member [*coronatum* biohorizon]

0.9 m: gray marl with *Actinostreon*, large belemnites and brachiopods.

0.1–0.12 m: gray-beige marly limestone.

0.45 m: gray marl, *Actinostreon*, belemnites.

0.1–0.12 m: gray-beige marly limestone. Ammonite: *Teloceras* sp.

3.4. Development area at Längenbergl (sewer trenches and construction pits)

In this development area, the ‘Oberer Blaukalk’ Member and max. 1.5 m of the overlying basal ‘Humphriesiolith’ Member were exposed. Higher parts of the ‘Humphriesiolith’ Member are eroded here.

Wedelsandstein Formation

‘Unterer Blaukalk’ Member, ‘Obere γ -Tone’ Member

In one of the sewer trenches a c. 1 m thick claystone and a 0.2 m thick limestone (‘Unterer Blaukalk’ Member) was exposed, followed by several metres of claystone and a thin marl bed. A connection to higher parts of the section was impossible. Starting with the ‘Oberer Blaukalk’ Member an exact section could be measured.

‘Oberer Blaukalk’ Member

0.35–0.4 m: bluish-gray calcareous sandstone, with light brown weathering crust, slightly micaceous. In the upper part a layer with flat, centimetre-sized phosphatic pebbles. At the base and at the top with abundant phosphatic body chamber fragments of *Kumatostephanus*. Ammonites: *Kumatostephanus* sp., *Emileia* sp., *Sonninia* sp.

0.18 m: gray brown clay marl, no macrofossils.

0.1–0.15 m: fossil-rich, strongly bioturbated finesandy calcareous marl with flaser bedding; on its top a hardground with belemnites, serpulids and bivalves.

Gosheim Formation**‘Humphriesioolith’ Member**

Max. c. 1.3–1.4 m exposed. Above a reddish iron-oolitic marl (c. 0.05 m), which often laterally disappeared, a single massive limestone bed (c. 1.3 m) followed which splitted into numerous layers in weathered state. In the basal 0.45 m a typical gray to reddish sparitic ‘Spathulatusbank’ lithology occurs with abundant, partly densely packed bivalve shells. Above, the first iron ooids appear; from 0.7 m above the basis onwards, the ooids become more abundant but are smaller than the deeper ones. The bivalve content decreases towards the top. The lithology of the limestone is still sparitic but becomes softer, with increasing ooid content.

Ammonites [from the ‘Spathulatusbank’]: *Emileia schassmanni* (Pl. 2, Figs. 4, 5), *Sonninia* sp.

3.5. Industrial area Sturmbühl (development trenches, construction pits)

Since the opening of this small industrial area several decades ago, sections of the ‘Oberer Blaukalk’ Member and basal parts of the ‘Humphriesioolith’ Member were repeatedly exposed. Despite intensive sampling in some of those very big outcrops, we did not find any ammonites there. However, we noticed a few specimens collected by amateurs in their private collections, among them *Sonninia* cf. *patella* (Pl. 7, Figs. 2, 4), *S. carinodiscus* (Pl. 8, Figs. 1, 2) and *Emileia arkelli* (Pl. 2, Figs. 2, 3) as well as two moderately preserved *Skirroceras macrum* (former BERNT † collection). In contrast, ammonites are quite abundant in the higher parts of the ‘Humphriesioolith’ Member in the industrial area Sturmbühl (see DIETZE et al. 2015).

In 2020, during the construction of an over 2.000 m² big area with commercial garages, we intensively searched for ammonites; however, only a single specimen came to light. Since the section in the lower part of the ‘Humphriesioolith’ Member is distinct from sections in the village Gosheim itself, we here provide the section of this outcrop. In the shredded excavation of a nearby small construction pit we found (in 2021) only two remains of *Skirroceras macrum* or *S. nodosum* preserved in a very hard gray, sparitic limestone.

Wedelsandstein Formation

‘Ober γ-Tone’ Member (min. 1.5 m): gray-blue micaceous silty to sandy clay marl.

‘Oberer Blaukalk’ Member (0.45 m): 0.35 m sandy limestone, followed by 0.1 m gray marl.

Gosheim Formation**‘Humphriesioolith’ Member (c. 4.2 m exposed)**

In the industrial area of Sturmbühl NE of Gosheim, the basal part of the ‘Humphriesioolith’ Member differs from all other outcrops in Gosheim. The basal part is made up here of thick gray, sparitic limestones.

0.6 m: massive gray, sparitic limestone, red-brown in weathered state.

0.25 m: hard flaser-bedded gray calcareous marl.

0.4 m: iron-oolitic calcareous marl splitting into several layers, with few *Entolium demissum*.

0.15 m: gray limestone with yellow-brown ooids.

0.17 m: soft, gray beige marl.

0.25 m: iron-oolitic marly limestone.

0.2 m: rusty red iron-oolitic marl with *Ctenostreon* and echinoid spines.

0.15 m: rusty red iron-oolitic limestone.

0.1 m: rusty red iron-oolitic limestone.

0.2 m: fine, gray beige marl.

0.15 m: hard gray sparitic limestone.

0.5 m: reddish, flaser-bedded ironoolithic calcareous marl/marly limestone.

0.4 m: rusty red soft ironoolithic marl with echinoid spines, *Ctenostreon*, “*Ostrea*” *eduliformis*, molds of endobenthic bivalves, and belemnites.

0.05–0.1 m: hard ironoolithic marly limestone.

0.15 m: gray clay marl.

0.15 m: gray-brown to beige, thinly splitting marly limestone.

0.4 m: soil with rock fragments and an ammonite (*Dorsetensia liostraca*).

3.6. Autunnel (FISCHER 1924, WEISERT 1932, DIETL 1978, railway cut)

Here we repeat the section taken by DIETL (1978) at the Autunnel. The ‘Humphriesioolith’ Member is more or less well-exposed in the railway cut from above the Autunnel to the village Gosheim.

Wedelsandstein Formation

‘Obere γ-Tone’ Member (c. 3.2 m exposed): dark gray to bluish gray, fine sandy micaceous claystone or clay marl; with a marly bed c. 2.5 m below the top.

‘Oberer Blaukalk’ Member (0.45 m): blue-gray, slightly micaceous calcareous sandstone followed by a fine sandy, micaceous clay marl bed (“Tonmergelhorizont”, c. 0.1 m).

Gosheim Formation

'Humphriesioolith' Member

(c. 0.15–0.2 m) ['Spathulatusbank']: blue-gray, fine sandy, sparitic, slightly iron-oolitic limestone; *Entolium demissum* (PHILLIPS) (= *Pecten spathulatus* ROEMER) very abundant in layers. The bed becomes more marly towards its top. Ammonite: *Sonninia* ex gr. *patella* (see DIETL 1978).

(c. 1.3 studied): 'Fe-oolithische Kalkmergel', rusty red calcareous marl to marl, remarkably sparitic in the basal part.

3.7. Construction of the underground car park at the Kreissparkasse

A section of this outcrop has not been documented. There, the *macrum* biohorizon was relatively rich in ammonites since most specimens of *Skirroceras macrum* and *S.* cf. *macrum* in various private collections originate from this site. During the excavations, the attentive operator recovered about ten large specimens of *Skirroceras* and a *Lytoceras* sp. and sold them to a fossil collector, who subsequently sold or swapped them to other collectors (F. FANZUTTI, Bisingen-Thenheim, pers. comm. September 2021). One of these specimens, a *Skirroceras* cf. *macrum*, is illustrated on Pl. 4, Fig. 2.

3.8. Denkinger Steige (FRANZ 1988; natural outcrop)

The section in FRANZ (1988) was taken along the Denkinger Steige (M. FRANZ, pers. comm. September 2021). The *Kumatostephanus triplicatus* specimen mentioned by FRANZ (1988) from the basis of the 'Oberer Blaukalk' Member is illustrated here on Pl. 1, Figs. 1, 2. At the Denkinger Steige the 'Humphriesioolith' Member has a thickness of c. 6.5 m.

3.9. Gosheim (OHMERT 1990)

OHMERT (1990: 129; fig. 7) published a continuous section from the 'Oberer Blaukalk' up to the basis of the 'Blagdenischichten' at Gosheim. It is not mentioned in his study, where exactly this section has been measured, or if it is a compound section from different places. Moreover, it is not explained on which ammonite material his biostratigraphy of the 'Humphriesioolith' Member was based. Unfortunately, no material from that section could be traced in the collection of the Geologisches Landesamt in Freiburg (M. FRANZ, pers. comm. October 2021). OHMERT's (1990) stratigraphy of the beds below the Romani Subzone (Humphriesianum Zone) is inconsistent and cannot be confirmed by our studies (see Fig. 2). Based on DIETL (1978), he drew the position of the Sauzei/Humphriesianum zonal boundary less than one metre above

the basis of the 'Humphriesioolith' Member. He assumed the type horizon of *Skirroceras macrum* and *S. plicatum* (after WEISERT 1932) to be the *frechi* biohorizon of the Pinguis Subzone ["In the section of Gosheim this horizon [= *frechi* biohorizon] may be comprised in the lower part of the Humphriesi-Oolith (4–4.5 m in fig. 7) with *Skirroceras macrum* (QUENSTEDT) and *Stephanoceras plicatum* (QUENSTEDT) (see WEISERT 1932: 184, 185)"]. In contrast, in the Lörrach section, he assigned the beds with *S. macrum* to the Sauzei Zone and the bed interval from 4–4.5 m in the Gosheim section following below the *frechi* biohorizon still to the *pinguis* biohorizon (OHMERT 1990: fig. 7).

3.10. Wassersteige (DIETL & RIEBER 1980; natural outcrop)

DIETL & RIEBER (1980) described the section at the Wassersteige, along the road connecting the villages Wilflingen and Gosheim:

Wedelsandstein Formation

'Oberer Blaukalk' Member (0.5 m): hard calcareous sandstone followed by a few centimetres of clay marl.

Gosheim Formation

Humphriesioolith Member (c. 6.5 m): At the basis c. 0.15 m 'Spathulatusbank'. The beds following above are more or less iron-oolitic. The ooids are fine in the basal part and become coarser up-section. A specimen of *Sonnina pseudofurciculata* from the BERNT † collection that originates from this outcrop is illustrated here (Pl. 9, Figs. 3, 4). The ammonites from the 'Humphriesioolith' Member mentioned by DIETL & RIEBER (1980) originate from another place.

4. Remarks on the ammonite fauna

Since we have only few biostratigraphically significant specimens, we preferred a purely morphospecific determination of the ammonite fauna.

4.1. Family Stephanoceratidae NEUMAYR, 1875

4.1.1. Subfamily Kumatostephaniinae CHANDLER, DIETZE & WHICHER, 2017

In the 'Oberer Blaukalk' Member, occasionally body chambers of *Kumatostephanus triplicatus* (RENZ, 1904) (Pl. 1, Figs. 1, 2) and *K.* cf. *perjucundus* BUCKMAN, 1927 (Pl. 2, Fig. 1) occur. Both species are very close to each other; however, *K. triplicatus* exhibits a somewhat broader and lower whorl section than *K. perjucundus*. A larger

ammonite (Pl. 1, Figs. 3, 4) with a rounded whorl section, slightly prorsiradiate and curved ribs is assigned to *K. aff. turgidulus* (QUENSTEDT, 1886). *K. turgidulus* s.str. differs by a more radiate ribbing style. *K. kumaterus* BUCKMAN, 1922 is denser ribbed than *K. aff. turgidulus* and has shovel-like primaries that begin radiate from the umbilical edge onwards and become convex at a deep point of the flank.

4.1.2. Subfamily Stephanoceratinae NEUMAYR, 1875

Stephanoceratids of the lower part of the 'Humphriesioolith' Member are solely represented by the genus *Skirroceras* MASCKE, 1907. A complete, precisely collected specimen of *Skirroceras macrum* (QUENSTEDT, 1886) (Pl. 3, Figs. 1, 2) corresponds perfectly to the lectotype of *Ammonites humphriesianus macer* QUENSTEDT, 1886 (pl. 65, fig. 11), which originates from an unknown Swabian locality. Despite the preoccupation of QUENSTEDT's third name, the taxon *S. macrum* has been validated (see ICZN 2005). The lectotype and all studied specimens of *S. macrum* and *S. cf. macrum* from Gosheim are well-preserved only on their stratinomic lower face, whereas the upper face is partly eroded. Sometimes the body chamber is more or less compressed (e.g., Pl. 4, Fig. 2). The description of *S. macrum* by WEISERT (1932) is very accurate; we can only add here that the maximum diameter of this species reaches at least 38 cm.

We concur with WEISERT's (1932) observation that morphological transients exist between *S. macrum* and *S. nodosum* (QUENSTEDT, 1857), although both taxa seem to be stratigraphically distinct from each other. The innermost whorls up to a diameter of 16–18 mm are almost identical in both species (WEISERT 1932): they are trapezoidal with hardly rounded ventrolateral margin. The whorls are arranged in a strongly involute style, with short, moderately deepened flanks, which are sculptured by sharp rectiradiate ribs. At mid-flank elevated pointed nodes with a broad basis are developed which gave rise to the secondary ribs. The broad-sectioned, strongly sculptured stage persists long in ontogeny of *S. nodosum* s. str., the whorl section of *S. macrum* becomes rounded in an earlier stage, the umbilicus widens and the conch becomes disc-shaped. The umbilical wall is rounded and relatively shallow. In contrast, the whorl section of *S. nodosum* (QUENSTEDT, 1857) is broad (Pl. 5, Fig. 1; Pl. 6, Figs. 3, 4) and the outer whorls still have relatively steep flanks, so that the conch has the appearance of a soup plate, whereas it is disc-shaped in *S. macrum*. In *S. nodosum* prominent high nodes occur which weaken towards the body chamber (Pl. 6, Figs. 3, 4) or persist until close to the aperture in some extreme variants (Pl. 5, Fig. 1), whereas in *S. macrum* the nodes become blunter in an earlier stage and form only shallow elevations on the body chamber. The adult

size of *S. nodosum* reaches little more than 30 cm; hence, it is much smaller than *S. macrum*.

The *Skirroceras* cf. *macrum* (QUENSTEDT, 1886) of Pl. 4, Fig. 2 retains the trapezoidal, strongly nodate stage longer than in *S. macrum* and even reiterates this character after the rounding of the umbilical edge. However, these characters are less prominently developed than in *S. nodosum*. Moreover, this specimen exhibits shovel-like nodes. The primary ribs are slightly retroradiate and continue after these nodes as rectiradiate secondaries. Although the whorl section of the ammonite illustrated on Pl. 4, Fig. 2 is less rounded in the area of the nodes than in the lectotype of *S. macrum*, this specimen is relatively evolute and does not show the typical deep umbilicus of *S. nodosum*. MAUBEUGE (1961, figs. on p. 121) determined a very similar specimen from the Sauzei Zone of northern Switzerland as *S. cf. macrum*.

Skirroceras cf. *nodosum* (QUENSTEDT, 1857) as illustrated on Pl. 6, Figs. 1, 2 shows inflated massive whorls and a smaller umbilicus like in *S. nodosum*, but it shares a shallowing of the nodes and the ribbing on the outer whorls with *S. macrum* without becoming disc-shaped. On the outer whorls, the umbilical wall is still rounded, but the umbilicus is depressed.

Skirroceras rochei MAUBEUGE, 1961 is another transitional form between *S. macrum* and *S. nodosum*. *S. thoralis* MAUBEUGE, 1961 is based on the badly preserved incomplete holotype (ROCHÉ, 1939, pl. 5, fig. 1) and thus hardly interpretable. Most likely, this species is just an extreme variant of *S. macrum*. Other nominal species of *Skirroceras* are hardly confusable with the material from the Sauzei Zone of Gosheim.

Two further specimens not collected bed-by-bed are determined as *Skirroceras* sp. (Pl. 5, Fig. 2) und *S. cf. freycineti* (Pl. 5, Figs. 3, 4). Both specimens originate from the lowermost 4.5 m of the 'Humphriesioolith' Member, since higher beds were not exposed at the finding site. After their mode of preservation, they originate from a different level than that of *S. macrum* or *S. nodosum*, supposedly from somewhat higher beds. *S. cf. freycineti* (Pl. 5, Figs. 3, 4) differs from true *S. freycineti* (BAYLE, 1878) by its wider-spaced ribbing. The nodes on the diverging points are more prominent than they appear in our illustration. The ammonite of Pl. 5, Fig. 2 exhibits a remarkably prorsiradiate ribbing with prominent nodes after a pathological stage. Due to this pathological sculpture, a specific determination of this *Skirroceras* is impossible.

4.2. Family Otoitidae MASCKE, 1907

Otoitids are extremely rare in the 'Oberer Blaukalk' and in the basal 'Humphriesioolith' Member, so that every record deserves our attention:

D. SCHREIBER (†) found the phosphatic body chamber of an *Emileia* sp. in the earth dump of Gosheim (extracted material from the development area at Längenbergraben). This ex-situ find originates from the 'Oberer Blaukalk' Member. The whereabouts of this specimen is unknown.

From the basal 'Humphriesioolith' Member (Sauzei Zone), we know of only four specifically identifiable specimens:

A specimen of *Emileia arkelli* MAUBEUGE, 1961 (Pl. 2, Figs. 2, 3) was collected from the screw during the development of the industrial area Sturmbühl. The specimen is preserved as a strongly iron-oolitic mold. This lithology occurs in the interval up to c. 2–3 m above the 'Spathulatusbank', which is not iron-oolitic and hence can be excluded as its finding level. In the *macrum* biohorizon of entire SW Germany, only a single large, densely ribbed *Emileia* specimen has been recorded from Ringsheim (Upper Rhinegraben) (see DIETZE et al. 2009). Hence, the *E. arkelli* specimen can be most likely assigned to the *pseudocontrahens* biohorizon, and all other biohorizons can be excluded.

A very large specimen of *Emileia vagabunda* BUCKMAN, 1927 (Pl. 1, Figs. 5, 6; maximum diameter 32 cm) is represented by a body chamber, which comprises a little more than one whorl and is slightly abraded in its anterior part. The similarly large-sized taxa *E. bulligera* BUCKMAN, 1927 and *E. greppini* MAUBEUGE, 1961, occurring in the same biostratigraphic level, are surely only variants of the same palaeobiospecies. Compared with *E. vagabunda*, *E. bulligera* is coarser ribbed and exhibits a broader whorl section; in *E. greppini* the ribbing is wider-spaced and the whorl height is slightly lower. Our *E. vagabunda* specimen illustrated on Pl. 1, Figs. 5, 6 was collected from just below the basis of the 'Spathulatusbank'. Some preserved rock matrix shows the typical lithology of the 'Spathulatusbank'; however, the strongly iron-oolitic filling of the body chamber differs from that lithology.

Another very large specimen of *Emileia schassmanni* MAUBEUGE, 1961 (Pl. 2, Figs. 4, 5; reconstructed final diameter c. 33 cm) originates from the 'Spathulatusbank'. This ammonite shows a remarkably broad whorl section. The primary ribs are densely spaced from the inner whorls onwards up to the body chamber. This ammonite is relatively involute because of its large whorl height. A short portion of the body chamber close to the aperture is still present but so badly preserved that we omitted this part from the illustration. *E. fuellinsdorfense* MAUBEUGE, 1961 and *E. pseudocontrahens* MAUBEUGE, 1961 are similarly fine-ribbed, but with a greater whorl section. Moreover, *E. pseudocontrahens* is significantly smaller than our specimen.

DIETL (1978) identified a well-preserved historical *Emileia* specimen from Gosheim as *E. (E.) cf. "polymera"* (WAAG.). After the rock matrix he assumed the lowest part

of the 'Humphriesioolith' Member as the finding horizon. We concur and can add that this ammonite must originate from the interval up to max. 2.5 m above the 'Spathulatusbank', even more likely from the lower 1.5 m of this member. Three further, badly preserved specimens of *Emileia* (priv. coll. V.D., SMNS) have not been collected bed-by-bed and thus cannot be determined specifically.

4.3. Family Sonninidae BUCKMAN, 1892

Only a single *Sonninia* (Pl. 3, Figs. 1, 2) was exactly sampled from an exposure of the basal 'Humphriesioolith' Member at Gosheim. All other specimens originate from the screw.

A *Sonninia* cf. *patella* (WAAGEN, 1867) (Pl. 4, Fig. 1) originates from a level c. 1.8 m above the basis des 'Humphriesioolith' Member; this is the same level as in the case of the *Skirroceras macrum* illustrated on Pl. 3, Figs. 1, 2. A further, much better preserved *Sonninia* cf. *patella* (Pl. 7, Figs. 2, 4) and a more depressed and more evolute *S. carinodiscus* (QUENSTEDT, 1886) (Pl. 8, Figs. 1, 2) as well as a specimen of *S. ("Sonninites") aff. felix* (BUCKMAN, 1923) (Pl. 7, Figs. 1, 3) originate from a not exactly known level within the interval above the 'Spathulatusbank' and max. 3–4 m higher up in the 'Humphriesioolith' Member.

Four additional sonniniids without precise information about their finding levels are determined as *S. pseudofur-ticarinata* MAUBEUGE, 1961 (Pl. 9, Figs. 1–4) and *S. alsatica* (HAUG, 1885) (Pl. 9, Figs. 5, 6), respectively.

A *Dorsetensia liostraca* BUCKMAN, 1892 (Pl. 8, Figs. 3, 4) was sampled from the lower part of the 'Humphriesioolith' Member at the Autunel. It originates from a bed located higher than c. 3.5 m above the basis of the member and which must be assigned already to the Romani Subzone (Humphriesianum Zone). QUENSTEDT (1886, pl. 63, fig. 7) illustrated a nicely preserved specimen of *D. liostraca* from the 'Humphriesioolith' at the nearby town Spaichingen.

The small-sized faunule questionably assigned to the *deltafalcata* biohorizon is not easy determinable both at genus and species level. We identified the ammonites of Figs. 3.3–3.4 as "*Dorsetensia punctatissima* (HAUG, 1893). A whorl fragment (Fig. 3.2) can be only determined as "*Dorsetensia*" sp. An ammonite from the screw (Figs. 3.1) which was found very close to this small-sized faunule represents *Pelekodites sulcatum* (BUCKMAN, 1893).

4.4. Family Hammatoceratidae BUCKMAN, 1887

A large-sized ammonite (Pl. 9, Figs. 7, 8) collected from the interval of c. 3 m up to max. 4.5 m above the basis

Lower Bajocian	Zone	Subzone	Biohorizons
	Humphriesianum	Blagdeni	<i>coronatum</i> <i>acuticostatum</i> <i>unnamed</i>
		Humphriesianum	<i>crassicosta</i> <i>umbilicum</i> <i>mutabile</i> <i>gerivillii/cycloides</i>
		Romani	<i>romani</i> <i>cf. complanata</i>
		Pinguis	? <i>deltafalcata</i> <i>frechi</i> <i>pinguis</i> ?
	Sauzei		? <i>carinodiscus</i> <i>macrum</i> <i>pseudocontrahens</i> <i>quenstedti</i> <i>dilatatus</i>

Fig. 4. Biohorizons of the Sauzei and Humphriesianum zones in SW Germany. The biohorizons recorded at Gosheim are marked in gray; uncertain records are indicated by a question mark.

of the ‘Humphriesioolith’ Member exhibits a fissilobate suture line and a high-oval whorl section with a rounded venter. Thus, we identified it as *Fissilobicerias furticarinum* (QUENSTEDT, 1856). The pyritic holotype of this species (re-figured in DIETZE et al. 2011, pl. 9, figs. 1, 3) originates from the Humphriesianum Zone (Pinguis or Romani Subzone) of Pfullingen near Reutlingen. A further specimen of *F. furticarinum* illustrated by QUENSTEDT (1886, pl. 68, fig. 5; re-figured in DORN 1935, pl. 20, fig. 1) from the same bed and locality shows bulging ribs like in the specimen from Gosheim (in the latter these ribs occur at the end of the phragmocone). Similar ribs occur in the lectotype of *F. fissilobatum* (WAAGEN, 1867; see photograph in SADKI & DIETZE 2020, fig. 10.A1, A2) from the Trigonalis Subzone of the Laeviuscula Zone. *F. fissilobatum* is the type species of *Fissilobicerias* BUCKMAN, 1919.

5. Bio-/chronostratigraphy and correlation of the Gosheim Formation

5.1. Biostratigraphy

The herein distinguished biohorizons are not recorded in every studied outcrop. It seems that some beds are not continuous over larger distances but form lenses. This is not surprising since the lower part of the ‘Humphriesioolith’ Member represents a locally diversified shallow water deposit (DIETZE et al. 2008). However, in cases where individual biohorizons are recorded, they occur in approximately the same vertical position within the ‘Humphriesioolith’ Member. As a result, although somewhat idealized, Fig. 4 shows an update of the biostratigraphical succession in the lower part of the ‘Humphriesioolith’ Member at Gosheim.

5.1.1. Laeviuscula Zone

It is unknown if the ‘Unterer Blaukalk’ Member at Gosheim (c. 4 m below the ‘Oberer Blaukalk’) with *Emileia polyschides*, *Otoites* sp. and *Sonninia* sp. still belongs to the Laeviuscula Zone or to the Sauzei Zone, since the specimens reported by FISCHER (1924) could not be traced anywhere.

5.1.2. Sauzei Zone

dilatus biohorizon – The ‘Oberer Blaukalk’ Member yields *Kumatostephanus triplicatus*, *K. cf. perjucundus*, *K. aff. turgidulus*, *Emileia* sp. and *Sonninia* sp. and is assigned to the *dilatus* biohorizon. In SW Germany, ammonites of the genus *Kumatostephanus* are only abun-

dant in this biohorizon (DIETZE et al. 2020). However, specimens of *Emileia* and *Sonninia* sampled from this level are not specifically determinable.

pseudocontrahens biohorizon – The *pseudocontrahens* biohorizon is proved by *Emileia vagabunda* and *E. schassmanni*. Both taxa occur in the interval of the only partly developed iron-oolitic basal marl layer of the ‘Humphriesioolith’ and in the ‘Spathulatusbank’. Most probably, this biohorizon expands up to 1.7 m above the basis of the ‘Humphriesioolith’ Member, since the oldest record of *Skirroceras macrum* appeared c. 1.8 m above this basis. Hitherto, large-sized species of *Emileia*, like *E. vagabunda* and *E. schassmanni*, have been reported from SW Germany only in the *pseudocontrahens* biohorizon. In the next-older *quenstedti* biohorizon, specimens of *Emileia* are significantly smaller and belong to different species (DIETZE et al. 2020). From the next-younger *macrum* biohorizon, only a single specimen of *Emileia* has been reported from the entire area of SW Germany (DIETZE et al. 2009).

macrum biohorizon – The bed interval c. 1.8–2 m above the basis of the ‘Humphriesioolith’ Member represents the *macrum* biohorizon. At Gosheim, this biohorizon is recorded by *Skirroceras macrum*, *S. cf. macrum* and *Sonninia patella*. However, it is unclear whether the herein illustrated sonniniids originate from this biohorizon as well, and we do not know if *Skirroceras nodosum* appears in this horizon or a little higher up in the section.

WEISERT (1932) mentioned *Skirroceras macrum* from a section of the ‘Humphriesioolith’ Member in the railroad cut at Autunnel in a position between c. 1.7 m up to max. 3.5 m above its basis. He named this part of the section “*nodosum-macrum-plicatum-Zone*“. We have some doubts about the correct specific identification of *S. macrum* from higher up than c. 2 m above the basis of the ‘Humphriesioolith’ Member in this section by WEISERT (1932). Despite many years of fieldwork, we never found this species at this high stratigraphic position. Moreover, WEISERT (1932) did not report *S. nodosum* – a species well-known to him – from the Autunnel section, which was well exposed in its time. Referring to WEISERT’s (1932) “*nodosum-macrum-plicatum-Zone*“, OHMERT et al. (1995) suggested to use a *Macrum* Subzone in the top of the Sauzei Zone; however, this view was not adopted since.

? carinodiscus biohorizon – The *carinodiscus* biohorizon is not safely recorded at Gosheim. In the adjacent area of the Zollernalb (DIETZE et al. 2020), this biohorizon is predominated by sonniniids of the variable nominal species *Sonninia carinodiscus*; only a single but typical *Skirroceras nodosum* has been recorded from this biohorizon. This is why we assign the rock interval around c. 2.3 m

		Ringsheim	Wutach	Gosheim	Hohen-zollern	Middle Swabian Alb	E Swabian Alb
Humphriesianum Z.	Pinguis Subzone		<i>deltafalcata</i>	? <i>deltafalcata</i> ² ? ¹	? <i>pinguis</i> ? ¹	<i>deltafalcata</i> <i>frechi</i> <i>pinguis</i>	<i>deltafalcata</i> ? ¹
Sauzei Zone		<i>macrum</i> <i>pseudoc.</i> <i>dilatus</i>	<i>carinodiscus</i> ? 	? <i>carinodiscus</i> ² <i>macrum</i> <i>pseudoc.</i> <i>dilatus</i>	<i>carinodiscus</i> <i>quenstedti</i> <i>dilatus</i>	? <i>macrum</i> <i>pseudoc.</i>	? <i>pseudoc.</i>

Fig. 5. Correlation of the Lower Bajocian biohorizons of SW Germany. Remarks:

¹In the Sauzei/Humphriesianum zonal boundary beds most likely a further biohorizon exists which yields *Sonninia alsatica* and related forms (see DIETZE et al. 2008, 2013, 2020).

²The *carinodiscus* and *deltafalcata* biohorizons are uncertain at Gosheim.

above the basis of the ‘Humphriesiolith’ Member yielding specimens of *S. nodosum* and *S. cf. nodosum* with some reservation to the *carinodiscus* biohorizon. However, we cannot exclude that this interval still belongs to the *macrum* biohorizon. We cannot decide whether some of the illustrated sonniniids collected from the screw originate from this interval. At least, this seems true for the specimen determined as *Sonninia carinodiscus*, the index of this biohorizon.

5.1.3. Humphriesianum Zone, Pinguis Subzone

Unfortunately, all ammonites illustrated on Pl. 9 have not been collected bed-by-bed; otherwise the Pinguis Subzone at Gosheim would become much better dividable. Then, it would become clear, whether *Sonninia alsatica* and *S. pseudofurticarinata* represent an individual biohorizon or belong to the *pinguis* biohorizon or even the Sauzei Zone. The basis of the Pinguis Subzone, however, must be located at least c. 3.6–4.2 m above the basis of the

‘Humphriesiolith’ Member, because the ammonites illustrated on Pl. 9, Figs. 3–10 originate already from the Pinguis Subzone.

?*deltafalcata* biohorizon – The interval of c. 3.6 to 4.2 m above the basis of the ‘Humphriesiolith’ Member with “*Dorsetensia*” sp. und “*D.*” *punctatissima* is tentatively assigned to the *deltafalcata* biohorizon. After OHMERT (1990: 123), *D. pinguis* ranges up to this biohorizon, and the type of *D. punctatissima* seems to originate from the *deltafalcata* biohorizon as well. Typical “*Dorsetensia*” specimens from the slightly older *pinguis* biohorizon (Fig. 4) show relatively narrow whorl sections (DIETZE et al. 2008, fig. 4). In contrast, the small-sized “*Dorsetensia*” from the *deltafalcata* biohorizon of the middle Swabian Alb exhibit broader whorl sections (see OHMERT 1990, pl. 1, figs. 4–7). *Nannina deltafalcata* (QUENSTEDT, 1857), the index of the *deltafalcata* biohorizon, has not been recorded from Gosheim. The few finds from Gosheim do not allow exact biostratigraphic interpretations, but they are definitely diagnostic for the Pinguis Subzone.

5.1.4. Humphriesianum Zone, Romani Subzone

The Romani Subzone is recorded by the *romani* biohorizon down to c. 0.6 m below the marlimestone bed containing the *gervillii/cycloides* biohorizon (Fig. 2; DIETZE et al. 2015). However, due to the absence of precisely collected ammonites, the exact basis of the Romani Subzone could not be drawn, and we cannot decide if the illustrated *Fissilobicerias furticarinum* originates from the Pinguis or from the Romani Subzone. After OHMERT (1990, fig. 9), the type horizon of *F. furticarinum* in the Middle Swabian Alb lies within the Romani Subzone (*furticarinum* biohorizon sensu OHMERT [= pars *deltafalcata* biohorizon and *romani* biohorizon in Fig. 2]). MORTON (1975) illustrated specimens termed as “*Sonninia*” aff. *furticarinata* from the Hebridica Subzone (Sauzei Zone) and the Cycloides Subzone (Humphriesianum Zone) of the Isle of Skye (Scotland) that are very close to our illustrated specimen. The Hebridica Subzone sensu MORTON correlates with uppermost parts of the Sauzei Zone and large parts of the Pinguis Subzone in SW Germany.

5.2. Age correlation

The correlation of several important localities in the upper Lower Bajocian of SW Germany with the Gosheim section is shown in Fig. 5.

Concerning the difficulties for age correlations with S England see DIETZE et al. (2020). The *dilatatus* biohorizon corresponds more or less to the *kumaterus/simulans* biohorizon, and the *pseudocontrahens* biohorizon is an approximate equivalent of the *kalum* biohorizon. The *macrum* and *carinodiscus* biohorizons are coeval with the *digbyi* and *rhytum* biohorizons in S England, respectively. The position of the *deltafalcata* biohorizon corresponds to that of the *delphinum* biohorizon (see CHANDLER 2019).

In France, the term ‘Propinquans Zone’ is used instead of the Sauzei Zone (RIOULT et al. 1997). It is subdivided into the older Patella Subzone and the younger Hebridica Subzone. A correlation with the French horizons is very difficult or even impossible (see DIETZE et al. 2020). The *macrum* and *carinodiscus* biohorizons partly correspond to the horizon à *Hebridica* (see DIETZE et al. 2020). The *deltafalcata* biohorizon might be coeval with the horizon à *Edouardiana*.

Acknowledgements

We thank E. BERNT †, F. FANZUTTI (Bisingen-Thenheim), F. NEUBAUER (Baltmannsweiler), T. OTT (Bisingen-Thenheim), and U. RYCK (Polling) for providing important data and several ammonites for this study and especially the late N. WANNENMACHER, with whom most of the sections were measured and the results were discussed. The manuscript benefitted from the thorough reviews by G. DIETL (Stuttgart) and A. GALÁČZ (Budapest).

6. References

- BAYLE, E. (1878): Atlas. Première Partie – Fossiles principaux des terrains. – Explication de la Carte Géologique de la France, 4: 158 pls.
- BERZ, K. C. (1933): Geologische Karte von Baden-Württemberg 1:25.000. Erläuterungen zu Blatt 7818 Wehingen. 91 pp.; Stuttgart.
- BUCKMAN, S. S. (1887–1907): A monograph of the ammonites of the “Inferior Oolite series”. – Monographs of the Palaeontographical Society: cclxii + 456 pp.
- BUCKMAN, S. S. (1909–1930): Yorkshire Type Ammonites/Type Ammonites, 1–7, 790 pls.; London (Wheldon & Wesley).
- CALLOMON, J. H. (1995): Time from fossils: S.S. Buckman and Jurassic high-resolution geochronology. In: LE BAS, M. J. (ed.): Milestones in geology. – Geological Society Memoirs, 16: 127–150.
- CHANDLER, R. B. (2019): Two new stephanoceratid ammonites from the Aalenian–Lower Bajocian (Middle Jurassic, Dorset, UK) and their phylogenetic significance. – Proceedings of the Geologists’ Association, 130: 307–325.
- CHANDLER, R. B., DIETZE, V. & WHICHER, J. (2017): A revision of some British Lower Bajocian stephanoceratid ammonites. – Proceedings of the Geologists’ Association, 128: 636–658.
- DIETL, G. (1977): The Braunjura (Brown Jurassic) in Southwest Germany. – Stuttgarter Beiträge zur Naturkunde, (B), 25: 1–41.
- DIETL, G. (1978): Zur Braunjura γ/δ -Grenze (Unter-Bajocium) im Westteil der Schwäbischen Alb (Baden-Württemberg). – Stuttgarter Beiträge zur Naturkunde, (B), 36: 1–15.
- DIETL, G. & RIEBER, H. (1980): Der Braune Jura der Westalb zwischen Balingen und Geisingen (Exkursion F1 am 10. April und Exkursion F2 am 11. April 1980). – Jahresberichte und Mitteilungen des oberrheinischen geologischen Vereins, Neue Folge, 62: 57–69.
- DIETZE, V., BOSCH, K., FRANZ, M., KUTZ, M., SCHWEIGERT, G., WANNENMACHER, N. & STUDER, S. (2013): Die Humphriesianum-Zone (Unter-Bajocium, Mitteljura) am Kahlenberg bei Ringsheim (Oberrheingraben, SW Deutschland). – Palaeodiversity, 6: 29–61.
- DIETZE, V., KUTZ, M., FRANZ, M. & BOSCH, K. (2009): Stratigraphy of the Kahlenberg near Ringsheim (Upper Rhine Valley, SW Germany) with emphasis on the Laeviuscula and Sauzei zones (Lower Bajocian, Middle Jurassic). – Palaeodiversity, 2: 19–65.
- DIETZE, V., SCHWEIGERT, G., FIDDER, U. & WANNENMACHER, N. (2011): The Giganteuston Member of Öschingen (Humphriesianum Zone, Lower Bajocian, Swabian Alb), with comments on the genera *Dorsetensia* BUCKMAN, 1892 and *Nannina* BUCKMAN, 1927. – Zitteliana, 51: 209–236.
- DIETZE, V., STAPPENBECK, G., WANNENMACHER, N. & SCHWEIGERT, G. (2008): Stratigraphie und Ammoniten-Faunenhorizonte im Grenzbereich Sauzei/Humphriesianum-Zone (Unter-Bajocium, Mitteljura) der westlichen Schwäbischen Alb (SW-Deutschland). – Palaeodiversity, 1: 141–165.
- DIETZE, V., WANNENMACHER, N., FRANZ, M. & SCHWEIGERT, G. (2015): Lithological section and biostratigraphy of the Gosheim Formation in its type area (Lower Bajocian, Swabian Alb, SW Germany). – Palaeodiversity, 8: 31–69.
- DIETZE, V., WANNENMACHER, N. & SCHWEIGERT, G. (2020): Die Sauzei-Zone (Mitteljura, Unter-Bajocium) im Gebiet des Hohenzollern (Schwäbische Alb, SW-Deutschland). – Zitteliana, 94: 37–83.

- DORN, P. (1935): Die Hammatoceraten, Sonninien, Ludwigien, Dorsetensien und Wichtellien des süddeutschen, insbesondere fränkischen Doggers. – *Palaeontographica*, **82**: 1–124.
- FISCHER, H. (1912): Beiträge zur Geologie von Rottweils Umgebung. – Wissenschaftliche Beilage zum Jahresbericht des königlichen Gymnasiums in Rottweil, **1912**: 1–63.
- FISCHER, H. (1924): Zur Stratigraphie des Doggers bei Gosheim (Rottweil – Spaichinger Gegend). – Jahresberichte und Mitteilungen des oberrheinischen geologischen Vereins, Neue Folge, **13**: 97–109.
- FRANZ, M. (1988): Makrofauna und Fazies der Oolithhorizonte im Bajocium Süddeutschlands. – Jahreshefte der Gesellschaft für Naturkunde in Württemberg, **143**: 79–105.
- GEYER, O. F. & GWINNER, M. P. (1984): Die Schwäbische Alb und ihr Vorland. – Sammlung geologischer Führer, **67**: 275 pp.; Berlin & Stuttgart (Borntraeger).
- GEYER, O. F. & GWINNER, M. P. (2011): Geologie von Baden-Württemberg (5th edition). X + 627 pp.; Stuttgart (Schweizerbart).
- HAUG, É. (1885): Beiträge zu einer Monographie der Ammonitengattung *Harpoceras*. – Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Beilage-Bände, **3**: 585–722.
- HAUG, É. (1893): Étude sur les ammonites des étages moyens du système Jurassique. – Bulletin de la Société géologique de France, **3**, **20**: 277–333.
- ICZN (2005): Opinion 2123. F.A. QUENSTEDT's trinomial nomenclature (1845–1888): usage of the third names of ammonites stabilized and 34 important QUENSTEDT names of ammonites placed on the Official List of Specific Names in Zoology (Cephalopoda, Ammonoidea). – Bulletin of Zoological Nomenclature, **62**: 158–163.
- MASCHE, E. (1907): Die *Stephanoceras*-Verwandten in den Coronatenschichten von Norddeutschland. Inaugural-Dissertation Universität Göttingen. 38 pp.; Alfeld (Dobler).
- MAUBEUGE, P. L. (1951): Les ammonites du Bajocien de la région frontière Franco-Belge (Bord septentrional du Bassin de Paris. – Mémoires de l'Institut Royal des sciences naturelles de Belgique, **2**(42): 1–104.
- MAUBEUGE, P. L. (1961): Catalogue des Ammonites du Jurassique inférieur et moyen (Hettangien à Bathonien) du Musée cantonal de Bâle-Campagne. – Tätigkeitsberichte der Naturforschenden Gesellschaft Baselland, **22**: 27–164.
- MORTON, N. (1975): Bajocian Sonniniidae and other ammonites from western Scotland. – *Palaeontology*, **18**(1): 41–91.
- NEUMAYR, M. (1875): Die Ammoniten der Kreide und die Systematik der Ammonitiden. – Zeitschrift der Deutschen Geologischen Gesellschaft, **27**: 854–892.
- OHMERT, W. (1990): The Humphriesianum zone in the type area. – In: CRESTA, S. & PAVIA, G. (eds.): Atti del meeting sulla stratigrafia del Baiociano. – Memorie descrittive della Carta geologica d'Italia, **40**: 117–140.
- OHMERT, W., GASSMANN, G., SCHATZ, R. H. & STETTER, L. (1995): Das höhere Unter-Bajocium (ehemals Mittel-Bajocium) von Lörrach. II. Die *humphriesianum*-Zone. – Jahreshefte des Geologischen Landesamtes Baden-Württemberg, **35**: 24–264.
- QUENSTEDT, F. A. (1856–1857): Der Jura. – 842 pp.; Tübingen (Laupp).
- QUENSTEDT, F. A. (1886–1887): Die Ammoniten des Schwäbischen Jura. 2. Der Braune Jura: 441–885; Stuttgart (Schweizerbart).
- RENZ, C. (1904): Der Jura von Daghestan. – Neues Jahrbuch für Mineralogie, Geologie und Palaeontologie, **1904**(2): 71–85.
- RIOULT, M., CONTINI, D., ELMI, S. & GABILLY, J. (1997): Bajocien. In: CARIOU, É. & HANTZPERGUE, P. (eds.): Biostratigraphie du Jurassique ouest-européen et méditerranéen. – Bulletin du Centre des Recherches Elf Aquitaine Exploration et Production, Memoirs, **17**: 41–54.
- ROCHÉ, M. (1939): Aalénien et Bajocien du Mâconnais et de quelques régions voisines. – Travaux des Laboratoires de la Faculté des Sciences de Lyon, fascicule 35, Mémoires, **29**: 1–355.
- SADKI, D. & DIETZE, V. (2020): New information on the Early Bajocian types of sonniniids and hammatoceratids (Ammonitina) described by W. WAAGEN (1867) from Gingen/Fils (SW Germany). – Boletín del Instituto de Fisiografía y Geología, **91**: 1–16.
- WAAGEN, W. (1867): Über die Zone des *Ammonites Sowerbyi*. – Benecke's geognostisch-paläontologische Beiträge, **1**(3)(2): 509–713.
- WANNENMACHER, N. (2017): Historische Fundstücke zur Geologie der Schwäbischen Alb. – Fossilien, **34**(5): 49–56.
- WEISERT, K. (1932): *Stephanoceras* im schwäbischen Braunen Jura Delta. – *Palaeontographica*, **76**: 121–191.

Addresses of the authors:

VOLKER DIETZE, Meraner Straße 41, D–86720 Nördlingen, Germany; e-mail: dietze.v@t-online.de

Dr. GÜNTER SCHWEIGERT, Staatliches Museum für Naturkunde in Stuttgart, Rosenstein 1, D–70191 Stuttgart, Germany; e-mail: guenter.schweigert@smns-bw.de

Manuscript received: 12 November 2021, revised version accepted: 27 December 2021.

Plate 1

(1–4) Ammonites of the Wedelsandstein Formation, ‘Oberer Blaukalk’ Member (topmost bed) at Gosheim, Lower Bajocian, Sauzei Zone, *dilatus* biohorizon.

(1, 2) *Kumatostephanus triplicatus* (QUENSTEDT, 1886), Denkinger Steige, SMNS 70635/5 [leg. M. FRANZ & E. KIEFER].

(3, 4) *Kumatostephanus* aff. *turgidulus* (QUENSTEDT, 1886), SMNS 70635/6 [leg. N. WANNENMACHER †].

(5, 6) *Emileia vagabunda* BUCKMAN, 1927; Gosheim Formation, ‘Humphriesoolith’ Member (basal 0.1 m), Lower Bajocian, Sauzei Zone, *pseudocontrahens* biohorizon, Gosheim, Industriestraße, SMNS 70635/7.

Scale bar equals 10 cm. Ammonites: x0.5. Beginning of body chamber is indicated by an asterisk.



Plate 2

(1) *Kumatostephanus* cf. *perjucundus* (BUCKMAN, 1927), Wedelsandstein Formation, 'Oberer Blaukalk' Member (topmost bed); Lower Bajocian, Sauzei Zone, *dilatus* biohorizon, Gosheim, SMNS 70635/8 [leg. K.-H. SPIETH].

(2–5) Ammonites of the Gosheim Formation, 'Humphriesioolith' Member at Gosheim; Lower Bajocian, Sauzei Zone, *pseudocontrahens* biohorizon.

(2, 3) *Emileia arkelli* MAUBEUGE, 1961, basal c. 2.5 m, Sturmbühl, SMNS 70635/9 [leg. F. NEUBAUER].

(4, 5) *Emileia schassmanni* MAUBEUGE, 1961, basal 0.45 m ("Spathulatusbank"), new development area at Längenberg, SMNS 70635/10 [leg. T. OTT].

Scale bar equals 10 cm. Ammonites x0.5. Beginning of body chamber is indicated by an asterisk.



Plate 3

(1, 2) *Skirroceras macrum* (QUENSTEDT, 1886), Gosheim Formation, 'Humphriesioolith' Member (c. 1.8 m above base), Lower Bajocian, Sauzei Zone, *macrum* biohorizon, Gosheim, Industriestraße, SMNS 70635/11.

Scale bar equals 10 cm. Ammonite x0.5. Beginning of body chamber is indicated by an asterisk.



Plate 4

Ammonites of the Gosheim Formation, 'Humphriesiolith' Member at Gosheim; Lower Bajocian, Sauzei Zone, *macrum* biohorizon.

(1) *Sonninia* cf. *patella* (WAAGEN, 1867), c. 1.8 m above base, Industriestraße, SMNS 70635/12.

(2) *Skirroceras* cf. *macrum* (QUENSTEDT, 1886), c. 1.8–2.5 m above base, *macrum* biohorizon [by preservation and matrix], Kreissparkasse, SMNS 70635/13.

Scale bar equals 10 cm. Ammonites x0.5. Beginning of body chamber is indicated by an asterisk.



Plate 5

Ammonites of the Gosheim Formation, 'Humphriesioolith' Member at Gosheim; Lower Bajocian.

(1) *Skirroceras* cf. *nodosum* (QUENSTEDT, 1857), c. 1.8–2.5 m above base; Sauzei Zone, *macrum* or *carinodiscus* biohorizon, SMNS 70635/14.

(2) *Skirroceras* sp. [pathogenic], between c. 2 m and max. 4.5 m above base, upper Sauzei or lowermost Humphriesianum Zone, Brücklestraße, SMNS 70635/15.

(3, 4) *Skirroceras* cf. *freycineti* (BAYLE, 1878), between c. 2 m and max. 4.5 m above base, upper Sauzei Zone or lowermost Humphriesianum Zone, Brücklestraße, SMNS 70635/16.

Scale bar equals 10 cm. Ammonites x0.5. Beginning of body chamber is indicated by an asterisk.



Plate 6

Ammonites of the Gosheim Formation, 'Humphriesiolith' Member at Gosheim (Brücklestraße), Lower Bajocian, Sauzei Zone, *?carinodiscus* biohorizon.

(1, 2) *Skirroceras* cf. *nodosum* (QUENSTEDT, 1857), c. 2.3 m above base, SMNS 70635/17.

(3, 4) *Skirroceras nodosum* (QUENSTEDT, 1857), c. 2.3 m above base as indicated by rock matrix and preservation, SMNS 70635/18 [leg. U. RYCK].

Scale bar equals 10 cm. Ammonites x0.5. Beginning of body chamber is indicated by an asterisk. The arrow indicates the beginning of the body chamber.



Plate 7

Ammonites of the Gosheim Formation, 'Humphriesioolith' Member at Gosheim, Lower Bajocian.

(1, 3) *Sonninia* ["*Sonninites*"] aff. *felix* (BUCKMAN, 1923), c. 1.5–max. 4.5 m above base, ?*carinodiscus* biohorizon, Brücklestraße, SMNS 70635/19 [leg. T. OTT].

(2, 4) *Sonninia* cf. *patella* (WAAGEN, 1867), c. 1.5–max. 4.5 m above base, ?*pseudocontrahens* biohorizon, Sturmbühl, SMNS 70635/20 [leg. E. BERNT †].

Scale bar equals 10 cm. Ammonites x0.5. Beginning of body chamber is indicated by an asterisk.



Plate 8

Ammonites of the Gosheim Formation, 'Humphriesioolith' Member at Gosheim, Lower Bajocian.

(1, 2) *Sonninia carinodiscus* (QUENSTEDT, 1886), c. 1.5–max. 4.5 m above base, ?*carinodiscus* biohorizon, Sturmbühl, SMNS 70635/21 [leg. E. BERNT].

(3, 4) *Dorsetensia liostraca* BUCKMAN, 1892, lower part of 'Humphriesioolith' Member, Romani Subzone (Humphriesianum Zone), Autunnel, SMNS 23697 [leg. K.-H. SPIETH; specimen mentioned in DIETL 1978: 9].

Scale bar equals 10 cm. Ammonites x0.5. Beginning of body chamber is indicated by an asterisk.



Plate 9

Ammonites of the Gosheim Formation, 'Humphriesioolith' Member at Gosheim; Lower Bajocian.

(1–4) *Sonninia pseudofurticarinata* MAUBEUGE, 1961, c. 1.5–max. 4.5 m above base, uppermost Sauzei Zone or lowermost Humphriesianum Zone, (1) SMNS 70635/22 [leg. E. GLÜCK], (2) Brücklestraße, SMNS 70635/23, (3, 4) Sturmbühl, SMNS 70635/24 [leg. E. BERNT].

(5, 6) *Sonninia alsatica* (HAUG, 1885), c. 1.5–max. 4.5 m above base, ?Pinguis Subzone (Humphriesianum Zone), Brücklestraße, SMNS 70635/25.

(7, 8) *Fissilobicerias furticarinatum* (QUENSTEDT, 1856), c. 2.5–max. 4.5 m above base, Pinguis Subzone or Romani Subzone (Humphriesianum Zone), Brücklestraße, SMNS 70635/26.

Scale bar equals 10 cm. Ammonites x0.5. Beginning of body chamber is indicated by an asterisk.

