



## **The genus *Psephellus* Cass. (Compositae, Cardueae) revisited with a broadened concept**

Authors: Wagenitz, Gerhard, and Hellwig, Frank H.

Source: *Willdenowia*, 30(1) : 29-44

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.30.30102>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

GERHARD WAGENITZ &amp; FRANK H. HELLWIG

## The genus *Psephellus* Cass. (*Compositae*, *Cardueae*) revisited with a broadened concept

### Abstract

Wagenitz, G. & Hellwig, F. H.: The genus *Psephellus* Cass. (*Compositae*, *Cardueae*) revisited with a broadened concept. – Willdenowia 30: 29-44. 2000. – ISSN 0511-9618.

A new concept of the genus *Psephellus* is presented on the basis of morphological, anatomical, palynological and caryological evidence. The few molecular data seem to confirm the monophyly of the genus. The following former sections of *Centaurea* are included: *C.* sect. *Psephelloideae*, *Psephellus*, *Hyalinella*, *Aetheopappus*, *Amblyopogon*, *Heterolophus*, *Czerniakovskya*, *Odontolophoideae*, *Odontolophus*, *Xanthopsis*, *Uralepis* and *Sosnovskya*. New combinations under *Psephellus* are provided for these sections and for 35 species, especially from Turkey and Iran. *Psephellus* in this broadened sense has 75-80 species and a distribution with a centre in E Anatolia, Caucasia and NW Iran; only few species occur outside this area. Close relationships exist between different sections despite considerable differences especially in the characters of the pappus.

### Introduction

In the *Centaureinae* the concept of genera varies enormously. This is clearly shown by the following list with the number of genera discerned by various authors:

Hoffmann (1894):	9 genera
Bobrov & Čerepanov (1963):	26 genera (only 'Flora SSSR area', but most genera occur there)
Dostál (1973):	51 genera
Dittrich (1977):	7 genera
Bremer (1994):	31 genera

Our aim is to establish moderately large genera which are monophyletic (see Wagenitz & Hellwig 1996). This is possible if morphological and molecular data are combined. One of these genera is presented here. It first emerged from the study of the pollen morphology (Wagenitz 1955). It could be shown that there is one group of sections of *Centaurea* s.l. (the *Psephellus* group of Wagenitz & Hellwig 1996) with a very conspicuous pollen type named *Centaurea dealbata* type. *C. dealbata* Willd. is the type of the genus *Psephellus* Cass. *Psephellus* has been accepted as a genus by Boissier (1875), Sosnovsky (1948), Dostál (1973), Gabrieljan (1995), but has been sunk into *Centaurea* in the 'Flora SSSR' (Bobrov & Čerepanov 1963). The same pollen type is found in several

sections of *Centaurea* centering in the area of E Turkey, Caucasia and NW Iran (with few representatives in the Carpathians and Russia). This group of taxa has some macromorphological features in common and is further united by micromorphological and karyological traits. The few molecular data seem to confirm the monophyly of the genus. Petit (1997) does not include any of the taxa concerning us here in his cladistic analysis of morphological data of the *Cardueae*. Besides the features common to the genus, it is held together by the geographical distribution. Remarkable is the absence of annuals or biennials and of spines or spinules on the phyllaries. *Psephellus dealbatus* and some allied species are decorative garden plants. The genus shows no weedy tendencies.

## Characters

**Macromorphological characters.** Herbaceous or suffruticose perennials. Leaves (at least the young ones) always tomentose with an indument of minute woolly hairs or with additional hairs with pedicel. Leaves variously divided, not decurrent. Appendage of phyllaries usually not decurrent, membranous, entire, dentate or ciliate, never with a spine. Receptacle with smooth bristles, persistent at maturity of the achenes. Marginal flowers with staminodes, often strongly radiant. Flowers purple, pink or pale yellow to whitish. Achenes medium-sized to large, laterally flattened to almost rounded in *Psephellus hedgei*. Pappus very variable, rarely lacking, when present always with biseriate glandular hairs at least on the inner pappus elements.

**Micromorphological characters of phyllaries and flowers.** Phyllaries without crystals. Corolla tube with small prismatic Ca-oxalate crystals or small prismatic crystals and small plates of Ca-oxalate. Cells at the base of the corolla tube strongly elongated with the walls evenly thickened (not thickened in *Psephellus bellus* and *P. marschallianus*). Apical appendages of the anthers obtuse and rather soft (with the exception of most species in *P.* sect. *Psephelloidei* and sect. *Amblyopogon*). Filaments with papillae or short hairs not exceeding half the width of the filaments. Style branches united, only in the uppermost part diverging.

**Anatomical characters of the achenes.** Dittrich (1968) investigated species belonging to four sections of *Psephellus* s.l. and united them in his “*Aetheopappus-Psephellus* branch”. Our studies on a much extended basis revealed the following features. Achenes elliptical in cross-section, almost circular in *P. hedgei*, usually with elaiosome (exceptions: see description of the sections), with unicellular hairs except in *P. pulcherrimus* (glabrous), in *P. trinervius* also with biseriate glandular hairs, surface smooth or with horizontal ridges, apically rounded (*P. xanthocephalus*) or crowned by a smooth or pinnated rim. Area of detachment in a lateral or (species of *P.* sect. *Psephelloidei*) basal-lateral position, surrounded by a hardened and often brighter zone, except in *P. xanthocephalus*. Cells of the fruit wall parenchyma lignified, except in *P. trinervius*. Outer layers of the pericarp without, inner layer with long prismatic Ca-oxalate crystals and with a dense layer of long prisms outside the testa epidermis. Seeds with hilum lateral.

**Pollen morphology.** This has first been investigated by Wagenitz (1955) as mentioned above. The “*dealbata* type”, or more correct “*Centaurea dealbata* type” is characterised by a spheroid to subprolate shape, a well developed inner row of bacula, a scarcely visible outer row of bacula and a minutely scabrate surface. It shows some similarity with the pollen types of *Centaurea montana* and *C. cyanus*, which, however differ markedly by their subprolate to prolate shape and the more or less triangular cross-section with the colpi on the planes. The “*Psephellus* type” established by Avetisjan (1964) included these two types and is thus less useful for taxonomic purposes. She gives, however, a list of additional taxa from *Psephellus* s.l. investigated by her.

**Chromosomes.** Results of karyological investigations in this group were rare and controversial for some time. Only recently reliable data for several of the sections have been available. From

these it clearly emerges that  $n = 15$  is the base number of the whole group. Deviating numbers have been published by some Russian and Armenian botanists but have not been confirmed by other authors. Polyploidy seems to be very rare, the only record is the count of  $2n = 50-60$  by Poddubnaja-Arnoldi (1931) for three closely allied species of *P.* sect. *Hyalinella*. B-chromosomes have been found in several species. The base number 15 is known from several other taxa (*Klasea*, *Centaurea* sect. *Centaurea*) in the *Centaureinae* and is probably primitive here.

**Molecular data.** So far only *Psephellus dealbatus* (Susanna & al. 1995), *P. hedgei*, *P. bellus* and *P. trinervius* (Hellwig 1996) have been included in phylogenetic analyses using molecular markers. In these studies the ITS regions of the nuclear ribosomal repeat were sequenced. As in Susanna's investigation *P. dealbatus* was the only representative of the *Psephellus* group, no conclusion on the monophyly of this group could be made. In Hellwig's study three species were included in the ITS data set and two of them (*P. hedgei* and *P. bellus*) were also included in an RFLP analysis of the chloroplast DNA (Hellwig 1996). In both cases *Psephellus* proved to be monophyletic. The bootstrap support for *Psephellus* in the ITS data set was moderately high (73 % in 1000 replicates) and comparable to the support for the *Cyanus/Protocyanus* group in the same analysis. *Psephellus bellus* and *P. hedgei* were grouped into a monophyletic clade with 100 % bootstrap support also in the restriction site analysis. Phylogenetic reconstruction in the subtribe *Centaureinae* (Hellwig 1996) using a data set combined from ITS sequence data and morphological data showed *Psephellus* as a monophyletic group which was well supported in a bootstrap analysis (94 %). The number of synapomorphies of *Psephellus* in the ITS sequence analysis was low (4) when compared to the respective values for *Plectocephalus* (9), *Acrocentron* (9) and *Cyanus/Protocyanus* (6). In contrast, there are 11 synapomorphic changes for the *Psephellus* group in the RFLP analysis of the cpDNA. This is the highest value for a clade within the subtribe *Centaureinae* and equals the number of synapomorphies for the whole subtribe (Hellwig 1996). The relatively high bootstrap value in the sequence analysis together with a low number of synapomorphies may be explained by the congruence of the changes in these few characters. Within *Centaureinae*, *Psephellus* can be qualified as a slowly evolving monophyletic clade when compared to the *Jacea* group, for example. This statement, however, is based only on the small amount of molecular information available so far. A detailed molecular study of *Psephellus* is under way.

### Affinities and phylogenetic position

There are many features shared by all species of *Psephellus*, admittedly however, several of these are plesiomorphic, such as the lack of crystals in the phyllaries, the staminodes and the chromosome number. The pollen type is clearly an apomorphic feature, probably the obtuse anther appendages are another one, but it is not found in all species. They are otherwise only known from few remote genera such as *Acroptilon*, *Callicephalus* and the *Carthamus* group.

*Centaurea* sect. *Cyanus* (better treated as a genus *Cyanus* Mill.) resembles *Psephellus* in some important characters. Both genera have similar pollen types differing markedly from all others by the well developed inner row of bacula in combination with a very inconspicuous outer row and loss of the spines (pollen grains only minutely scabrate). It is remarkable too that in both groups the phyllary appendages are never spiny. There are, however, important differences: in *Cyanus* the appendages are strongly decurrent, crystals are present in the phyllaries, the base chromosome numbers are  $n = 8, 10, 11, 12$  and the style branches are much shorter and completely free. Further features of *Cyanus* absent in *Psephellus* are the curved apical anther appendages and the characteristic trichomes at the detachment area of the achenes.

*Psephellus* has several features in common with *Centaurea* s.str. and *Cheirolophus*, e.g. the occurrence of staminodes and the chromosome base number ( $n = 15$  in *Centaurea*, 15 and 16 in *Cheirolophus*), but the pollen types and the apical anther appendages are very different.

Extrafloral nectaries, which can be found on the outer face of the phyllaries in *Cyanus* and on the inner surface in *Centaurea*, are lacking in *Psephellus*.

### Comparison with previous systems

Two extreme positions are possible in the system of the *Centaureinae* with respect to the group we are concerned with. Several authors (Hoffmann 1894, Dittrich 1977, Bremer 1994) include all the sections of *Psephellus*, as treated here, in *Centaurea*, while Dostál (1973) recognizes four genera with pollen of the *Centaurea dealbata* type: *Aetheopappus*, *Psephellus*, *Heterolophus* and *Odontolophus*. It is not clear if the other relevant taxa were not known to him or are to be included in these. As Table 1 shows, there are also several authors who have treated some of the taxa as separate genera, others as subgenera of *Centaurea*. Boissier (1875), who relied heavily on the pappus when delimiting *Centaurea* (see Wagenitz 1963), dispersed the species of our group in several genera:

1. typical double *Centaurea* pappus: *Centaurea* sect. *Jacea* § *Psephelloideae* Boiss., *C.* sect. *Phalolepis* (later *C.* sect. *Hyalinella* Tzvelev and *C.* sect. *Uralepis* (DC.) Wagenitz)
2. pappus simple or double with plumose bristles: *Aetheopappus* (this pappus type seems to vary even within *C. pulcherrima* s.l.)
3. pappus very short and deciduous: *Psephellus* Cass.
4. pappus short and not distinctly double: *Phaeopappus* Boiss. (species now belonging to *Psephellus* sect. *Odontolophus* and sect. *Sosnovskya*).

His system is certainly unnatural and unacceptable in the view of our present knowledge.

The systems by Bobrov & Čerepanov (1963) and Tahtadžjan (1995) fail, however, likewise in creating a phylogenetically sound system, as they distribute the members of the *Psephellus* group in several subgenera inside and one or two genera outside *Centaurea* (Table 1.).

Table 1. Status of the sections of *Psephellus* in previous systems (*C.* = *Centaurea*, — = taxon not treated).

<i>P.</i> sect.	Boissier (1875)	Bobrov & Čerepanov (1963)	Tahtadžjan (1995)
<i>Psephelloidei</i>	<i>C.</i> sect. <i>Jacea</i> § <i>Psephelloideae</i> & <i>C.</i> sect. <i>Phalolepis</i> p.p.	<i>C.</i> subg. <i>Pseudohyalea</i> p.p.	
<i>Psephellus</i>	<i>Psephellus</i>	<i>C.</i> subg. <i>Psephellus</i>	<i>Psephellus</i>
<i>Hyalinella</i>	<i>C.</i> sect. <i>Phalolepis</i> p.p.	<i>C.</i> subg. <i>Hyalinella</i>	<i>C.</i> subg. <i>Hyalinella</i>
<i>Aetheopappus</i>	<i>Aetheopappus</i>	<i>Aetheopappus</i>	<i>Aetheopappus</i>
<i>Amblyopogon</i>	<i>Psephellus</i> p.p. & <i>C.</i> sect. <i>Jacea</i> § <i>Psephelloideae</i>	<i>C.</i> subg. <i>Amblyopogon</i>	—
<i>Heterolophus</i>	<i>C.</i> sect. <i>Jacea</i> § <i>Psephelloideae</i>	<i>C.</i> subg. <i>Heterolophus</i>	—
<i>Czerniakovskya</i>	—	<i>C.</i> subg. <i>Czerniakovskya</i>	—
<i>Odontolophoidei</i>	<i>Psephellus</i> p.p.	<i>C.</i> subg. <i>Odontolophopsis</i>	
<i>Odontolophus</i>	<i>Phaeopappus</i> p.p.	<i>C.</i> subg. <i>Odontolophus</i>	
<i>Xanthopsis</i>	<i>Psephellus</i> p.p.	<i>C.</i> subg. <i>Xanthopsis</i>	
<i>Uralepis</i>	<i>C.</i> sect. <i>Phalolepis</i> p.p.	—	—
<i>Sosnovskya</i>	<i>Phaeopappus</i> p.p.	<i>C.</i> subg. <i>Sosnovskya</i>	

### Subdivision of the genus

The main differences within *Psephellus* s.l. concern the habit, the marginal flowers and the characters of the pappus. According to the habit of the plants two groups of sections can be distinguished, but there is no clear-cut division between them (Table 2). In the sections with a “*Psephellus* habit” the plants are mesomorphic, the stems single or few, erect or ascending to prostrate, they have usually pinnatisect leaves and relatively large heads with showy marginal flowers. With one exception the flowers are pink to purple (in Table 2 “purple” denotes different shades from pink and rose-purple to purple). Several species in this group live in subalpine to alpine habitats. In the second group (“*Xanthopsis* habit”) the plants have numerous stems from a woody base, the leaves are predominantly undivided and linear-lanceolate, the small heads have inconspicuous marginal flowers and the flowers are purple or yellowish. The plants of this second group have a more xeromorphic habit and inhabit mountain steppe habitats or even desert-like places. As can be seen from Table 2, there are intermediate groups which do not fit this pattern: *P.* sect. *Czerniakovskya* has the *Psephellus* habit but yellowish not radiant flowers, while *P.* sect. *Odontolophus* with the *Xanthopsis* habit has radiant flowers. This distribution of characters shows the close phylogenetic connections between all these taxa. It is worth mentioning that some authors have already considered an amplification of the genus *Psephellus*. This is shown by the names or combinations *Psephellus aucherianus* (DC.) Boiss., *P. erivanensis* Lipsky, *P. incanescens* (DC.) Boiss., *P. integrifolius* (C. A. Mey.) C. Koch, *P. marschallianus* (Spreng.) C. Koch and *P. xanthocephalus* (DC.) Boiss. & Buhse. The similarities in habit are also shown by the names *Psephelloides* or *Psephelloideae* given to two different sections.

Table 2. Main features of the sections of *Psephellus* Cass. s.l.  
(h = habit, 1 = “*Psephellus* habit”, 2 = “*Xanthopsis* habit”, see text for further explanations).

<i>P.</i> sect.	h	leaves	phyllary appendage	marginal flowers	flower colour	pappus	pappus bristles
<i>Psephelloidei</i>	1	tomentose	entire/dentate	large	purple	double	scabrous
<i>Psephellus</i>	1	bi-coloured	dentate	large	purple	short, deciduous	scabrous
<i>Hyalinella</i>	1	bi-coloured	denticulate	large	purple	double	scabrous
<i>Aetheopappus</i>	1	tomentose	dentate	large	purple/yellow	double/simple	plumose
<i>Amblyopogon</i>	1	tomentose	ciliate	not radiant	purple	short or 0	scabrous
<i>Heterolophus</i>	1	bi-coloured	ciliate/dentate	radiant	purple	double	scabrous
<i>Czerniakovskya</i>	1	tomentose	ciliate	not radiant	yellowish	double	scabrous
<i>Odontolophoidei</i>	2	tomentose	ciliate	not radiant	purple	double, short	scabrous
<i>Odontolophus</i>	2	tomentose	ciliate/lacerate	radiant	pink	double/short	scabrous
<i>Xanthopsis</i>	2	tomentose	ciliate	not radiant	yellowish	short or 0	scabrous
<i>Uralepis</i>	2	tomentose/ glabrous	entire/ dentate	not radiant	yellowish	double, long	scabrous
<i>Sosnovskya</i>	2	tomentose/ glabrous	ciliate	slightly radiant	pink or whitish	double/simple	scabrous

**Conspectus of the genus *Psephellus* s.l.**

*Psephellus* Cass. in Cuvier, Dict. Sci. Nat. 43: 488. 1826.

Type: *Centaurea dealbata* Willd. ≡ *Psephellus dealbatus* (Willd.) C. Koch

*Psephellus* sect. *Psephelloidei* (Boiss.) Wagenitz & Hellwig, **comb. nova**

Type: *Centaurea pyrrhoblephara* Boiss. ≡ *Psephellus pyrrhoblepharus* (Boiss.) Wagenitz

≡ *Centaurea* sect. *Jacea* § *Psephelloideae* Boiss., Fl. Orient. 3: 616. 1875 ≡ *Centaurea* sect. *Psephelloideae* (Boiss.) Sosn. in Věstn. Tiflissk. Bot. Sada, ser. 2, 5: 29. 1931.

= *Centaurea* sect. *Pseudohyalea* Tzvelev in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 428. 1959 ≡ *Centaurea* subg. *Pseudohyalea* (Tzvelev) Tzvelev, in Bobrov & Čerepanov, Fl. SSSR 28: 486. 1963. – Type: *Centaurea pergamacea* DC.

*Perennial herbs* with woody rootstock. *Stems* erect or ascending, simple or sparsely branched. *Leaves* tomentose, rarely glabrescent, undivided, lyrate or pinnatifid, not decurrent. *Involucre* 15-30 mm long and 10-30 mm broad, ovoid to subglobose. *Phyllaries* in several series, appendages membranous, usually large and semicircular to ovate or rhombic, sometimes shortly decurrent, margin entire, denticulate or regularly ciliate, the terminal mucro often shorter than the cilia, rarely slightly longer or totally lacking. *Flowers* rose-purple, the marginal radiant, often strongly so, with staminodes, the central flowers numerous. *Achenes* 5-8 mm long, detachment area sublateral. *Pappus* (2-)5-13 mm, of scabrous bristles, the inner row of short scales.

The section shows considerable variation in the structure of the appendages, which may be entire, denticulate or with long cilia and decurrent or well differentiated. As the plate in Wagenitz (1975: 561) shows, there are many transitions in this character and the separation of *C.* sect. *Pseudohyalea* is scarcely justified. The appendages of the anthers are also variable, usually acute but obtuse in *Psephellus gilanicus* and *P. mucroniferus*.

Species and distribution: 12 species in central, S and E Anatolia and three additional in Iran, one of these also in Transcaucasia.

*Psephellus bornmuelleri* (Hausskn. ex Bornm.) Wagenitz, **comb. nova** ≡ *Centaurea bornmuelleri* Hausskn. ex Bornm. in Mitt. Thüring. Bot. Vereins 20: 21. 1905.

*P. brevifimbriatus* (Hub.-Mor.) Wagenitz, **comb. nova** ≡ *Centaurea brevifimbriata* Hub.-Mor. in Bauhinia 3: 315. 1967.

*P. gilanicus* (Bornm.) Wagenitz, **comb. nova** ≡ *Centaurea gilanica* Bornm. in Mitt. Thüring. Bot. Vereins 20: 23. 1905.

*P. gracillimus* (Wagenitz) Wagenitz, **comb. nova** ≡ *Centaurea gracillima* Wagenitz in Notes Roy. Bot. Gard. Edinburgh 33: 225. 1974.

*P. hadimensis* (Wagenitz, Ertugrul & Dural) Wagenitz, **comb. nova** ≡ *Centaurea hadimensis* Wagenitz, Ertugrul & Dural in Willdenowia 28: 157. 1998.

*P. holtzii* (Wagenitz) Wagenitz, **comb. nova** ≡ *Centaurea holtzii* Wagenitz in Notes Roy. Bot. Gard. Edinburgh 33: 226. 1974.

*P. huber-morathii* (Wagenitz) Wagenitz, **comb. nova** ≡ *Centaurea huber-morathii* Wagenitz in Notes Roy. Bot. Gard. Edinb. 33: 226. 1974.

*P. leuzeoides* (Jaub. & Spach) Wagenitz, **comb. nova** ≡ *Hyalea leuzeoides* Jaub. & Spach, Ill. Pl. Orient. 3: 21, 216. 1847.

*P. mucroniferus* (DC.) Wagenitz, **comb. nova** ≡ *Centaurea mucronifera* DC., Prodr. 6: 569. 1838.

*P. oltensis* Wagenitz, **nom. nov.** ≡ *Centaurea taochia* Sosn. in Věstn. Tiflissk. Bot. Sada, ser. 2, 5: 30. 1931.

*P. pergamaceus* (DC.) Wagenitz, **comb. nova** ≡ *Centaurea pergamacea* DC., Prodr. 6: 569. 1838.

*P. psephelloides* (Freyn & Sint.) Wagenitz, **comb. nova** ≡ *Centaurea psephelloides* Freyn & Sint. in Österr. Bot. Z. 42: 237. 1892.

*P. pyrrhoblepharus* (Boiss.) Wagenitz, **comb. nova** ≡ *Centaurea pyrrhoblephara* Boiss., Diagn. Pl. Orient., ser. 2, 3: 66. 1856.



*P. schischkinii* (Tzvelev) Wagenitz, **comb. nova**  $\equiv$  *Centaurea schischkinii* Tzvelev in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 20: 399. 1960.

*P. xeranthemoides* (Rech. f.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea xeranthemoides* Rech. f. in Anz. Österr. Akad. Wiss., Math.-Naturwiss. Kl. 88: 264. 1951.

Systematic position: Allied to *Psephellus* sect. *Aetheopappus* and sect. *Psephellus*.

Lit.: Sosnovsky 1931; Wagenitz 1975: 559-564, 1980: 399-402.

### *Psephellus* sect. *Psephellus*

$\equiv$  *Centaurea* sect. *Psephellus* (Cass.) DC., Prodr. 6: 575. 1838  $\equiv$  *Centaurea* subg. *Psephellus* (Cass.) Spach, Hist. Nat. Végét. (Phan.) 6: 11. 1841  $\equiv$  *Psephellus* sect. *Dealbati* Sosn. in Zаметki Sist. Geogr. Rast. 14: 12. 1948. – Type: *Centaurea dealbata* Willd.

$\equiv$  *Psephellus* sect. *Hypoleuci* Sosn. in Zаметki Sist. Geogr. Rast. 14: 6. 1948  $\equiv$  *Centaurea* sect. *Hypoleucae* (Sosn.) Sosn. in Bobrov & Čerepanov, Fl. SSSR 28: 424. 1963. – Type: *Centaurea hypoleuca* DC.

$\equiv$  *Psephellus* sect. *Leucophylli* Sosn. in Zаметki Sist. Geogr. Rast. 14: 7. 1948  $\equiv$  *Centaurea* sect. *Leucophyllae* (Sosn.) Sosn. in Bobrov & Čerepanov, Fl. SSSR 28: 428. 1963. – Type: *Centaurea leucophylla* M. Bieb.

*Perennial herbs*, often with ascending or prostrate stems with few branches and a central rosette of leaves. *Leaves* lyrate to pinnatisect or more rarely undivided, arachnoid on the upper side, densely tomentose on the lower. *Involucre* c. 14-21 mm long and 12-20 mm broad, cup-shaped. *Phyllaries* in several series, nearly glabrous; appendages membranous, triangular to ovate, ciliate, without prominent terminal mucro or spinule, not decurrent. *Flowers* purple or the central whitish, the marginal strongly to moderately radiant, infundibuliform, with 5-6 segments, with staminodes; central flowers numerous. *Achenes* 5-6 mm long, only slightly laterally compressed, detachment area lateral, with elaiosome. *Pappus* 1-1.5 mm, of deciduous scabrous bristles, inner row shorter.

Chromosome numbers: 12 counts, all with  $2n = 30$ , have been reported for different species (Matveeva & Tichonova ined. in Fedorov 1969; Žukova 1967, cited from Moore 1973; Čuksanova & al. 1968; Davlianidze 1985; Ghaffari & Chariat-Panahi in Taxon 34: 549. 1985). For *Psephellus somcheticus*  $2n = 30$  is known too, only Tonjan (1968) gave  $2n = 28$  for this species.

Species and distribution: 30 species are enumerated by Sosnovsky (1963) for the 'Flora SSSR area', where nearly all species of the section occur. Some have been added later (Galuško & Alieva 1976, Gabrieljan & Tahtadžjan 1992, Gabrieljan 1994), but species delimitation is very critical in this section and several of these may better be looked at as subspecies. The distribution centres in Caucasia, extending to the Crimea, E Anatolia and N Iran.

Systematic position: Allied to *Psephellus* sect. *Psephelloidei*, *Aetheopappus* and others, mainly distinguished by characters of the pappus.

Lit.: Sosnovsky 1948, 1963; Gabrieljan 1995.

### *Psephellus* sect. *Hyalinella* (Tzvelev) Wagenitz & Hellwig, **comb. nova**

Type: *Centaurea simplicicaulis* Boiss. & Huet  $\equiv$  *Psephellus simplicicaulis* (Boiss. & Huet) Wagenitz  $\equiv$  *Centaurea* sect. *Hyalinella* Tzvelev in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 426. 1959  $\equiv$  *Centaurea* subg. *Hyalinella* (Tzvelev) Tzvelev in Bobrov & Čerepanov, Fl. SSSR 28: 488. 1963.

$\equiv$  *Centaurea* sect. *Albovia* Tzvelev in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 421. 1959. – Type: *Centaurea pecho* Albov.



*Perennial herbs* with branched rootstock and erect or ascending stems, simple or with few branches in or below middle. *Leaves* arachnoid and glabrescent above, densely appressed-tomentose below, the lower pinnatisect to lyrate with elliptic to ovate lateral segments, sometimes undivided, middle leaves similar but with narrower segments or undivided like the few upper leaves. *Involucre* 12-19 mm long, (6-)8-14 mm broad, ovoid. *Phyllaries* in several series, nearly glabrous, the appendages nearly circular to broadly ovate, brown in the centre with hyaline or straw-coloured, denticulate or ciliate margin, not or very slightly decurrent. *Flowers* rose-coloured, the marginal scarcely to distinctly radiant, with staminodes, central flowers numerous. *Achenes* 4-5 mm long, laterally compressed, with lateral detachment area, without elaiosome. *Pappus* 5-8(-10) mm long, of scabrous bristles, inner row short, scaly.

Chromosome numbers: Three species belonging to the *Psephellus simplicicaulis* group have been counted with  $2n = 50-60$  (Poddubnaja-Arnoldi 1931).

Species and distribution: Seven species have been described, but several of these seem to belong to the polymorphous *Psephellus simplicicaulis*. The small species group is distributed in NE Anatolia and Transcaucasia.

*Psephellus bellus* (Trautv.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea bella* Trautv. in Bull. Acad. Imp. Sci. Saint-Pétersbourg 10: 394. 1866.

*P. pecho* (Albov) Wagenitz, **comb. nova**  $\equiv$  *Centaurea pecho* Albov in Bull. Herb. Boissier, ser. 1, 2: 640. 1894.

*P. simplicicaulis* (Boiss. & Huet) Wagenitz, **comb. nova**  $\equiv$  *Centaurea simplicicaulis* Boiss. & Huet in Boiss., Diagn. Pl. Orient., ser. 2, 3: 67. 1856.

Systematic position: Allied to *Psephellus* sect. *Psephelloidei* and sect. *Psephellus*.

Lit.: Cvelev 1963; Wagenitz 1975: 565-566.

***Psephellus* sect. *Aetheopappus*** (Cass.) Wagenitz & Hellwig, **comb. nova**

Type: *Centaurea pulcherrima* Willd.  $\equiv$  *Psephellus pulcherrimus* (Willd.) Wagenitz  $\equiv$  *Aetheopappus* Cass. in Cuvier, Dict. Sci. Nat. 50: 250. 1827  $\equiv$  *Centaurea* sect. *Aetheopappus* (Cass.) DC., Prodr. 6: 574. 1838  $\equiv$  *Centaurea* ser. *Aetheopappus* (Cass.) Benth. in Bentham & Hooker, Gen. Pl. 2: 480. 1873.

*Perennial herbs* with erect stem, usually unbranched or with 1-3 branches. *Leaves* tomentose on both sides or glabrescent above, very variable, ovate to narrowly lanceolate in outline, the lower entire, denticulate, lyrate or pinnatifid, the median and upper usually undivided, the upper with an appendage similar to that of the phyllaries. *Involucre* 18-28 mm long and 15-30 mm broad, subglobose. *Phyllaries* in several series, the appendages very large, membranous, brownish, triangular to lanceolate or ovate with numerous regular cilia, not decurrent. *Flowers* rose-purple or pale yellow, the marginal strongly radiant with more than 5 segments, with staminodes, the central numerous. *Achenes* 6-6.5 mm long, flattened, with small sublateral detachment area without elaiosome. *Pappus* 11-16 mm, of barbellate to plumose bristles, the inner row short in *Psephellus appendicigerus*, not distinct in *P. pulcherrimus*.

Species and distribution: Two main species, *Psephellus appendicigerus* in NE Anatolia, the polymorphous *P. pulcherrimus* in E Anatolia and Transcaucasia. The latter may actually comprise several distinct species.

*Psephellus appendicigerus* (C. Koch) Wagenitz, **comb. nova**  $\equiv$  *Centaurea appendicigera* C. Koch in Linnaea 24: 425. 1851.

*P. pulcherrimus* (Willd.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea pulcherrima* Willd., Sp. Pl. 3(3): 2298. 1803.

Systematic position: Allied to *Psephellus* sect. *Psephelloidei* and sect. *Psephellus*.

Lit.: Sosnovsky 1953, 1963: 353-356; Wagenitz 1975: 567-569.

***Psephellus* sect. *Amblyopogon* (DC.) Wagenitz & Hellwig, **comb. nova****Type: *Amberboa incanescens* DC.  $\equiv$  *Psephellus incanescens* (DC.) Boiss.

$\equiv$  *Amberboa* sect. *Amblyopogon* Fisch. & C. A. Mey. ex DC., Prodr. 6: 561. 1838  $\equiv$  *Amblyopogon* (DC.) Jaub. & Spach, Ill. Pl. Orient. 3: 23. 1847  $\equiv$  *Centaurea* sect. *Amblyopogon* (DC.) Sosn. in Věstn. Tiflissk. Bot. Sada, ser. 2, 5: 21. 1931  $\equiv$  *Centaurea* subg. *Amblyopogon* (DC.) Tzvelev in Bobrov & Čerepanov, Fl. SSSR 28: 472. 1963.

*Herbaceous perennials* with decumbent or erect and very short stems, simple or sparsely branched, stems (always?) arising from the base of the rosette. *Leaves* tomentose on both sides, lyrate, pinnatisect or undivided. *Involucre* 13-30 mm long, 10-20 mm broad, ovoid. *Phyllaries* in several series, appendages membranous, lanceolate or triangular, not or shortly decurrent, ciliate. *Flowers* rose-purple, the marginal not or scarcely radiant, with staminodes, the central rather numerous. *Achenes* 4-8 mm long, detachment area lateral, large (in *Psephellus karduchorum*), with elaiosome. *Pappus* usually short, 1-2.5 mm, of scabrous bristles or slender scales, lacking in *P. poluninii*, 8 mm long in *P. congestus*, inner row shortest, sometimes indistinctly so.

Chromosome numbers: Unknown.

Species and distribution: Five species are known, two of them only from the type, all with a restricted distribution in E Anatolia, Talysh and W Iran.

*Psephellus congestus* (Wagenitz) Wagenitz, **comb. nova**  $\equiv$  *Centaurea congesta* Wagenitz in Rechinger, Fl. Iran. 139b: 405. 1980.

*P. incanescens* (DC.) Boiss., Fl. Orient. 3: 610. 1875.

*P. integrifolius* C. Koch in Linnaea 24: 438. 1851  $\equiv$  *Centaurea integrifolia* C. A. Mey., Verz. Pfl. Casp. Meer.: 64. 1831, nom. illeg. [non Tausch 1828]  $\equiv$  *C. meyeranus* Tzvelev in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 418. 1959.

*P. karduchorum* (Boiss.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea karduchorum* Boiss., Fl. Orient. 3: 633. 1875.

*P. poluninii* (Wagenitz) Wagenitz, **comb. nova**  $\equiv$  *Centaurea poluninii* Wagenitz in Notes Roy. Bot. Gard. Edinburgh 33: 228. 1974.

Systematic position: Probably related to *Psephellus* sect. *Psephelloidei*, but with reduced marginal flowers and the “mymecochory syndrom”: short or prostrate stems, large achenes with short pappus and with elaiosome. *P. congesta* is aberrant with its long pappus.

Lit.: Sosnovsky 1931; Cvelev 1963: 472 ff.; Wagenitz 1975: 569-570, 1980: 404-406.

***Psephellus* sect. *Heterolophus* (Cass.) Wagenitz & Hellwig, **comb. nova****Type: *Centaurea sibirica* L.  $\equiv$  *Psephellus sibiricus* (L.) Wagenitz

$\equiv$  *Heterolophus* Cass. in Cuvier, Dict. Sci. Nat. 50: 250. 1827  $\equiv$  *Centaurea* subg. *Heterolophus* (Cass.) Spach, Hist. Nat. Végét. (Phaner.) 6: 11. 1841  $\equiv$  *Centaurea* sect. *Heterolophus* (Cass.) M. Dittrich in Bot. Jahrb. Syst. 88: 156. 1968.

*Perennials* with woody base and several simple or sparsely branched ascending or prostrate stems, arising from the base of the rosette. *Leaves* tomentose, especially on the lower side, partly pinnatipartite, partly undivided. *Involucre* c. 14-22 mm long and 10-20 mm broad, cup-shaped to subglobose. *Capitula* heterogamous. *Phyllaries* in several series, slightly woolly, appendages membranous, not decurrent, very variable from suborbicular to lanceolate or triangular or even linear-lanceolate with distinct long cilia or lacerate or dentate. *Flowers* rose-purple, the marginal distinctly radiant with 4-7 broad segments, with staminodes, central flowers numerous. *Achenes* 4-7 mm long, laterally compressed, detachment area lateral-basal, with elaiosome. *Pappus* 1-2.5(-3) mm long, of scabrous bristles, inner row short.

Chromosome numbers: Three counts with  $2n = 30$  have been reported for *Psephellus sibiricus* (Rostovzeva 1979, Pulkina 1988 [two provenances]). For *P. marschallianus*  $2n = 18$  has been re-

corded from cultivated material; especially in the case of cultivated plants misidentification may be involved, another count by Kuzmen & Georgieva (in *Taxon* 32: 665. 1983) revealed  $2n = 20$ . Checking of these numbers is necessary.

Species and distribution: According to Klokov (1963) there are six species distributed from Bulgaria, Ukraine and S Russia to W Siberia and Central Asia. They form evidently a closely knit group as all of them have been assigned either to *Centaurea sibirica* or *C. marschalliana*. The variability of the appendages is very marked in this section.

*Psephellus marschallianus* (Spreng.) C. Koch in *Linnaea* 24: 438  $\equiv$  *Centaurea marschalliana* Spreng., *Syst. Veg.* 3: 398. 1826.

*P. sibiricus* (L.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea sibirica* L., *Sp. Pl.* 2: 913. 1753.

Systematic position: Closely related to *Psephellus* sect. *Amblyopogon* and *Czerniakovskya*. *P.* sect. *Amblyopogon* is distinguished mainly by its inconspicuous, not radiant marginal flowers.

Lit.: Klokov 1963.

***Psephellus* sect. *Czerniakovskya* (Czerep.) Wagenitz & Hellwig, **comb. nova****

Type: *Centaurea kopet-daghensis* Iljin  $\equiv$  *Psephellus kopet-daghensis* (Iljin) Wagenitz  $\equiv$  *Centaurea* subg. *Czerniakovskya* Czerep. in Bobrov & Čerepanov, *Fl. SSSR* 28: 609. 1963  $\equiv$  *Centaurea* sect. *Czerniakovskya* (Czerep.) Wagenitz in Rechinger, *Fl. Iran.* 139b: 407. 1980.

*Perennial* with woody rhizome with several ascending to nearly prostrate stems. *Stems* simple or with few branches. *Leaves* tomentose, lanceolate. *Involucre* 15-20 mm long and 8-15 mm broad, cup-shaped. *Phyllaries* in several series, woolly, appendages triangular, ciliate, the terminal mucro scarcely differing from the cilia, not decurrent (*Psephellus kopet-daghensis*) or decurrent (*P. iljinii*). *Capitula* heterogamous. *Flowers* pale yellow to rose-coloured, the sterile marginal not or slightly radiant, with staminodes, the central hermaphrodite numerous. *Achenes* 7-8 mm long, laterally compressed, detachment area lateral, with(?) elaiosome. *Pappus* 3-8 mm long, of scabrous bristles, the inner row of shorter scales.

Chromosome numbers: Unknown.

Species and distribution: Only two species known from Turkmenia (Kopet-Dagh) and N Iran. The species are similar in habit but differ remarkably in the appendages.

*Psephellus iljinii* (Czerniak.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea iljinii* Czerniak. in *Repert. Spec. Nov. Regni Veg.* 27: 285. 1930.

*P. kopet-daghensis* (Iljin) Wagenitz, **comb. nova**  $\equiv$  *Centaurea kopet-daghensis* Iljin in *Izv. Bot. Sada Akad. Nauk SSSR* 30: 359. 1932.

Systematic position: Closely allied to *Psephellus* sect. *Amblyopogon* and perhaps not really distinct, as the length of the pappus is the main distinguishing character.

Lit.: Čerepanov 1963: 418-420; Wagenitz 1980: 407-409.

***Psephellus* sect. *Odontolophoidei* (Tzvelev) Wagenitz & Hellwig, **comb. nova****

Type: *Centaurea aucheriana* DC.  $\equiv$  *Psephellus aucherianus* (DC.) Boiss. ( $\equiv$  *Phaeopappus microcephalus* Jaub. & Spach, nom. illeg.)

$\equiv$  *Centaurea* sect. *Odontolophoideae* Tzvelev in *Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR* 19: 422. 1959  $\equiv$  *Centaurea* subg. *Odontolophopsis* Tzvelev in Bobrov & Čerepanov, *Fl. SSSR* 28: 485. 1963  $\equiv$  *Phaeopappus* sect. *Psephelloides* Jaub. & Spach, *Ill. Pl. Orient.* 3: 16. 1847, non *Centaurea* sect. *Psephelloideae* Boiss. 1875, nec *Psephellus* sect. *Psephelloidei* (Boiss.) Wagenitz & Hellwig 2000.

*Perennial herbs*, woody at base, with several stems branched from near the base or in the lower half. *Leaves* floccose or appressed tomentose, the lower (and median) petiolate and lyrate,

pinnatifid or pinnatipartite or undivided, the median and upper sessile, lanceolate to nearly linear. *Involucre* 10-20 mm long, 6-22 mm broad, ovoid to nearly globose. *Phyllaries* in several series, appendages membranous, straw-coloured, concealing the basal part of the phyllaries or nearly so, ovate to circular, ciliate, not decurrent. *Capitula* heterogamous. *Flowers* pink, the marginal not or slightly radiant, with 5-6 segments, with staminodes. *Achenes* 4-6 mm long, laterally compressed, smooth, detachment area lateral, relatively large. *Pappus* 1-4.5 mm long, of scabrous bristles, the inner row short.

Chromosome numbers: Only one count,  $2n = 30$  for *Psephellus phaeopappoides* (Garcia-Jacas & al. 1998), has been reported.

Nomenclature: The oldest name at the rank of a section is *Phaeopappus* sect. *Psephelloides* Jaub. & Spach, but this must be looked at as an orthographic variant of *Centaurea* sect. *Psephelloideae* Boiss.  $\equiv$  *Psephellus* sect. *Psephelloidei* (Boiss.) Wagenitz & Hellwig and cannot be used in *Psephellus* at the same time.

Species and distribution: Three species in E Anatolia, Transcaucasia and NW Iran.

*Psephellus aucherianus* (DC.) Boiss., Fl. Orient. 3: 611. 1875.

*P. eugenii* (Sosn.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea eugenii* Sosn. in Věstn. Tiflissk. Bot. Sada, ser. 2, 5: 32. 1931.

*P. phaeopappoides* (Bordzil.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea phaeopappoides* Bordzil. in Žurn. Inst. Bot. Vseukrajins'k. Akad. Nauk 3: 82. 1935.

Systematic position: Closely allied to *Psephellus* sect. *Xanthopsis*. Leaves similar to those in *P.* sect. *Odontolophus*.

Lit.: Cvelev 1959, 1963: 485-486; Wagenitz 1975: 570.

***Psephellus* sect. *Odontolophus*** (Cass.) Wagenitz & Hellwig, **comb. nova**

Type: *Centaurea trinervia* Willd.  $\equiv$  *Psephellus trinervius* (Willd.) Wagenitz

$\equiv$  *Odontolophus* Cass. in Cuvier, Dict. Sci. Nat. 50: 252. 1827  $\equiv$  *Centaurea* sect. *Odontolophus* (Cass.) DC., Prodr. 6: 579. 1838  $\equiv$  *Centaurea* subg. *Odontolophus* (Cass.) Spach, Hist. Nat. Vég. (Phaner.) 6: 11. 1841.

*Perennial* with woody rhizome, stems strongly branched near the base with numerous long branches with one capitulum. *Leaves* narrowly lanceolate, thinly tomentose, indumentum of woolly hairs and hairs with pedestal. *Involucre* 15-17 mm long, c. 10 mm broad, cup-shaped to subcylindric. *Phyllaries* in several series, glabrous, appendages hyaline, ovate to suborbicular, shortly ciliate or lacerate, decurrent. *Capitula* heterogamous. *Marginal flowers* pink, radiant, with 7-8 broad segments; *central flowers* numerous, whitish to pink, anther tube rose-violet. *Achenes* 4-6 mm, laterally compressed, with relatively large basal-lateral detachment area, with biseriate glandular hairs, with elaiosome. *Pappus* 1-2.5 mm long, of scabrous bristles, inner row similar, but shorter.

Chromosome numbers: Only one count is available:  $2n = 32$  (Guinochet & Foissac 1962). This may be a real deviation from the usual number or due to B-chromosomes.

Species and distribution: The best known species, *Psephellus trinervius*, is distributed from Romania (Carpathian) through S Russia (Ukraine?) and the Crimea to Caucasia. Two other species are from Caucasia.

*Psephellus avaricus* (Tzvelev) Wagenitz, **comb. nova**  $\equiv$  *Centaurea avarica* Tzvelev in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 20: 27. 1960.

*P. kobstanicus* (Tzvelev) Wagenitz, **comb. nova**  $\equiv$  *Centaurea kobstanica* Tzvelev in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 425. 1959.

*P. trinervius* (Willd.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea trinervia* Steph. ex Willd., Sp. Pl. 3(3): 2301. 1803.

Systematic position: Allied to *Psephellus* sect. *Uralepis* and *Sosnovskya*.

Lit.: Janka 1882; Cvelev 1963: 481-485.

***Psephellus* sect. *Xanthopsis* (DC.) Wagenitz & Hellwig, **comb. nova****

Type: *Amberboa xanthocephala* DC.  $\equiv$  *Psephellus xanthocephalus* (DC.) Boiss. & Buhse  
 $\equiv$  *Amberboa* sect. *Xanthopsis* DC., Prodr. 6: 561. 1838  $\equiv$  *Xanthopsis* (DC.) C. Koch in *Linnaea*  
 24: 422. 1851  $\equiv$  *Amblyopogon* sect. *Xanthopsis* (DC.) Grossh. in *Trudy Prikl. Bot. Ser.1. Sist.*  
*Rast. Obščie Vopr. Rasteniev.* 2: 243. 1937  $\equiv$  *Centaurea* sect. *Xanthopsis* (DC.) Tzvelev in  
*Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR* 19: 418. 1959  $\equiv$  *Centaurea* subg.  
*Xanthopsis* (DC.) Tzvelev in *Bobrov & Čerepanov, Fl. SSSR* 28: 474. 1963.

*Perennials* with woody base, branched from near the base, the branches with one to several capitula, tomentose especially in the lower part. *Leaves* appressed-tomentose, more or less glabrescent, the lower petiolate, the upper sessile, all undivided and linear-lanceolate to lanceolate or the lower pinnatipartite to lyrate. *Involucre* 11-18 mm long, 6-11 mm broad, ovoid. *Phyllaries* in several series, their glabrous basal part totally concealed by the large appendages. *Appendages* membranous, whitish to straw-coloured, ovate or hemispherical to nearly circular, not decurrent, denticulate to ciliate. *Capitula* heterogamous. *Flowers* whitish to sulphur-yellow, the marginal with staminodes, nearly filiform, not radiant, about as long or slightly shorter than the central ones, with 4-5 segments, the central rather few, hermaphrodite. *Achenes* 5-7 mm long, laterally compressed, detachment area lateral, small, probably without elaiosome. *Pappus* 0.5-1 mm (7-8 mm in *Psephellus straminicephalus*), of scabrous bristles, inner row scarcely different (shorter or equal in *P. straminicephalus*).

Chromosome numbers: Two counts are available for *Psephellus xanthocephalus*:  $n = 15$  (Bakshi Khaniki 1995) and  $2n = 30 + 0-3 B$  (Garcia Jacas & al. 1998). Tonjan (1968) gives  $2n = 32$  for *P. erivanensis*.

Species and distribution: Three or four species in E Anatolia, the Aras valley and NW Iran. *Psephellus straminicephalus* differs by the long pappus.

*Psephellus erivanensis* Lipsky, *Fl. Caucasi*, suppl. 1 [= *Acta Horti Tiflis* 6]: 64. 1902

*P. straminicephalus* (Hub.-Mor.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea straminicephala* Hub.-Mor. in *Bauhinia* 3: 316. 1967.

*P. xanthocephalus* (DC.) Fisch. & C. A. Mey. ex Boiss. & Buhse in *Nouv. Mém. Soc. Imp. Naturalistes Moscou* 12: 129. 1860  $\equiv$  *Amberboa xanthocephala* DC., Prodr. 6: 561. 1838.

Systematic position: Close to *Psephellus* sect. *Sosnovskya*. *P.* sect. *Amblyopogon* has some affinities too, but differs in habit.

Lit.: Gabrieljan 1988; Grossgejm 1937; Cvelev 1963: 474-477; Wagenitz 1975: 571-572, 1980: 411-412.

***Psephellus* sect. *Uralepis* (DC.) Wagenitz & Hellwig, **comb. nova****

Type: *Microlonchus persicus* DC.  $\equiv$  *Psephellus persicus* (DC.) Wagenitz  
 $\equiv$  *Microlonchus* sect. *Uralepis* DC., Prodr. 5: 563. 1838  $\equiv$  *Centaurea* sect. *Uralepis* (DC.) Wagenitz in *Bot. Jahrb. Syst.* 82: 195. 1963.

*Perennial* with woody base and several erect to ascending stems, branched from the base with long branches with one capitulum. Lower *leaves* tomentose, lanceolate to rhomboid, sometimes lyrate, denticulate with cartilaginous teeth, median and upper nearly glabrous, linear-lanceolate or linear. *Involucre* 13-16 mm long and (5-)8-12 mm broad, cup-shaped. *Phyllaries* in several series, coriaceous, glabrous, appendage crescent-shaped, hyaline, entire or denticulate, decurrent. *Flowers* ochroleucous, the marginal not radiant, sterile, with staminodes and sometimes a rudimentary style, with filiform segments. *Achenes* 6-7 mm long, laterally compressed, detachment

area small, lateral, probably without elaiosome. *Pappus* 7-9 mm long, of scabrous bristles, the inner row of scales 3-4(-5) mm long.

Chromosome numbers: Unknown.

Species and distribution: Only one species, endemic in W and central Iran.

*Psephellus persicus* (DC.) Wagenitz, **comb. nova**  $\equiv$  *Microlonchus persicus* DC., Prodr. 6: 563. 1838 = *Centaurea gaubae* (Bornm.) Wagenitz in Bot. Jahrb. Syst. 82: 195. 1963.

Nomenclature: In *Psephellus* the oldest available epithet must be used again. For details see Wagenitz (1963: 196).

Systematic position: Although superficially similar, *Psephellus persicus* has surely no affinity with *Mantisalca* (*Microlonchus*) as supposed by Candolle. It clearly belongs to the group of sections with pollen of the *Centaurea dealbata* type, but is rather isolated. The habit is similar to *P.* sect. *Odontolophus* and *Sosnovskya*

Lit.: Wagenitz 1963: 195-197, 1980: 410-411.

***Psephellus* sect. *Sosnovskya* (Takht.) Wagenitz & Hellwig, **comb. nova****

Type: *Centaurea amblyolepis* Ledeb.  $\equiv$  *Psephellus amblyolepis* (Ledeb.) Wagenitz

$\equiv$  *Sosnovskya* Takht. in Sovetsk. Bot. 1936(5): 98. 1936  $\equiv$  *Centaurea* sect. *Sosnovskya* (Takht.) Wagenitz in Bot. Jahrb. Syst. 82: 193. 1963  $\equiv$  *Centaurea* subg. *Sosnovskya* (Takht.) Czerep. in Bobrov & Čerepanov, Fl. SSSR 28: 477. 1963.

*Perennials* with woody base, stems branched near the base with numerous long branches with one capitulum, woolly at base. *Leaves* thinly tomentose, glabrescent, narrowly lanceolate to entire, the lower sometimes with few distant lobes or teeth or lyrate, the upper linear, nearly needle-like. *Involucre* c. 15-17 mm long, 8-13 mm broad, cup-shaped. *Phyllaries* in few series, glabrous, the appendages triangular to crescent-shaped, not or slightly decurrent, ciliate. *Capitula* heterogamous. *Flowers* whitish or pink, the marginal slightly radiant, with 5-6 narrow segments, with staminodes, the central flowers rather numerous. *Achenes* 5.5-6 mm long, detachment area small (only 1.5 mm in *P. amblyolepis*), lateral-basal. *Pappus* 3-5 mm long, of scabrous bristles, the inner row slightly shorter or not differentiated.

Chromosome numbers: Unknown.

Species and distribution: Four species occur in the Caucasian region (Čerepanov 1963). They are apparently closely related and two of them may be only subspecies of *Psephellus amblyolepis*. A further species, *P. galactochrous* (Rech. f.) Parsa, occurs in N Iran.

*Psephellus amblyolepis* (Ledeb.) Wagenitz, **comb. nova**  $\equiv$  *Centaurea amblyolepis* Ledeb., Fl. Ross. 2: 703. 1845  $\equiv$  *Phaeopappus amblyolepis* (Ledeb.) Boiss., Fl. Orient. 3: 601. 1875  $\equiv$  *Sosnovskya amblyolepis* (Ledeb.) Takht. in Sovetsk. Bot. 1936(5): 99. 1936 & in Repert. Spec. Nov. Regni Veg. 41: 191. 1936.

*P. arvensis* (Czerep.) Wagenitz, **comb. nova**  $\equiv$  *Sosnovskya arvensis* Czerep. in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. SSSR 20: 484. 1960  $\equiv$  *Centaurea arvensis* (Czerep.) Czerep. in Bobrov & Čerepanov, Fl. SSSR 28: 481. 1963.

*P. galactochrous* (Rech. f.) Parsa, Fl. Iran 3: 633. 1948  $\equiv$  *Centaurea galactochroa* Rech. f. in Repert. Spec. Nov. Regni Veg. 48: 153. 1940.

Systematic position: Closely related to *Psephellus* sect. *Odontolophus* and probably better included in it, differing mainly in the less radiant marginal flowers and not or scarcely decurrent appendages.

Lit.: Wagenitz 1963: 193-195; Čerepanov 1963: 477-481.



## Acknowledgements

This study is partly based on investigations of micromorphological and molecular characters made possible by a grant of the Deutsche Forschungsgemeinschaft (DFG). This is gratefully acknowledged. D. Petit and an unknown reviewer made valuable suggestions for improving the text.

## References

- Avetisjan, E. M. 1964: K palinosistematike nekotoryh rodov triby *Centaureinae* semejstva *Asteraceae*. – Trudy Bot. Inst. Akad. Nauk Armjansk. S.S.R. **14**: 31-47.
- Bakshi Khaniki, G. 1995: Meiotic studies on some Iranian *Centaurea* (*Compositae*). – [Cytologia](#) **60**: 341-346.
- Boissier, E. 1875: Flora orientalis **3**. – Basel & Genève.
- Bobrov, E. G. & Čerepanov S. K. (ed.) 1963: Flora SSSR **28**. – Moskva & Leningrad.
- Bremer, K. 1994: *Asteraceae*: Cladistics and classification. – Portland.
- Čerepanov, S. K. 1963: *Centaurea* podrod 9. *Sosnovskya* (Takht.) Czer. comb. nova. – Pp. 477-481 in: Bobrov, E. G. & Čerepanov S. K. (ed.), Flora SSSR **28**. – Moskva & Leningrad.
- Čuksanova, N. A., Svešnikova, L. I. & Alexandrova, T. V. 1968: Novyje dannye o čislah hromosom u vidov semejstva sloznocvetnyh. [Data on the karyology of the family *Compositae* Giseke]. – Citologija [Tsitologija] **10**: 198-206.
- Cvelev, N. 1959: Kritičeskie zametki o nekotoryh sekczijah roda *Centaurea* L. [Notulae criticae de sectionibus nonnullis generis *Centaureae* L.] – Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR **19**: 409-441.
- 1963: *Centaurea* podrod 13. *Hyalinella* (Tzvel.) Tzvel. comb. nova. – Pp. 488-493 in: Bobrov, E. G. & Čerepanov S. K. (ed.), Fl. SSSR **28**. – Moskva & Leningrad.
- Davlianidze, M. T. 1985: Čisla hromosom predstavitelej semejstv *Asteraceae*, *Boraginaceae*, *Brassicaceae*, *Liliaceae*, *Fabaceae*, *Paeoniaceae*, *Poaceae*, *Primulaceae*, *Ranunculaceae*, *Rosaceae* flory Gruzinskoj SSR. [Chromosome numbers in the representatives of the flora from Georgia]. – Bot. Žurn. (Moscow & Leningrad) **70**: 698-700.
- Dittrich, M. 1968: Karpologische Untersuchungen zur Systematik von *Centaurea* und verwandten Gattungen. – Bot. Jahrb. Syst. **88**: 79-162.
- 1977: *Cynareae* – systematic review. – Pp. 999-1015 in: Heywood, V. H., Harborne, J. B. & Turner, B. L. (ed.), The biology and chemistry of the *Compositae* **2**. – London, etc.
- Dostál, J. 1973: Preliminary notes on the subtribe *Centaureinae*. – Acta Bot. Acad. Sci. Hung. **19**: 73-79.
- Fedorov, A. A. (ed.) 1969: Hromosomnye čisla cvetkovykh rastenij. [Chromosome numbers of flowering plants]. – Leningrad.
- Gabrieljan, E. 1988: Obzor vidov podroda *Xanthopsis* (DC.) Tzvel. roda *Centaurea* L. (*Asteraceae*). [Synopsis specierum generis *Centaurea* L. (*Asteraceae*) subgeneris *Xanthopsis* (DC.) Tzvelev]. – Novosti Sist. Vysš. Rast. **25**: 160-171.
- 1994: Novye taksony podroda *Psephellus* roda *Centaurea* (*Asteraceae*) iz Armenii i Irana [New taxa of the subgenus *Psephellus* of the genus *Centaurea* from Armenia and Iran]. – Bot. Žurn. (Moscow & Leningrad) **79**(4):120-129.
- 1995: *Psephellus* Cass. – Pp. 336-352 in: Tahtadžjan, A. (ed.), Flora Armenii **9**. – Havlickuv Brod.
- & Tahtadžjan, A. L. 1992: Novyj vid *Centaurea cronquistii* (*Asteraceae*) iz Armenii. [A new species of [sic] *Centaurea cronquistii* (*Asteraceae*) from Armenia.]. – Bot. Žurn. (Moscow & Leningrad) **77**(9): 65-67.
- Galuško, A. & Alieva, A. 1976: Novyj vid roda *Psephellus* Cass. (*Asteraceae*) e severnogo



- Kavkaza. [Nova species generis *Psephellus* Cass. (Asteraceae) e Caucasi boreali]. – *Novosti Sist. Vysš. Rast.* **13**: 246-248.
- Garcia-Jacas, N., Susanna, A. & Mozaffarian, V. 1998: New chromosome counts in the subtribe *Centaureinae* (Asteraceae, Cardueae) from West Asia III. – *Bot. J. Linn. Soc.* **128**: 413-422.
- Grossgejm, A. A. 1937: Materialy k poznannju dekorativnyh “suhocvetov” Zakavkaz’ja. Rod *Amblyopogon* (F. et. Mey.) J. & Sp. [A contribution to the knowledge of the ornamental xerophytes of Transcaucasia. The genus *Amblyopogon* (F. & M.) J. & Sp.]. – *Trudy Prikl. Bot. Ser. 1. Sist. Rast. Obščie Vopr. Rasteniev.* **2**: 241-248.
- Guinochet, M. & Foissac, J. 1962: Sur les caryotypes de quelques espèces du genre *Centaurea* L. – *Rev. Cytol. Biol. Végét.* **25**: 373-389.
- Hellwig, F. H. 1996: Untersuchungen zur Phylogenie der *Cardueae-Centaureinae* (Compositae) unter Verwendung molekularer und morphologisch-anatomischer Merkmale. – *Habilitationsschrift Univ. Göttingen.* – Göttingen.
- Hoffmann, O. 1894: *Compositae*. – Pp. 87-387 in: Engler, A. & Prantl, K., *Die natürlichen Pflanzenfamilien* **4(5)**. – Leipzig.
- Janka, V. v. 1882: *Odontolophus*, eine ausgezeichnete Gattung. – *Österr. Bot. Z.* **32**: 280-281.
- Klokov, M. V. 1963: *Centaurea* podrod 6. *Heterolophus* (Cass.) Dobrocz. – Pp. 463-472 in: Bobrov, E. G. & Čerepanov S. K. (ed.), *Flora SSSR* **28**. – Moskva & Leningrad.
- Moore, R. J. (ed.) 1973: Index to plant chromosome numbers 1967-1971. – *Regnum Veg.* **90**.
- Petit, D. P. 1997: Generic relationships of the *Cardueae* (Compositae): a cladistic analysis of morphological data. – *Pl. Syst. Evol.* **207**: 173-203.
- Poddubnaja-Arnoldi, W. 1931: *Vergleichende embryologisch-zytologische Untersuchungen über die Gruppe Cynareae, Fam. Compositae.* – *Beih. Bot. Centralbl., Abt. II*, **48**: 141-237.
- Pulkina, S. V. 1988: Čisla hromosom nekotoryh vidov semejstva Asteraceae. [Chromosome numbers in some species of the Asteraceae]. – *Bot. Žurn. (Moscow & Leningrad)* **73**: 607-608.
- Rostovzeva, T. S. 1979: Čisla hromosom nekotoryh vidov sem. Asteraceae Dumort. [Chromosome numbers of some species of the family Asteraceae Dumort.]. – *Bot. Žurn. (Moscow & Leningrad)* **64**: 582-589.
- Sosnovsky, D. 1931: O dvuh sekcijah roda *Centaurea* L. s. str. [On two sections of the genus *Centaurea* L. s. str.]. – *Věstn. Tiflissk. Bot. Sada [= Monit. Jard. Bot. Tiflis N. S.]*, ser. 2, **5**: 21-34.
- 1948: Obzor predstavitelej roda *Psephellus* (Cass.) D. Sosn. [Specierum caucasicarum generis *Psephelli* (Cass.) D. Sosn. emend. revisio]. – *Zametki Sist. Geogr. Rast.* **14**: 5-22.
- 1953: Predvaritel’nye rezul’taty revizii *Aetheopappus* Cass. [Prodromus generis *Aetheopappi* Cass.] – *Zametki Sist. Geogr. Rast.* **17**: 4-16.
- 1963: *Centaurea* podrod 4. *Psephellus* (Cass.) Schmalh. – Pp. 420-440 in: Bobrov, E. G. & Čerepanov S. K. (ed.), *Flora SSSR* **28**. – Moskva & Leningrad.
- Susanna, A., Garcia Jacas, N., Soltis, D. E. & Soltis, P. S. 1995: Phylogenetic relationships in tribe *Cardueae* (Asteraceae) based on ITS sequences. – *Amer. J. Bot.* **82**: 1056-1068.
- Tahtadžjan, A. (ed.) 1995: *Flora Armenii* 9. *Campanulaceae, Asteraceae.* – Havlickuv Brod.
- Tonjan, Z. R. 1968: Čisla hromosom nekotoryh vidov roda *Centaurea*. [The chromosome numbers of some species of the genus *Centaurea*]. – *Biol. Žurn. Armenii* **21**: 86-96.
- Wagenitz, G. 1955: *Pollenmorphologie und Systematik in der Gattung Centaurea L. s.l.* – *Flora* **142**: 213-279.
- 1963: Die Eingliederung der “*Phaeopappus*”-Arten in das System von *Centaurea*. – *Bot. Jahrb. Syst.* **82**: 137-215. 1963.
- 1975: *Centaurea* L. – Pp. 465-585 in: Davis, P. H. (ed.), *Flora of Turkey and the East Aegean Islands* **5**. – Edinburgh.
- 1980: *Centaurea* L. – Pp. 313-420 in: Rechinger, K. H. (ed.), *Flora iranica* **139b**. – Graz.

- & Hellwig, F. H. 1996: Evolution of characters and phylogeny of the *Centaureinae*. – Pp. 491-510 in: Hind, D. J. N. & Beentje, H. J. (ed.): *Compositae: Systematics*. Proc. Int. *Compositae* Conf., Kew, 1994, **1**. – Kew.

Addresses of the authors:

Prof. em. Dr Gerhard Wagenitz, Albrecht-von-Haller Institut für Pflanzenwissenschaften, Abt. Systematische Botanik, Untere Karspüle 2, D-37073 Göttingen; e-mail: gwageni@gwdg.de

Prof. Dr F. H. Hellwig, Institut für Spezielle Botanik, Universität Jena, Philosophenweg 16, D-07743 Jena; e-mail: hellwig@otto.biologie.uni-jena.de