

1 **Vegetation history of central Chukotka deduced from permafrost paleoenvironmental**
2 **records of the El'gygytgyn Impact Crater**

3
4 **A.A. Andreev¹, E. Morozova², G. Fedorov², L. Schirrmeister³, A.A. Bobrov⁴, F. Kienast⁵,**
5 **G. Schwamborn³**

6
7 ¹Institute of Geology and Mineralogy, University of Cologne, Zùlpicher St. 49a, D-50674, Cologne,
8 Germany

9 ²Arctic and Antarctic Research Institute, Bering St. 38, St. Petersburg, 199397 Russia

10 ³Alfred Wegener Institute for Polar and Marine Research, Department of Periglacial Research,
11 Telegrafenberg A43, D-14473 Potsdam, Germany

12 ⁴Faculty of Soil Science, Moscow State University, Vorobievsky Gory, 119899 Moscow, Russia

13 ⁵Senckenberg, Research Institute and Natural History Museum, Research Station for
14 Quaternary Paleontology, am Jakobskirchhof 4, 99423 Weimar, Germany

15
16 *Correspondence to:* A. Andreev

17 (aandreev@uni-koeln.de)

18
19 **Abstract.** Frozen sediments from three cores bored in the permafrost surrounding of the El'gygytgyn
20 Impact Crater Lake have been studied for pollen, non-pollen palynomorphs, plant macrofossils and
21 rhizopods. The palynological study of these cores contributes to a higher resolution of time intervals
22 presented in a poor temporal resolution in the lacustrine sediments; namely the Allerød and succeeding
23 periods. Moreover, the permafrost records better reflect local environmental changes, allowing a more
24 reliable reconstruction of the local paleoenvironments. The new data confirm that shrub tundra with
25 dwarf birch, shrub alder and willow dominated the lake surroundings during the Allerød warming.
26 Younger Dryas pollen assemblages reflect abrupt changes to grass-sedge-herb dominated
27 environments reflecting significant climate deterioration. Low shrub tundra with dwarf birch and
28 willow dominate the lake vicinity at the onset of the Holocene. The find of larch seeds indicate its

29 local presence around 11,000 cal. yr BP and, thus a northward shift of treeline by about 100 km during
30 the early Holocene thermal optimum. Forest tundra with larch and shrub alder stands grew in the area
31 during the early Holocene. After ca. 3500 cal. yr BP similar-to-modern plant communities became
32 common in the lake vicinity.

33

34 **1 Introduction**

35 El'gygytgyn Impact Crater is located in central Chukotka, approximately 100 km north of the
36 Arctic Circle (Fig. 1). The crater was formed 3.6 Myr ago (Gurov and Gurova, 1979; Layer,
37 2000). As inferred from geomorphologic research, the study area was never glaciated after the
38 time of the impact ca. 3.6 Myr ago (e.g. Brigham-Grette et al., 2007 and references therein),
39 and thus, the lake is probably the longest archive for Arctic terrestrial environmental and
40 climate history. Elgygytgyn Late Quaternary lacustrine palynological records were first
41 reported by Shilo et al. (2001) followed by more continuous and detailed records published by
42 Lozhkin et al. (2007) and Matrosova (2009). The studied sediments comprise the oldest
43 continuous Quaternary pollen record in the Arctic, which provides history of vegetation and
44 climate changes since ca. 350 kyr.

45 Generally, sediments from large and deep lakes are valuable paleoenvironmental archives
46 which contain pollen data reflecting vegetation and climate history of surrounding areas.
47 However, such pollen records reflect predominately regional environmental changes because
48 of the large input of long distance wind-transported pollen into the spectra. The Lake
49 El'gygytgyn sediments, where the pollen from a several thousand square-kilometer source
50 area is trapped, also provide a reliable record of extra-regional vegetation and climate changes
51 (Lozhkin et al., 2007; Matrosova 2009; Lozhkin and Anderson 2011). The importance of such
52 continuous long-term regional records is obvious. Nevertheless, short-term palynological
53 records reflecting local paleoenvironmental dynamics are also highly desired. These records
54 document predominate changes in local vegetation and may be compared with extra-regional

55 variations in order to better understand the role of local and regional vegetation in the
56 paleobotanical records, resulting in more reliable environmental reconstructions. Moreover,
57 these records often have better temporal resolution for some abrupt changes such as Younger
58 Dryas providing unique possibilities for high-resolution environmental studies.

59 Palynological studies of surface samples from the study area complement reliable
60 reconstructions. A total of 56 surface sediment samples from Lake El'gygytyn and 26
61 surface soil samples from the crater slopes have been recently studied (Matrosova et al., 2004;
62 Matrosova, 2006, 2009; Glushkova et al., 2009). These studies demonstrate that pollen of
63 trees and shrubs may reach up to 82% of the recent lacustrine spectra although the only
64 willow and dwarf birch stands grow in the crater in protected locations. Although soil pollen
65 spectra reliably reflect the local vegetation, pollen of long-distance-transported taxa dominate
66 even there (Matrosova, 2006; Glushkova et al., 2009). It is characteristic that pollen contents
67 of *Pinus pumila* and *Alnus fruticosa*, species not growing in the crater vicinity, may reach up
68 to 15 and 37% of the spectra consequently. Thus, by interpretations of fossil pollen
69 assemblages it has to be taken in a consideration that a significant part of the pollen may have
70 originated from some dozens and even hundreds of kilometers away.

71 This paper presents palaeoenvironmental and palaeoclimatic changes during the Lateglacial
72 and Holocene inferred from permafrost pollen, plant macrofossil, and rhizopod records from
73 the permafrost surrounding of the El'gygytyn Crater Lake. The Lateglacial/Holocene
74 transition is considered as a unique period of intensive glaciation and deglaciation events
75 accompanied by remarkable changes in global temperature, atmospheric circulation, air
76 humidity, precipitation and vegetation (Johnsen et al., 1995, Stuiver et al., 1995, Blunier and
77 Brook, 2001). Our studies of three permafrost cores add to a better understanding of
78 palaeoenvironmental changes during these time intervals which are not well represented in a
79 high temporal resolution in the lacustrine archive. A comparison of the palynological data
80 from the new permafrost cores and previously studied exposures and lake cores were used to

81 make a local chronostratigraphy scheme because of the partly insufficient geochronological
82 datasets. Such comparison resulted in a more reliable reconstruction of vegetation and climate
83 changes, especially during the transitional intervals from cold to warm periods.

84

85 **2 Geographical setting**

86 The El'gygytgyn Impact Crater is 18 km in diameter and holds a ca 170 m deep lake that has
87 a bowl-shaped morphology ca 12 km in diameter (Fig. 1). The crater is superimposed on the
88 Anadyr lowland and was formed in an Upper Cretaceous volcanic plateau (Belyi, 1998). The
89 crater rim comprises peaks between 600 and 930 m above sea level (a.s.l.), and the lake level
90 is situated at 492 m a.s.l. Unconsolidated Quaternary permafrost deposits cover the crater
91 bottom surrounding the lake. They show a distinctly asymmetrical distribution with a broad
92 fringe of loose sediment that is 500 to 600 m wide to the north and west, and only 10 to 20 m
93 elsewhere around the lake (Fig. 1).

94 The study area belongs to the continuous permafrost zone with a mean annual ground
95 temperature of -10 °C at 12.5 m depth (Schwamborn et al., 2008). In 2003, the active layer
96 was about 40 cm deep in peaty silts and reached 50 to 80 cm in sand, pebbles, and gravels.
97 The region is characterized by extremely harsh climate with average annual air temperature
98 ca. -10 °C, mean July temperatures of 4 to 8 °C and mean January temperatures of -32 to -36
99 °C. The precipitation consists of 70 mm summer rainfall (June-September) and ca. 110 mm
100 water equivalent of snowfall (Nolan and Brigham-Grette 2007). Climate variables are
101 strongly depending on oceanic influence expressed in decreasing summer temperatures
102 (Kozhevnikov, 1993). According to Kozhevnikov (1993), long-distance atmospheric
103 convection bringing air masses from the south and north, dominates at the lake area. These air
104 masses bring tree and shrub pollen grains playing an important role in the recent pollen
105 assemblages from long distances. This situation may also have occurred the past.

106 The study area belongs to the subzone of southern shrub and typical tundra (Galanin et al,
 107 1997). The modern treeline for larch (*Larix cajanderi*) and stone pine (*Pinus pumila*) is
 108 positioned roughly 100 km to the south and west of the lake (Galanin et al, 1997 and
 109 references therein). Although the northern boundary of shrub alder is reportedly much to the
 110 north of the lake, the only shrub alder stands grow approximately 10 km from the lake, in the
 111 Enmyvaam River valley (P. Minyuk, personal com). The local vegetation has been well
 112 studied during the last decades (e.g. Belikovich, 1988, 1989, 1994; Kozhevnikov, 1993;
 113 Belikovich and Galanin, 1994 and references therein).

114 According to Belikovich (1994), ca. 40% of the area (low parts of smooth crater slopes and
 115 low lake terraces) are covered by hummock tundra with *Eriophorum vaginatum*, *E. callitrix*,
 116 *E. polystachion*, *Pedicularis pennellii*, *P. albolabiata*, *Carex rotundata*, *C. lugens*, *Salix*
 117 *fuscescens*, *S. reticulata*, *Senecio atropurpureus*, *Ledum decumbens*, *Andromeda polifolia*,
 118 and *Vaccinium uliginosum*; ca. 20% (low-middle parts of crater slopes) by moss-lichens
 119 tundra with *Cassiope tetragona*, *Rhododendron parvifolium*, *Senecio resedifolus*, *Ermania*
 120 *parryoides*, *Silene stenophylla*, *Dryas octopetala*, *Crepis nana*, *Potentilla elegans*, and
 121 *Androsace ochotensis*; ca. 15% (upper mountain plains) - by tundra with rare beds with *Salix*
 122 *phlebophylla*, *Pedicularis lanata*, *Artemisia furcata*, *Potentilla elegans*, *Eritrichium*
 123 *aretioides*, *Minuartia arctica*, *Potentilla uniflora*, *Arenaria capillaris*, *Poa pseudoabbreviata*,
 124 *Cardamine bellidifolia*, *Saxifraga serpyllifolia*, *Kobresia myosuroides*, and *Crepis nana*; ca.
 125 10% - by nival vegetation with *Salix polaris*, *Cassiope tetragona*, *Carex tripartita*, *Phippsia*
 126 *algida*, *Koenigia islandica*, *Saxifraga hyperborea*, *Eritrichium villosum*, *Primula*
 127 *tschuktschorum*, *Hierochloe pauciflora*; ca. 10% - by meadow and shrubby tundra with
 128 *Artemisia arctica*, *Aconitum delphiniphodium*, *Arctagrostis arundinacea*, *Carex podocarpa*,
 129 *Festuca altaica*, *Luzula multiflora*, *Senecio tundricola*, *Thalictrum alpinum*, *Veratrum*
 130 *oxysepalum*. Rare steppe-like communities with *Potentilla stipularis*, *Artemisia kruhseana*,
 131 *Myosotis asiatica*, *Saxifraga eschscholtzii*, *Papaver lapponicum*, *Senecio jacuticus*, *Woodsia*

132 *ilvensis*, *Dianthus repens* can be found in rocky habitats. Along the Enmyvaam River and
133 alongside large creeks, grow low willow stands with *Salix tschuktschorum*, *S. saxatilis*,
134 *Androsace ochotensis*, *Empetrum subholarcticum*, *Pleuropogon sabinii*, *Polemonium boreale*,
135 *Beckwithia chamissonis*, *Saussurea tilesii*, *Lagotis minor*, *Pedicularis hirsuta* and meadow-
136 shrub willow communities with *Salix alaxensis*, *S. krylovii*, *Deschampsia borealis*,
137 *Chamerion latifolium*, *Equisetum variegatum*, *Stellaria fischerana*, *Potentilla hyparctica*,
138 *Eutrema edwardsii*, *Cardamine blaisdellii*, *Trollius membranostylus*, *Polemonium*
139 *acutiflorum*, *Parnassia kotzebuei*, *Poa paucispicula*.

140

141 **3 Methods**

142 A standard HF technique was used for pollen preparation (Berglund and Ralska-
143 Jasiewiczowa, 1986). A tablet of *Lycopodium* marker spores was added to each sample for
144 calculating total pollen and spore concentrations following Stockmarr (1971). Water-free
145 glycerol was used for sample storage and preparation of the microscopic slides. Pollen and
146 spores were identified at magnifications of 400×, with the aid of published pollen keys and
147 atlases (Kupriyanova et al., 1972, 1978; Bobrov et al., 1983; Reille, 1992, 1995, 1998; Beug,
148 2004). In addition to pollen and spores a number of non-pollen-palynomorphs such as fungi
149 spores, remains of algae and invertebrate, were also identified when possible and counted.
150 These non-pollen-palynomorphs are also valuable indicators of past environments (e.g. van
151 Geel, 2001 and references therein).

152 At least 250 pollen grains were counted in each sample. The relative frequencies of pollen
153 taxa were calculated from the sum of the terrestrial pollen taxa. Spore percentages are based
154 on the sum of pollen and spores. The relative abundances of reworked taxa (mineralized
155 pollen and spores of Tertiary and early Quaternary age) are based on the sum of pollen and
156 redeposited taxa, the percentages of non-pollen palynomorphs are based on the sum of the
157 pollen and non-pollen palynomorphs, and the percentages of algae are based on the sum of

158 pollen and algae. TGVView software (Grimm, 2004) was used for the calculation of
159 percentages and for drawing the diagrams (Figs 3-5). The diagrams were zoned by a
160 qualitative inspection of significant changes in pollen associations, pollen concentrations and
161 occurrence of particularly indicative taxa. CorelDraw software was used for preparation of the
162 final pollen diagrams.

163 At a depth of 146.5-151 cm in core P2, we detect a number of well-preserved plant
164 remains, picked using a stereomicroscope and identified by comparison with a modern
165 reference material from the Herbarium Senckenbergianum (IQW). Additionally, a *Carex*
166 identification key (Egorova, 1999) has been used.

167 The core sediments were also studied for testate amoebae tests. The samples were sieved
168 through a 0.5 mm mesh and testate amoebae tests were concentrated with a centrifuge. A drop
169 of suspension was placed on the slide, and then glycerol was added. Normally, 5 slides were
170 examined at x200-400 magnification with a light microscope.

171 A total of 33 AMS ^{14}C ages were obtained from the studied deposits (Tables 1). Plant
172 macrofossils (i.e. grass remains) were picked from the cores P1 and P2 and the uppermost
173 segment of 5011-3 for AMS radiocarbon dating. Because of the lack of plant remains in the
174 lower part of core 5011-3 only bulk organic was dated. AMS datings were done at the Leibniz
175 Laboratory for Radiometric Dating and Stable Isotope Research (Christian Albrechts
176 University, Kiel, Germany) and the Poznan Radiocarbon Laboratory (Adam Mickiewicz
177 University, Poznan, Poland). Calibrated ages (cal. yr BP) were calculated using "CALIB 5.0"
178 (Reimer et al., 2004).

179

180 **4 Results**

181 **4.1 P1 core**

182 The first permafrost core (P1) has extracted from a piedmont terrace about 1.7 km southeast
183 of the lake (67°22'26''N, 172°13'10''E, Fig. 1) during field work in summer 2003 (for details

184 see Schwamborn et al., 2006). The study site is located on a slope exposed to the southwest
185 with the angle of 5°. The vegetation cover at core site was relatively dense (ca. 80%).

186 The 5 m slope debris core consists mostly of a silty-to-sandy diamicton interpreted as a
187 result from proluvial, colluvial and solifluctional deposition (Schwamborn et al., 2006).
188 Prominent peaty layers interrupt the section between 330 and 220 cm core depth, which is
189 also reflected in maximum values of total organic carbon (TOC on Fig. 2). Non-identified
190 plant remains from several layers have been dated and show a correct depth-to-age
191 relationship (Table 1, Fig. 2). The oldest date from 463 cm depth shows that the oldest core
192 sediments are around 13,000 cal. yr BP old or slightly older.

193 Generally the P1 sequence is very rich in pollen and palynomorphs (Fig. 3). The studied
194 pollen spectra can be subdivided into 5 pollen zones (PZ). PZ-I (ca. 495-430 cm) is dominated
195 by Cyperaceae, Poaceae, and *Betula* sect. *Nanae* and *Salix* pollen. PZ-II (ca. 430-380 cm)
196 shows the significant increase of Cyperaceae pollen content, while *Betula* sect. *Nanae* content
197 is decreased. PZ-III (ca. 380-330 cm) is notable for an increase in *Betula* sect. *Nanae* and
198 appearance of small amounts of *Alnus fruticosa*. Pollen concentration is also increased in the
199 upper part of the zone. The amounts of tree and shrub pollen (predominantly *Alnus fruticosa*)
200 have a maximum in PZ-IV (ca. 330-265 cm). The pollen concentration is the highest in PZ-V
201 (ca. 265-50 cm), which is notable for high amounts of *Betula* sect. *Nanae*, *Alnus fruticosa* and
202 Cyperaceae pollen. Single pollen grains of *Pinus*, *Larix*, and *Picea* are also characteristic for
203 this zone. PZ-V can be subdivided into 2 subzones, the upper one (50-0 cm) showing the
204 higher contents of *Salix* pollen.

205 P1 has also been studied for rhizopods (Table 2). The only sphagnobiotic/hygrophilic
206 *Heleopera petricola* v. *amethystea*, pointing to a very wet environment, has been found at
207 463-473 cm depth. Mostly soil-eurybiotic (e.g. *Centropyxis aerophila*, *C. constricta*, *C.*
208 *sylvatica*) and hydrophilic (*Diffflugia* and *Lagenodifflugia*.) species dominated the sediments
209 between 334 and 223 cm. However, sphagnobiotic taxa (*Arcella*, *Heleopera*, *Nebela*,

210 *Centropyxis aculeata*) are also common. The role of soil-eurybiotic species gradually
211 increases in the upper part.

212

213 **4.2 P2 core**

214 The core was retrieved 12.5 km away from P1 across the lake, to the north (67°32'50''N,
215 172°07'31''E), Fig. 1). The site is placed on a gently inclined (<3°) surface about 100 m from
216 the north lake shoreline (for details see Schwamborn et al., 2008). The surface is characterized
217 by a boggy environment composed of a loamy substrate covered by grass tundra. Similarly
218 like core P1 deposits, core P2 is composed of a silty-to-sandy diamicton deposition
219 (Schwamborn et al., 2008b). The lower part of the core (510-250 cm) is interpreted as
220 weathering debris of the local volcanic basement. The upper 250 cm consisted of proluvial
221 slope wash out deposits. The lithological transition between the units is also very
222 distinguishable by an increase of TOC contents (Fig. 2).

223 Non-identified plant remains found in the P2 deposits have also been dated and show a
224 rather reliable depth-age relationship (Table 1, Fig. 2). Three radiocarbon dates from the
225 sediments between 205 and 226 cm depth demonstrates that these sediments might have
226 accumulated about 14,000-12,400 cal. yr BP. Taking in consideration the comparison with
227 other dated pollen records from the area (Lozhkin et al., 2007; Matrosova 2009; Glushkova et
228 al., 2009) the youngest date seems to be the most reliable.

229 P2 core sediments are rich in pollen and palynomorphs except of the lowermost 170 cm.
230 The studied pollen spectra can be subdivided into 6 PZ (Fig. 4). Sediments from PZ-I (ca.
231 510-350 cm) contain only single pollen grains of Pinaceae, *Betula* sect. *Nanae*, *Alnus*
232 *fruticosa*, and Cyperaceae. Pollen concentration is slightly higher in the lowermost sample
233 which contains few pollen of *Betula* sect. *Nanae*, *Alnus fruticosa*, *Pinus* s/g *Haploxylon* and
234 Cyperaceae. Pollen concentration is higher (up to 2650 grains/g) in PZ-II (ca. 350-330 cm),
235 which is also notable for high content of *Lycopodium* and *Botrychium* spores. The lowermost

236 PZ-I and PZ-II were not used for paleoenvironmental reconstructions because of very low
237 pollen concentration in many samples which may lead to over-representing some taxa due to
238 possible contamination or selective preservation of palynomorphs (e.g. abnormal presence of
239 spores may indirectly point to it). Pollen concentration is much higher (up to 5800 grains/g) in
240 PZ-III (ca. 330-265 cm), which is characterized by high pollen contents of *Betula* sect. *Nanae*,
241 *Alnus fruticosa*, Cyperaceae, Poaceae. Rather high amounts of *Pinus* s/g *Haploxyton* and
242 Pinaceae are also notable in this zone. The pollen concentration increases significantly (up to
243 35.700 grains/g) in PZ-IV (ca. 265-180 cm). *Betula* sect. *Nanae* and *Alnus fruticosa* pollen
244 contents decreased dramatically at the beginning of the zone and gradually increased in the
245 upper part. The zone can be subdivided in two subzones based on the shrub pollen contents.
246 Pollen concentration is highest (up to 83.600 grains/g) in PZ-V (ca. 180-40 cm), which is
247 dominated by pollen of *Betula* sect. *Nanae*, *Alnus fruticosa*, *Salix*, Cyperaceae, and Poaceae.
248 Additionally, on the 146.5-151 cm depth seeds and short spurs of *Larix dahurica* as well as
249 numerous utricle and nutlets of *Carex rostrata* were found. The uppermost PZ-VI (ca. 40-0
250 cm) is characterized by decreasing *Betula* sect. *Nanae* and *Alnus fruticosa* pollen contents,
251 while Cyperaceae, *Pinus* s/g *Haploxyton* and *Salix* pollen contents increased.

252 The P2 core has been also studied for rhizopods, but no tests were found there.

253

254 **4.3 5011-3 core**

255 The core was drilled on the western margin of the crater (67°29'04''N, 171°56'40''E)
256 approximately 300 m west from the lake shore (Fig. 1). This 141.5 m long core was recovered
257 during a drilling campaign in winter 2008 in the frame of the international ICDP funded
258 project "El'gygytgyn Drilling Project" (Melles et al., 2011). The main objective of the coring
259 was to extend the permafrost record back in time in order to better understand the interaction
260 between catchment processes and lake sedimentation. The sediment core drilled in an alluvial
261 fan consists of sediment layers of sandy gravel to gravelly sand, which is interpreted to

262 represent alternating subaerial and subaquatic parts of the fan. Occasionally intercalations of
263 sandy beds occur, e.g. at 7, 9, 14.5, 18-19.5, 24, and 26 m. The modern setting of the coring
264 site is placed in an alluvial-proluvial sediment fan, and from aerial imagery it is obvious that
265 the fan has a subaquatic prolongation into the lake. In total, 12 samples from the core were
266 AMS ^{14}C dated (Table 1). Although the non-identifiable plant remains (possibly grass roots)
267 were picked throughout the upper meter of the core and expected to provide reliable age
268 control for studied sediments, the ages appeared to be modern reflecting the presence of
269 modern plant roots in the active layer. The bulk AMS ^{14}C dates from some selected horizons
270 (Table 1) did not provide reliable ages either. These ages are not in a chronological order
271 reflecting the reworked character of TOC in the samples. The ages are obviously too old,
272 taking into consideration the comparison with other dated pollen records from the area (e.g.
273 Matrosova 2009; Glushkova et al., 2009; P1 and P2 records). Therefore, age estimations for
274 the 5011-3 core are based on a comparison with the dated pollen sequences from the area.

275 Generally, the upper 9 m of 5011-3 sediments are rich in pollen and palynomorphs, but
276 only single pollen grains were found below this depth, except in sediments between 19.8 and
277 19.3 m (Fig. 5). The studied pollen spectra can be subdivided into 7 PZ. PZ-I (ca. 1980-1930
278 cm) is dominated by *Betula* sect. *Nanae*, *Alnus fruticosa*, *Salix*, Cyperaceae, Poaceae, and
279 Ericales pollen. The presence of *Larix* pollen and high contents of *Sphagnum* and *Lycopodium*
280 spores is also characteristic for the zone, where pollen concentration is rather low (up to 3500
281 grains/g). No pollen has been found between ca. 1930 and 1400 cm and only few pollen
282 grains of *Betula* sect. *Nanae*, *Alnus fruticosa*, *Salix*, Cyperaceae, Poaceae and single spores of
283 *Sphagnum* and *Lycopodium* have been found in PZ-II (ca. 1400-900 cm). PZ-III (ca. 900-330
284 cm) is notable for much higher pollen concentration (up to 101,330 grains/gr). The spectra are
285 dominated by *Betula* sect. *Nanae*, *Alnus fruticosa*, *Salix*, Cyperaceae, Poaceae, Ericales and
286 spores of *Sphagnum*. Contents of *Sphagnum* as well as pollen concentration reduced
287 significantly in the upper PZ-IV (ca. 330-250 cm). PZ-V (ca. 250-180 cm) is notable for the

288 significant increase of Poaceae pollen content, while contents of *Betula* sect. *Nanae*, *Alnus*
289 *fruticosa*, *Salix*, Ericales and *Sphagnum* are dramatically decreased. The pollen concentration
290 is the highest in the zone (up to 829,400 grains/g). The contents of *Betula* and *Alnus* pollen
291 increased again in PZ-VI (ca. 180-100 cm), which is also notable for high content of
292 *Artemisia*. The pollen concentration significantly (up to 15,000 grains/g) reduced in this zone.
293 The uppermost PZ-VII (100-0 cm) is dominated by of *Betula* sect. *Nanae*, *Alnus fruticosa*,
294 Cyperaceae, Poaceae, and Ericales, where pollen concentration is very high (up to 770,000
295 grains/g). Single pollen of long-distance transported *Pinus* and *Picea* are also characteristic
296 for this zone.

297 The 5011-3 core has been also studied for rhizopods, but no tests were found.

298

299 **5 Discussion: paleoenvironmental reconstructions**

300 **5.1 MIS 7(?) environment**

301 The oldest pollen spectra are presented in the lower part (1980 to 1930 cm) of the studied
302 section of the core 5011-3 (PZ-I, Fig. 5). The pollen assemblages are dominated by *Alnus*
303 *fruticosa*, *Betula* sect. *Nanae* and Poaceae. However, pollen of *Larix*, *Salix*, Cyperaceae,
304 Ericales, Caryophyllaceae and spores of *Sphagnum*, *Lycopodium* and *Huperzia* are also
305 important components of the revealed spectra. They are not dated but the comparison with
306 lacustrine pollen records shows that spectra of our PZ-I are similar to those from the zone E14
307 of the TL-dated lacustrine core LZ1024 (Matrosova, 2009) and to those from the zone EG2 of
308 the core PG1357 (Lozhkin et al, 2007). Based on the comparison of our record with the
309 lacustrine records, we may suggest a MIS 7 age for our PZ-I zone. However, an older age for
310 the revealed interglacial interval cannot be completely excluded.

311 According to the pollen spectra, shrub alder, dwarf birch, and willows grew in the lake
312 catchment. Relatively high content of larch pollen in the spectra (up to 4.5%) requires the

313 movement of northern boundary of larch forest at least 100 km to the north. Our conclusion is
314 also supported by the lacustrine pollen records (Lozhkin et al., 2007, Matrosova, 2009).
315 However, the cores drilled in the center of the El'gygytgyn Lake do not contain larch pollen at
316 all and show low presence of *Salix*, Ericales, Caryophyllaceae pollen and *Sphagnum*,
317 *Lycopodium* and *Huperzia* spores. This difference most likely reflects the larger presence of
318 the local components in the 5011-3 core pointing to the importance of studying of the
319 terrestrial (non-lacustrine) sediments in addition to the lacustrine ones. Taking in consideration
320 all El'gygytgyn pollen records, we assume that open larch forest with shrub alder, dwarf birch
321 and willows dominated the local vegetation during the revealed warm interval. However,
322 grass-sedge dominated communities with other herbs and *Sphagnum* and *Lycopodium* growing
323 in mesic habitats were also common in lake vicinity.

324

325 **5.2 Lateglacial**

326 Lateglacial sediments are revealed in both radiocarbon dated slope cores (P1 and P2) and in
327 the long permafrost 5011-3 core. Unfortunately, we do not have a good age control for the
328 lowermost part of the core P1. Taking into consideration the P1 bottom age of $11,160 \pm 70$ ^{14}C
329 yr BP (12,283-13,424 cal. yr BP), the most reliable P2 age of $10,450 \pm 60$ ^{14}C yr BP (12,124-
330 12,654 cal. yr BP), and pollen-based correlation with lacustrine pollen records (zone E4 of
331 LZ1024 in Matrosova 2009) we may assume that our PZ-I of P1 (Fig. 3), PZ-III of P2 (Fig. 4)
332 and PZ-III and PZ-IV of 5011-3 (Fig. 5) accumulated during the Allerød, before 13 cal. kyr.

333 Sediments attributed to the Allerød are dominated by pollen of *Betula* sect. *Nanae*, *Alnus*
334 *fruticosa*, *Salix*, Cyperaceae, Poaceae, Ericales and spores of *Sphagnum*. The relatively high
335 pollen concentration is also characteristic for the sediments. However, a number of samples
336 show very low pollen concentrations or do not contain pollen at all. Most likely, this reflects a
337 very high accumulation rate during the sedimentation. This conclusion is in a good agreement
338 with thicknesses of Allerød-attributed deposits of about 2.5 m in the P2 core, and at least 6.5

339 m in the 5011-3 core. Warmer and wetter climate conditions in the Allerød may have
340 intensified erosion and, therefore, produced higher influx of terrestrial material. The absence
341 or very low thickness of the underlying Late Pleistocene sediments might also be connected
342 with these erosion processes.

343 The main pollen taxa in the spectra slightly differ from site to site. For example 5011-3
344 sediments contain large amounts of *Salix* and Ericales pollen and *Sphagnum* spores; P1 and P2
345 sediments contain numerous pollen of Cyperaceae; lacustrine pollen records contain larger
346 amounts of long-distance pollen (including *Betula* and *Alnus*). However, the PG1351
347 lacustrine record also contains large amounts of *Sphagnum* spores in the late glacial sediments
348 confirming wet habitats in the lake vicinity (Lozhkin et al., 2007). The sphagnobiotic
349 rhizopod, *Heleopera petricola*, found in the Allerød-dated P1 sediments is in good agreement
350 with numerous *Sphagnum* spores in the pollen records. Such habitats were probably common
351 along the creeks as today.

352 Our interpretation of the studied sediments is very similar to those from the PG1351
353 lacustrine record (Lozhkin et al., 2007; Matrosova 2009) and from LZ1024 (Glushkova et
354 al., 2009; Matrosova 2009). Glushkova et al., (2009) also have reported pollen spectra with
355 dominance of shrub pollen taxa from the undated terrace sediments (sections GS-10 and GS-
356 12/1 in Fig 1) attributed to a Late Glacial warm interval. Similar paleoenvironmental records
357 are also known from adjacent regions (e.g. Brubaker et al., 2005; Anderson and Lozhkin 2006;
358 Shilo et al., 2006, 2007; Kokorowski et al., 2008; Andreev et al., 2009 and references
359 therein). Lozhkin et al., (2007) based on their PG1351 lacustrine pollen record, have
360 suggested that birch was regionally present at about 12,800 yr ¹⁴C BP (15,300 cal. yr BP),
361 while alder established in the area around 10,700 yr ¹⁴C BP (12,700 cal. yr BP).

362 There are plant macrofossil data from the sediments of section GS-37 (Fig 1) ¹⁴C dated to
363 12,215±40 yr BP (14,027-14,491 cal. yr BP). The studied sediments do not contain any shrub
364 remains. Glushkova et al. (2009) interpreted this as the absolute absence of any shrub stands in

365 the lake vicinity and very severe climate conditions. Thus, it seems that Allerød pollen and
366 plant macrofossil data are contradictory. However, the conclusion about herb dominated
367 tundra vegetation around 14,250 cal. yr BP is based on the single studied sample, which reflect
368 very wet, but not a typical tundra habitat. Moreover, they interpret the sediments containing
369 numerous pebbles and eggs of *Daphnia* as the lake terrace periodically overflowed by the lake
370 (Glushkova et al., 2009). It is obvious that shrubs cannot survive in such flooded habitats.
371 Therefore, the found plant macrofossils reflect a very local, flooded habitat, which cannot be
372 extrapolated to the whole lake vicinity.

373 Thus, according to the pollen spectra shrub alder, dwarf birch, and willows grew in the
374 lake surrounding during the Allerød interstadial with relatively warm and wet climate (Melles
375 et al., 2012). We can reconstruct shrub tundra vegetation with dwarf birch, shrub alder and
376 willow around the lake.

377 Pollen spectra from PZ-II of P1 (Fig. 3), PZ-IVa of the core P2 (Fig. 4), and PZ-V of 5011-3
378 (Fig. 5) are dominated mostly by Cyperaceae and Poaceae pollen and reflect disappearance of
379 shrubs from the area pointing to climate deterioration which can be attributed to the Younger
380 Dryas. The most reliable ^{14}C dates from core P2 and P1 (Table 1) confirm this conclusion.
381 Pollen spectra with a significant increase in herbs (mostly Poaceae) and *Selaginella rupestris*
382 have also been revealed in the lacustrine sediments (E3 of LZ1024 in Matrosova 2009), and
383 are interpreted as reflecting the Younger Dryas event (Glushkova et al., 2009; Matrosova;
384 2009; Melles et al., 2012). Thus, grass-herb tundra dominated the area during the Young
385 Dryas cooling. Younger Dryas dated pollen records from the adjacent regions (e.g. Anderson
386 et al., 2002; Kokorowski et al., 2008; Andreev et al., 2009, 2011 and references therein) reflect
387 the similar environmental changes.

388

389 **5.3 Holocene**

390 Pollen spectra of the PZ-IVb of P2 (Fig. 4) accumulated before 9640 ± 60 ^{14}C yr BP (11,200-
391 10,780 cal. yr BP) show a gradual increase of *Alnus fruticosa* and *Betula* sect. *Nanae* pollen
392 contents reflecting early Holocene climate amelioration. The early Holocene pollen
393 assemblages are also well represented in the undated PZ-III of P1 (Fig. 3), where they are
394 dominated mostly by pollen of *Betula* sect. *Nanae*, Cyperaceae and Poaceae with few *Alnus*
395 *fruticosa* and *Salix*. Four ^{14}C dates (Table 1) confirm that these sediments were accumulated
396 before 9000 ^{14}C yr BP (10,200 cal. yr BP). Similar pollen assemblages have been revealed in
397 the lowermost pollen zone of the so-called Olga Creek section (OC on Fig. 1, Shilo et al.,
398 2008; Glushkova et al., 2009), situated ca 100 m from P1 coring site. These lowermost spectra
399 are also not ^{14}C dated, however two ^{14}C dates: 9250 ± 90 and 9125 ± 30 yr BP, from overlain
400 sediments confirm that these sediments were accumulated before 9300 ^{14}C yr BP (10,550 cal.
401 yr BP). Similar undated early Holocene pollen assemblages are also reported by Glushkova et
402 al., (2009) from the section GS-12/1 (Fig. 1). Thus, we may assume that the earliest shrub
403 tundra, with dwarf birches and willows and probably a few shrub alder, dominated the lake
404 vicinity at the onset of the Holocene. The early Holocene pollen records from adjacent regions
405 (e.g. Anderson et al., 2002; Lozhkin and Anderson, 2002; Kokorowski et al., 2008; Andreev
406 et al., 2009 and references therein) have revealed similar environmental changes.

407 The contents of *Alnus fruticosa* are significantly higher in the PZ-V of the core P2 (up to
408 30%) ^{14}C dated to ca 9600 yr BP (11,200-10,780 cal. yr BP) and PZ-IV of the core P1 (up to
409 50%) ^{14}C dated around 8900-8800 yr BP (9940-9700 cal. yr BP). Most likely, this increase
410 reflects the further distribution of shrub alder stands in the area during the early Holocene.
411 Pollen spectra of the PZ-V of P2 (Fig. 4) radiocarbon dated to about 7200-7300 cal. yr BP,
412 PZ-VI of 5011-3 (Fig. 5) and bottom spectra from the terrace section GS-8403 (Glushkova et
413 al., 2009) and the section OC in the Enmyvaam River valley (Glushkova and Smirnov 2007;
414 Shilo et al., 2008; Glushkova et al., 2009) also demonstrate high amounts of *Alnus fruticosa*
415 pollen in the early Holocene sediments. Moreover, the lacustrine sediments (Matrosova

416 2009; Melles et al., 2012) accumulated above sediments attributed to the Younger Dryas, are
417 also contain very high amounts of *Alnus* (up to 60%). Large shrub alder trunks and smaller
418 twig fragments ^{14}C dated to 9250 ± 90 and 9125 ± 30 yr BP respectively, as well as numerous
419 undated alder nuts from the same layers well confirm that shrub alder grew in the lake
420 vicinity at least 10,550 cal. yr BP (Shilo et al., 2008). Thus, it is likely that shrub alder stands
421 were well established in the El'gygytgyn Lake Crater at about 11,200 cal. yr BP or even
422 slightly earlier.

423 The well-preserved larch seeds (Fig 6) found in peaty layer in the core P2 prove the local
424 presence of trees directly at the lake crater as early as 11,200-10,780 cal. yr BP. Larch
425 remains were also found in the sediments accumulated shortly before 9300 ^{14}C yr BP
426 (10,550 cal. yr BP) from the OC section (Fig. 1, Shilo et al., 2008; Glushkova et al., 2009),
427 thus, also confirming local presence of the larch trees at the area during the early Holocene.
428 Such forest (tundra-forest) environments are also good habitats for the shrub alder stands.
429 The local presence of *Larix* indicates a treeline shift of about 100 km northward (CAVM-
430 Team, 2003) as result of the early Holocene climate amelioration. Larch requires a mean
431 temperature of the warmest month of at least 10 °C, thus such climate conditions must have
432 existed at the lake crater during the early Holocene.

433 The studied early Holocene pollen assemblages slightly differ from site to site. For
434 example, the early Holocene 5011-3 spectra (PZ-VI) show high contents of *Artemisia* (up to
435 25%), while GS-8403 spectra reported by Glushkova et al., (2009) contain up to 23% of
436 Ericales. The difference may reflect the mosaic character of the local vegetation cover and/or
437 different age of the revealed pollen assemblages. The lacustrine record (Matrosova 2009;
438 Melles et al., 2012) accumulated above the sediments attributed to the Younger Dryas shows
439 very high amounts of *Alnus* (up to 60%), which might have been transported from a distance
440 and, thus, reflect the regionally dominated vegetation.

441 Rhizopod tests of soil-eurybiotic *Centropyxis* and hydrophilic *Diffflugia* taxa (Table 2) are
442 numerous in the P1 early Holocene sediments, however, sphagnobiotic *Arcella*, *Heleopera*,
443 and *Nebela* are also common. The high contents of hydrophilic and sphagnophilic taxa point
444 to wet oligotrophic and mesotrophic soil environment at the core site. Later, after ca 6300 cal.
445 yr BP, the role of soil-eurybiotic species increased reflecting drier soil environment.

446 Thus, pollen and macrofossil data show that forest and/or forest-tundra communities
447 with larches, shrub alder, dwarf birches, and willows were well distributed the low
448 elevations around the lake during the early Holocene at least between 11,200 and 9100 cal.
449 yr BP. It is most likely that larch and shrub alder grew in the close vicinity to the lake only
450 before ca. 8200 cal. yr BP. Similar changes in the high Arctic vegetation cover are also
451 characteristic for coastal areas of the Laptev and East Siberian Seas (e.g. Andreev et al.,
452 2009, 2011 and references therein). Recovered larch remains document that larch grew
453 approximately 100 km from its modern northern distribution limit. The mean July
454 temperatures were at least 10-12 °C (Lozhkin & Anderson, 1995), ca 4-5 °C higher than
455 modern July temperatures (Shilo et al., 2008). This is in agreement with the early Holocene
456 pollen-based paleoclimate reconstruction from the El'gygytgyn lacustrine record (Lozhkin et
457 al., 2007; Melles et al., 2012) and other high arctic sites (e.g. Andreev et al., 2009, 2011 and
458 references therein).

459 A number of ¹⁴C dates (Table 1) from P1 (PZ-Va) and P2 (PZ-V) cores confirm that
460 permafrost sediments containing relatively high amounts of *Alnus fruticosa* pollen were
461 accumulated until ca. 3500 cal. yr BP. Therefore we may assume that shrub alder might
462 grow around the lake in more protected habitats or very close to the lake vicinity before this
463 time. This conclusion is in good agreement with pollen and plant macrofossil data from
464 adjacent regions, documenting the presence of shrubs and trees to the north from modern
465 distribution areas (e.g. MacDonald et al., 2000; Andreev et al., 2009, 2011; Binney et al.,
466 2009 and references therein). However, the dated woody remains from the Enmyvaam River

467 valley (Glushkova and Smirnov 2007; Lozhkin et al., 2011) confirm the presence of high
468 shrubs in the area only until ca. 7400 ¹⁴C yr BP (8200 cal. yr BP). The studied deposits also
469 contain the rather high amounts (up to 35%) of *Alnus fruticosa* pollen in the sediments
470 accumulated after 7400 ¹⁴C yr BP (Shilo et al., 2008; Lozhkin et al., 2011), pointing to a
471 possible local presence of shrub alder, however, the age of the pollen assemblages is
472 unknown.

473 Generally, late Holocene pollen spectra from the uppermost sediments (upper spectra of
474 PZ-Vb of the core P1, Fig. 3; PZ-VI of the core P2, Fig. 4; LZ-1024 record in Matrosova
475 2009 and Melles et al., 2012) show a decrease in contents of *Alnus fruticosa* (mean values
476 are up 20% and less) and some increases of contents of *Salix*, *Pinus*, *Betula*, Ericales, and
477 Cyperaceae. These changes can be interpreted as disappearance of shrub alder from the lake
478 vicinity. The main components of pollen assemblages slightly change from site to site
479 reflecting local vegetation cover at coring sites.

480 The late Holocene pollen assemblages are characterized by higher amounts of *Pinus* s/g
481 *Haploxylon*. The modern boundary of the stone pine (*Pinus pumila*) is about 80 km from the
482 study area (Vas'kovskiy 1958), thus, it is most likely that all *Pinus* pollen grains are of long
483 distance origin. Its pollen presence is especially remarkable in the uppermost lake sediments
484 (Lozhkin et al., 2007; Matrosova 2009) and the modern spectra (Matrosova 2006) reflecting
485 the extra-regional vegetation pollen influx. Taking into consideration all pollen records from
486 the study area we may assume that stone pine did not grow around the lake during the
487 Holocene.

488 Late Holocene sediments dated between ca. 900 and 450 cal. yr BP (Glushkova et al.,
489 2009) contain pollen spectra similar to those revealed in this study. They also show lower
490 contents of *Alnus* pollen in many spectra and high fluctuations in *Betula*, Ericales, *Thalictrum*,
491 and *Selaginella rupestris* reflecting local environments. Thus, pollen data show that herb
492 tundra communities started to dominate in the lake catchment after ca. 3000 cal. yr BP.

493

494 **6 Conclusions**

495 New permafrost records document vegetation and climate changes in the El'gygytgyn Lake
496 Crater during the Late Quaternary. The studied records reflect the local vegetation changes
497 that result in a better understanding of the possible role of local and regional components in
498 the fossil pollen spectra, and in more reliable environmental reconstructions. It is evident that
499 terrestrial records better reflect the local environments than the lacustrine ones where long-
500 distance transported pollen overshadows the local components.

501 The oldest pollen spectra of the studied sections of the core 5011-3 are possibly of the
502 MIS 7 age. They document that open larch forest with shrub alder, dwarf birch and willows
503 dominate vegetation suggesting the northern movement of larch forests. Treeless grass-sedge
504 dominated communities with other herbs and *Sphagnum* and *Lycopodium* growing in mesic
505 habitats were also common in lake vicinity.

506 Lateglacial pollen records show that shrub tundra with dwarf birch, shrub alder and willow
507 dominated in the lake surroundings during the relatively warm Allerød interstadial. Rather low
508 pollen concentrations in many samples of Allerød age reflect very high accumulation rate
509 during the sedimentation.

510 Younger Dryas pollen records reflect dramatic changes in the vegetation cover. Grass-
511 sedge-herb tundra dominated the area pointing to significant climate deterioration.

512 Forest-tundra with larches, dwarf birches and willows dominate the lake vicinity at the
513 onset of the Holocene between ca. 11,200 and 9100 cal yr BP. Shrub alder stands might grow
514 at the low elevations around the lake during the Holocene, between ca. 11.200 and 3500 cal yr
515 BP. Later, similar-to-modern herb tundra communities dominated the Elgygytgyn Impact
516 Crater.

517 **Acknowledgments:** Drilling operations for the ICDP 5011-3 core was funded by the
518 International Continental Scientific Drilling Program (ICDP), the U.S. National Science

519 Foundation (NSF), the German Federal Ministry of Education and Research (BMBF), Alfred
 520 Wegener Institute (AWI) and Helmholtz Centre Potsdam (GFZ), the Russian Academy of
 521 Sciences Far East Branch (RAS FEB), the Russian Foundation for Basic Research (RFBR),
 522 and the Austrian Federal Ministry of Science and Research (BMWF). Funding of core
 523 analyses was provided by BMBF (grant no. 03G0642) . We are grateful to M. Edwards and an
 524 anonymous reviewer for their constructive comments and suggestions. Special thanks also go
 525 to Dr. Alison McAnena for reviewing the English.

526

527 **References**

- 528 Anderson, P.M. and Lozhkin, A.V.: Palynological and radiocarbon data from Quaternary deposits of
 529 northeastern Siberia, in: Quaternary vegetation and climate of Siberia and Russian Far East: a
 530 palynological and radiocarbon database, edited by Anderson, P.M. and Lozhkin, A.V., NOAA
 531 Paleoclimatology and North-East Science Center, Magadan, 27-34, (in Russian and English),
 532 2002.
- 533 Anderson, P.M., Lozhkin, A.V., and Brubaker, L.B.: Implications of a 24,000-yr palynological record
 534 for a Younger Dryas cooling and for boreal forest development in Northeastern Siberia, *Quatern.*
 535 *Res.*, 57, 325-333, 2002.
- 536 Andreev, A.A., Grosse, G., Schirrmeyer, L., Kuznetsova, T.V., Kuzmina, S.A., Bobrov, A.A.,
 537 Tarasov, P.E., Novenko, E.Yu., Meyer, H., Derevyagin, A.Yu., Kienast, F., Bryantseva, A., and
 538 Kunitsky, V.V.: Weichselian and Holocene paleoenvironmental history of the Bol'shoy
 539 Lyakhovsky Island, New Siberian Archipelago, Arctic Siberia, *Boreas*, 38, 72-110, 2009.
- 540 Andreev, A., Schirrmeyer, L., Tarasov, P., Ganopolski, A., Brovkin, V., Siebert, C., and Hubberten,
 541 H.-W.: Vegetation and climate history in the Laptev Sea region (arctic Siberia) during Late
 542 Quaternary inferred from pollen records, *Quatern. Sci. Rev.*, 30, 2182-2199, 2011.
- 543 Belikovich, A.V.: Permanent monitoring in the El'gygytyn Lake area, IBPS FEB RAS, Magadan. (in
 544 Russian), 1988.
- 545 Belikovich, A.V.: Graf method by analyses of habitats in a landscape unit using environment
 546 characteristics (El'gygytyn Lake, central Chukotka as a model), in: Graf method in ecology,
 547 edited by Galanin, A.V., FEB RAS, Vladivostok, 104-111, (in Russian), 1989.
- 548 Belikovich, A.V.: Recreation sources of planned "El'gygytyn Lake Park", *Vestnik FEB RAS*, 3,
 549 57-63, (in Russian), 1994.
- 550 Belikovich A.V. and Galanin, A.V.: "El'gygytyn Lake" Reservation (central Chukotka). *Vestnik*
 551 *FEB RAS*, 4, 22-24, (in Russian). 1994.
- 552 Belyi, F.V.: Impactogenesis and volcanism of El'gygytyn depression, *Petrology*, 6, 86-99, 1998.
- 553 Berglund, B.E. and Ralska-Jasiewiczowa, M.: Pollen analysis and pollen diagrams, in: *Handbook of*
 554 *Holocene palaeoecology and palaeohydrology*, edited by Berglund, B.E., Wiley, Chichester, 455-
 555 484, 1986.
- 556 Binney, H.A., Willis, K.J., Edwards, M.E., Bhagwat, S.A., Anderson, P., Andreev, A.A., Blaauw, M.,
 557 Damblon, F., Haesaerts, P., Kienast, F.W., Kremenetski, K.V., Krivonogov, S.K., Lozhkin, A.V.,
 558 MacDonald, G.M., Novenko, E.Y., Oksanen, P., Sapelko, T.V., Valiranta, M. and Vazhenina, L.,

- 559 The distribution of late-Quaternary woody taxa in Eurasia: evidence from a new macrofossil
560 database, *Quatern. Sci. Rev.*, 28, 2445-2464, 2009.
- 561 Brigham-Grette, J., Melles, M., Minyuk, P. and Scientific Party, Overview and significance of a
562 250-ka paleoclimate record from El'gygytyn Crater Lake, NE Russia, *J. Paleolim.*, 37, 1-16,
563 2007.
- 564 Blunier, T., Brook, E.J., Timing of Millennial-Scale Climate change in Antarctica and
565 Greenland during the last glacial period. *Science*, 291, 109-112. 2001.
- 566 Bobrov, A.E., Kupriyanova, L.A., Litvintseva, M.V., and Tarasevich, V.F.: Spores and pollen
567 of gymnosperms from the flora of the European part of the USSR. Nauka, Leningrad, (in Russian),
568 1983.
- 569 Brubaker, L.B., Anderson, P.M., Edwards, M.E., and Lozhkin, A.V.: Beringia as a glacial refugium for
570 boreal trees and shrubs: new perspectives from mapped pollen data, *J. Biogeogr.*, 32, 833-848,
571 2005.
- 572 CAVM-Team: Circumpolar Arctic Vegetation Map. Conservation of Arctic Flora and Fauna (CAFF)
573 Map No. 1. U.S. Fish and Wildlife Service, Anchorage, Alaska, 2003.
- 574 Chardez, D.: Ecologie generale des Thecamoebiens (Rhizopoda, Testacea), *Bulletin de l'Institut*
575 *Agronomique et des Stations de Recherches Gembloux*, 2, 306-341, 1965.
- 576 Egorova, T.V.: The sedges (*Carex* L.) of Russia and adjacent states, St. Petersburg State Chemical-
577 Pharmaceutical Academy, St. Petersburg, 1999.
- 578 Fradkina, A.F., Alekseev, M.N., Andreev, A.A., and Klimanov, V.A., East Siberia, Cenozoic climatic
579 and environmental changes in Russia, *Geol. Society of America Spec. Paper*, 382, 89-103. 2005a.
- 580 Fradkina, A.F., Grinenko, O.V., Laukhin, S.A., Nechaev, V.P., Andreev, A.A., and Klimanov, V.A.,
581 North-eastern Asia, Cenozoic climatic and environmental changes in Russia, *Geol. Society of*
582 *America Spec. Paper*, 382, 105-120, 2005b.
- 583 Galanin A.V., Belikovich, A.V., Galanin A.A., Tregubov, O.D.: *Priroda i resursy Chukotki (Nature*
584 *and sources of Chukotka)* IBPS FEB RAS, Magadan, (in Russian), 1997.
- 585 Glushkova, O.Y. and Smirnov, V.N.: Pliocene to Holocene geomorphic evolution and paleogeography
586 of the El'gygytyn Lake region, NE Russia, *J. Paleolim.*, 37, 37-47, 2007.
- 587 Glushkova, O.Y., Smirnov, V.N., Matrosova, T.V., Vazhenina, L.N., and Braun, T.A.: Climatic-
588 stratigraphic characteristic and radiocarbon dates of terrace complex in El'gygytyn Lake basin,
589 *Vestnik FEB RAS*, 2, 31-43, (in Russian), 2009.
- 590 Grimm, E.C., 2004. TGView. Illinois State Museum, Research and Collections Center, Springfield.
- 591 Gurov, E.P. and Gurova, E.P.: Stages of shock metamorphism of silica volcanic rocks in the
592 El'gygytyn meteorite Crater, Chukotka, *Doklady Akademii Nauk SSSR*, 249, 121-123, (in
593 Russian), 1979.
- 594 Johnsen, S.J., Dahl-Jensen, D., Dansgaard, W., and Gundestrup, N.: Greenland palaeotemperatures
595 derived from GRIP borehole temperature and ice core isotope profiles, *Tellus B* 47, 624-629, 1995.
- 596 Kokorowski, H.D., Anderson, P.M., Sletten, R.S., Lozhkin, A.V., and Brown, T.A.: Late Glacial
597 and early Holocene climatic changes based on a multiproxy lacustrine sediment records from
598 Northeast Siberia. *Arct., Antarct. Alp. Res.*, 40, 497-505, 2008.
- 599 Kozhevnikov, Yu.P.: Vascular plants around El'gygytyn Lake, in: Natural depression El'gygytyn
600 Lake (problems of study and preservation). NEISRI FEB RAS, Magadan, 62-82, (in Russian),
601 1993.
- 602 Kupriyanova, L.A., and Alyoshina, L.A.: Pollen and spores of plants from the flora of European part
603 of USSR. Vol. I., Academy of Sciences USSR, Komarov Botanical Institute, Leningrad, (in
604 Russian), 1972.

- 605 Kupriyanova, L.A. and Alyoshina, L.A.: Pollen and spores of plants from the flora of European part of
606 USSR. Academy of Sciences USSR, Komarov Botanical Institute, Leningrad (in Russian). 1978.
- 607 Layer, P.: Argon-40/argon-39 age of the Elgygytyn impact event, Chukotka, Russia, Meteor. Planet.
608 Sci., 35, 591-599, 2000.
- 609 Lozhkin, A.V. and Anderson, P.M.: The Last Interglaciation in Northeast Siberia, Quatern. Res., 43,
610 147-158, 1995.
- 611 Lozhkin, A.V. and Anderson, P.M.: A reconstruction of the climate and vegetation of Northeastern
612 Siberia based on lake sediments. Paleontol. J., 40, 622-628. 2006.
- 613 Lozhkin, A.V., Anderson, P.M., Matrosova, T.V. and Minyuk, P.S. The pollen record from
614 El'gygytyn Lake: implications for vegetation and climate histories of northern Chukotka since the
615 late middle Pleistocene. J. Paleolim., 37, 135-153, 2007.
- 616 Lozhkin, A.V., Anderson, P.M., and Vazhenina, L.N. Younger Dryas and early Holocene peats from
617 northern Far East Russia. Quatern. Int., 237, 54-64, 2011.
- 618 MacDonald, G.M., Velichko, A.A., Kremenetski, C.V., Borisova, O.K., Goleva, A.A., Andreev, A.A.,
619 Cwynar, L.C., Riding, R.T., Forman, S.L., Edwards, T.W. D., Aravena, R., Hammarlund, D.,
620 Szeicz, J.M., Gataullin, V.N. Holocene treeline history and climate change across Northern Eurasia.
621 Quatern. Res., 53, 302-311. 2000.
- 622 Matrosova, T.V.: Modern spore-pollen spectra of Anadyr Plateau (El'gygytyn Lake), in: Geology,
623 geography and biodiversity of North-East Russia, Materials of Far-East regional conference in
624 memory of A.P. Vas'kovskiy, edited by Chereshev, I.A., NEISRI FEB RAS, Magadan, 159-162,
625 (in Russian). 2006.
- 626 Matrosova, T.V.: Vegetation and climate change in northern Chukotka during the last 350 ka (basing
627 on lacustrine pollen records from El'gygytyn Lake, Vestnik FEB RAS, 2, 23-30. (in Russian).
628 2009.
- 629 Matrosova, T.V., Anderson, P.M., Lozhkin, A.V., and Minyuk, P.S.: Climate history of Chukotka
630 during the last 300,000 years from the Lake El'gygytyn pollen record, in: Climate records from
631 Quaternary sediments of Beringia, edited by Lozhkin, A.V., NEISRI FEB RAS, Magadan, 26-42,
632 (in Russian), 2004.
- 633 Melles, M., Brigham-Grette, J., Glushkova, O., Minyuk, P., Nowaczyk, N.R., and Hubberten, H.-W.:
634 Sedimentary geochemistry of a pilot core from Elgygytyn Lake - a sensitive record of climate
635 variability in the East Siberian Arctic during the past three climate cycles, J. Paleolim., 37, 89-104.
636 2007.
- 637 Melles, M., Brigham-Grette, J., Minyuk, P., Koeberl, C., Andreev, A., Cook, T., Fedorov, G.,
638 Gebhardt, C., Haltia-Hovi, E., Kukkonen, M., Nowaczyk, N., Schwamborn, G., Wennrich, V., and
639 the El'gygytyn Scientific Party: The Lake El'gygytyn Scientific Drilling Project - Conquering
640 Arctic Challenges in Continental Drilling, Scientific Drilling, 11, 29-40, 2011.
- 641 Melles, M., Brigham-Grette, J., Minyuk, P.S., Nowaczyk, N.R., Wennrich, V., DeConto, R.M.
642 Anderson, P.M., Andreev, A., Coletti, A., Cook, T., Haltia-Hovi, E., Kukkonen, Lozhkin, A.V.,
643 Rosén, P., Tarasov, P., Vogel, H., Wagner, B.: 2.8 Million Years of Arctic Climate Change from
644 Lake El'gygytyn, NE Russia, Science. 2012.
- 645 Nolan, M. and Brigham-Grette, J.: Basic hydrology, limnology, and meteorology of modern
646 El'gygytyn Lake, Siberia. J. Paleolim., 37, 17-35, 2007.
- 647 Reille, M.: Pollen et spores d'Europe et d'Afrique du nord, Laboratoire de Botanique Historique et
648 Palynologie, Marseille, 1992.
- 649 Reille, M.: Pollen et spores d'Europe et d'Afrique du nord, supplement 1. Laboratoire de Botanique
650 Historique et Palynologie, Marseille, 1995.
- 651 Reille, M.: Pollen et spores d'Europe et d'Afrique du nord, supplement 2. Laboratoire de Botanique
652 Historique et Palynologie, Marseille, 1998.

- 653 Reimer, P.J., Baillie, M.G.L., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C.,
654 Buck, C.E., Burr, G.S., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I.,
655 Heaton, T.J., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., McCormac, F.G., Manning,
656 S.W., Reimer, R.W., Richards, D.A., Southon, J.R., Talamo, S., Turney, C.S.M., van der Plicht, J.,
657 and Weyhenmeyer, C.E., *Radiocarbon* 51, 1111-1150, 2009.
- 658 Schwamborn, G., Meyer, H., Fedorov, G., Schirrmeister, L., and Hubberten, H.-W.: Ground ice and
659 slope sediments archiving Late Quaternary paleoenvironment and paleoclimate signals at the
660 margins of Elgygytgyn Impact Crater, NE Siberia, *Quatern. Res.*, 66, 259-272, 2006.
- 661 Schwamborn, G., Förster, A., Diekmann, B., Schirrmeister, L., and Fedorov, G.: Mid to late
662 Quaternary cryogenic weathering conditions in Chukotka, northeastern Russia: Inference from
663 mineralogical and microtextural properties of the El'gygytgyn Crater Lake sediment record, in:
664 Ninth International Conference on Permafrost, Institute of Northern Engineering, University of
665 Alaska Fairbanks, 1601-1606, 2008a.
- 666 Schwamborn, G., Fedorov, G., Schirrmeister, L., Meyer, H., and Hubberten, H.-W.: Periglacial
667 sediment variations controlled by late Quaternary climate and lake level change at El'gygytgyn
668 Crater, Arctic Siberia, *Boreas*, 37, 55-65, 2008b.
- 669 Shilo, N.A., Lozhkin, A.V., Anderson, P.M., Belaya, B.V., Stetsenko, T.V., Glushkova, O.Yu.,
670 Brigham-Grette, J., Melles, M., Minyuk, P.S., Nowaczyk, N., and Forman, S.: First continuous
671 pollen record of vegetation and climate change in Beringia during the last 300 thousand years,
672 *Doklady Akademii Nauk*, 376, 231-234, (in Russian), 2001.
- 673 Shilo, N.A., Lozhkin, A.V., Anderson, P.M., Brown, T.A., Matrosova, T.V., and Kotov, A.T.:
674 Radiocarbon dates and palynological characteristics of sediments in Lake Melkoe, Anadyr River
675 Basin, Chukotka, *Doklady Akademii Nauk*, 407, 235-237, 2006.
- 676 Shilo, N.A., Lozhkin, A.V., Anderson, P.M., Brown, T.A., Pakhomov, A.Yu. and Solomatkina, T.B.:
677 Glacial refugium of *Pinus pumila* (Pall.) Regel in northeastern Siberia, *Doklady Akademii Nauk*,
678 412, 401-403, 2007.
- 679 Shilo, N.A., Lozhkin, A.V., Anderson, P.M., Vazhenina, L.N., Stetsenko, T.V., Glushkova, O.Yu. and
680 Matrosova, T.V.: First data on the expansion of *Larix gmelinii* (Rupr.) Rupr. into arctic regions of
681 Beringia during the early Holocene, *Doklady Akademii Nauk*, 423, 680-682, 2008.
- 682 Stockmarr, J.: Tablets with spores used in absolute pollen analysis, *Pollen et Spores*, 13, 614-621,
683 1971.
- 684 Stuiver, M., Grootes, P.M., Braziunas, T.F.: The GISP2 $\delta^{18}\text{O}$ climate record of the past 16,500 years
685 and the role of the sun, ocean, and volcanoes, *Quatern. Res.*, 44, 341-354, 1995.
- 686 Van Geel, B.: Non-pollen palynomorphs, in: Tracking environmental change using lake sediments.
687 Vol. 3: Terrestrial, algal and siliceous indicators, edited by Smol, J. P., Birks, H. J. B., Last, W.
688 M., Bradley, R. S., and Alverson, K., Kluwer, Dordrecht, 99-119, 2001.
- 689 Vas'kovskiy, A.P. New data about distribution limits of trees and shrubs in extreme North-East of
690 USSR, *Materialy po geologii i poleznym iskopaemym severo-vostoka SSSR*, 13, 187-204,
691 Magadan Publishing House, (in Russian), 1958.
- 692 Yurtsev, B.A.: Botanical-geographical zonation and flora regionalization tundra on Chukchi Peninsula.
693 *Botanicheskiy Zhurnal*, 58, 945-964, (in Russian), 1973.
- 694
- 695
- 696
- 697
- 698

699 Table 1: Radiocarbon and calibrated ages enclose the two-sigma range of highest probability.
 700 The ages have been calibrated using CALIB Rev 6.1.0. (Reimer, P.J. et al., 2009). The
 701 obviously inversed ages were *rejected dates* and marked with *

702

Depth core (cm),	Dated material	¹⁴ C ages (yr BP)	Calibrated age intervals (cal. yr BP)	Lab. number	Reference
20, P1	plant remains	3000±30	3078-3268	KIA25979	Schwamborn et al., 2006
43, P1	plant remains	3095±45	3209-3403	KIA25980	Schwamborn et al., 2006
114, P1	plant remains	3670±30	3906-4087	KIA23976	Schwamborn et al., 2006
150, P1	plant remains	3665±35	3890-4090	KIA25981	Schwamborn et al., 2006
207, P1	plant remains	8145±45*		KIA28241	Schwamborn et al., 2006
233, P1	plant remains	5585±40	4493-6447	KIA23977	Schwamborn et al., 2006
265, P1	plant remains	8760±45	9558-9914	KIA23978	Schwamborn et al., 2006
292, P1	plant remains	8830±55	9695-10,159	KIA23979	Schwamborn et al., 2006
314, P1	plant remains	8885±40	9887-10,182	KIA24865	Schwamborn et al., 2006
325, P1	plant remains	8920±110	9660-10,249	KIA28242	Schwamborn et al., 2006
463, P1	plant remains	11,160±70	12,801-13,243	KIA23980	Schwamborn et al., 2006
46, P2	grass remains	1675±25	1526-1626	KIA24866	Schwamborn et al., 2008
52, P2	grass remains	3365±35	3553-3692	KIA27258	Schwamborn et al., 2008
95, P2	grass remains	4400±110	4812-5320	KIA27259	Schwamborn et al., 2008
119, P2	grass remains	5350±45	5998-6218	KIA27260	Schwamborn et al., 2008
132 P2	grass remains	6345±35	7171-7330	KIA24867	Schwamborn et al., 2008
146-151, P2	<i>Larix</i> seeds	9640±60	10,775-11,193	Poz-42874	this study
170-184, P2	bulk organic	1890±100*		Poz-42875	this study
205, P2	grass remains	10,450±60	12,116-12,560	KIA24868	Schwamborn et al., 2008
210, P2	grass remains	11,180±147	12,706-13,320	KIA28243	Schwamborn et al., 2008
226, P2	grass remains	11,790±242	13,113-14,220	KIA28244	Schwamborn et al., 2008
0-40, 5011-3	plant remains	modern		Poz-33404	this study
40-50, 5011-3	plant remains	modern		Poz-33406	this study

50-60, 5011-3	plant remains	modern		Poz-33407	this study
60-70, 5011-3	plant remains	modern		Poz-33408	this study
70-100, 5011-3	plant remains	modern		Poz-33409	this study
100-110, 5011-3	plant remains	modern		Poz-33410	this study
173-183, 5011-3	bulk organic	$27,690 \pm 200^*$		Poz-35975	this study
208-230, 5011-3	bulk organic	$20,860 \pm 170^*$		Poz-35977	this study
315-325, 5011-3	bulk organic	$18,800 \pm 120^*$		Poz-35978	this study
395-400, 5011-3	bulk organic	$24,070 \pm 320^*$		Poz-35979	this study
845-852, 5011-3	bulk organic	$24,590 \pm 220^*$		Poz-35980	this study
899-910, 5011-3	bulk organic	$28,440 \pm 320^*$		Poz-35981	this study

703

704

705 **Figure captures**

706 Figure 1. Location map of the study sites and mentioned cores and sections. OC – Olga Creek
707 terrace section from Enmyvaam River valley (Glushkova and Smirnov 2007; Shilo et al.,
708 2008; Glushkova et al., 2009).

709 Figure 2. Lithological, geochronological, grain size and TOC data from P1, P2, and 5011-3
710 cores.

711 Figure 3. Percentage pollen diagram of core P1. Dots are <2% pollen contents.

712 Figure 4. Percentage pollen diagram of core P2. Dots are <2% pollen contents.

713 Figure 5. Percentage pollen diagram of core 5011-3. Dots are <2% pollen contents.

714 Figure 6. Seeds of *Larix* found in core P2.

715

716

717

718

719

720

721


722

723

724


725

726 Figure 1
727




728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753

754 Figure 2




755
756
757
758
759
760

761 Figure 3




Pollen analysts: E. Morozova & A. Andreev

764 Figure 4



Analysts: E. Morozova and A. Andreev

767 Figure 5



Analysts: A. Andreev & E. Morozova

5111-3

769 Figure 6
770



771