

Research Article

Screening of α -Glucosidase Inhibitory Activity from Some Plants of Apocynaceae, Clusiaceae, Euphorbiaceae, and Rubiaceae

Berna Elya, Katrin Basah, Abdul Mun'im, Wulan Yuliasuti, Anastasia Bangun, and Eva Kurnia Septiana

Department of Pharmacy, Faculty of Mathematics and Sciences, University of Indonesia, Depok 16424, Indonesia

Correspondence should be addressed to Berna Elya, elya64@yahoo.com

Received 30 July 2011; Revised 8 September 2011; Accepted 9 September 2011

Academic Editor: Munekazu Inuma

Copyright © 2012 Berna Elya et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Diabetes mellitus (DM) is recognized as a serious global health problem that is characterized by high blood sugar levels. Type 2 DM is more common in diabetic populations. In this type of DM, inhibition of α -glucosidase is a useful treatment to delay the absorption of glucose after meals. As a megabiodiversity country, Indonesia still has a lot of potential unexploited forests to be developed as a medicine source, including as the α -glucosidase inhibitor. In this study, we determine the α -glucosidase inhibitory activity of 80% ethanol extracts of leaves and twigs of some plants from the Apocynaceae, Clusiaceae, Euphorbiaceae, and Rubiaceae. Inhibitory activity test of the α -glucosidase was performed *in vitro* using spectrophotometric methods. Compared with the control acarbose (IC₅₀ 117.20 μ g/mL), thirty-seven samples of forty-five were shown to be more potent α -glucosidase inhibitors with IC₅₀ values in the range 2.33–112.02 μ g/mL.

1. Introduction

Diabetes mellitus (DM) is the most common endocrine disease worldwide. About 173 million people suffer from diabetes mellitus. The number of people with diabetes mellitus will more than double over the next 25 years to reach a total of 366 million by 2030 [1]. In 2000, Indonesia is ranked the fourth largest number of people with DM, after India, China, and the United States, which is about 8.4 million people. The amount is expected to rise to 21.3 million in 2030 [2].

DM consists of several types, one of which is noninsulin-dependent diabetes mellitus (type 2 DM). This type of DM is more common, reaching 90–95% of the population with DM [3]. This increasing trend in type 2 DM has become a serious medical concern worldwide that prompts every effort in exploring for new therapeutic agents to stem its progress.

In type 2 DM, inhibition of α -glucosidase therapy is beneficial to delay absorption of glucose after a meal [4]. α -

glucosidase plays a role in the conversion of carbohydrates into glucose. By inhibiting α -glucosidase, glucose levels in the blood can be returned within normal limits [5].

Natural resources provide a huge and highly diversified chemical bank from which we can explore for potential therapeutic agents by bioactivity-targeted screenings [6]. As a megabiodiversity country, Indonesia still has a lot of potential unexploited forests to be developed as a source of phytopharmaca or modern medicine [7]. Opportunity exploration of medicinal plants is still very wide open in line with the development of herbal industry, herbal medicine, and phytopharmaca. Therefore, researchers try to explore the potential antidiabetic agents with the mechanism of action of α -glucosidase inhibition in several plant species from four families: Apocynaceae, Clusiaceae, Euphorbiaceae, and Rubiaceae. The four families were chosen because members of some species have been scientifically proven to have antidiabetic activity. Based on the theory of kinship through a systematic approach to plant (chemotaxonomy), plants with the same family generally have similar chemical content,

TABLE 1: Phytochemical screening of 80% ethanol extracts from some plants of Apocynaceae, Clusiaceae, Euphorbiaceae, and Rubiaceae.

Simplicia	Chemical contents						
	Alkaloid	Flavonoid	Terpenoid	Tannin	Glycoside	Saponin	Anthraquinone
Apocynaceae							
<i>Beaumontia multiflora</i> Teijsm. & Binn. Folium	+	+	-	+	+	+	+
<i>Beaumontia multiflora</i> Teijsm. & Binn. Cortex	-	-	-	-	+	+	-
<i>Carissa carandas</i> L. Folium	-	+	+	+	+	+	+
<i>Carissa carandas</i> L. Cortex	-	-	-	+	+	+	+
<i>Ochrosia citrodora</i> Lauterb. & K. Schum. Folium	+	-	+	+	+	+	+
<i>Rauvolfia sumatrana</i> Jack Folium	+	-	+	-	+	+	-
<i>Strophanthus caudatus</i> (Blume.f.) Kurz Folium	-	-	+	-	+	+	+
<i>Strophanthus caudatus</i> (Blume.f.) Kurz Cortex	+	-	+	+	+	+	-
<i>Strophanthus gratus</i> Baill. Folium	-	-	+	-	-	+	-
<i>Strophanthus gratus</i> Baill. Cortex	+	-	+	-	+	+	-
<i>Tabernaemontana sphaerocarpa</i> Blume Folium	+	-	+	-	+	+	-
<i>Willughbeia tenuiflora</i> Dyer ex Hook.f Folium	-	-	+	+	+	+	-
<i>Willughbeia tenuiflora</i> Dyer ex Hook.f Cortex	+	-	+	+	+	+	-
Clusiaceae							
<i>Calophyllum tomentosum</i> Wight. Folium	+	+	+	+	+	+	-
<i>Garcinia bancana</i> Miq. Folium	+	-	+	+	+	+	-
<i>Garcinia daedalanthera</i> Pierre. Folium	-	+	+	+	+	+	+
<i>Garcinia daedalanthera</i> Pierre. Cortex	-	-	+	+	+	+	+
<i>Garcinia hombroniana</i> Pierre. Folium	+	-	+	+	+	+	+
<i>Garcinia kydia</i> Roxb. Folium	-	+	+	+	+	+	+
<i>Garcinia rigida</i> Miq. Folium	+	+	+	+	+	+	+
Euphorbiaceae							
<i>Antidesma bunius</i> (L.) Spreng Folium	-	-	+	+	+	+	+
<i>Antidesma bunius</i> (L.) Spreng Cortex	+	-	+	+	+	+	-
<i>Antidesma celebicum</i> Cortex	-	-	-	+	+	+	-
<i>Antidesma celebicum</i> Folium	-	+	-	+	+	+	+
<i>Antidesma montanum</i> (Blume) Folium	+	-	+	+	+	+	-
<i>Antidesma neurocarpum</i> Miq. Folium	+	+	-	+	+	-	+
<i>Blumeodendron toksbrai</i> (Blume.) Kurz. Cortex	+	-	-	-	+	+	-
<i>Blumeodendron toksbrai</i> (Blume.) Kurz. Folium	+	-	+	-	+	-	-
<i>Croton argyratus</i> Blume. Folium	-	-	+	-	+	-	-

TABLE 1: Continued.

Simplicia	Chemical contents						
	Alkaloid	Flavonoid	Terpenoid	Tannin	Glycoside	Saponin	Anthraquinone
<i>Cephalomappa mallotica</i> J.J.Sm. Cortex	–	–	+	+	+	+	+
<i>Cephalomappa mallotica</i> J.J.Sm. Folium	–	–	+	+	+	–	+
<i>Galearia filiformis</i> Blume. Folium	+	–	+	+	+	–	+
<i>Sumbaviopsis albicans</i> (Blume) J.J.Sm. Cortex	–	–	+	–	+	+	–
<i>Sumbaviopsis albicans</i> (Blume) J.J.Sm. Folium	–	–	+	–	+	+	–
<i>Suregada glomerulata</i> (Blume) Baill. Folium	+	–	+	–	+	–	–
Rubiaceae							
<i>Adina trichotoma</i> Zoll. & Moritzi. Folium	+	–	+	–	+	–	+
<i>Amaracarpus pubescens</i> Blume. Folium	+	+	+	–	+	–	+
<i>Canthium glabrum</i> Blume. Folium	+	+	+	+	+	–	+
<i>Chiococca javanica</i> Blume. Folium	+	+	+	–	+	–	+
<i>Hydnophytum formicarum</i> Folium	+	–	+	+	+	+	–
<i>Hydnophytum formicarum</i> Cortex	+	+	+	–	+	–	–
<i>Nauclea calycina</i> (Batr.) ex DC.) Merr. Folium	+	–	–	+	+	–	–
<i>Nauclea calycina</i> (Batr.) ex DC.) Merr. Cortex	–	+	+	+	+	–	+
<i>Posoqueria latifolia</i> (Lam.) Roem. & Schult. Folium	–	–	+	+	+	–	–

Key: +: present; -: absent.

so it may just have the same potential for the treatment of a disease [8].

2. Method and Material

2.1. Plant Material. The stem bark and leaves of plants material were collected in November 2010 and identified by Center for Plant Conservation-Bogor Botanical Garden.

2.2. Extraction. Each dried powdered of wood bark, twig and leaves (20 g) were extracted by reflux with ethanol 80% then evaporated.

2.3. Inhibition Assay for α -Glucosidase Activity. The inhibition of α -glucosidase activity was determined using the modified published method [9]. One mg of α -glucosidase (*Saccharomyces cerevisiae*, Sigma-Aldrich, USA) was dissolved in 100 mL of phosphate buffer (pH 6.8) containing 200 mg of bovine serum albumin (Merck, German). The reaction mixture consisting 10 μ L of sample at varying concentrations (0.52 to 33 μ g/mL) was premixed with 490 μ L phosphate buffer pH 6.8 and 250 μ L of 5 mM *p*-nitrophenyl α -D-glucopyranoside (Sigma-Aldrich, Switzerland). After preincubating at 37°C for 5 min, 250 μ L α -glucosidase (0.15 unit/mL) was added and incubated at 37°C for 15 min.

The reaction was terminated by the addition of 2000 μ L Na₂CO₃ 200 mM. α -glucosidase activity was determined spectrophotometrically at 400 nm on spectrophotometer UV-Vis (Shimadzu 265, Jepang) by measuring the quantity of *p*-nitrophenol released from *p*-NPG. Acarbose was used as positive control of α -glucosidase inhibitor. The concentration of the extract required to inhibit 50% of α -glucosidase activity under the assay conditions was defined as the IC₅₀ value.

2.4. Kinetics of Inhibition against α -Glucosidase. Inhibition modes of sample that had the best α -glucosidase inhibiting activity in Clusiaceae, Euphorbiaceae, and Rubiaceae were measured with increasing concentration of *p*-nitrophenyl α -D-glucopyranoside as a substrate in the absence or presence of ethanolic extract at different concentrations. Inhibition type was determined by the Lineweaver-Burk plots analysis of the data, which were calculated from the result according to the Michaelis-Menten kinetics.

2.5. Phytochemistry Test. In this research we performed phytochemistry test which consists of alkaloid test with Mayer, Dragendorff, and Bouchardat reagents; Flavonoid test with Shinoda and Wilson Töubock reaction; tannin test with

TABLE 2: IC₅₀ values of rude extracts against α -glucosidase.

Number Sample		IC ₅₀ (μ g/mL)
(1)	Acarbose	117.20
Apocynaceae		
(2)	<i>Beaumontia multiflora</i> Teijsm. & Binn. Folium	79.80
(3)	<i>Beaumontia multiflora</i> Teijsm. & Binn. Cortex	130.20
(4)	<i>Carissa carandas</i> L. Folium	21.14
(5)	<i>Carissa carandas</i> L. Cortex	20.44
(6)	<i>Ochrosia citrodora</i> Lauterb. & K. Schum. Folium	112.02
(7)	<i>Rauvolfia sumatrana</i> Jack Folium	174.27
(8)	<i>Strophanthus caudatus</i> (Blume.f.) Kurz Folium	706.81
(9)	<i>Strophanthus caudatus</i> (Blume.f.) Kurz Cortex	13.93
(10)	<i>Strophanthus gratus</i> Baill. Folium	50.61
(11)	<i>Strophanthus gratus</i> Baill. Cortex	202.17
(12)	<i>Tabernaemontana sphaerocarpa</i> Blume Folium	554.32
(13)	<i>Willughbeia tenuiflora</i> Dyer ex Hook.f Folium	8.16
(14)	<i>Willughbeia tenuiflora</i> Dyer ex Hook.f Cortex	42.11
Clusiaceae		
(15)	<i>Calophyllum tomentosum</i> Wight. Folium	15.83
(16)	<i>Garcinia bancana</i> Miq. Folium	22.41
(17)	<i>Garcinia daedalanthera</i> Pierre. Folium	2.33
(18)	<i>Garcinia daedalanthera</i> Pierre. Cortex	3.71
(19)	<i>Garcinia hombroniana</i> Pierre. Folium	11.30
(20)	<i>Garcinia kydia</i> Roxb. Folium	3.88
(21)	<i>Garcinia rigida</i> Miq. Folium	24.48
Euphorbiaceae		
(22)	<i>Antidesma buniis</i> (L.) Spreng Folium	7.94
(23)	<i>Antidesma buniis</i> (L.) Spreng Cortex	3.90
(24)	<i>Antidesma celebicum</i> Cortex	3.93
(25)	<i>Antidesma celebicum</i> Folium	2.34
(26)	<i>Antidesma montanum</i> (Blume) Folium	2.83
(27)	<i>Antidesma neurocarpum</i> Miq. Folium	4.22
(28)	<i>Blumeodendron toksbrai</i> (Blume.) Kurz. Cortex	22.82
(29)	<i>Blumeodendron toksbrai</i> (Blume.) Kurz. Folium	64.78
(30)	<i>Croton argyratus</i> Blume. Folium	366.07
(31)	<i>Cephalomappa mallotica</i> J.J.Sm. Cortex	12.22
(32)	<i>Cephalomappa mallotica</i> J.J.Sm. Folium	2.66
(33)	<i>Galearia filiformis</i> Blume. Folium	21.54
(34)	<i>Sumbaviopsis albicans</i> (Blume) J.J.Sm. Cortex	42.66
(35)	<i>Sumbaviopsis albicans</i> (Blume) J.J.Sm. Folium	43.40
(36)	<i>Suregada glomerulata</i> (Blume) Baill. Folium	57.46
Rubiaceae		
(37)	<i>Adina trichotoma</i> Zoll. & Moritz. Folium	28.22
(38)	<i>Amaracarpus pubescens</i> Blume. Folium	3.64
(39)	<i>Canthium glabrum</i> Blume. Folium	117.85
(40)	<i>Chiococca javanica</i> Blume. Folium	23.86
(41)	<i>Hydnophytum formicarum</i> Folium	181.90
(42)	<i>Hydnophytum formicarum</i> Cortex	11.04

TABLE 2: Continued.

Number Sample		IC ₅₀ (μ g/mL)
(43)	<i>Nauclea calycina</i> (Batrl.ex DC.) Merr. Folium	18.81
(44)	<i>Nauclea calycina</i> (Batrl.ex DC.) Merr. Cortex	25.99
(45)	<i>Posoqueria latifolia</i> (Lam.) Roem. & Schult. Folium	80.27

gelatin test, gelatin-salt test, and test with ferrous (III) chloride; glycoside test with Molisch reaction; saponin test with honeycomb froth test; anthraquinone test with Bornträger reaction; terpenoid test with Liebermann-Burchard reagent.

3. Results and Discussion

3.1. Phytochemistry Test. Compounds with α -glucosidase inhibitory activity were preliminary identified by the existence of alkaloid, terpene, saponin, tannin, glycoside, flavonoid, and quinone (Table 1).

3.2. Assay for α -Glucosidase Inhibitory Activity. The α -glucosidase of *S. cerevisiae* is used to investigate the inhibitory activity of the rude extracts. α -glucosidase inhibitory activity of rude extracts compounds against α -glucosidases were determined using *p*-nitrophenyl- α -D-glucopyranoside (*p*-NPG) as a substrate and these were compared with acarbose (Table 2). The IC₅₀ values of compounds range from 2.33 μ g/mL to 706.81 μ g/mL. There are thirty-seven of samples which have IC₅₀ lower than acarbose. Extracts derived from leaves of *Garcinia daedalanthera* showed inhibitory activity against α -glucosidase enzyme significantly, with IC₅₀ value of 2.33 μ g/mL. Inhibitory activity of the enzyme α -glucosidase at forty-five extracts may be due to the glycoside content in each extract. Glycosides consist of sugars that may be structurally similar to carbohydrate which is a substrate of the enzyme α -glucosidase [10]. IC₅₀ value of samples of plant extracts are lower than acarbose because their active chemical compounds have no further fractionation and may have a synergistic effect in inhibiting α -glucosidase [11].

Inhibition mode of leaves extract of *Antidesma celebicum* from Euphorbiaceae was investigated. Inhibition mode of 80% ethanol extract showed competitive inhibitory mode. This mode may have been due because the structure is similar with glucose. This result is similar with inhibition mode of Nojirimycin which has a competitive inhibition against α -glucosidase [9] (Figure 1).

Inhibition mode of leaves extract of *Garcinia kydia* from Clusiaceae was investigated. Inhibition mode of 80% ethanol extract showed noncompetitive inhibitory mode [12] (Figure 2).

Inhibition mode of 80% ethanol extract from *Amaracarpus pubescens* Blume. leaves had a combination of competitive and uncompetitive inhibition. Combination of competitive and noncompetitive may have been due to the extract having more than one compound that has α -glucosidase inhibitory activity [13] (Figure 3).




FIGURE 1: Lineweaver-Burk plot of 80% ethanol extract of leaves of *Antidesma celebicum* with concentration of 2.08 $\mu\text{g/mL}$ with pNPG substrate concentration of 1.25, 2.5, 5, 10, and 20 mM.




FIGURE 2: Lineweaver-Burk plot of the reaction α -glucosidase in the presence of 80% ethanol extract from *Garcinia kydia* Roxb. Leaves.

4. Conclusion

In vitro assays of α -glucosidase activity showed thirty-seven of forty-five samples had IC_{50} values of between 2.33 $\mu\text{g/mL}$ and 112.02 $\mu\text{g/mL}$, which were lower than that of acarbose (117.20 $\mu\text{g/mL}$). Based on family, 80% ethanol extract from *Garcinia daedalanthera* Pierre. leaves (Clusiaceae), *Antidesma celebicum* leaves (Euphorbiaceae), *Amaracarpus pubescens* Blume. leaves (Rubiaceae), and *Willughbeia tenuiflora* Dyer ex Hook.f leaves (Apocynaceae) had the highest




FIGURE 3: Lineweaver-Burk plot of the reaction α -glucosidase in the presence of 80% ethanol extract from *Amaracarpus pubescens* Blume.

α -glucosidase inhibiting activity with IC_{50} of 2.33 $\mu\text{g/mL}$, 2.34 $\mu\text{g/mL}$, 3.64 $\mu\text{g/mL}$, and 8,16 $\mu\text{g/mL}$. Meanwhile, types of enzyme inhibition mechanism from *Garcinia kydia* Roxb. leaves (Clusiaceae), *Antidesma celebicum* leaves (Euphorbiaceae), and *Amaracarpus pubescens* Blume. leaves (Rubiaceae) were noncompetitive inhibitor, competitive inhibitor, and mixed inhibitor. Currently attempts to purify the active compound from leaves extract of *Garcinia kydia* Roxb. (Clusiaceae), *Antidesma celebicum* (Euphorbiaceae), and *Amaracarpus pubescens* Blume. (Rubiaceae) are conducted to understand the inhibitory mechanisms more clearly. Moreover, further *in vivo* study is also required.

Acknowledgment

The authors would like thank to Department of Pharmacy, Faculty of Mathematics and Natural Sciences, University of Indonesia for supporting this project.

References

- [1] I. Funke and M. F. Melzig, "Traditionally used plants in diabetes therapy—phytotherapeutics as inhibitors of α -amylase activity," *Revista Brasileira de Farmacognosia Brazilian Journal of Pharmacognosy*, vol. 16, no. 1, pp. 1–5, 2006.
- [2] S. Wild, G. Roglic, A. Green, R. Sicree, and H. King, "Global prevalence of diabetes: estimates for the year 2000 and projections for 2030," *Diabetes Care*, vol. 27, no. 5, pp. 1047–1053, 2004.
- [3] Department of Health Republic of Indonesia, *Pharmaceutical Care for Diabetes Mellitus*, Department of Health Republic of Indonesia, Kurihara, Indonesia, 2008.
- [4] K. Y. Kim, K. A. Nam, H. Kurihara, and S. M. Kim, "Potent α -glucosidase inhibitors purified from the red alga *Grateloupia elliptica*," *Phytochemistry*, vol. 69, no. 16, pp. 2820–2825, 2008.
- [5] L. H. Bösenberg and D. G. Van Zyl, "The mechanism of action of oral antidiabetic drugs: a review of recent literature," *Journal of Endocrinology, Metabolism and Diabetes of South Africa*, vol. 13, no. 3, pp. 80–88, 2008.

- [6] S. H. Lam, J. M. Chen, C. J. Kang, C. H. Chen, and S. S. Lee, " α -Glucosidase inhibitors from the seeds of *Syagrus romanzoffiana*," *Phytochemistry*, vol. 69, no. 5, pp. 1173–1178, 2008.
- [7] M. S. H. Wahyuningsih, S. Wahyuono, D. Santosa et al., "Plants exploration from central kalimantan forest as a source of bioactive compounds," *Biodiversitas*, vol. 9, pp. 169–172, 2008.
- [8] L. S. De Padua, N. Bunyapiaphatsara, and R. H. M. J. Lemmens, *Plant Resources of South-East Asia: Medicinal and Poisonous Plants 1*, Prosea Foundation, Bogor, Indonesia, 1999.
- [9] R. T. Dewi, Y. M. Iskandar, M. Hanafi et al., "Inhibitory effect of Koji *Aspergillus terreus* on α -glucosidase activity and postprandial hyperglycemia," *Pakistan Journal of Biological Sciences*, vol. 10, no. 18, pp. 3131–3135, 2007.
- [10] S. Sugiwati, S. Setiasi, and E. Afifah, "Antihyperglycemic activity of the mahkota dewa [*Phaleria macrocarpa* (scheff.) boerl.] leaf extracts as an alpha-glucosidase inhibitor," *Makara Kesehatan*, vol. 13, no. 2, pp. 74–78, 2009.
- [11] A. Andrade-Cetto, J. Becerra-Jiménez, and R. Cárdenas-Vázquez, "Alfa-glucosidase-inhibiting activity of some Mexican plants used in the treatment of type 2 diabetes," *Journal of Ethnopharmacology*, vol. 116, no. 1, pp. 27–32, 2008.
- [12] D. S. Lee and S. H. Lee, "Genistein, a soy isoflavone, is a potent α -glucosidase inhibitor," *FEBS Letters*, vol. 501, no. 1–3, pp. 84–86, 2001.
- [13] K. B. Storey, *Functional Metabolism: Regulation and Adaptation*, Wiley-Interscience, Hoboken, NJ, USA, 2004.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

