

NO. 2203

for. 282.24. (72)
G.R.E.

A CONSPECTUS OF THE MOSSES OF ANTARCTICA, SOUTH GEORGIA, THE FALKLAND ISLANDS AND SOUTHERN SOUTH AMERICA

By

DOROTHY M. GREENE

Institute of Terrestrial Ecology

Natural Environment Research Council



LIBRARY
BRITISH ANTARCTIC SURVEY
MADINGLEY ROAD
CAMBRIDGE CB3 0ET



0057792

CAMBRIDGE: PUBLISHED BY THE BRITISH ANTARCTIC SURVEY: 1986

NATURAL ENVIRONMENT RESEARCH COUNCIL

Front Cover: Tortula robusta, Cumberland Bay, South Georgia. Photo: I Everson



£7.00

Authors address:

Institute of Terrestrial Ecology
Monkswood Experimental Station
Abbots Ripton
Huntingdon PE17 2LS
United Kingdom
Telephone 04873 381/8

Copies available from:

The Distribution Centre
Blackhorse Road
Letchworth
Herts SG6 1HN
United Kingdom
Telephone 04626 72555

A CONSPECTUS OF THE MOSSES OF ANTARCTICA, SOUTH GEORGIA, THE FALKLAND ISLANDS AND SOUTHERN SOUTH AMERICA

By

DOROTHY M. GREENE

Institute of Terrestrial Ecology

Natural Environment Research Council

ABSTRACT

A SURVEY is presented of the taxa of mosses which have been reported from Antarctica, the islands of the Scotia Ridge, South Georgia, the Falkland Islands and southern South America, i.e. Argentina, Chile and Uruguay, including Juan Fernandez, from earliest times up to the end of 1977. The results show that 1727 species and 254 subspecies or varieties are currently accepted for the area; of these 254 species and 40 subspecies or varieties are believed to be endemic.

Nomenclature is based on that in *Index muscorum* (van der Wijk and others, 1959, 1962, 1964, 1967, 1969) or in later treatments where appropriate but in four cases, viz. *Racomitrium crispulum*, the Polytrichaceae, *Bryum antarcticum* and *Sphagnum*, reasons are given for adopting a different approach. The list of synonyms includes the basionym and those epithets which have been used for material from the area but is not nomenclaturally exhaustive. Some 60 *nomina nuda* are also listed since specimens bearing these names are frequently encountered in herbaria.

Distribution follows the worldwide scheme adopted in *Index muscorum* grouped here under six headings, viz. Juan Fernandez, southern South America, the Falkland Islands, South Georgia, Antarctic Peninsula region and Antarctic continental region, the records for the South Orkney Islands being included with the peninsula region. It was found that 41 of the taxa listed in *Index muscorum* as occurring in the area could not be substantiated from any other source and so have been deleted from the list. As genera and species are revised, further reductions in the totals of currently accepted taxa can be expected, although further field work may add others.

The list of references is extensive and includes about 900 entries, all of which have been examined by the author. Included in the list is the literature relating to the basionyms. Where appropriate, both bibliographic and effective dates have been given to facilitate literature retrieval and taxonomic requirements.

I. INTRODUCTION

FACILITIES provided during the International Geophysical Year (1957–58) offered bryologists their first opportunities to work in Antarctica. As a result of visits by S. W. Greene in 1960–61, R. E. Longton in 1963–65, R. I. L. Smith in 1964–67, 1969–71, the author in 1967–68, G. C. S. Clarke in 1967–68 and B. G. Bell in 1971–72, substantial collections were assembled in the British Antarctic Survey Herbarium (Greene, 1972) to be used for revisionary work on Antarctic bryophytes. Most pre-IGY collections had been made from ship-borne expeditions by collectors who were almost entirely non-botanists and who sent their material to European specialists for examination. As a consequence there developed a very scattered literature published in a variety of journals and expedition reports. The first attempt to draw this together was made by Steere (1961a, b), who produced check lists for Antarctica and South Georgia, respectively, which were later updated by Greene (1968a).

As the collections made by the above-mentioned bryologists contained so much that was new for the region, it soon became clear that the work of identification and preparation of taxonomic

revisions could not proceed satisfactorily without detailed knowledge of the taxa, and records of their distribution, in neighbouring areas. This applied in particular to southern South America and New Zealand, since it appeared that the Antarctic Peninsula and continental Antarctic moss floras were most closely related to the flora of Magellanic and Nova Zealandic regions. For the latter region Sainsbury's (1955) *A handbook of the New Zealand mosses* and Dixon's earlier *Studies in the bryology of New Zealand with special reference to the herbarium of Robert Brown, Christchurch, New Zealand* (Dixon, 1913, 1914, 1923, 1926, 1927, 1929a) provided a reasonable background. Kühnemann (1938) had compiled from the literature a check list of mosses for Argentina, which included the Falkland Islands and later he gave a list of genera for the neighbourhood of Buenos Aires (Kühnemann, 1944). Looser (1952) reviewed the contribution made by Thériot to knowledge of the Chilean moss flora, while Herter (1927–28, 1953) listed many relevant works for Uruguay. For the Magellanic region, where the first moss recorded as having been collected was obtained in 1690 (Middleton, 1909; Gunckel, 1971), Cardot (1908a) in *La flore bryologique des terres Magellaniques, de la Géorgie du Sud et de l'Antarctide* drew together the work of the early expeditions. But all these accounts remained unsynthesized, a different nomenclature was used in each and much of relevance had been published subsequently.

So, in response to the needs of the British Antarctic Survey's bryologists, a synthesis of the literature was undertaken to update Cardot (1908a) and to amalgamate into a single consolidated account the reports and records of mosses from southern South American and Antarctic regions based on the nomenclature of van der Wijk and others (1959, 1962, 1964, 1967, 1969) in *Index muscorum*. This synthesis is provided here in the form of a conspectus of taxa for almost the whole of the areas referred to in *Index muscorum* as Antarctica (Ant.) and America area No. 6 (Am. 6), i.e.:

<i>Antarctica</i>	Antarctic Peninsula and continental region together with all off-shore islands.
<i>America 6</i>	Chile, Argentina, Uruguay, Juan Fernandez and Easter Island, Falkland Islands, South Georgia, South Sandwich Islands.

The one exception is the records for the Easter Island flora which have been excluded since they are not believed to have any real affinity with the flora of Antarctica. It should also be appreciated that the moss floras of the sub-Antarctic Marion and Prince Edward Islands, îles Crozet, îles Kerguelen, Heard and Macdonald Islands and Macquarie Island, as well as oceanic island groups to the north of these, e.g. the islands of the Tristan–Gough group in the South Atlantic, New Amsterdam and St. Paul Islands in the South Indian Ocean and the islands normally referred to as the New Zealand Shelf islands, are not treated as they form parts of other geographical areas, *sensu Index muscorum*.

The present work lists 1959 currently accepted taxa, 1728 species and 232 subspecies or varieties, of which 254 species and 40 subspecies or varieties are, according to *Index muscorum*, endemic; in addition 60 *nomina nuda* are included. It covers the period from the earliest records up to 1977, although a few papers from 1978 and 1979 have been included. Purely ecological papers have been excluded except where these works contain lists of species from an area which is not covered in any taxonomic work.

Nomenclature follows that in *Index muscorum*, Crosby (1977a), or the most recent treatment, in which case the authority is given. Exceptions are the treatment of *Racomitrium crispulum* by Clifford (1955) and the Polytrichaceae by Smith (1971), because it was found that contemporary opinion did not accept the conclusions in these works. Another exception is the list of synonyms given for *Bryum antarcticum* by Dixon and Watts (1918). In recent years considerable doubt has been cast on the validity of these reductions by Horikawa and Ando (1967), Greene (1968a), and others. In view of this, and the fact that doubt exists as to whether *B. antarcticum* really belongs to the genus *Bryum*, it was decided to list all the species involved under their pre-Dixon and Watts (1918) nomenclature. The taxonomic status of *Bryum inconnexum* Card. is also in considerable doubt in spite of a recent paper by Ochi (1976) in Japanese.

The treatment of the genus *Sphagnum* has been presented in a simplified form. It proved too time-consuming and beyond the competence of the author to treat the nomenclature for *Sphagnum* in the same way as has been done for other genera. On the advice of Mr A. Eddy, of the British Museum (Nat. Hist.), the *Sphagnum* taxa which have been reported for the region are listed with their geographical distribution but no attempt has been made to quote basionyms or any other synonyms, although *Index muscorum* has been followed as to status. This is indicated in the same way as in the rest of the text.

Apart from the foregoing cases, the author has not made any taxonomic decisions, so obvious anomalies or mistakes remain, although some attempt has been made to draw attention to them.

The list is arranged alphabetically in genera and species and, as in *Index muscorum*, nothing under varietal rank has been given, although forms are sometimes mentioned in the distribution. Validly published, currently accepted names are given in bold italics, synonyms in italics, *nomina illegitima*, *homonyma illegitima* and *nomina nuda* are in Roman type (Table I) with the status indicated in each case. Synonyms are cross-referenced to their final names. The list of synonyms is not exhaustive but always includes the basionym and those names which have been used for the geographical regions covered by the list, as well as those which were considered necessary to make the nomenclature clear.

TABLE I
ABBREVIATIONS AND CONVENTIONS

Abbreviations

<i>comb. inval.</i>	invalid combination
distr.	distribution
err.	in error
f.	filius or son
<i>fide</i>	according to
forma	lowest-ranked infraspecific category
<i>hom. illeg.</i>	later and thus illegitimate homonym
<i>nom. illeg.</i>	illegitimate name
<i>nom. ival.</i>	invalidly published name
<i>nom. nud.</i>	name lacking a description or reference to a description
<i>nom. rejic.</i>	name rejected in favour of a later conserved name
<i>non</i>	not
pro parte	in part
<i>sensu</i>	as understood by
ssp.	subspecies or highest-ranked infraspecific category
syn.	synonym
<i>synon. excl.</i>	excluding synonyms in original publication
var.	variety or intermediate-ranked infraspecific category

The foregoing terms are to be understood in accordance with usage in botanical nomenclature, as explained in the introductory matter to volume one of *Index muscorum* (van der Wijk and others, 1959).

Conventions

Names in bold italics	(e.g. <i>Achrophyllum dentatum</i>) are those currently accepted.
Names in italics	(e.g. <i>Hookeria dentata</i>) are synonyms.
Names in Roman type	(e.g. <i>Hookeria denticulata</i>) are those which, for one reason or another contravene the rules of <i>The international code of botanical nomenclature</i> (Stafleu, 1978).

Although *nomina nuda* have no nomenclatural significance, they have been treated in the same way as valid species because both Mueller and Dusén created a significant number and, in the case of Dusén, specimens bearing these names are frequently encountered in world herbaria. It will be noted that on a number of occasions a name has been used by an author prior to its valid publication by the authority cited. For example, Paris frequently published names attributed to Dusén a year before Dusén himself published his descriptions. Camp (1915, 1921) and Costes (1921) both cited Thériot names before the latter validly published them. In those cases where it has been possible to be sure that the record pre-dating the valid publication of the taxon is based on the same type as the later valid publication, it has been cited without further comment in the distribution, although technically these names are all *nomina nuda*. Where the literature has revealed a misidentification, this has been shown by giving the reference followed by *fide* and a reference to the appropriate source. Orthographical errors not listed in *Index muscorum* are mentioned in the distribution lists after a reference as orthographical variants.

In a number of cases an author has used a specific epithet which is not given in *Index muscorum* or elsewhere and although an authority was cited it has proved impossible to trace the original publication. These names are included in the list as *nomina nuda*. One other situation deserves comment. A number of authors have reported *Hypnum liliiputeanum* from Chile but, as Thériot (1916) pointed out, Montagne (1842) described the taxon on material from Cuba. Mitten (1869) erroneously listed it for Chile and this error has been faithfully copied by later authors.

The distribution lists are an amalgamation of all the papers in which a taxon or its synonyms have been reported as occurring in the area. This extended method of citation has been adopted as an aid to specimen recovery for the taxonomic work. Despite what appears to be an extensive literature, the specimen base is surprisingly small, the same specimen, quite often the type, being cited by several authors as the only one available.

All taxa reported in *Index muscorum* as occurring in America 6 or Antarctica have been abstracted, but not all of these appear in the list of taxa. In most cases, this is due to the fact that synonyms irrelevant to the region have been deleted. There remain the 41 taxa cited in Table II where *Index muscorum* appears to be in error as the author has been unable to find any other reference to the occurrence of any of these taxa in the region.

The abbreviations for the author citations follow those used in *Index muscorum*; in the case of more recent authors, not cited in that work, the full name is used. The abbreviations of standard nomenclatural terms used in the text are summarized in Table I.

TABLE II
TAXA REPORTED ERRONEOUSLY FROM SOUTHERN SOUTH AMERICA OR ANTARCTICA IN
Index muscorum

Southern South America

- Aulacomnium androgynum* (Hedw.) Schwaegr.
- Bartramia micrangia* Herz.
- Breutelia karsteniana* (C. Muell.) Jaeg.
- Breutelia wainioi* Broth.
- Bryum cirrhatum* Hoppe et Hornsch. *hom. illeg.*
- Bryum incacorralis* Herz.
- Bryum incrasso-limbatum* Card.
- Bryum micro-laevigatum* C. Muell.
- Bryum pungens* Tayl.
- Bryum urbanskyi* Broth.
- Campylopus acervatus* Mitt.
- Campylopus bogotensis* Thér.
- Coscinodon trinervis* (Williams) Broth.
- Cryphaea patens* Hornsch. ex C. Muell.
- Cryphaea patens* Hornsch. ex C. Muell. var. *dilatata* Herz.
- Fissidens brevifolius* Hook. f. et Wils.
- Fissidens mollis* Mitt.
- Fissidens subcrispus* Besch.
- Fissidens taylorii* C. Muell.
- Funaria megalostoma* Mitt.
- Leptodontium subacutifolium* Besch. *nom. nud.*
- Meesia uliginosa* Hedw.
- Microdus densus* (Hook.) Besch.
- Mielichhoferia campylocarpa* (Hook. et Arnott) Mitt.
- Octoblepharum ampullaceum* Mitt.
- Orthotrichum albidum* Hedw.
- Papillaria pseudo-funalis* C. Muell.
- Philonotis hastata* (Dub.) Wijk et Marg.
- Pilotrichum fendleri* C. Muell.
- Polytrichum alpinum* var. *polare* (C. Muell.) Hag.
- Pseudocrossidium apiculatum* Williams
- Ptychomitrium serratum* B.S.G.
- Racopilum cuspidigerum* (Schwaegr.) Aongstr.
- Schlotheimia rugifolia* (Hook.) Schwaegr.
- Sematophyllum contiguum* (Mitt.) Mitt.
- Tortula austro-ruralis* (C. Muell.) Broth.
- Verrucidens macrosporus* Reim.

Antarctica

- Brachythecium pallidoflavens* Card.
- Isopterygium ambiguum* Card.
- Isopterygium brownii* Card.
- Rhynchostegium pseudo-murale* (Hamp.) Jaeg.

The list of references has been made as complete as possible and is arranged under the bibliographic (i.e. title page) date. In taxonomic work it is the effective date, i.e. the date of issue or availability, that is important. In *Index muscorum* only the effective dates are cited, which presents difficulty in locating many works since card indices and other retrieval systems in libraries are based exclusively on bibliographic dates. To facilitate retrieval, the references are arranged in the bibliography according to bibliographic dates with the effective date given within square brackets where it differs. The same policy has been adopted for author citations but elsewhere in the text only the bibliographic date is used. If the evidence for an effective date is available internally in a publication, e.g. in those volumes of the *Revue Bryologique et Lichénologique*, where the effective date is given on the cover, the date itself is cited without a source; in all other cases the authority for the effective date is given.

Extensive use has been made of the following works when determining effective dates of publication:

- van Steenis-Krauseman and Stearn* (1954),
- Sayre* (1959, 1969, 1971, 1975),
- Stafleu* (1967) and
- Stafleu and Cowan* (1976, 1979).

Considerable difficulty was experienced in finding effective dates of publication for the first edition of *Index bryologicus* (Paris, 1894, 1895, 1896, 1897, 1898), *Index muscorum* being found to be inconsistent in the dates quoted for the different parts. The dates used here are those given by Paris (1900b), but the complicated inter-relationships between the dates of publication of the fascicules and the order in which pages appeared in *Actes de la Société linnéenne de Bordeaux* have been dealt with by Stafleu and Cowan (1983).

Every reference has been checked personally by the author. This includes the literature of the basionyms, since many of these works are often cited inaccurately and a lot of them are rare and not readily available outside of large institutional libraries. It is hoped that the trouble taken to trace these works will facilitate their recovery by working taxonomists.

Inevitably a number of papers came to hand too late to be incorporated into the body of the text. Grassi's (1975a, b, 1976) introduction to Argentine bryophytes gives lists of species for the north-west of the country, while Mahu's (1979) paper presents a list of genera of Chilean mosses based on *Index muscorum*. The existence of an early paper by Porter (1900), dealing with Chilean mosses which is, as its author clearly states, a résumé of the appropriate parts of Cardot's (1900) preliminary account of the mosses collected by the Belgian Antarctic Expedition, has only recently come to the author's attention. Zander (1978) has proposed a number of new combinations in *Didymodon* and the relevant nomenclatural changes involving taxa treated here in *Barbula* and *Trichostomopsis* have been shown as footnotes in the text. This has also been done for Ochi (1977) and for his provisional revision of the genus *Bryum* in Antarctica (Ochi, 1979) in which some new synonyms are proposed. However, there is, as yet, no answer to the vexed question of the identity and synonymy of *Bryum antarcticum* Hook.f. et Wils.

Acanthocladium hom. illeg.

A. breviflagellosum Broth. (Brotherus, 1908) *nom. nud.*

Syn. Heterophyllum breviflagellosum C. Muell. (Mueller in Brotherus, 1908) *nom. nud.*

Distr. Southern South America (Hosseus, 1940).

Acaulon

A. lorentzii C. Muell.

Astomum lorentzii

***A. vesiculosum* C. Muell. (Mueller, 1888).**

Syn. Sphaerangium vesiculosum (C. Muell.) Kindb. (Kindberg, 1889).

Distr. Southern South America (Mueller, 1888; Paris, 1905b; Roth, 1911; Brotherus, 1924a; Kühnemann, 1938).

Achrohypnella

A. subnervis Herz.

Vesiculariopsis spirifolium

Achrophyllum***A. anomalam* (Schwaegr.) H. Robinson (Robinson, 1974).**

Syn. Racopilum anomalum Schwaegr. (Schwaegrichen, 1830).

Hookeria anomala (Schwaegr.) C. Muell. (Mueller, 1851a).

Pterygophyllum anomalum (Schwaegr.) Mitt. (Mitten, 1869).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1974, 1975).

Southern South America (Mitten, 1869; Mueller, 1885; Paris, 1905b; Cardot, 1905a, 1908a; Brotherus, 1925; Kühnemann, 1938; Welch, 1969).

***A. dentatum* (Hook. f. et Wils.) Vitt et Crosby (Vitt and Crosby, 1972).**

Syn. Hookeria dentata Hook. f. et Wils. (Hooker and Wilson, 1844).

Pterygophyllum dentatum (Hook. f. et Wils.) Dix. (Dixon, 1912).

Hookeria denticulata Hook. f. et Wils. (Hooker and Wilson in Wilson and Hooker, 1845) *hom. illeg.*

Pterygophyllum denticulatum Mitt. (Mitten, 1869) *nom. illeg.*

Hookeria nigella Hook. f. et Wils. (Hooker and Wilson in Wilson, 1854).

Pterygophyllum nigellum (Hook. f. et Wils.) Mitt. (Mitten, 1856).

Distr. Juan Fernandez (Mitten, 1885; Johow, 1896; Paris, 1905b; Brotherus, 1924b, 1925; Bartram, 1959; Robinson, 1975).

Southern South America (Wilson and Hooker, 1847; Mueller, 1851a, 1885; Mitten, 1869; Jaeger, 1877; Paris, 1905b; Cardot, 1908a; Thériot, 1917b; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Herzog, 1939, 1954).

Falkland Islands (Wilson and Hooker, 1847; Mueller, 1851a, 1885; Jaeger, 1877; Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Welch, 1969).

***A. tenuinerve* (Broth.) H. Robinson (Robinson, 1974).**

Syn. Pterygophyllum tenuinerve Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Welch, 1969; Matteri, 1972; Robinson, 1974, 1975).

Southern South America (Matteri, 1972).

Acidodontium

A. kunthii Schwaegr.

Acidodontium megalocarpum

***A. macropoma* (C. Muell.) Broth. (Brotherus, 1903).**

Syn. Bryum macropoma C. Muell. (Mueller, 1879a).

Brachymenium macropoma (C. Muell.) Kindb. (Kindberg, 1889).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1903; Brotherus, 1903, 1924a; Kühnemann, 1938).

A. megalocarpum (Hook.) Ren. et Card. (Renauld and Cardot, 1892 [1893]).

Syn. *Bryum megalocarpum* Hook. (Hooker *in Kunth*, 1822).

Acidodontium kunthii Schwaegr. (Schwaegrichen, 1827a) *nom. illeg.*

Distr. Southern South America (Hornschuch, 1840; Herter, 1928, 1933a).

Acroceratium nom. nud.

A. auriculatum (Mont.) Mitt.

Acrocladium auriculatum

A. politum (Hook. f. et Wils.) Mitt.

Catagonium politum

A. sarmentosum (Wahlenb.) Mitt.

Calliergon sarmentosum

Acrocladiopsis

A. complanata Card.

Catagonium politum

A. myura Card.

Catagonium myurum

A. nitida (Hook. f. et Wils.) Card.

Catagonium nitidum

Acrocladium

A. auriculatum (Mont.) Mitt. (Mitten, 1869).

Syn. *Hypnum auriculatum* Mont. (Montagne, 1843).

Stereodon auriculatus (Mont.) Mitt. (Mitten, 1859).

Coelidium auriculatum (Mont.) Jaeg. (Jaeger, 1878).

Acroceratium auriculatum (Mont.) Mitt. (Mitten, 1883).

Lembophyllum auriculatum (Mont.) Par. (Paris, 1896).

Hypnum chlamydophyllum Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Acrocladium chlamydophyllum (Hook. f. et Wils.) C. Muell. et Broth. (Mueller and Brotherus, 1900).

Hypnum encalyptratum Schimp. (Schimper *in Ångström*, 1872) *hom. illeg.*

Eurhynchium encalyptratum Schimp. (Schimper *in Jaeger*, 1878) *nom. nud.*

Eurhynchium eucalyptratum Schimp. ex Par. (Schimper *ex Paris*, 1895) *nom. nud.*

Hypnum eucalyptratum Lindb. ex Card. (Lindberg *ex Cardot*, 1908a) *err.*

Plagiothecium magellanicum Schimp. (Schimper *in Jaeger*, 1878) *nom. nud.*

Distr. Juan Fernandez (Robinson, 1975).

Southern South America (Montagne, 1843, 1845b, 1850, 1856; Wilson and Hooker, 1847 (and as *Hypnum chlamydophyllum* *fide* Karczmarz, 1966); Mueller, 1851a, 1885, 1889; Sullivant, 1859 (as *Hypnum chlamydophyllum* *fide* Karczmarz, 1966); Mitten, 1869; Ångström, 1872; Jaeger, 1878; Bescherelle, 1889; Paris, 1896, 1905a; Neger, 1899; Cardot, 1900, 1901, 1905a, 1908a (and as *A. chlamydophyllum*), 1913a; Dusén, 1903a, 1905d; Spegazzini, 1922; Cardot and Brotherus, 1923; Herzog, 1923, 1939, 1940, 1954, 1957, 1960; Brotherus, 1925; Roivainen, 1934; Thériot, 1934b; Kühnemann, 1938; Andrews, 1949a; Bartram, 1952; Karczmarz, 1966; Seki, 1974).

A. chlamydophyllum (Hook. f. et Wils.) C. Muell. et Broth.

Acrocladium auriculatum

A. cuspidatum (Hedw.) Lindb.

Calliergonella cuspidata

A. morenoi (C. Muell.) Dus.

Pleurozium schreberi

A. phyllogenium Jaeg.

Catagonium politum var. *phyllogenium*

A. politum (Hook. f. et Wils.) Mitt.

Catagonium politum

A. sarmentosum (Wahlenb.) Richs. et Wall.

Calliergon sarmentosum

A. stramineum (Brid.) Richs. et Wall.

Calliergon stramineum

Acrocryphaea *nom. illeg.*

A. julacea (Mitt.) Jaeg. (Jaeger, 1876).

Syn. *Cryphaea julacea* Mitt. (Mitten, 1869).

Grimmia julacea Hornsch. (Hornschuch, 1840) *hom. illeg.*

Distr. Southern South America (Brotherus, 1925; Kühnemann, 1938; Herzog, 1952a).

Acroschisma

A. wilsonii (Hook. f.) Jaeg.

Andreaea wilsonii

Aloina

A. catillum (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula catillum* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Paris, 1903; Kühnemann, 1938; Delgadillo, 1975).

A. cucullatifolia (C. Muell.) Broth.

Aloinella cucullatifolia

A. galeata (C. Muell.) Broth.

Aloinella galeata

A. recurvipatula (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula recurvipatula* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Brotherus, 1902, 1924a; Paris, 1903; Hosseus, 1935b, 1937; Kühnemann, 1938).

A. roseae (Williams) Delgadillo (Delgadillo, 1973).

Syn. *Pterygoneurum roseae* Williams (Williams, 1915).

Crossidium roseae (Williams) Bartr. (Bartram, 1942).

Distr. Southern South America (Bartram, 1942, 1943, 1952).

A. sedifolia (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula sedifolia* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Paris, 1903; Kurtz, 1904; Hosseus, 1935b, 1937; Kühnemann, 1938).

Aloinella

A. cucullatifolia (C. Muell.) Broth. (Brotherus, 1924a).

Syn. *Barbula cucullatifolia* C. Muell. (Mueller, 1879a).

Aloina cucullatifolia (C. Muell.) Broth. (Brotherus, 1902).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Paris, 1903; Kühnemann, 1938).

A. galeata (C. Muell.) Broth. (Brotherus, 1924a).

Syn. *Barbula galeata* C. Muell. (Mueller, 1879a).

Aloina galeata (C. Muell.) Broth. (Brotherus, 1902).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Paris, 1903; Kühnemann, 1938).

Amblystegiella nom. illeg.

A. densissima (Card.) Broth. (Brotherus, 1908).

Syn. *Amblystegium densissimum* Card. (Cardot, 1900).

Distr. Antarctic, peninsula region (Cardot, 1900, 1901, 1908a; Paris, 1903; Brotherus, 1908, 1925; Robinson, 1972 (as *Platydictya densissima*)).

A. tenuifolia (Hedw.) Fleisch.

Rhynchostegium tenuifolium

Amblystegium

A. aduncum (Hedw.) De Not.

Drepanocladus aduncus

ssp. *orthothecoides* (Lindb.) Lindb.

Drepanocladus uncinatus var. *subjulaceus*

var. *plumosum* (Schimp.) Hult

Drepanocladus uncinatus

A. austro-fluviatile (C. Muell.) Kindb.

Hygroamblystegium austro-fluviatile

A. austro-riparium Broth. (Brotherus *in* Kurtz, 1904) *nom. nud.*

Distr. Southern South America (Kurtz, 1904; Hosseus, 1936; Kühnemann, 1938).

A. austro-serpens (C. Muell.) Kindb.

Haplocladium microphyllum

A. brachiatum (Mitt.) Broth.

Drepanocladus brachiatus

<i>A. brachypelmatum</i> C. Muell.	<i>Leptodictyum brachypelmatum</i>
<i>A. campicola</i> (C. Muell.) Kindb.	<i>Leptodictyum campicola</i>
<i>A. chilense</i> Lor.	<i>Hygroamblystegium chilense</i>
<i>A. Clarazii</i> (Dub.) Jaeg.	<i>Helicodontium clarazii</i>
<i>A. densissimum</i> Card.	<i>Amblystegiella densissima</i>
<i>A. dusenii</i> Broth. (<i>Brotherus in Paris</i> , 1906) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (<i>Paris</i> , 1906).	
<i>A. exannulatum</i> (B.S.G.) De Not.	<i>Drepanocladus exannulatus</i>
<i>A. excurrens</i> Card. et Broth.	<i>Hygroamblystegium fuegianum</i> var. <i>excurrens</i>
<i>A. fluitans</i> (Hedw.) De Not.	<i>Drepanocladus fluitans</i>
ssp. <i>exannulatum</i> (B.S.G.) Arn. et C. Jens.	<i>Drepanocladus exannulatus</i>
var. <i>exannulatum</i> (B.S.G.) Lindb.	<i>Drepanocladus exannulatus</i>
var. <i>falcifolium</i> (Ren.) Arn. et C. Jens.	<i>Drepanocladus fluitans</i>
<i>A. fuegianum</i> Mitt.	<i>Drepanocladus fuegianus</i>
<i>A. intermedium</i> Lindb. var. <i>revolvens</i> (Sw.) Vent. et Bott.	<i>Drepanocladus revolvens</i>
<i>A. jaffuelii</i> Thér.	<i>Leptodictyum riparium</i> var. <i>jaffuelii</i>
<i>A. kneiffii</i> B.S.G. var. <i>polycarpum</i> (Voit) Braithw.	<i>Drepanocladus aduncus</i> var. <i>polycarpus</i>
<i>A. kochii</i> B.S.G.	<i>Leptodictyum kochii</i>
<i>A. longifolium</i> Mitt.	<i>Drepanocladus brachiatus</i>
<i>A. lucidum</i> (Hook. f. et Wils.) Jaeg.	<i>Plagiothecium lucidum</i>
<i>A. megachaete</i> Dus. (<i>Dusén</i> , 1903a) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (<i>Dusén</i> , 1903a).	
<i>A. orthocarpus</i> Broth. (<i>Brotherus in Kurtz</i> , 1904) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (<i>Kurtz</i> , 1904; <i>Hosseus</i> , 1936, 1937; <i>Kühnemann</i> , 1938).	
<i>A. orthothecoides</i> (Lindb.) Lindb. ex Par.	<i>Drepanocladus uncinatus</i> var. <i>subjulaceus</i>
<i>A. philonotis</i> Dus. (<i>Dusén in Paris</i> , 1906) <i>nom. nud.</i>	
<i>Distr.</i> Southern America (<i>Paris</i> , 1906; <i>Dixon</i> , 1929b).	
<i>A. polycarpum</i> (Voit) Vent. et Bott.	<i>Drepanocladus aduncus</i> var. <i>polycarpus</i>
<i>A. polygamum</i> B.S.G.	<i>Campylium polygamum</i>
<i>A. pulchellum</i> C. Muell.	<i>Amblystegium varium</i>
<i>A. radicale</i> (P. Beauv.) Mitt.	<i>Campylium radicale</i>
<i>A. revolvens</i> (Sw.) De Not.	<i>Drepanocladus revolvens</i>
<i>A. riparium</i> (Hedw.) B.S.G. ssp. <i>kochii</i> (B.S.G.) Ren. et Card.	<i>Leptodictyum kochii</i>
var. <i>kochii</i> (B.S.G.) Buyss.	<i>Leptodictyum kochii</i>
<i>A. rivulare</i> (C. Muell.) Broth. (<i>Brotherus</i> , 1908) <i>hom. illeg.</i>	
<i>Syn.</i> <i>Dimerodontium rivulare</i> C. Muell. (<i>Mueller</i> , 1897a).	
<i>Distr.</i> Southern South America (<i>Mueller</i> , 1897a; <i>Paris</i> , 1900a, 1904b; <i>Brotherus</i> , 1908, 1925; <i>Fleischer</i> , 1920; <i>Kühnemann</i> , 1938).	
<i>A. sarmentosum</i> (Wahlenb.) De Not.	<i>Calliergon sarmentosum</i>
<i>A. serpens</i> (Hedw.) B.S.G. (Bruch and others, 1853).	
<i>Syn.</i> <i>Hypnum serpens</i> Hedw. (<i>Hedwig</i> , 1801).	
<i>Hypnum oligorrhizone</i> Guemb. (<i>Guembel in Austin</i> , 1877) <i>nom. nud.</i>	
<i>Distr.</i> Juan Fernandez (<i>Robinson</i> , 1975).	
Southern South America (<i>Gibert</i> , 1873; <i>Rovainen</i> , 1934; <i>Piovano</i> , 1954).	
Falkland Islands (<i>Cardot</i> and <i>Brotherus</i> , 1923; <i>Kühnemann</i> , 1938).	

A. sparsifolium (Hamp.) Jaeg. (Jaeger in Jaeger and Sauerbeck, 1879a).

Syn. *Hypnum sparsifolium* Hamp. (Hampe, 1870).

Distr. Southern South America (Felippone, 1928; Herter, 1933a).

A. stramineum (Brid.) De Not.

Calliergon stramineum

A. subvarium Broth. (Brotherus, 1906a).

Distr. Antarctic, peninsula region (Darbshire, 1923; Greene, 1968a).

A. tenellum Card. et Broth.

Hygroamblystegium tenellum

A. trichopodium (Schultz) Hartm. ssp. *kochii* (B.S.G.) Loesk.

Leptodictyum kochii

var. *kochii* (B.S.G.) Lindb.

Leptodictyum kochii

A. uncinatum (Hedw.) De Not.

Drepanocladus uncinatus

var. *orthothecoides* (Lindb.) C. Jens.

Drepanocladus uncinatus var. *subjulaceus*

var. *plumosum* (Schimp.) Moell.

Drepanocladus uncinatus

var. *subjulaceum* (B.S.G.) C. Jens.

Drepanocladus uncinatus var. *subjulaceus*

var. *symmetricum* (Ren. et Card.) Dus.

Drepanocladus uncinatus var. *symmetricus*

A. varium (Hedw.) Lindb. (Lindberg, 1879).

Syn. *Leskea varia* Hedw. (Hedwig, 1801).

Hypnum varium (Hedw.) P. Beauv. (Palisot de Beauvois, 1805a).

Stereodon varius (Hedw.) Mitt. (Mitten, 1864).

Rigodium varium (Hedw.) Kindb. (Kindberg, 1883).

Amblystegium pulchellum C. Muell. (Mueller in Ule, 1899) nom. nud.

Hygroamblystegium varium (Hedw.) Moenk. (Mönkemeyer, 1911).

Distr. Southern South America (Williams, 1930).

var. *patagonicum* Card.

Drepanocladus aduncus var. *polycarpus*

Amphidium

A. cyathicarpum (Mont.) Broth.

Amphidium tortuosum

A. tortuosum (Hornschr.) H. Robinson (Robinson, 1975).

Syn. *Syrrhopodon tortuosus* Hornsch. (Hornschuch, 1841).

Zygodon cyathicarpus Mont. (Montagne, 1845a).

Didymodon cyathicarpus (Mont.) Mitt. (Mitten, 1860).

Rhabdoweisia cyathicarpa (Mont.) Mitt. (Mitten, 1869).

Amphoridium cyathicarpum (Mont.) Jaeg. (Jaeger, 1874).

Oncophorus cyathicarpus (Mont.) Mitt. (Mitten, 1885).

Amphidium cyathicarpum (Mont.) Broth. (Brotherus, 1902).

Weisia sphaerothecia C. Muell. (Mueller, 1882).

Rhabdoweisia sphaerothecia (C. Muell.) Kindb. (Kindberg, 1889).

Distr. Juan Fernandez (Mitten, 1885; Johow, 1896; Paris, 1900a, 1903; Skottsberg, 1914;

Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849, 1882; Jaeger, 1874; Brotherus, 1902, 1924a; Dusén, 1903a; Paris, 1903; Cardot, 1908a; Thériot, 1917b, 1921a; Costes, 1921; Kühnemann, 1938; Herzog, 1939; Lawton, 1961; Seki, 1974).

South Georgia (Cardot and Brotherus, 1923; Steere, 1961b).

Amphoridium hom. illeg.

A. cyathicarpum (Mont.) Jaeg.

Amphidium tortuosum

Amphoritheca

A. bonplandii (Hook.) Jaeg.

Funaria bonplandii

A. clavellata (Mitt.) Jaeg.

Funaria clavellata

A. papillosa (C. Muell.) Kindb.

Funaria papillosa

Anacalypta

<i>A. brunnea</i> (C. Muell.) Kindb.	<i>Didymodon brunneus</i>
<i>A. glauco-viridis</i> C. Muell.	<i>Didymodon glauco-viridis</i>
<i>A. gracillima</i> (C. Muell.) C. Muell.	<i>Trichostomum gracillimum</i>
<i>A. gymna</i> (C. Muell.) Kindb.	<i>Didymodon gymnus</i>
<i>A. imperfecta</i> (C. Muell.) Kindb.	<i>Barbula imperfecta</i>
<i>A. microthecium</i> C. Muell.	<i>Didymodon microthecius</i>
<i>A. nuda</i> (C. Muell.) Kindb.	<i>Didymodon brunneus</i>
<i>A. spathulato-linearis</i> (C. Muell.) Kindb.	<i>Didymodon spathulato-linearis</i>

Anacolia

<i>A. aurea</i> Dus. (Dusén, 1903a) <i>nom. nud.</i>	
Distr. Southern South America (Dusén, 1903a).	
<i>A. auricolor</i> Dus.	<i>Breutelia integrifolia</i>
<i>A. laevisphaera</i> (Tayl.) Flow. (Flowers, 1935).	

Syn. *Glyphocarpa laevisphaera* Tayl. (Taylor, 1846).
Bartramia laevisphaera (Tayl.) C. Muell. (Mueller, 1849).
Glyphocarpus laevisphaerus (Tayl.) Jaeg. (Jaeger, 1875).
Bartramia subsessilis Tayl. (Taylor, 1847).
Anacolia subsessilis (Tayl.) Broth. (Brotherus, 1904).
Anacolia subsessilis (Tayl.) Broth. var. *brevifolia* Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Flowers, 1952; Bartram, 1959;
Robinson, 1975).
Southern South America (Thériot, 1936; Hosseus, 1938c, d, 1939; Flowers, 1952;
Herzog, 1960).

<i>A. subsessilis</i> (Tayl.) Broth.	<i>Anacolia laevisphaera</i>
var. <i>brevifolia</i> Broth.	<i>Anacolia laevisphaera</i>

Ancistrodes

<i>A. genuflexa</i> (C. Muell.) Crosby (Crosby, 1976).	
Syn. <i>Pilotrichum genuflexum</i> C. Muell. (Mueller, 1844b [1845]).	
<i>Meteoriump genuflexum</i> (C. Muell.) Mitt. (Mitten, 1869).	
<i>Duseniella genuflexa</i> (C. Muell.) Broth. (Brotherus, 1906b).	
<i>Bryodusenia genuflexa</i> (C. Muell.) H. Robinson (Robinson, 1974).	
<i>Hookeria ancistrodes</i> Mont. (Montagne, 1845a).	
<i>Pilotrichum ancistrodes</i> (Mont.) Lor. (Lorentz, 1866).	
<i>Meteoriump ancistrodes</i> (Mont.) Par. (Paris, 1896) <i>hom. illeg.</i>	
Distr. Southern South America (Mueller, 1844b, 1850; Montagne, 1845a, 1850, 1856; Lorentz, 1866; Mitten, 1869; Jaeger, 1877; Paris, 1897, 1900a, 1905a; Brotherus, 1906b, 1925; Cardot, 1908a; Herzog, 1923, 1939, 1954, 1960; Reimers, 1926; Thériot, 1934b; Kühnemann, 1938; Seki, 1974).	

Andreaea

<i>A. acuminata</i> Mitt. (Mitten in Wilson, 1859).	
Syn. <i>Andreaea opaca</i> Card. ex Roth (Cardot ex Roth, 1911) <i>fide</i> Greene, 1968b.	
Distr. South Georgia (Roth, 1911, 1913; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b; Greene, 1968b).	
<i>A. acutifolia</i> Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).	
Distr. Southern South America (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Mitten, 1869; Jaeger, 1870; Bescherelle, 1889; Dusén, 1903a; Paris, 1903; Cardot, 1905a, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Seki, 1974).	

Falkland Islands (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a; Jaeger, 1870; Paris, 1903; Cardot, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

var. *rufescens* Hook. f. et Wils. (Hooker and Wilson *in Wilson and Hooker*, 1847).

Distr. Southern South America (Wilson and Hooker, 1847).

A. *alpina* Hedw. (Hedwig, 1801).

Syn. *Andreaea squamata* C. Muell. (Mueller, 1884 [1883]) *fide* Greene, 1968a.

Andreaea loricata Dus. (Dusén, 1903b).

Andreaea brevifolia Dus. (Dusén, 1903b) *fide* Roivainen and Bartram, 1937.

Andreaea parallela C. Muell. var. *brevifolia* (Dus.) Card. (Cardot, 1908a) *fide* Roivainen and Bartram, 1937.

Andreaea lanceolata Dus. ex Roth. (Roth, 1910) *hom. illeg.*

Andreaea ruficaulis Dus. (Dusén *in Roivainen and Bartram*, 1937) *nom. nud.*

Distr. Juan Fernandez (Robinson, 1975).

Southern South America (Wilson and Hooker, 1847; Sullivant, 1859; Mitten, 1969; Jaeger, 1870; Dusén, 1903a, b; Paris, 1906; Cardot, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Greene, 1968b; Seki, 1974).

Falkland Islands (Cardot and Brotherus, 1923; Roivainen and Bartram, 1937; Kühnemann, 1938; Greene, 1968b).

South Georgia (Cardot, 1906a, 1908a; Roth, 1910; Steere, 1961b (and as *A. regularis* var. *brevifolia*); Greene, 1968a, b).

A. *appendiculata* B.S.G. (Bruch and others, 1855).

Distr. Southern South America (Wilson and Hooker, 1847 (as *A. rupestris* L.); Bruch and others, 1855; Mueller, 1864, 1885; Mitten, 1869; Jaeger, 1870; Dusén, 1903a; Paris, 1903; Cardot, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

A. *arachnoidea* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1903; Roth, 1910; Brotherus, 1924a; Kühnemann, 1938).

A. brevifolia Dus.

Andreaea alpina

A. *depressinervis* Card. (Cardot, 1900).

Syn. *Andreaea depressinervis* Card. var. *compacta* Card. (Cardot, 1901) *fide* Greene, 1968b.

Distr. South Georgia (Greene, 1968b; Schultze-Motel, 1970).

Antarctic, peninsula region (Cardot, 1900, 1901, 1906a, 1907c, 1908a, 1911b, c, d, 1912b, 1913a; Paris, 1903; Wright, 1905 (as *Andreaea* sp.); Roth, 1910; Dixon, 1935; Steere, 1961a; Greene, 1968b; Greene and others, 1970; Robinson, 1972).

var. *compacta* Card.

Andreaea depressinervis

A. *flabellata* C. Muell. (Mueller, 1884 [1883]).

Syn. *Andreaea grimmoides* Dus. (Dusén, 1903b) *fide* Greene, 1968b.

Andreaea pumila Card. (Cardot, 1906a) *fide* Greene, 1968b.

Distr. Southern South America (Dusén, 1903b; Cardot, 1905a, 1908a; Roth, 1910; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1954; Greene, 1968b).

Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938; Greene, 1968b).

South Georgia (Cardot, 1906a, 1908a; Roth, 1910; Brotherus, 1924a; Steere, 1961b; Greene, 1968b).

A. *fragilis* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1903; Roth, 1910; Kühnemann, 1938).

A. fuegiana (Card.) S.W. Greene (Greene, 1972 [1973]).

Syn. *Neuroloma fuegiana* Card. (Cardot, 1911a) *hom. illeg. non Neuroloma Cand.*

Distr. Southern South America (Cardot, 1911a; Roth, 1913; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Seki, 1974). South Georgia (Greene, 1968b, 1972).

A. gainii Card. (Cardot, 1911b).

Distr. South Georgia (Greene, 1968b).

Antarctic, peninsula region (Cardot, 1911b, d, 1913a; Roth, 1913; Brotherus, 1924a; Bartram, 1957 (as *A. rupestris forma fide* Greene, 1968b); Steere, 1961a; Greene, 1968b; Greene and others, 1970; Robinson, 1972).

var. parallela (C. Muell.) S.W. Greene

Andreaea parallela

A. grimmiooides Dus.

Andreaea flabellata

A. heterophylla Card.

Andreaea regularis

A. hookeri Schimp. (Schimper in Bruch and others, 1855).

Distr. Southern South America (Bruch and others, 1855).

A. lanceolata Dus. ex Roth.

Andreaea alpina

A. latinervis Bartr. (Bartram in Roivainen and Bartram, 1937).

Distr. Southern South America (Roivainen and Bartram, 1937).

A. laxifolia Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).

Distr. Southern South America (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Mitten, 1869; Jaeger, 1870; Dusén, 1903a; Paris, 1903; Cardot, 1908a; Roth, 1910; Brotherus, 1924a).

var. minor Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).

Distr. Southern South America (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Paris, 1903; Cardot, 1908a).

A. leiophylla Card. ex Roth (Cardot ex Roth, 1913).

Distr. Southern South America (Roth, 1913; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938).

A. liliputana C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

A. lorentziana C. Muell. (Mueller, 1879a).

Syn. *Andreaea lorentzii* C. Muell. ex Kindb. (Kindberg, 1889) *nom. illeg.*

Distr. Southern South America (Mueller, 1879a; Paris, 1903; Roth, 1910; Brotherus, 1924a; Kühnemann, 1938).

A. lorentzii C. Muell. ex Kindb.

Andreaea lorentziana

A. loricata Dus.

Andreaea alpina

A. marginata Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).

Distr. Southern South America (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Mitten, 1869; Jaeger, 1870; Dusén, 1903a, b; Paris, 1903; Cardot, 1908a; Roth, 1910; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1954; Seki, 1974).

A. mutabilis Hook. f. et Wils.

Andreaea rupestris

var. microphylla Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

var. subsecunda Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).

Distr. Falkland Islands (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a; Paris, 1903; Cardot, 1908a; Kühnemann, 1938).

- var. *uncinata*** Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).
Distr. Falkland Islands (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a; Paris, 1903; Cardot, 1908a; Kühnemann, 1938).
- A. opaca* Card. ex Roth *Andreaea acuminata*
- A. pachyphylla* (C. Muell.) Broth. (Brotherus, 1901).**
Syn. *Grimmia pachyphylla* C. Muell. (Mueller, 1885).
Andreaea pachyphylla (C. Muell.) Broth. var. *acutifolia* Dus. (Dusén, 1903b) *fide* Schultze-Motel, 1970.
Distr. Southern South America (Mueller, 1885; Brotherus, 1901, 1924a; Dusén, 1903a, b; Paris, 1903; Cardot, 1908a; Roth, 1910; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1954; Schultze-Motel, 1970; Seki, 1974).
var. *acutifolia* Dus. *Andreaea pachyphylla*
- A. parallela* C. Muell. (Mueller, 1884 [1883]).**
Syn. *Andreaea gainii* Card. var. *parallela* (C. Muell.) S.W. Greene (Greene, 1968b) *hom.* *illeg.*
Distr. South Georgia (Cardot, 1906a, 1908a; Roth, 1910; Steere, 1961b; Greene, 1968b). Antarctic, peninsula region (Greene and others, 1970).
var. *brevifolia* (Dus.) Card. (Cardot, 1908a). *Andreaea alpina*
- A. patagonica* Dus. (Dusén, 1903b).** *Andreaea alpina*
Distr. Southern South America (Dusén, 1903b; Cardot, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).
- A. perichaetialis* (Hook. f. et Wils.) Jaeg. *Andreaea subulata*
A. petrophila Fuernr. *Andreaea rupestris*
- A. pilifera* Herz. et Thér. (Herzog and Thériot *in Herzog*, 1937).**
Distr. Southern South America (Herzog, 1937).
- A. pseudo-alpina* C. Muell. (Mueller, 1859).**
Distr. Southern South America (Mueller, 1859, 1885; Jaeger, 1870; Dusén, 1903a, b; Paris, 1903; Cardot, 1908a; Roth, 1910; Brotherus, 1924a).
- A. pseudomutabilis* Dus. (Dusén, 1903b).**
Distr. Southern South America (Dusén, 1903b; Cardot, 1908a; Roth, 1910; Brotherus, 1924a).
A. pseudo-subulata C. Muell. *Andreaea subulata*
- A. pulvinatula* Dus. (Dusén, 1903a) *nom. nud.*
Distr. Southern South America (Dusén, 1903a).
- A. pumila* Card. *Andreaea flabellata*
- A. purpurascens* (Dus.) Roth (Roth, 1910).**
Syn. *Andreaea remotifolia* Dus. var. *purpurascens* Dus. (Dusén, 1903b).
Distr. Southern South America (Dusén, 1903a, b; Cardot, 1908a; Roth, 1910; Brotherus, 1924a).
A. pycnotyla Card. *Andreaea regularis*
A. pygmaea Card. *Andreaea regularis*
- A. regularis* C. Muell. (Mueller, 1890a).**
Syn. *Andreaea pycnotyla* Card. (Cardot, 1900) *fide* Greene, 1968b.
Andreaea regularis C. Muell. var. *pycnotyla* (Card.) Card. (Cardot, 1908a) *fide* Greene, 1968b.
Andreaea heterophylla Card. (Cardot, 1906a) *fide* Greene, 1968b.
Andreaea pygmaea Card. (Cardot, 1900) *fide* Greene, 1968b.
Andreaea verruculosa Card. (Cardot, 1905a) *fide* Greene, 1968b.

Andreaea viridis C. Muell. (Mueller, 1890a) *fide* Greene, 1968b.
Andreaea willii C. Muell. (Mueller, 1890a) *fide* Greene, 1968b.

Distr. Southern South America (Cardot, 1905a, 1908a (including *A. mutabilis* *fide* Greene, 1968b); Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937 (including *A. mutabilis* p.p. *fide* Greene, 1968b); Kühnemann, 1938; Bartram, 1952; Greene, 1968b; Seki, 1974).
Falkland Islands (Cardot and Brotherus, 1923 (as *A. mutabilis* *fide* Greene, 1968b); Greene, 1968b).
South Georgia (Mueller, 1890a; Paris, 1903; Cardot, 1906a, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b; Greene, 1968b).
Antarctic, peninsula region (Cardot, 1900, 1901, 1906a, b, 1907a, c, 1908a, 1911b, d, 1913a; Paris, 1903; Roth, 1910; Brotherus, 1924a; Bartram, 1957 (as *A. rupestris* *fide* Greene, 1968b); Steere, 1961a (and as *A. rupestris* *fide* Greene, 1968b); Greene, 1968b; Greene and others, 1970; Robinson, 1972).

var. *pycnotyla* (Card.) Card.

Andreaea regularis

A. remotifolia Dus. (Dusén, 1903b).

Distr. Southern South America (Dusén, 1903b; Cardot, 1908a; Roth, 1910; Brotherus, 1924a).

var. *purpurascens* Dus.

Andreaea purpurascens

A. rothii Web. et Mohr (Weber et Mohr, 1807).

Syn. *Andreaea rupestris* A. Roth (Roth, 1802) *hom. illeg.*

Distr. Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

A. ruficaulis Dus.

Andreaea alpina

A. rupestris Hedw. (Hedwig, 1801).

Syn. *Andreaea mutabilis* Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).

Andreaea petrophila Fuernr. (Fürnrohr, 1827).

Distr. Southern South America (Mueller, 1851a; Jaeger, 1870; Dusén, 1903a; Paris, 1903; Cardot, 1905a, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952).
Falkland Islands (Jaeger, 1870; Dusén, 1903a; Paris, 1903; Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

A. rupestris A. Roth

Andreaea rothii

A. semi-squarrosa C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1903; Roth, 1910; Brotherus, 1924a; Kühnemann, 1938).

A. squamata C. Muell.

Andreaea alpina

A. squarrosa Mitt. (Mitten, 1869).

Distr. Southern South America (Wilson and Hooker, 1847 (as *A. alpina* Hedw. var. 1 *fide* Mitten, 1879); Donat, 1936a; Herzog, 1954; Seki, 1974)).

A. subremotifolia Dix. (Dixon, 1920).

Distr. South Georgia (Dixon, 1920; Greene, 1968b).

A. subulata Harv. (Harvey in Hooker, 1839).

Syn. *Andreaea subulata* Harv. var. *perichaetialis* Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).

Andreaea subulata Harv. var. *rigida* Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).

Andreaea pseudo-subulata C. Muell. (Mueller, 1864).

Andreaea perichaetialis (Hook. f. et Wils.) Jaeger (Jaeger, 1870) *hom. illeg.*

Distr. Southern South America (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1864, 1885; Mitten, 1869; Jaeger, 1870; Dusén, 1903a, b; Paris, 1903; Cardot, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Roivainen and Bartram, 1937; Kühnemann, 1938; Seki, 1974).

Falkland Islands (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a; Jaeger, 1870; Dusén, 1903a; Paris, 1903; Cardot, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Kühnemann, 1938).

var. *perichaetialis* Hook. f. et Wils. *Andreaea subulata*
 var. *rigida* Hook. f. et Wils. *Andreaea subulata*

A. vaginalis Herz. (Herzog in Donat, 1936a).

Distr. Southern South America (Donat, 1936a, b; Seki, 1974).

A. vermicularis Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a).

A. verruculosa Card.

Andreaea regularis

A. viridis C. Muell.

Andreaea regularis

A. willii C. Muell.

Andreaea regularis

A. wilsonii Hook. f. (Hooker in Hooker, J.D. and Wilson, 1844).

Syn. *Acroschisma wilsonii* (Hook. f.) Jaeg. (Jaeger, 1870).

Distr. Southern South America (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1851a, 1885; Mitten, 1869; Jaeger, 1870; Dusén, 1903a, b; Paris, 1903; Cardot, 1908a; Roth, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952; Herzog, 1954; Seki, 1974).

Anictangium *nom. rejic.*

A. ciliatum Hedw.

Hedwigia ciliata

A. coronatum Hook. et Wils.

Erpodium coronatum

A. humboldtii Hook.

Rhacocarpus purpurascens

A. imberbe (Sm.) Hook. et Tayl.

Hedwigidium integrifolium

A. orthotrichoides Gill. ex Grev. (Gillies ex Greville, 1830).

Distr. Southern South America (Greville, 1830).

A. subsessile (Brid.) Grev. et Arnott

Pterygoneurum subsessile

Anisothecium

A. capituligerum (C. Muell.) Thér. (Thériot, 1934b [1935]).

Syn. *Aongstroemia capituligera* C. Muell. (Mueller, 1879a).

Dicranella capituligera (C. Muell.) Kindb. (Kindberg, 1889).

Dichodontium capituligerum (C. Muell.) Par. (Paris, 1900a).

Distr. Southern South America (Mueller, 1879a; Paris, 1895, 1900a, 1903; Brotherus, 1901, 1924a; Kurtz, 1904; Hosseus, 1935b, 1937; Kühnemann, 1938).

A. elegans (Dub.) Thér. (Thériot, 1934b [1935]).

Syn. *Dicranum elegans* Dub. (Duby, 1870).

Dichodontium elegans (Dub.) Jaeg. (Jaeger in Jaeger and Sauerbeck, 1879b).

Aongstroemia elegans (Dub.) Broth. (Brotherus, 1901).

Distr. Southern South America (Duby, 1870; Jaeger and Sauerbeck, 1879b; Brotherus, 1901, 1924a; Paris, 1903; Herzog, 1923, 1938; Thériot, 1934b).

A. flexisetum Dus. (Dusén in Herzog, 1938) *nom. nud.*

Distr. Southern South America (Herzog, 1938).

A. hookeri (C. Muell.) Broth.

Dicranella hookeri

var. minus Card. et Thér. (Cardot and Thériot in Thériot, 1929c [1930]).

Distr. Southern South America (Thériot, 1929c).

A. jamesonii Mitt.

Anisothecium vaginatum

A. krausei (Lor.) Herz. (Herzog, 1939).

Syn. *Dicranella krausei* Lor. (Lorentz, 1866).

Dichodontium krausei (Lor.) Jaeg. (Jaeger, 1872).

Distr. Southern South America (Lorentz, 1866; Paris, 1904b; Herzog, 1939, 1954; Seki, 1974).

A. *lorentzii* (C. Muell.) Broth. (Brotherus, 1924a).

Syn. *Aongstroemia lorentzii* C. Muell. (Mueller, 1882).
Dicranella lorentzii (C. Muell.) Kindb. (Kindberg, 1889).
Dichodontium lorentzii (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1882; Paris, 1894, 1904b; Brotherus, 1901, 1924a; Kühnemann, 1938).

A. *paludella* (Besch.) Broth. (Brotherus, 1924a).

Syn. *Dichodontium paludella* Besch. (Bescherelle, 1885a).
Aongstroemia paludella (Besch.) C. Muell. (Mueller, 1901a [1900]).
Dicranella paludella (Besch.) Dus. (Dusén, 1905a).

Distr. Southern South America (Bescherelle, 1885a, 1889; Mueller, 1901a; Paris, 1904b; Dusén, 1905a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1954; Seki, 1974).

A. *perpusillum* Dus.

Dicranella hookeri

A. *persquarrosum* (Dus.) Broth.

Dicranella cardotii

var. *falklandicum* (Card.) Wijk et Marg. (van der Wijk and Margadant, 1960b).

Syn. *Dichodontium dicranelloides* Card. var. *falklandicum* Card. (Cardot, 1905a).
Dichodontium persquarrosum (Dus.) Card. var. *falklandicum* (Card.) Card. (Cardot, 1908a).

Distr. Falkland Islands (Cardot, 1905a, 1908a).

A. *pseudorufescens* (Card. et Broth.) Broth. (Brotherus, 1924a).

Syn. *Dicranella pseudorufescens* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

var. *leptocladus* (Card. et Broth.) Kühn. (Kühnemann, 1938).

Syn. *Dicranella pseudorufescens* Card. et Broth. var. *leptocladus* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

A. *rufipes* (C. Muell.) Dus. (Dusén, 1903a).

Syn. *Aongstroemia rufipes* C. Muell. (Mueller, 1882).
Dicranella rufipes (C. Muell.) Kindb. (Kindberg, 1889).
Dichodontium rufipes (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1882; Paris, 1894, 1904b); Brotherus, 1901, 1924a; Dusén, 1903a; Kühnemann, 1938).

A. *skottsbergii* (Card. et Broth.) Broth. (Brotherus, 1924a).

Syn. *Dicranella skottsbergii* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1954).

A. *vaginatum* (Hook.) Mitt. (Mitten, 1869).

Syn. *Dicranum vaginatum* Hook. (Hooker, 1816).
Aongstroemia vaginata (Hook.) C. Muell. (Mueller, 1851a).
Dichodontium vaginatum (Hook.) Jaeg. (Jaeger, 1872).
Dicranella vaginata (Hook.) Card. (Cardot, 1908a).
Dicranum jamesonii Tayl. (Taylor, 1847) hom. illeg.
Leptotrichum jamesonii Mitt. (Mitten, 1859).
Anisothecium jamesonii Mitt. (Mitten, 1869).
Dichodontium jamesonii (Mitt.) Jaeg. (Jaeger, 1872).
Aongstroemia jamesonii (Mitt.) C. Muell. (Mueller, 1901a [1900]).
Dicranella jamesonii (Mitt.) Broth. (Brotherus, 1901).

Distr. Southern South America (Montagne, 1839b (as *Dicranum vaginatum* Hook. ?); Wilson and Hooker, 1847; Mitten, 1869; Jaeger, 1872; Brotherus, 1901, 1924a; Paris, 1904b; Dusén, 1905a; Thériot, 1934b; Hosseus, 1938c, d, 1939; Herzog, 1938, 1939, 1940, 1954, 1960 (as *Anisodontium jamesonii* Tayl.)).

Anoectangium

- | | |
|---|------------------------------------|
| <i>A. acaule</i> (Web. et Mohr) Roehl. | <i>Pterygoneurum subsessile</i> |
| <i>A. ciliatum</i> (Hedw.) Roehl. | <i>Hedwigia ciliata</i> |
| <i>A. euchloron</i> (Schwaegr.) Mitt. | <i>Gymnostomum aestivum</i> |
| <i>A. excelsum</i> (C. Muell.) Kindb. | <i>Molendoa sendtneriana</i> |
| <i>A. fernandezianum</i> Card. (Cardot in Skottsberg, 1914) <i>nom. nud.</i> | |
| Distr. Juan Fernandez (Skottsberg, 1914). | |
| <i>A. humboldtii</i> (Hook.) Brid. | <i>Rhacocarpus purpurascens</i> |
| var. <i>australe</i> Hook. f. et Wils. | <i>Rhacocarpus purpurascens</i> |
| <i>A. integrifolium</i> (P. Beauv.) Schwaegr. | <i>Hedwigidium integrifolium</i> |
| <i>A. laxifolium</i> Par. | <i>Gymnostomum tenerrimum</i> |
| <i>A. lechlerianum</i> Mitt. | <i>Molendoa sendtneriana</i> |
| var. <i>limbatulum</i> Bartr. (Bartram, 1936). | |
| Distr. Southern South America (Bartram, 1936; Hosseus, 1938c, 1939). | |
| <i>A. lineare</i> (C. Muell.) Kindb. (Kindberg, 1889). | |
| Syn. <i>Zygodon linearis</i> C. Muell. (Mueller, 1879a). | |
| <i>Anoectangium mandonianum</i> (C. Muell.) Par. (Paris, 1894) <i>nom. inval.</i> | |
| <i>Zygodon mandonianus</i> Schimp. ex C. Muell. (Schimper ex Mueller, 1897b). | |
| Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1903; Brotherus, 1924a; Hosseus, 1938d; Kühnemann, 1938). | |
| <i>A. mandonianum</i> (C. Muell.) Par. | <i>Anoectangium lineare</i> |
| <i>A. patagonicum</i> Card. et Broth. (Cardot and Brotherus, 1923). | |
| Distr. Southern South America (Cardot and Brotherus, 1923). | |
| <i>A. sendtnerianum</i> B.S.G. | <i>Molendoa sendtneriana</i> |
| <i>A. subsessile</i> (Brid.) Spreng. | <i>Pterygoneurum subsessile</i> |
| <i>A. tenerrimum</i> C. Muell. | <i>Gymnostomum tenerrimum</i> |

Anomobryum

- | | |
|---|--|
| <i>A. brachymeniopsis</i> (C. Muell.) Broth. (Brotherus, 1903). | |
| Syn. <i>Bryum brachymeniopsis</i> C. Muell. (Mueller, 1879a). | |
| Distr. Southern South America (Mueller, 1879a; Paris, 1903; Brotherus, 1903, 1924a; Kühnemann, 1938). | |
| <i>A. conicum</i> (Hornschr.) Broth. (Brotherus, 1903). | |
| Syn. <i>Bryum conicum</i> Hornsch. (Hornschuch, 1840). | |
| Distr. Southern South America (Felippone, 1909; Herter, 1933a). | |
| <i>A. cygnicollum</i> (C. Muell.) Jaeg. (Jaeger, 1875). | |
| Syn. <i>Bryum cygnicollum</i> C. Muell. (Mueller, 1858). | |
| Mielichhoferia cygnicolla Hamp. (Hampe in Mueller, 1858) <i>nom. nud.</i> | |
| Climacodontium cygnicollum (C. Muell.) Hamp. (Hampe in Jaeger and Sauerbeck, 1879b) <i>comb. inval.</i> | |
| Bryum climacodontium Par. (Paris, 1894) <i>nom. illeg.</i> | |
| Distr. Southern South America (Mueller, 1858; Paris, 1894, 1904a). | |

A. felipponei Thér. (Thériot *in Felippone*, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933a).

A. filiforme (Dicks.) Husn. (Husnot, 1888) *hom. illeg.*

Syn. *Bryum filiforme* Dicks. (Dickson, 1801).

Pohlia filiformis (Dicks.) Andrews (Andrews, 1935).

Bryum julaceum Schrad. ex Gaertn., Meyer et Scherb. (Schrader *in Gärtner and others*, 1802).

Distr. Southern South America (Montagne, 1850; Mitten, 1869; Gibert, 1873; Paris, 1903; Campo, 1921; Cardot and Brotherus, 1923; Brotherus, 1924c; Herter, 1928, 1933a, b, 1943; Kühnemann, 1938; Herzog, 1938, 1939, 1954; Espinosa B., 1941; Seki, 1974).

A. laxirete Card. et Broth. (Cardot et Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

A. semireticulatum (C. Muell.) Broth. (Brotherus, 1903).

Syn. *Bryum semi-reticulatum* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a, 1882; Paris, 1903; Brotherus, 1903, 1924a; Hosseus, 1938c, d, 1939; Kühnemann, 1938).

A. terminale (C. Muell.) Broth. (Brotherus, 1903).

Syn. *Bryum terminale* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1903; Brotherus, 1903, 1924a; Hosseus, 1938c, d, 1939; Kühnemann, 1938).

Anomodon

A. curtipendulus (Hedw.) Hook. et Tayl.

Antitrichia curtipendula

A. pellicola C. Muell.

Leskea pellicula

A. pellicula C. Muell.

Leskea pellicula

A. toccae Sull. et Lesq.

Herpetineuron toccae

Antitrichia

A. curtipendula (Hedw.) Brid. (Bridel, 1819).

Syn. *Neckera curtipendula* Hedw. (Hedwig, 1801).

Hypnum curtipendulum (Hedw.) L. ex With. (Linnaeus ex Withering, 1801).

Anomodon curtipendulus (Hedw.) Hook. et Tayl. (Hooker and Taylor, 1818).

Cyrtopus curtipendulus Hedw. Spruc. (Spruce, 1849).

Haptymenium curtipendulum (Hedw.) Fuernr. (Fürnrohr, 1827).

Leucodon curtipendulus (Hedw.) T. Jens. (Jensen, 1856).

Distr. Southern South America (Bridel, 1801b, 1817, 1827b; Paris, 1903).

Aongstroemia

A. argentinica C. Muell.

Dicranella argentinica

A. aulacocarpa (Mont.) C. Muell.

Dicranella aulacocarpa

A. capituligera C. Muell.

Anisothecium capituligerum

A. elegans (Dub.) Broth.

Anisothecium elegans

A. euchlora (Mont.) C. Muell.

Microdus euchlorus

A. gayana (Mont.) C. Muell. (Mueller, 1848a).

Syn. *Dicranum gayanum* Mont. (Montagne, 1845a).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Lorentz, 1866; Mitten, 1869; Jaeger, 1872; Brotherus, 1901, 1924a; Paris, 1903; Dusén, 1905a; Campo, 1921; Costes, 1921; Thériot, 1921a; Herzog, 1938, 1939, 1954; Seki, 1974).

A. hilariana (Mont.) C. Muell.

Dicranella hilariana

<i>A. hookeri</i> C. Muell.	<i>Dicranella hookeri</i>
<i>A. jamesonii</i> (Mitt.) C. Muell.	<i>Anisothecium vaginatum</i>
<i>A. leiocarpa</i> C. Muell. (Mueller, 1901a [1900]) <i>nom. nud.</i>	
<i>Syn.</i> <i>Cynodontium leiocarpum</i> Lor. (Lorentz in Mueller, 1901a [1900]) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Mueller, 1901a).	
<i>A. lorentzii</i> C. Muell.	<i>Anisothecium lorentzii</i>
<i>A. paludella</i> (Besch.) C. Muell.	<i>Anisothecium paludella</i>
<i>A. patagonica</i> C. Muell.	<i>Dicranella patagonica</i>
<i>A. persquarrosa</i> Dus.	<i>Dicranella cardotii</i>
<i>A. rufipes</i> C. Muell.	<i>Anisothecium rufipes</i>
<i>A. subclathrata</i> (Lor.) C. Muell.	<i>Dicranella hookeri</i>
<i>A. tenuirostris</i> (Schwaegr.) C. Muell.	<i>Dicranella hilariana</i>
<i>A. vaginata</i> (Hook.) C. Muell.	<i>Anisothecium vaginatum</i>

Apalodium

<i>A. australe</i> (Hook. f. et Wils.) Mitt.	<i>Orthodontium lineare</i>
<i>A. confine</i> (Hamp.) Mitt.	<i>Orthodontium pellucens</i>
<i>A. lineare</i> (Schwaegr.) Mitt.	<i>Orthodontium lineare</i>
<i>A. pellucens</i> (Hook.) Mitt.	<i>Orthodontium pellucens</i>
<i>A. tenue</i> Mitt.	<i>Pohlia humilis</i>

Aptychella***A. chilensis* Herz. (Herzog, 1954).***Distr.* Southern South America (Herzog, 1954; Seki, 1974; Tixier, 1977).***Aptychus***

<i>A. ampullatus</i> C. Muell. <i>ex Par.</i>	<i>Sematophyllum ampullulatum</i>
<i>A. ampullulatus</i> C. Muell.	<i>Sematophyllum ampullulatum</i>
<i>A. aureo-viridis</i> C. Muell.	<i>Sematophyllum aureo-viride</i>
<i>A. catilliformis</i> C. Muell.	<i>Sematophyllum bracteatum</i>
<i>A. cochleatulus</i> C. Muell.	<i>Rhaphidostegium cochleatum</i>
<i>A. condensatulus</i> C. Muell.	<i>Rhapidostegium condensatum</i>
<i>A. diaphanodictyus</i> C. Muell.	<i>Rhaphidostegium diaphanodictyon</i>
<i>A. grandicellulosus</i> C. Muell.	<i>Sematophyllum grandicellulosum</i>
<i>A. laxo-alaris</i> C. Muell.	<i>Entodon laxo-alaris</i>
<i>A. macrocytus</i> C. Muell.	<i>Sematophyllum macrocytus</i>
<i>A. micrangius</i> C. Muell.	<i>Sematophyllum micrangium</i>
<i>A. micropyxis</i> C. Muell.	<i>Trichosteleum micropyxidium</i>
<i>A. nanocephalus</i> C. Muell.	<i>Sematophyllum nanocephalum</i>
var. <i>subglaucus</i> C. Muell.	<i>Sematophyllum nanocephalum</i> var. <i>subglaucus</i>
<i>A. serifolius</i> C. Muell.	<i>Sematophyllum serifolium</i>
<i>A. temperatus</i> C. Muell.	<i>Sematophyllum temperatum</i>
<i>A. tenerifolius</i> C. Muell.	<i>Sematophyllum tenerifolium</i>

Archidium***A. amplexicaule* C. Muell. (Mueller, 1882).**

Distr. Southern South America (Mueller, 1882; Paris, 1903; Roth, 1911; Brotherus, 1924a; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939).

var. *brevi-solonaceum* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1903; Roth, 1911).

A. *arechavaletae* C. Muell. (Mueller, 1888).

Distr. Southern South America (Mueller, 1888; Paris, 1903; Roth, 1911; Felippone, 1912; Brotherus, 1924a; Herter, 1933a).

A. *ephemeroides* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1903; Roth, 1911; Brotherus, 1924a; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939).

A. *gibertii* Mitt. (Mitten, 1886).

Distr. Southern South America (Mitten, 1886; Paris, 1903; Roth, 1910; Brotherus, 1924a; Herter, 1933a).

A. *julaceum* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1903; Roth, 1911; Thériot, 1918, 1924; Campo, 1921; Brotherus, 1924a; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939).

A. *lorentzii* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1903; Roth, 1911; Brotherus, 1924a; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939; Herzog, 1952a).

Aschistodon *nom. rejic.*

A. *conicus* Mont.

Ditrichum conicum

Asteriscium *hom. illeg.*

A. *australasiae* (Hook. et Grev.) Hilp.

Trichostomopsis australasiae

A. *curvipes* (C. Muell.) Hilp.

Trichostomopsis curvipes

A. *deciduum* Hilp.

Barbula uruguensis

A. *flavisetum* Herz.

Trichostomopsis umbrosa

A. *geniculatum* (Mont.) Herz.

Barbula geniculata

A. *graminicola* (C. Muell.) Hilp.

Barbula graminicolor

A. *poeppigianum* (C. Muell.) Hilp.

Trichostomopsis australasiae

A. *riparium* (C. Muell.) Hilp.

Barbula riparia

A. *subtophaceum* (Williams) Hilp.

Trichostomopsis australasiae

A. *umbrosum* (C. Muell.) Herz.

Trichostomopsis umbrosa

Astomiopsis

A. *amblyocalyx* C. Muell. (Mueller, 1882).

Syn. Astomiopsis amblyocarpa C. Muell. ex Par. (Mueller ex Paris, 1894) *nom. inval.*
Pleuridiopsis amblyocalyx (C. Muell.) Par. (Paris, 1897) *comb. inval.*

Distr. Southern South America (Mueller, 1882; Paris, 1897, 1905b; Brotherus, 1924a; Kühnemann, 1938).

A. *amblyocarpa* C. Muell. ex Par.

Astomiopsis amblyocalyx

A. *pacifica* Buck et Landrum (Buck and Landrum, 1977).

Distr. Juan Fernandez (Buck and Landrum, 1977).

A. *subulata* C. Muell. (Mueller, 1882).

Syn. Pleuridiopsis subulata (C. Muell.) Par. (Paris, 1897) *comb. inval.*

Distr. Southern South America (Mueller, 1882; Paris, 1897, 1905b; Brotherus, 1924a; Kühnemann, 1938).

Astomum

A. chilense Williams

Tetrapherum recurvirostre

A. fruchartii (C. Muell.) Broth. (Brotherus, 1900).

Syn. *Phascum fruchartii* C. Muell. (Mueller, 1888).

Systemium fruchartii (C. Muell.) Kindb. (Kindberg, 1889).

Distr. Southern South America (Mueller, 1888; Paris, 1905b; Roth, 1911; Felippone, 1912; Brotherus, 1924a; Herter, 1933a, 1943).

A. lorentzii (C. Muell.) Broth. (Brotherus, 1901).

Syn. *Acaulon lorentzii* C. Muell. (Mueller, 1882).

Sphaerangium lorentzii (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1882; Paris, 1897, 1905b; Brotherus, 1901, 1924a; Roth, 1911, 1914; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939).

A. nervosum (Hook.) C. Muell.

Pleuridium nervosum

A. recurvirostre (C. Muell.) Roth

Tetrapherum recurvirostre

A. robinsonii (Mont.) C. Muell.

Pleuridium robinsonii

A. subnervosum C. Muell.

Pleuridium subnervosum

Atractylocarpus

A. patagonicus Herz. et Thér. (Herzog and Thériot in Herzog and others, 1933).

Distr. Southern South America (Herzog and others, 1933).

Atrichopsis

A. compressa (Hook. f. et Wils.) G.L. Smith (Smith, 1969a).

Syn. *Polytrichum compressum* Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).

Catharinea compressa (Hook. f. et Wils.) C. Muell. (Mueller, 1848a).

Psilotilum compressum (Hook. f. et Wils.) Mitt. (Mitten, 1869).

Oligotrichum compressum (Hook. f. et Wils.) Kindb. (Kindberg, 1888).

Atrichopsis magellanica Card. (Cardot, 1912a) *fide* Smith, 1969a.

Distr. Southern South America (Hooker, J.D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Mitten, 1869; Jaeger, 1875; Paris, 1905b; Cardot, 1908a, 1912a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Herzog, 1939; Smith, 1969a).

Falkland Islands (Cardot, 1912a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Smith, 1969a).

A. magellanica Card.

Atrichopsis compressa

Atrichum

A. planifolium (C. Muell.) Jaeg.

Atrichum polycarpum

A. polycarpum (C. Muell.) Mitt. (Mitten, 1869).

Syn. *Catharinea polycarpa* C. Muell. (Mueller, 1851a).

Catharinea planifolia C. Muell. (Mueller in Lorentz, 1868).

Atrichum planifolium (C. Muell.) Jaeg. (Jaeger, 1875).

Distr. Southern South America (Lorentz, 1868; Jaeger, 1875; Frye and Duckering, 1948).

A. rigidum Lor.

Oligotrichum rigidum

Aulacomnium

A. chilense C. Muell.

Rhizogonium mnioides

A. gaudichaudii (Schwaegr.) Mitt.

Leptotheca gaudichaudii

A. palustre (Hedw.) Schwaegr. (Schwaegrichen, 1827b).

Syn. *Mnium palustre* Hedw. (Hedwig, 1801).

Bryum palustre (Hedw.) Gaertn., Meyer et Scherb. (Gärtner and others, 1802) hom. illeg.

Hypnum palustre (Hedw.) Web. et Mohr (Weber and Mohr, 1803) hom. illeg.

Bartramia palustris (Hedw.) P. Beauv. (Palisot de Beauvois, 1805b).

Orthopyxis palustris (Hedw.) P. Beauv. (Palisot de Beauvois, 1805a).

Gymnocybe palustris (Hedw.) Fries (Fries, 1825).

Limnobryum palustre (Hedw.) Rabenh. (Rabenhorst, 1863).

Sphaerocephalus palustris (Hedw.) Lindb. (Lindberg, 1879).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938; Herzog, 1957).

A. pentastichum Mont.

Zygodon pentastichus

Aulacopilum

A. glaucum Wils. (Wilson, 1848).

Syn. *Aulacopilum intermedium* Broth. (Brotherus, 1918) *fide* Crum, 1972.

Distr. Southern South America (Brotherus, 1918, 1925; Kühnemann, 1938; Crum, 1972).

A. intermedium Broth.

Aulacopilum glaucum

Barbula

B. aculeonervis C. Muell.

Tortula aculeonervis

B. aestiva (Hedw.) Schultz

Tortula muralis var. *aestiva*

B. alpicola C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

B. amblyophylla (Hook.) Jaeg.

Didymodon amblyophyllus

B. amphidiifolia C. Muell.

Tortula amphidiifolia

B. anacamptophylla C. Muell.

Tortula lepto-syntrichia

B. anastomosans C. Muell. (Mueller, 1879a).

Syn. *Barbula anostomosans* Par. (Paris, 1894) err.

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904a; Brotherus, 1918, 1924a; Kühnemann, 1938).

B. anderssonii (Aongstr.) Jaeg.

Tortula anderssonii

B. annulus C. Muell.

Tortula muralis

B. anostomosans Par.

Barbula anastomosans

B. antarctica Hamp.

Tortula princeps

B. apoclada Par.

Barbula perrevoluta

B. appressa (Mitt.) Jaeg.

Tortula appressa

B. arechavaleta C. Muell. (Mueller, 1901a [1900]) nom. nud.

Distr. Southern South America (Mueller, 1901a).

B. arenae Besch.

Tortula arenae

B. arenicola Dus. (Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Brotherus, 1924a; Kühnemann, 1938).

B. argentinica Par.

Didymodon argentinicus

B. atrovirens (Sm.) Schimp.

Desmatodon convolutus

B. australasiae (Hook. et Grev.) Brid.

Trichostomopsis australasiae

B. australis Par.

Tortula fuegiana

B. austrogracilis Dus. (Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Brotherus, 1924a).

<i>B. berteroana</i> C. Muell.	<i>Tortula muralis</i>
<i>B. berteroi</i> C. Muell. ex Kindb.	<i>Tortula muralis</i>
<i>B. brachychaete</i> Dus.	<i>Tortula brachychaete</i>
<i>B. breviseta</i> (Mont.) C. Muell.	<i>Tortula breviseta</i>
<i>B. brunnea</i> (C. Muell.) Broth.	<i>Didymodon brunneus</i>

B. byrdii Bartr. (Bartram, 1938).

Distr. Antarctic, continental region (Bartram, 1938, 1957; Steere, 1961a).

<i>B. caespitosa</i> Schwaegr.	<i>Tortella humilis</i>
<i>B. calycina</i> Schwaegr.	<i>Tortella calycina</i>

B. calyculosa (Mitt.) Jaeg. (Jaeger, 1873).

Syn. *Tortula calyculosa* Mitt. (Mitten, 1869).

Distr. Southern South America (Herzog, 1954).

B. cardotii Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b, 1906; Brotherus, 1924a; Kühnemann, 1938).

<i>B. catillum</i> C. Muell.	<i>Aloina catillum</i>
<i>B. cavifolia</i> Schimp.	<i>Pterygoneurum ovatum</i>
<i>B. characodonta</i> C. Muell.	<i>Tortula characodonta</i>
<i>B. chilensis pilifera</i> C. Muell. (Mueller, 1901a [1900]) <i>nom. nud.</i>	

Distr. Southern South America (Mueller, 1901a).

<i>B. chrysopila</i> C. Muell.	<i>Tortula chrysopila</i>
<i>B. conotricha</i> C. Muell.	<i>Tortula anderssonii</i>
var. <i>fagicola</i> C. Muell.	<i>Tortula anderssonii</i> var. <i>fagicola</i>
<i>B. contorta</i> Hamp. ex C. Muell.	<i>Tortula contorta</i>

B. costessii Thér. (Thériot, 1921a).

Distr. Southern South America (Thériot, 1921a; Costes, 1921; Brotherus, 1924a).

<i>B. crinita</i> Schultz	<i>Tortula pilifera</i>
<i>B. crispatula</i> C. Muell. ex Par.	<i>Calyptopogon mnioides</i>
<i>B. cucullatifolia</i> C. Muell.	<i>Aloinella cucullatifolia</i>
<i>B. curvipes</i> C. Muell.	<i>Trichostomopsis curvipes</i>
<i>B. curvirostris</i> Lindb.	<i>Hymenostylium recurvirostre</i>
<i>B. decidua</i> C. Muell.	<i>Barbula uruguensis</i>

B. declivium C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

<i>B. densifolia</i> Hook. f. et Wils.	<i>Tortula densifolia</i>
<i>B. depressa</i> Sull.	<i>Tortula pilifera</i>
var. <i>denticulata</i> (Dus.) Thér.	<i>Tortula pilifera</i> var. <i>denticulata</i>
var. <i>gracilis</i> (Dus.) Thér.	<i>Tortula pilifera</i> var. <i>gracilis</i>
var. <i>oliviensis</i> (Card.) Thér.	<i>Tortula pilifera</i> var. <i>oliviensis</i>
<i>B. dioica</i> (R. Brown ter.) Par.	<i>Tortula robusta</i>

<i>B. filaris</i> C. Muell.	<i>Tortula filaris</i>
<i>B. flagellaris</i> Schimp.	<i>Tortula flagellaris</i>
var. <i>denticulata</i> Dus.	<i>Tortula pilifera</i> var. <i>denticulata</i>
var. <i>gracilis</i> Dus.	<i>Tortula pilifera</i> var. <i>gracilis</i>
<i>B. flavidо-pilosa</i> Dus.	<i>Tortula flavidо-pilosa</i>
<i>B. flaviseta</i> (Herz.) Wijk et Marg.	<i>Trichostomopsis umbrosa</i>
<i>B. fontana</i> C. Muell.	<i>Tortula fontana</i>
<i>B. fuegiana</i> (Mitt.) Jaeg.	<i>Tortula fuegiana</i>
 <i>B. fusca</i> C. Muell. (Mueller, 1849).	
Syn. <i>Tortula fusca</i> (C. Muell.) Mont. (Montagne, 1850).	
Distr. Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Paris, 1904a; Dusén, 1906; Cardot, 1908a; Thériot, 1918, 1921a, 1928; Campo, 1921; Costes, 1921; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1938; Bartram, 1952; Seki, 1974).	
 <i>B. fuscinervia</i> (Mitt.) Jaeg. (Jaeger, 1873).	
Syn. <i>Tortula fuscinervia</i> Mitt. (Mitten, 1869).	
Distr. Southern South America (Mitten, 1869; Jaeger, 1873; Paris, 1904a; Brotherus, 1924a, c, Skottsberg, 1950; Bartram, 1952; Herzog, 1954, 1960; Seki, 1974).	
 <i>B. fusco-viridis</i> Broth. ex Thér. (Brotherus ex Thériot, 1921a).	
Distr. Southern South America (Paris, 1906; Campo, 1921; Thériot, 1921a).	
 <i>B. fuscula</i> C. Muell.	<i>Erythrophyllopsis fuscula</i>
 <i>B. galeata</i> C. Muell.	<i>Aloinella galeata</i>
 <i>B. geniculata</i> (Mont.) C. Muell. (Mueller, 1849).	
Syn. <i>Tortula geniculata</i> Mont. (Montagne, 1845a). <i>Asteriscium geniculatum</i> (Mont.) Herz. (Herzog, 1938).	
Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Mitten, 1869; Jaeger, 1873; Paris, 1904a; Brotherus, 1924a; Herzog, 1938, 1954).	
 <i>B. glacialis</i> Kunz. ex C. Muell.	<i>Tortula glacialis</i>
 <i>B. glaucula</i> C. Muell. (Mueller, 1882).	
Distr. Southern South America (Mueller, 1882).	
 <i>B. graminicolor</i> C. Muell. (Mueller, 1849).	
Syn. <i>Tortula graminicolor</i> (C. Muell.) Mont. (Montagne, 1850). <i>Asteriscium graminicolor</i> (C. Muell.) Hilp. (Hilpert, 1933).	
Distr. Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Paris, 1904a; Thériot, 1921b; Brotherus, 1924a).	
 ssp. <i>subgraminicolor</i> Thér. (Thériot, 1917b).	
Syn. <i>Barbula subgraminicolor</i> Thér. (Thériot, 1917b) nom. inval.	
Distr. Southern South America (Thériot, 1917b; Costes, 1921).	
 <i>B. grimiacea</i> C. Muell. (Mueller, 1882).	
Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Hilpert, 1933; Kühnemann, 1938).	
 <i>B. hosseusii</i> Thér. (Thériot, 1935b).	
Distr. Southern South America (Bauer, 1934; Thériot, 1935b; Hosseus, 1936, 1937, 1938a, b, c, d, 1939; Kühnemann, 1938).	

<i>B. humilis</i> Hedw.	<i>Tortella humilis</i>
<i>B. hyperborea</i> Mont. ex Par.	<i>Tortula mucronifolia</i> var. <i>arctica</i>
<i>B. imperfecta</i> (C. Muell.) Broth. (Brotherus, 1902).	
<i>Syn.</i> <i>Trichostomum imperfectum</i> C. Muell. (Mueller, 1879a).	
<i>Anacalypta imperfecta</i> (C. Muell.) Kindb. (Kindberg, 1889).	
<i>Pottia imperfecta</i> (C. Muell.) Par. (Paris, 1898).	
<i>Distr.</i> Southern South America (Mueller, 1879a; Paris, 1898; Brotherus, 1902, 1924a; Warnstorf, 1916; Kühnemann, 1938).	
<i>B. inundata</i> (Mitt.) Jaeg.	<i>Didymodon inundatus</i>
<i>B. juniperoides</i> C. Muell. (Mueller, 1882).	
<i>Distr.</i> Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).	
<i>B. kunzeana</i> C. Muell.	<i>Tortula kunzeana</i>
<i>B. kunzei</i> C. Muell. ex Kindb.	<i>Tortula kunzeana</i>
<i>B. laevigata</i>* (Mitt.) Jaeg. (Jaeger, 1873).	
<i>Syn.</i> <i>Tortula laevigata</i> Mitt. (Mitten, 1869).	
<i>Distr.</i> Southern South America (Dusén, 1907; Cardot, 1908a; Seki, 1974).	
<i>B. laevipila</i> (Brid.) Garov.	<i>Tortula laevipila</i>
<i>B. lamprocalyx</i> C. Muell. (Mueller, 1849).	
<i>Syn.</i> <i>Tortula lamprocalyx</i> (C. Muell.) Mont. (Montagne, 1850).	
<i>Distr.</i> Southern South America (Mueller, 1849; Montagne, 1850; Jaeger, 1873; Paris, 1904a; Brotherus, 1924a).	
<i>B. lechleri</i> C. Muell.	<i>Tortula lechleri</i>
<i>B. lepto-syntrichia</i> C. Muell.	<i>Tortula lepto-syntrichia</i>
<i>B. leucocalyx</i> (Mont.) C. Muell.	<i>Pseudocrossidium leucocalyx</i>
<i>B. lonchodonta</i> C. Muell. (Mueller, 1879a).	
<i>Distr.</i> Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935a, 1937; Kühnemann, 1938).	
<i>B. lorentzii</i> C. Muell.	<i>Tortula lorentzii</i>
<i>B. lurida</i> Hornsch. (Hornschuch, 1840).	
<i>Distr.</i> Southern South America (Hosseus, 1938c).	
<i>B. magellanica</i> (Mont.) C. Muell.	<i>Tortula magellanica</i>
<i>B. magellanica</i> C. Muell.	<i>Tortula fuegiana</i>
<i>B. marginato-serrata</i> Dus.	<i>Tortula marginato-serrata</i>
<i>B. mendozensis</i> (Mitt.) Jaeg. (Jaeger, 1873).	
<i>Syn.</i> <i>Tortula mendozensis</i> Mitt. (Mitten, 1869).	
<i>Distr.</i> Southern South America (Mitten, 1869; Jaeger, 1873; Paris, 1904a; Brotherus, 1924a).	
<i>B. micro-runcinata</i> Dus.	<i>Leptodontium longicaule</i> var. <i>microruncinatum</i>
<i>B. minutirosetula</i> C. Muell.	<i>Tortula minutirosetula</i>
<i>B. mniadelphus</i> C. Muell.	<i>Tortula mniadelphus</i>
<i>B. mnioides</i> Schwaegr.	<i>Calyptopogon mnioides</i>

* = *Didymodon laevigatus* (Mitt.) Zander *fide* Zander (1978).

<i>var. prostrata</i> (Mont.) C. Muell.	<i>Tortula prostrata</i>
<i>B. muelleri</i> B.S.G.	<i>Tortula princeps</i>
<i>B. muelleri</i> Kindb.	<i>Barbula perrevoluta</i>
<i>B. muralis</i> (Hedw.) Crom.	<i>Tortula muralis</i>
ssp. <i>aestiva</i> (Hedw.) Kindb.	<i>Tortula muralis</i> var. <i>aestiva</i>
var. <i>aestiva</i> (Hedw.) Roehl.	<i>Tortula muralis</i> var. <i>aestiva</i>
var. <i>australis</i> Hamp.	<i>Tortula muralis</i>
var. <i>rupestris</i> Schultz	<i>Tortula muralis</i>
<i>B. muricola</i> C. Muell.	<i>Tortula muralis</i>
<i>B. nanophylla</i> Dus. (Dusén, 1903a) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Dusén, 1903a).	
<i>B. oliviensis</i> Card.	<i>Tortula pilifera</i> var. <i>oliviensis</i>
<i>B. ovata</i> (Hedw.) Schimp.	<i>Pterygoneurum ovatum</i>
<i>B. pachyneura</i> Dus.	<i>Pseudocrossidium leucocalyx</i>
<i>B. patagonica</i> (Mitt.) Jaeg.	<i>Didymodon patagonicus</i>
<i>B. patagonica</i> C. Muell.	<i>Tortula fuegiana</i>
<i>B. perangusta</i> Dus. (Dusén, 1903a) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Dusén, 1903a).	
<i>B. percarnosa</i> C. Muell.	<i>Tortula percarnosa</i>
<i>B. perfirma</i> Dus. (Dusén in Paris, 1906) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Paris, 1906; Rövainen, 1934).	
<i>B. perflaccida</i> Broth.	<i>Tortula flagellaris</i>
<i>B. pernana</i> C. Muell. (Mueller, 1879a).	
<i>Distr.</i> Southern South America (Mueller, 1879a; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).	
<i>B. perpusilla</i> C. Muell.	<i>Tortula perpusilla</i>
<i>B. perrevoluta</i> C. Muell. (Mueller, 1882).	
Syn. <i>Barbula subrevoluta</i> C. Muell. (Mueller, 1879a) <i>hom. illeg.</i>	
<i>Barbula muelleri</i> Kindb. (Kindberg, 1889) <i>hom. illeg.</i>	
<i>Barbula apoclada</i> Par. (Paris, 1900a) <i>nom. illeg.</i>	
<i>Distr.</i> Southern South America (Mueller, 1879a, 1882; Paris, 1900a, 1904a; Kurtz, 1904; Brotherus, 1924a; Hosseus, 1935a, b, 1937; Kühnemann, 1938).	
var. <i>acutifolia</i> (C. Muell.) Par. (Paris, 1894).	
Syn. <i>Barbula subrevoluta</i> C. Muell. var. <i>acutifolia</i> C. Muell. (Mueller, 1879a).	
<i>Distr.</i> Southern South America (Mueller, 1879a; Paris, 1894, 1904a; Hosseus, 1935b, 1937; Kühnemann, 1938).	
var. <i>linearifolia</i> (C. Muell.) Par. (Paris, 1894).	
Syn. <i>Barbula subrevoluta</i> C. Muell. var. <i>linearifolia</i> C. Muell. (Mueller, 1879a).	
<i>Distr.</i> Southern South America (Mueller, 1879a; Paris, 1894, 1904a; Hosseus, 1935a, 1937; Kühnemann, 1938).	
<i>B. perrubiginosa</i> Dus.	<i>Tortula anderssonii</i>
<i>B. perrufula</i> C. Muell.	<i>Tortella perrufula</i>
<i>B. pertorquescens</i> Broth. (Brotherus, 1918).	
<i>Distr.</i> Southern South America (Brotherus, 1918, 1924a; Kühnemann, 1938).	
<i>B. pilifera</i> (Hook.) Brid.	<i>Tortula pilifera</i>

B. plebeja C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

<i>B. podocarpi</i> C. Muell.	<i>Tortula podocarpi</i>
<i>B. poeppigiana</i> C. Muell.	<i>Trichostomopsis australisiae</i>
<i>B. princeps</i> (De Not.) C. Muell.	<i>Tortula princeps</i>
<i>B. prostrata</i> (Mont.) Jaeg.	<i>Tortula prostrata</i>
<i>B. pseudo-caespitosa</i> C. Muell.	<i>Tortella pseudo-caespitosa</i>
var. <i>brachybasis</i> C. Muell.	<i>Tortella pseudo-caespitosa</i> var. <i>brachybasis</i>
var. <i>pungens</i> C. Muell.	<i>Tortella pseudo-caespitosa</i> var. <i>pungens</i>
<i>B. punctulata</i> (Brid.) Jaeg.	<i>Tortula robusta</i>

B. purpurascens Dus. (Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1938, 1960; Seki, 1974).

<i>B. pusilla</i> Jaeg.	<i>Tortula brachypelma</i>
-------------------------	----------------------------

B. pycnophylla Card. (Cardot, 1906a).

Distr. Southern South America (Dusén, 1906; Cardot, 1908a; Brotherus, 1924a; Bartram, 1952; Seki, 1974).

South Georgia (Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b).

<i>B. recurvipatula</i> C. Muell.	<i>Aloina recurvipatula</i>
-----------------------------------	-----------------------------

B. replicata Tayl. (Taylor, 1846).

Syn. *Tortula replicata* (Tayl.) Wils. (Wilson, 1846).

Distr. Southern South America (Bartram, 1942, 1943, 1952).

<i>B. rigidula</i> (Hedw.) Mitt.	<i>Didymodon rigidulus</i>
----------------------------------	----------------------------

B. riparia C. Muell. (Mueller, 1882).

Syn. *Asteriscium riparium* (C. Muell.) Hilp. (Hilpert, 1933).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Hosseus, 1938a, d; Kühnemann, 1938).

<i>B. ripicola</i> Card.	<i>Hymenostylium recurvirostre</i>
<i>B. robusta</i> (Hook. et Grev.) Brid.	<i>Tortula robusta</i>
<i>B. rubiginosa</i> C. Muell.	<i>Didymodon argentinicus</i>
<i>B. rubra</i> (Mitt.) Jaeg.	<i>Tortula robusta</i>
<i>B. rubricaulis</i> Kindb. (Kindberg, 1891) <i>nom. nud.</i>	

Distr. Southern South America (Mueller, 1901a).

B. rubriseta Bartr. (Bartram, 1936).

Distr. Southern South America (Bartram, 1936; Hosseus, 1938c, d, 1939).

<i>B. runcinata</i> C. Muell.	<i>Tortula robusta</i> var. <i>runcinata</i>
<i>B. ruralis</i> Hedw.	<i>Tortula ruralis</i>

B. santessonii Bartr. (Bartram, 1952).

Distr. Southern South America (Bartram, 1952; Seki, 1974).

B. santiagensis Broth. (Brotherus *in* Dusén, 1907).

Distr. Southern South America (Dusén, 1907; Brotherus, 1924a; Thériot, 1934b; Herzog, 1954).

var. <i>piligera</i> Broth. (<i>Brotherus in Dusén</i> , 1907).	
<i>Distr.</i> Southern South America (<i>Dusén</i> , 1907).	
B. scabrella Dus.	<i>Tortula scabrella</i>
B. scabrinervis C. Muell.	<i>Tortula scabrinervis</i>
B. schimperi (Mont.) Mitt.	<i>Didymodon schimperi</i>
B. schnyderi C. Muell.	<i>Tortula schnyderi</i>
B. sedifolia C. Muell.	<i>Aloina sedifolia</i>
B. serripungens Lor. et C. Muell.	<i>Tortula serripungens</i>
var. <i>exesa</i> C. Muell.	<i>Tortula serripungens</i> var. <i>exesa</i>
B. serrulata (Hook. et Grev.) Brid.	<i>Tortula robusta</i>
B. speciosa Hook. f. et Wils.	<i>Tortula robusta</i>
B. spiralis Schimp. (<i>Schimper in Mueller</i> , 1849).	
<i>Distr.</i> Southern South America (<i>Hosseus</i> , 1938a, d).	
B. streptostega C. Muell. (<i>Mueller</i> , 1901a [1900]) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (<i>Mueller</i> , 1901a).	
B. subdecidua Kindb.	<i>Barbula uruguensis</i>
B. subgraminicolor Thér.	<i>Barbula graminicolor</i> ssp. <i>subgraminicolor</i>
B. subgrimiacea Thér. (<i>Thériot in Felippone</i> , 1929 [1930]).	
<i>Distr.</i> Southern South America (<i>Felippone</i> , 1929; <i>Herter</i> , 1933a).	
B. subrevoluta C. Muell.	<i>Barbula perrevoluta</i>
var. <i>acutifolia</i> C. Muell.	<i>Barbula perrevoluta</i> var. <i>acutifolia</i>
var. <i>linearifolia</i> C. Muell.	<i>Barbula perrevoluta</i> var. <i>linearifolia</i>
B. subrubiginosa Kindb.	<i>Didymodon argentinicus</i>
B. subruncinata C. Muell. (<i>Mueller</i> , 1901a [1900]).	
<i>Distr.</i> Southern South America (<i>Mueller</i> , 1901a).	
B. tenella (Broth.) Par.	<i>Tortula tenella</i>
B. tessellata C. Muell. (<i>Mueller</i> , 1901a [1900]) <i>nom. nud.</i>	
<i>Distr.</i> Am. 6 in <i>Index muscorum</i> , 1, 156, but not found in <i>Mueller</i> (1901a).	
B. tortelloides C. Muell. (<i>Mueller</i> , 1879a).	
<i>Distr.</i> Southern South America (<i>Mueller</i> , 1879a; <i>Paris</i> , 1904a; <i>Brotherus</i> , 1924a; <i>Kühnemann</i> , 1938).	
B. tortuosa (Hedw.) Web. et Mohr	<i>Tortella tortuosa</i>
B. umbrosa C. Muell.	<i>Trichostomopsis umbrosa</i>
B. uncinicoma C. Muell. (<i>Mueller</i> , 1879a).	
<i>Distr.</i> Southern South America (<i>Mueller</i> , 1879a; <i>Kurtz</i> , 1904; <i>Paris</i> , 1904a; <i>Brotherus</i> , 1924a; <i>Williams</i> , 1930; <i>Hosseus</i> , 1935a, 1937; <i>Kühnemann</i> , 1938).	
B. unguiculata Hedw. (<i>Hedwig</i> , 1801).	
<i>Syn.</i> <i>Bryum unguiculatum</i> (Hedw.) With. (<i>Withering</i> , 1801).	
<i>Tortula unguiculata</i> (Hedw.) A. Roth ex P. Beauv. (A. Roth ex <i>Palisot de Beauvois</i> , 1805a) <i>hom. illeg.</i>	
<i>Streblotrichum unguiculatum</i> (Hedw.) Loesk. (<i>Loeske</i> , 1910).	
<i>Distr.</i> Southern South America (<i>Felippone</i> , 1917; <i>Herter</i> , 1933a).	
var. <i>patagonica</i> C. Muell. (<i>Mueller</i> , 1843a).	
<i>Syn.</i> <i>Syntrichia ericetorum</i> (With.) Brid. (<i>Bridel</i> , 1826).	

Distr. Southern South America (Bridel, 1826; Mueller, 1843a, 1849; Paris, 1904a).

B. unguiculatula C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904a; Brotherus, 1924a; Hosseus, 1938c, d; Kühnemann, 1938).

B. uruguensis Broth. (Brotherus, 1900).

Distr. Southern South America (Brotherus, 1900; Paris, 1904a; Felippone, 1912; Herter, 1933a; Hosseus, 1938d).

B. uruguensis Par. (Paris, 1894).

Syn. *Barbula decidua* C. Muell. (Mueller, 1882) *hom. illeg.*
Barbula subdecidua Kindb. (Kindberg, 1889) *nom. nud.*
Asteriscium deciduum Hilp. (Hilpert, 1933).

Distr. Southern South America (Mueller, 1882; Paris, 1894, 1904a; Felippone, 1912; Brotherus, 1924a; Herter, 1933a; Hosseus, 1938a, c; Herzog, 1938; Kühnemann, 1938).

B. vaginata Warnst. (Warnstorf, 1915).

Distr. Southern South America (Warnstorf, 1915).

B. vahliana Schultz

B. ventanae C. Muell. ex Par.

B. ventanae C. Muell.

B. ventanica C. Muell. (Mueller, 1897a).

Syn. *Barbula ventanae* C. Muell. ex Par. (Mueller ex Paris, 1900a) *err.*

Distr. Southern South America (Mueller, 1897a; Paris, 1900a; Kühnemann, 1938).

B. vinealis* Brid. var. *flaccida* B.S.G. (Bruch and others, 1842).

Syn. *Tortula vinealis* (Brid.) Spruc. var. *flaccida* (B.S.G.) Spruc. (Spruce, 1845).

Distr. Southern South America (Bruch and others, 1842; Montagne, 1850).

Bartramia

B. abbreviata Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

B. acicularis C. Muell.

Philonotis acicularis

B. acutissima C. Muell.

Philonotis acutissima

B. ambigua Mont. (Montagne, 1845a).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Mitten, 1869; Jaeger, 1875; Paris, 1904a; Thériot, 1915, 1917b, 1921b; Campo, 1921; Costes, 1921; Cardot and Brotherus, 1923; Brotherus, 1924a, c; Reimers, 1926; Roivainen, 1934; Herzog, 1938, 1939, 1940, 1954, 1957, 1960; Kühnemann, 1938; Espinosa B., 1941; Bartram, 1952; Seki, 1974).

var. horrida Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

var. penicillatula (C. Muell.) Herz. (Herzog, 1920).

Syn. *Bartramia penicillatula* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. aristata Schimp. ex C. Muell. (Schimper ex Mueller, 1862).

Syn. *Bartramia lechleri* Schimp. (Schimper in Mueller, 1901a [1900]) *nom. illeg.*

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Robinson, 1975).

Southern South America (Lechler, 1857; Mueller, 1862; Lorentz, 1866; Jaeger, 1874; Mueller, 1901a; Dusén, 1903a; Paris, 1904a; Cardot, 1905a, 1908a; Thériot, 1918, 1934b; Campo, 1921; Brotherus, 1924a; Reimers, 1926; Herzog, 1938, 1957; Kühnemann, 1938).

B. aureola Besch. ex C. Muell.

Breutelia aureola

B. australis Mitt.

Conostomum pentastichum

B. austro-arcuata C. Muell.

Breutelia austro-arcuata

B. austro-georgica Par.

Bartramia patens

B. austro-stricta C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

B. carinata Mitt.

Breutelia plicata

B. chilensis Lor.

Breutelia dumosa

B. chrysura C. Muell.

Breutelia integrifolia

B. comiramea C. Muell.

see **Breutelia dumosa**

B. comosa Mitt.

Philonotis crenatula

B. crenatula C. Muell.

Bartramia pomiformis

B. crispa Brid. var. *pomiformis* (Hedw.) Lindb.

Philonotis curvata

B. curvata Hamp.

Breutelia cuspidatissima

B. cuspidatissima C. Muell.

Bartramidula cygnea

B. cygnea Mont.

Philonotis pomangium

B. declivium C. Muell.

Philonotis glaucescens

B. defecta C. Muell.

B. delicatula C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

B. diminutiva C. Muell. (Mueller, 1884 [1883]).

Distr. South Georgia (Cardot, 1906a, 1908a; Dixon, 1920; Steere, 1961b).

Antarctic, peninsula region (Cardot, 1907c, 1908a, 1911b, d, 1913a; Steere, 1961a).

B. dimorpha Schimp.

Philonotis vagans

B. dumosa (Mitt.) Mitt.

Breutelia dumosa

B. elegantula Tayl.

Philonotis rufiflora

B. elegantula Par.

Philonotis elongatula

B. elongatula C. Muell.

Philonotis elongatula

var. *tenella* C. Muell. (Mueller, 1882) *nom. nud.*

Distr. Southern South America (Mueller, 1882).

B. exigua Sull.

Philonotis scabrifolia

B. fernandeziana Card. (Cardot in Thériot, 1921b).

Distr. Juan Fernandez (Thériot, 1921b; Brotherus, 1924a, b).

B. flaviuscula C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

B. flexipes (C. Muell.) C. Muell.

Philonotis flexipes

B. fontana (Hedw.) Turn.

Philonotis fontana

var. *magellanica* (Brid.) P. Beauv.

Philonotis fontana var. magellanica

B. fontanoides Gill. et Grev.

Philonotis fontanoides

* = *Didymodon vinealis* (Brid.) Zander var. *flaccida* (B.S.G.) Zander *fide* Zander (1978).

- B. fragilifolia** C. Muell. (Mueller, 1897b).
Distr. Southern South America (Bartram, 1965).
- B. glaucescens** Hornsch.
B. graminicola C. Muell.
- B. halleriana** Hedw. (Hedwig, 1801).
Distr. Southern South America (Mitten, 1869; Bartram, 1952; Seki, 1974).
- var. brachydonta** Kab. (Kabiersch, 1937).
Syn. *Bartramia mossmaniana* C. Muell. (Mueller, 1851b).
Bartramia magellanica Aongstr. (Ångström, 1872) *hom. illeg.*
Bartramia mossmanii C. Muell. ex Kindb. (Mueller ex Kindberg, 1888) *nom. illeg.*
Distr. Southern South America (Ångström, 1872; Jaeger, 1875; Mueller, 1885; Neger, 1899; Dusén, 1903a, 1905d; Paris, 1904a; Cardot, 1905a, 1908a, 1913a; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1940, 1957; Brotherus 1924a; Roivainen, 1934; Thériot, 1935a; Kabiersch, 1937; Kühnemann, 1938).
- B. hariotiana** Besch. ex C. Muell.
B. integrifolia Tayl.
- B. ithyphylla** Brid. (Bridel, 1803).
Syn. *Plagiopus ithyphylloides* (Brid.) Mach. (Machado, 1919).
Distr. Southern South America (Montagne, 1850; Campo, 1921 (as *Bartramia ithyla*)).
var. arenae Besch. (Bescherelle, 1889).
Distr. Southern South America (Bescherelle, 1889; Dusén, 1903a; Paris, 1904a; Cardot, 1908a; Kühnemann, 1938).
- B. ithyphylloides** Schimp. ex C. Muell. (Schimper ex Mueller, 1859).
Distr. Southern South America (Lechler, 1857; Mueller, 1859; Jaeger, 1875; Neger, 1899; Paris, 1904a; Brotherus, 1924a; Roivainen, 1934; Thériot, 1934b; Herzog, 1939, 1954; Seki, 1974).
- B. jacquenotiana** Schimp. (Schimper in Paris, 1904a) *nom. nud.*
Distr. Southern South America (Paris, 1904a).
- B. joannae** C. Muell. (Mueller, 1901a [1900]) *nom. nud.*
Distr. Southern South America (Mueller, 1901a).
- B. krausei** C. Muell.
B. laevisphaera (Tayl.) C. Muell.
- B. lechleri** Schimp.
B. leptophylla Card. et Broth. (Cardot and Brotherus, 1923).
Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a).
- B. leptophylla** Card. (Cardot in Skottsberg, 1914) *nom. nud. fide* (Brotherus, 1924b).
Distr. Juan Fernandez (Skottsberg, 1914).
Southern South America (Campo, 1921).
- B. leucocolea** Card.
var. brevifolia Broth. et Card.
- var. glaucoflava** Broth. et Card. (Brotherus and Cardot in Cardot and Brotherus, 1923).
Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).
- B. leucolomacea** C. Muell.
B. lorentzii C. Muell.
- B. magellanica** (Sull.) Mitt.
B. magellanica Aongstr.
- Philonotis glaucescens**
Breutelia integrifolia
- B. magellanica** Brid.
B. microbasis C. Muell. (Mueller, 1882).
Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938; Bizot, 1967).
- B. mossmaniana** C. Muell.
B. mossmanii C. Muell. ex Kindb.
- B. nevadensis** C. Muell. (Mueller, 1882).
Distr. Southern South America (Mueller, 1882; Brotherus, 1924a; Hosseus, 1938c, d, 1939; Kühnemann, 1938).
- B. oreadella** C. Muell.
var. microphylla Card.
- B. palustris** (Hedw.) P. Beauv.
- B. patens** Brid. (Bridel, 1803).
Syn. *Bartramia reticulata* P. Beauv. (Palisot de Beauvois, 1805a).
Bartramia magellanica Brid. (Bridel in Bescherelle, 1889) *nom. nud.*
Bartramia leucolomacea C. Muell. (Mueller, 1890a) *fide* Newton, 1974a.
Bartramia oreadella C. Muell. (Mueller, 1890a) *fide* Newton, 1974a.
Bartramia oreadella C. Muell. var. *microphylla* Card. (Cardot, 1905a) *fide* Newton, 1974a.
Bartramia pycnocolea C. Muell. (Mueller, 1890a) *fide* Newton, 1974a.
Bartramia subpatens C. Muell. (Mueller, 1890a) *hom. illeg.*
Bartramia austro-georgica Par. (Paris, 1894).
Bartramia leucocolea Card. (Cardot, 1905a) *fide* Newton, 1974a.
Bartramia leucocolea Card. var. *brevifolia* Broth. et Card. (Brotherus and Cardot in Cardot and Brotherus, 1923) *fide* Newton, 1974a.
- Distr. Juan Fernandez (Brotherus, 1924b; Robinson, 1975).
Southern South America (Bridel, 1803, 1827b; Palisot de Beauvois, 1805a; Schwaegrichen, 1816; Wilson and Hooker, 1847; Mueller, 1849, 1885, 1889; Montagne, 1850; Sullivant, 1859; Mitten, 1869; Jaeger, 1875; Bescherelle, 1889; Dusén, 1903a; Paris, 1904a; Cardot, 1905a, 1908a, 1913a; Cardot and Brotherus, 1923; Herzog, 1923, 1954, 1957, 1960; Brotherus, 1924a; Roivainen, 1934; Kühnemann, 1938; Bartram, 1952; Seki, 1974).
Falkland Islands (Wilson and Hooker, 1847; Jaeger, 1875; Mueller, 1885; Bescherelle, 1889; Dusén, 1903a; Paris, 1904a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a).
South Georgia (Mueller, 1890a; Paris, 1894, 1904a; Cardot, 1906a, 1908a; Dixon, 1920, 1934; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b; Newton, 1974a; Bell, 1977b).
Antarctic, peninsula region (Cardot, 1906a, 1908a, 1911b, d, 1913a; Bartram, 1957; Steere, 1961a; Greene, 1968a; Robinson, 1972).
- var. arenae** Card. (Cardot in Cardot and Brotherus, 1923) *nom. nud.*
Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).
Falkland Islands (Cardot and Brotherus, 1923).
- var. intermedia** Hook. f. et Wils. (Hooker and Wilson in Wilson and Hooker, 1847).
Distr. Falkland Islands (Wilson and Hooker, 1847; Kühnemann, 1938).
- var. minor** Dus. (Dusén, 1903a).
Distr. Southern South America (Dusén, 1903a; Kühnemann, 1938).
- B. pendula** (Sm.) Hook. see *Breutelia dumosa*
B. penicillatula C. Muell. *Bartramia ambigua* var. *penicillatula*
B. pentasticha Brid. *Conostomum pentastichum*
- B. perpumila** C. Muell. (Mueller, 1882).
Distr. Southern South America (Mueller, 1882; Paris, 1904a; Kühnemann, 1938; Brotherus, 1924a; Bartram, 1965).
- Conostomum magellanicum**
Bartramia halleriana var. **brachydonta**

<i>B. plicata</i> (Mitt.) Mitt.	<i>Breutelia plicata</i>
<i>B. pomangium</i> C. Muell.	<i>Philonotis pomangium</i>
<i>B. pomiformis</i> Hedw. (Hedwig, 1801).	
Syn. <i>Bryum pomiforme</i> (Hedw.) With. (Withering, 1801).	
<i>Bartramia crispa</i> Brid. var. <i>pomiformis</i> (Hedw.) Lindb. (Lindberg, 1872) nom. illeg.	
<i>Plagiopus pomiformis</i> (Hedw.) Mach. (Machado, 1919).	
Distr. Southern South America (Bridel, 1803, 1827b; Montagne, 1845b, 1850; Mitten, 1869; Mueller, 1885; Dusén, 1903a; Paris, 1904a; Cardot, 1908a).	
var. <i>crispa</i> (Brid.) B.S.G.	<i>Bartramia pomiformis</i> var. <i>elongata</i>
var. <i>elongata</i> Turn. (Turner, 1805).	
Syn. <i>Bartramia pomiformis</i> Hedw. var. <i>crispa</i> (Brid.) B.S.G. (Bruch and others, 1842). nom. illeg.	
Distr. Southern South America (Wilson and Hooker, 1847; Rowlee, 1916).	
<i>B. potosica</i> Mont. (Montagne, 1838).	
Distr. Southern South America (Mitten, 1869; Jaeger, 1875; Paris, 1904a; Brotherus, 1924a).	
<i>B. pseudo-cryptopodium</i> C. Muell. (Mueller, 1882).	
Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).	
<i>B. pusilla</i> Sull.	<i>Philonotis scabrifolia</i>
<i>B. pycnocolea</i> C. Muell.	<i>Bartramia patens</i>
<i>B. reticulata</i> P. Beauv.	<i>Bartramia patens</i>
<i>B. robusta</i> Hook. f. et Wils. (Hooker and Wilson in Wilson and Hooker, 1845).	
Syn. <i>Breutelia robusta</i> (Hook. f. et Wils.) Jaeg. (Jaeger, 1875).	
Distr. Southern South America (Sullivan, 1859; Jaeger, 1875; Mueller, 1885; Dusén, 1903a; Paris, 1904a; Cardot, 1908a; Brotherus, 1924a).	
<i>B. rufiflora</i> Hornsch.	<i>Philonotis rufiflora</i>
<i>B. rupestris</i> Mitt.	<i>Breutelia plicata</i>
<i>B. rupta</i> C. Muell. (Mueller, 1882).	
Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935b, 1937, 1939; Kühnemann, 1938).	
<i>B. scaberrima</i> Card. et Broth. (Cardot and Brotherus, 1923).	
Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).	
<i>B. scabrifolia</i> (Hook. f. et Wils.) C. Muell.	<i>Philonotis scabrifolia</i>
<i>B. simplicissima</i> C. Muell.	<i>Philonotis simplicissima</i>
<i>B. striata</i> C. Muell.	<i>Philonotis striata</i>
<i>B. stricta</i> Brid. (Bridel, 1803).	
Syn. <i>Plagiopus strictus</i> (Brid.) Mach. (Machado, 1919).	
Distr. Juan Fernandez (Montagne, 1835; Johow, 1896).	
Southern South America (Bridel, 1803, 1827b; Schwaegrichen, 1816; Montagne, 1850; Paris, 1904a).	
<i>B. strictiuscula</i> (C. Muell.) C. Muell.	<i>Philonotis strictiuscula</i>
<i>B. subpatens</i> C. Muell.	<i>Bartramia patens</i>
<i>B. subsecunda</i> C. Muell. (Mueller, 1901a [1900]) nom. nud.	
Distr. Southern South America (Mueller, 1901a).	

<i>B. subsessilis</i> Tayl.	<i>Anacolia laevisphaera</i>
<i>B. subsymmetrica</i> Card. (Cardot, 1906a).	
Syn. <i>Exodokidium subsymmetricum</i> (Card.) Card. (Cardot, 1908a) fide Newton, 1974a.	
Distr. South Georgia (Cardot, 1906a, 1908a; Brotherus, 1924a; Newton, 1974a; Bell, 1977b).	
<i>B. tenella</i> (Brid.) C. Muell.	<i>Philonotis glaucescens</i>
<i>B. thrausta</i> Schimp. ex Britt. (Schimper ex Britton, 1896).	
Distr. Southern South America (Paris, 1904a).	
<i>B. vagans</i> (Hook. f. et Wils.) Mitt.	<i>Philonotis vagans</i>
<i>B. ventanae</i> C. Muell.	<i>Philonotis scabrifolia</i>
<i>B. wilkesiana</i> Par. (Paris, 1894) nom. nud.	
Distr. Southern South America (Paris, 1894, 1904a).	
<i>B. willii</i> C. Muell.	<i>Philonotis scabrifolia</i>
Bartramidula	
<i>B. cygnea</i> (Mont.) Par. (Paris, 1894).	
Syn. <i>Bartramia cygnea</i> Mont. (Montagne, 1845a).	
<i>Glyphocarpus cygneus</i> (Mont.) Jaeg. (Jaeger, 1875).	
Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Lorentz, 1866; Mitten, 1869; Jaeger, 1875; Paris, 1894, 1904a; Herzog, 1923; Brotherus, 1924a; Seki, 1974).	
<i>B. defecta</i> (C. Muell.) Broth.	<i>Philonotis glaucescens</i>
<i>B. exigua</i> (Sull.) Par.	<i>Philonotis scabrifolia</i>
<i>B. krausei</i> (C. Muell.) Par.	<i>Philonotis krausei</i>
Blindia	
<i>B. acuta</i> (Hedw.) B.S.G. (Bruch and others, 1846).	
Syn. <i>Weisia acuta</i> Hedw. (Hedwig, 1801).	
Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1848a; Paris, 1904a).	
Falkland Islands (Wilson and Hooker, 1847; Kühnemann, 1938).	
<i>B. antarctica</i> C. Muell.	<i>Dicranoweisia antarctica</i>
<i>B. arcuata</i> Mitt.	<i>Blindia magellanica</i>
<i>B. auriculata</i> C. Muell.	<i>Holodontium inerme</i>
<i>B. austro-crispula</i> C. Muell.	<i>Dicranoweisia austro-crispula</i>
<i>B. brevipes</i> C. Muell.	<i>Dicranoweisia brevipes</i>
<i>B. capillifolia</i> Card. (Cardot, 1906a).	
Distr. South Georgia (Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b).	
<i>B. churuccana</i> Besch.	<i>Holodontium pumilum</i>
<i>B. consimilis</i> Card.	<i>Blindia magellanica</i>
<i>B. coniecta</i> (Hook. f. et Wils.) C. Muell. (Mueller, 1848a).	
Syn. <i>Weisia coniecta</i> Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).	
<i>Dicranoweisia coniecta</i> (Hook. f. et Wils.) Par. (Paris, 1896).	
Distr. Southern South America (Ångström, 1872; Paris, 1896; Dusén, 1903a; Paris, 1904a; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).	
<i>B. curviseta</i> Mitt. (Mitten, 1869).	
Syn. <i>Dicranum gracilescens</i> Dus. (Dusén in Paris, 1904b) nom. nud.	
<i>Blindia praticola</i> Card. et Broth. (Cardot and Brotherus, 1923).	

Distr. Southern South America (Mitten, 1869; Mueller, 1885; Paris, 1900a, 1904a, b; Dusén, 1903a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen, 1934; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1939).
Falkland Islands (Dusén, 1903a; Paris, 1904a; Cardot, 1908a).
South Georgia (Cardot and Brotherus, 1923).

B. dicranellacea C. Muell. (Mueller, 1890a).

Distr. South Georgia (Mueller, 1890a; Paris, 1904a; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b).

B. globularis Dus.

Blindia magellanica

B. grimmiae C. Muell.

Dicranoweisia grimmiae

B. humilis C. Muell.

Holodontium pumilum

B. inundata (Card.) Card.

Blindia magellanica var. *inundata*

B. leptotrichacea C. Muell. ex. Par.

Ditrichum hookeri

B. leptotrichocarpa C. Muell.

Ditrichum hookeri

var. *strictiuscula* C. Muell.

Ditrichum hookeri var. *strictiusculum*

B. lygodipoda C. Muell. (Mueller, 1885).

Syn. *Blindia lygopodina* C. Muell. (Mueller in Paris, 1894) nom. illeg.

Distr. Southern South America (Mueller, 1885; Paris, 1894, 1904a; Dusén, 1903a; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

B. lygopodina C. Muell.

Blindia lygodipoda

B. magellanica Schimp. (Schimper in Mueller, 1862).

Syn. *Blindia arcuata* Mitt. (Mitten, 1869).

Blindia consimilis Card. (Cardot, 1905a).

Blindia globularis Dus. (Dusén, 1905a).

Blindia microdicranum Dus. (Dusén in Roivainen and Bartram, 1937) nom. nud.

Distr. Southern South America (Mueller, 1862, 1885; Mitten, 1869; Jaeger, 1873; Dusén, 1903a, 1905a; Paris, 1904a, 1906; Cardot, 1908a; Campo, 1921; Thériot, 1921a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Herzog, 1938, 1939; Kühnemann, 1938; Seki, 1974).

Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938).

var. *inundata* (Card.) Herz. (Herzog, 1939).

Syn. *Ditrichum inundatum* Card. (Cardot, 1905a).

Blindia inundata (Card.) Card. (Cardot, 1908a).

Distr. Southern South America (Cardot, 1905a, 1908a; Brotherus, 1924a; Roivainen and Bartram, 1937; Herzog, 1939, 1940, 1960).

B. microdicranum Dus.

Blindia magellanica

B. pallidifolia C. Muell.

Dicranoweisia subinclinata

B. praticola Card. et Broth.

Blindia curviseta

B. pseudolygodipoda Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).
Falkland Islands (Cardot and Brotherus, 1923).

B. pseudorobusta Dus.

Blindia robusta

B. robusta Hamp. (Hampe, 1860).

Syn. *Dicranum tenuifolium* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844) nom. illeg.
Blindia tenuifolia Mitt. (Mitten, 1869).
Blindia pseudorobusta Dus. (Dusén, 1905a).

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Sullivant, 1859; Mitten, 1969; Jaeger, 1872, 1873; Dusén, 1903a, 1905a; Paris, 1904a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Herzog, 1938, 1954; Kühnemann, 1938; Seki, 1974).

B. savatieri C. Muell.

Holodontium sp.

B. savatieriana C. Muell.

Holodontium sp.

B. skottsbergii Card. (Cardot, 1906a).

Distr. South Georgia (Cardot, 1906a, 1908a; Dixon, 1920; Brotherus, 1924a; Steere, 1961b). Antarctic, peninsula region (Wright, 1905 (as *Campylopus vesticaulis*); Cardot, 1906a, 1908a; 1911c, 1912b; Brotherus, 1924a; Steere, 1961a; Greene, 1968a; Robinson, 1972).

B. subinclinata C. Muell.

Dicranoweisia subinclinata

B. sulphurea Dus.

Dicranoweisia austro-crispula

B. tenuifolia Mitt.

Blindia robusta

B. torrentium Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. turpis Card.

Verrucidens turpis

Brachymenium

B. ellipticum Jaeg.

Brachymenium meyenianum var. *ellipticum*

B. erubescens (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Bryum erubescens* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. fabroniellum Par. (Paris, 1906) nom. nud.

Syn. *Dicranobryum fabroniellum* Dus. (Dusén in Paris, 1906) nom. nud.

Distr. Southern South America (Paris, 1906).

B. fabronioides (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Bryum fabronioides* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935b; Kühnemann, 1938).

var. *densirete* Thér. (Thériot, 1935b).

Distr. Southern South America (Thériot, 1935b; Hosseus, 1938c, d).

B. gilliesii (Hook.) Jaeg.

Bryum gilliesii

B. hyalinum Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933a).

B. indicum (Doz. et Molk.) Bosch et Lac. (van den Bosch and van der Sande Lacoste, 1860).

Syn. *Bryum indicum* Doz. et Molk. (Dozy and Molkenboer, 1845).

Peromnion magellanicum Sull. (Sullivant, 1850).

Bryum magellanicum (Sull.) C. Muell. (Mueller, 1851a).

Brachymenium magellanicum (Sull.) Par. (Paris, 1894).

Distr. Southern South America (Sullivant, 1850, 1859; Mueller, 1851a, 1885; Mitten, 1869; Jaeger, 1875; Paris, 1894, 1904a; Dusén, 1903a; Spegazzini, 1922; Kühnemann, 1938).

B. macropoma (C. Muell.) Kindb.

Acidodontium macropoma

B. magellanicum (Sull.) Par.

Brachymenium indicum

- B. malachiticum** (C. Muell.) Kindb. (Kindberg, 1889).
 Syn. *Bryum malachiticum* C. Muell. (Mueller, 1882).
 Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).
- B. meyenianum** (C. Muell.) Jaeg. (Jaeger, 1875).
 Syn. *Webera meyeniana* Hamp. (Hampe, 1837c).
Bryum meyenianum (Hamp.) C. Muell. (Mueller, 1848a).
Brachymenium meyenii Hamp. ex Kindb. (Hampe ex Kindberg, 1888) nom. illeg.
Dicranobryum subfabroniellum Dus. (Dusén in Paris, 1906) nom. nud.
Brachymenium subfabroniellum Par. (Paris, 1906) nom. nud.
 Distr. Southern South America (Hampe, 1837c; Mueller, 1848a; Montagne, 1850; Mitten, 1869; Jaeger, 1875; Paris, 1904a, 1906; Brotherus, 1924a; Thériot, 1928; Hosseus, 1936, 1937; Kühnemann, 1938; Herzog, 1939, 1954).
- var. *ellipticum* (Jaeg.) Thér. (Thériot, 1928).
 Syn. *Bryum ellipticum* Mitt. (Mitten, 1869) hom. illeg.
Brachymenium ellipticum Jaeg. (Jaeger, 1875).
 Distr. Southern South America (Mitten, 1869; Jaeger, 1875; Paris, 1904a; Costes, 1921; Thériot, 1921b, 1928; Brotherus, 1924a).
- B. meyenii Hamp. ex Kindb.
 B. ovatum Hook. f. et Wils.
 B. papillosum Jaeg.
 B. porteri Thér. (Thériot, 1925a).
 Distr. Southern South America (Thériot, 1925a, 1926).
- B. robertii** Broth. (Brotherus, 1918).
 Distr. Southern South America (Brotherus, 1918, 1924a; Kühnemann, 1938).
- B. subfabroniellum Par.
 B. wilsonii (Mitt.) Jaeg.
 Brachypodium hom. illeg.
 B. crispatum (Hedw.) Brid.
- Ptychomitrium**
- Ptychomitrium crispatum**
Ptychomitrium balansae
Ptychomitrium chimborazense var. *obtusifolium*
Ptychomitrium crispatum
Ptychomitrium fernandesianum
Ptychomitrium deltorii
Ptychomitrium emersum
Ptychomitrium ligulatum
Ptychomitrium sellowianum
Ptychomitrium sellowianum
Ptychomitrium balansae
Ptychomitrium vernicosum
- Brachythecium**
 B. alari-decurrans Dus.
- Brachythecium subplicatum**

- B. antarcticum Card.
 var. *cavifolium* Card.
- B. arenarium** Card. et Broth. (Cardot and Brotherus, 1923).
 Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).
- B. argyrocaulon Dus. (Dusén in Paris, 1906) nom. nud.
 Distr. Southern South America (Paris, 1906).
- B. austro-glareosum** (C. Muell.) Kindb. (Kindberg, 1891).
 Syn. *Hypnum austro-glareosum* C. Muell. (Mueller, 1884 [1883]).
Brachythecium austro-glareosum (C. Muell.) Kindb. var. *diffusum* Card. (Cardot, 1911b) fide Newton, 1979a.
 Distr. South Georgia (Newton, 1979a, b).
 Antarctic, peninsula region (Cardot, 1906b, 1907c, 1908a, 1911b, d, 1913a; Brotherus, 1924a; Steere, 1961a; Robinson, 1972; Newton, 1979a).
 var. *diffusum* Card. **Brachythecium austro-glareosum**
- B. austro-salebrosum** (C. Muell.) Kindb. (Kindberg, 1889).
 Syn. *Hypnum austro-salebrosum* C. Muell. (Mueller, 1884 [1883]).
Hypnum georgico-glareosum C. Muell. (Mueller, 1890a) fide Newton, 1979a.
Brachythecium georgico-glareosum (C. Muell.) Kindb. (Kindberg, 1891) fide Newton, 1979a.
Brachythecium antarcticum Card. (Cardot, 1900) fide Newton, 1979a.
Brachythecium antarcticum Card. var. *cavifolium* Card. (Cardot, 1900) fide Newton, 1979a.
Brachythecium skottsbergii Card. (Cardot, 1906a) fide Newton, 1979a.
 Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Bartram, 1952; Seki, 1974).
 Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).
 South Georgia (Mueller, 1890a; Paris, 1894, 1904a; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Dixon, 1932; Steere, 1961b; Newton 1979a, b).
 Antarctic, peninsula region (Cardot, 1900, 1901, 1906a, b, 1907c, 1908a, 1911b, d, 1913a; Paris, 1904a; Dixon, 1920, 1935; Brotherus, 1925; Bartram, 1957; Steere, 1961a; Robinson, 1972; Newton, 1979a).
- B. calerae** (C. Muell.) Kindb. (Kindberg, 1888).
 Syn. *Hypnum calerae* C. Muell. (Mueller, 1879a).
 Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904a; Kurtz, 1904; Brotherus, 1925; Hosseus, 1935b, 1937; Kühnemann, 1938).
- B. clinocarpum (Tayl.) Jaeg. **Oxyrrhynchium clinocarpum**
 B. confluens (C. Muell.) Kindb. **Sciaromium confluens**
- B. conostomum** (Tayl.) Jaeg. (Jaeger, 1878).
 Syn. *Hypnum conostomum* Tayl. (Taylor, 1848a).
Hypnum taylorii Mitt. (Mitten, 1869) nom. illeg.
 Distr. Southern South America (Reimers, 1926; Herzog, 1938).
- B. cuspidarioides Dus. **Brachythecium subplicatum**
- B. fasciculato-caudatum** C. Muell. (Mueller, 1897a).
 Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1925; Kühnemann, 1938).
- B. filirameum** C. Muell. (Mueller, 1897a).
 Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1925; Kühnemann, 1938).

B. filirepens Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b; Paris, 1906; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938; Seki, 1974).

B. flagellare (Hedw.) Jenn.

Brachythecium plumosum***B. fuegianum*** Broth. (Brotherus, 1908).

Syn. *Stereophyllum fuegianum* Besch. (Bescherelle, 1885a *syn. excl.*) *nom. nud.*
Stereophyllum fuegianum Besch. (Bescherelle, 1889) *hom. illeg.*
Brachythecium stereophyllum C. Muell. (Mueller *in* Bescherelle, 1889) *nom. nud.*

Distr. Southern South America (Bescherelle, 1885a, 1889; Brotherus, 1908, 1925).

B. georgico-glareosum (C. Muell.) Kindb.

Brachythecium austro-salebrosum***B. glaciale*** B.S.G. (Bruch and others, 1853).

Distr. Southern South America (Newton, 1979a, b).
South Georgia (Newton, 1979a, b).

B. grandirete C. Muell. (Mueller, 1897b).

Syn. *Brachythecium pumilum* Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b; Brotherus, 1925).

B. lepidopilooides (C. Muell.) Kindb.

Stereophyllum lepidopilooides

B. longidecurrens Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

B. longidens (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum longidens* C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Dusén, 1903a; Paris, 1904a, 1905b; Cardot, 1908a; Kühnemann, 1938; Bartram, 1946).

B. macroconum (C. Muell.) Kindb.

Brachythecium plumosum

var. ***robustius*** (C. Muell.) Par. (Paris, 1894).

Syn. *Hypnum macroconum* C. Muell. var. *robustius* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904a).

B. macrogynum Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Brotherus, 1925; Kühnemann, 1938; Seki, 1974).

B. majusculum M. Newton (Newton, 1974d).

Syn. *Brachythecium majusculum* Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Newton, 1974d, 1979a, b).
South Georgia (Newton, 1974d, 1979a, b).

B. majusculum Dus.

Brachythecium majusculum

B. microcollinum Bartr. (Bartram, 1946).

Distr. Southern South America (Rovainen, 1934; Bartram, 1946).

B. minusculifolium C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1925; Kühnemann, 1938).

B. mollirameum C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Kühnemann, 1938).

B. morenoi C. Muell.

Brachythecium sericeo-virens***B. nematogonium*** (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum nematogonium* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904a; Brotherus, 1925; Hosseus, 1935b, 1937; Kühnemann, 1938).

B. paradoxum (Hook. f. et Wils.) Jaeg. (Jaeger, 1878).

Syn. *Hypnum paradoxum* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Rhynchostegium mollissimum Dus. (Dusén 1903a) *nom. nud.*

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1851a, 1885; Mitten, 1869; Jaeger, 1878; Bescherelle, 1889; Cardot, 1901, 1905a, 1908a; Dusén, 1903a; Paris, 1904a, 1905b; Cardot and Brotherus, 1923; Brotherus, 1925; Rovainen, 1934; Kühnemann, 1938; Herzog, 1954; Seki, 1974).

B. parvulum Dus. (Dusén *in* Paris, 1904a) *nom. nud.*

Distr. Southern South America (Paris, 1904a).

B. patagonicum Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

B. plumosum (Hedw.) B.S.G. (Bruch and others, 1853).

Syn. *Hypnum plumosum* Hedw. (Hedwig, 1801).
Hypnum macroconum C. Muell. (Mueller, 1879a).
Brachythecium macroconum (C. Muell.) Kindb. (Kindberg, 1888).
Hypnum flagellare Hedw. (Hedwig, 1801).
Brachythecium flagellare (Hedw.) Jenn. (Jennings, 1913) *nom. illeg.*

Distr. Southern South America (Mueller, 1879a; Paris, 1904a; Brotherus, 1925; Rovainen, 1934; Kühnemann, 1938).

B. pumilum Dus.***Brachythecium grandirete******B. rivulare*** B.S.G. (Bruch and others, 1853).

Distr. Southern South America (Thériot, 1934b).

B. rivularioides (C. Muell.) Par. (Paris, 1894).

Syn. *Hypnum rivularioides* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904a; Kurtz, 1904; Brotherus, 1925; Hosseus, 1935a, 1937; Kühnemann, 1938).

B. rutabulum (Hedw.) B.S.G. (Bruch and others, 1853).

Syn. *Hypnum rutabulum* Hedw. (Hedwig, 1801).
Pancovia rutabula (Hedw.) Kickz (Kickz, 1867).

Distr. Southern South America (Wilson and Hooker, 1847; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938).

B. sericeo-virens (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum sericeo-virens* C. Muell. (Mueller, 1885).
Brachythecium morenoi C. Muell. (Mueller, 1879a) *fide* Seki, 1974.

Distr. Southern South America (Mueller, 1885, 1897a; Paris, 1900a, 1904a; Dusén, 1903a; Cardot, 1905a, 1908a; Brotherus, 1925; Kühnemann, 1938; Seki, 1974).

B. seribracteatum Dix.***Brachythecium subpilosum******B. skottsbergii*** Card.***Brachythecium austro-salebrosum******B. spegazzinii*** (C. Muell.) Kindb.***Hypnum spegazzinii******B. spurio-albicans*** C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Dusén, 1903a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

B. squalidissimum (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum squalidissimum* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904a; Brotherus, 1925; Hosseus, 1935b, 1937; Kühnemann, 1938).

B. stereophyllum C. Muell.

Brachythecium fuegianum

B. subpilosum (Hook. f. et Wils.) Jaeg. (Jaeger, 1878).

Syn. *Hypnum subpilosum* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Brachythecium seribracteatum Dix. (Dixon in Newton, 1979a) *nom. nud.*

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1851a, 1885; Mitten, 1869; Jaeger, 1878; Paris, 1904a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Herzog, 1923; Brotherus, 1925; Kühnemann, 1938; Newton, 1979a).

Falkland Islands (Dusén, 1903a; Paris, 1904a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

South Georgia (Cardot, 1906a, 1908a; Dixon, 1920; Cardot and Brotherus, 1923; Brotherus, 1925; Steere, 1961b; Newton, 1979a, b).

Antarctic, peninsula region (Cardot, 1906a, 1908a; Brotherus, 1925; Steere, 1961a).

B. subplicatum (Hamp.) Jaeg. (Jaeger, 1878).

Syn. *Hypnum subplicatum* Hamp. (Hampe in Mueller, 1851a).

Hypnum rutabulum Hedw. var. 5 (in Wilson and Hooker, 1847) *fide* Mueller, 1851a.

Brachythecium cuspidarioides Dus. (Dusén, 1905b) *fide* Cardot, 1908a.

Brachythecium turgens Dus. (Dusén, 1905b) *fide* Seki, 1974.

Brachythecium alari-decurrens Dus. (Dusén in Newton, 1979a) *nom. nud.*

Distr. Southern South America (Mueller, 1851a, 1885; Dusén, 1903a, 1905b; Paris, 1906; Cardot, 1908a; Herzog, 1923, 1939, 1954; Brotherus, 1925; Thériot, 1929c; Rövainen, 1934; Kühnemann, 1938; Bartram, 1952; Seki, 1974; Newton, 1979a).

Falkland Islands (Wilson and Hooker, 1847; Mueller, 1851a; Jaeger, 1878; Cardot, 1905a, 1908a; Brotherus, 1925; Rövainen, 1934; Kühnemann, 1938; Newton, 1979a).

South Georgia (Cardot, 1906a, 1908a; Brotherus, 1925; Steere, 1961b; Newton, 1979a, b).

var. dilaceratum Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Kühnemann, 1938).

var. minus Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

B. subtilicaule Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1904a).

B. sulphureum (Geh. et Hamp.) Par. (Paris, 1894).

Syn. *Hypnum sulphureum* Geh. et Hamp. (Geheebe and Hampe in Hampe and Geheebe, 1881).

Distr. Southern South America (Paris, 1894; Brotherus, 1918, 1925; Hosseus, 1938a (as *Brachystegium*), b, c, d, 1939; Kühnemann, 1938).

B. tenui-prostratum C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1925; Kühnemann, 1938).

B. trachychaete Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b; Cardot, 1908a; Brotherus, 1925; Newton, 1979a).

B. turgens Dus.

Brachythecium subplicatum

B. turquetii Card. (Cardot, 1906b).

Distr. Antarctic, peninsula region (Cardot, 1906b, 1907c, 1908a; Brotherus, 1925; Steere, 1961a).

Braunia

B. argentinica C. Muell.

Braunia cirrhifolia

B. canescens Schimp. ex Britt.

Braunia cirrhifolia

B. chilensis Broth.

Hedwigidium chilense

B. cirrhifolia (Mitt.) Wils. ex Jaeg. (Wilson ex Jaeger, 1876).

Syn. *Harrisonia secunda* (Hook.) Spreng. var. *cirrhifolia* Mitt. (Mitten, 1851) *nom. nud.* *Hedwigia cirrhifolia* Mitt. (Mitten, 1869).

Braunia incana C. Muell. (Mueller, 1879a).

Braunia argentinica C. Muell. (Mueller, 1882).

Braunia canescens Schimp. ex Britt. (Schimper ex Britton, 1896).

Distr. Southern South America (Mueller, 1879a, 1882; Kurtz, 1904; Paris, 1904a; Brotherus, 1918, 1925; Hosseus, 1936, 1937, 1938c, d, 1939; Kühnemann, 1938).

B. cochlearifolia C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1925; Hosseus, 1936, 1937, 1938c, d, 1939; Kühnemann, 1938).

B. exserta C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904a; Brotherus, 1925; Kühnemann, 1938; Bartram, 1965).

var. *reflexifolia* C. Muell.

Braunia reflexifolia

B. gibertii Besch.

Hedwigidium gibertii

B. humboldtii (Hook.) Hook. f.

Rhacocarpus purpurascens

B. incana C. Muell.

Braunia cirrhifolia

B. patagonica C. Muell.

Hedwigidium patagonicum

B. reflexifolia (C. Muell.) Bartr. (Bartram, 1965).

Syn. *Braunia exserta* C. Muell. var. *reflexifolia* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904a; Bartram, 1965).

B. rhabdocarpa (Hamp.) C. Muell.

Hedwigidium integrifolia

Braunia Hamp. hom. illeg.

B. cuspidata (Sull.) Hamp.

Dendrocyphaea cuspidata

Breutelia

B. angustiretis Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946; Matteri, 1973a; Seki, 1974).

B. areola Besch.

Breutelia aureola

B. aureola (C. Muell.) Besch. (Bescherelle, 1885a).

Syn. *Bartramia aureola* Besch. ex C. Muell. (Bescherelle ex Mueller, 1885).

Breutelia areola Besch. (Bescherelle in Cardot and Brotherus, 1923) err.

Breutelia rupestris (Mitt.) Jaeg. var. *gracilior* Card. (Cardot, 1908a) *fide* Matteri, 1973a.

Distr. Southern South America (Mueller, 1885; Bescherelle, 1885a, 1889; Dusén, 1903a (and as *B. elongata* *fide* Matteri, 1973a); Paris, 1904a; Cardot, 1905a, 1908a (and as *B. elongata* *fide* Matteri, 1973a); Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1954; Matteri, 1973a; Seki, 1974).

B. auricolor Herz.

Breutelia integrifolia

B. austro-arcuata (C. Muell.) Kindb. (Kindberg, 1889).

Syn. **Bartramia austro-arcuata** C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1894, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. brachycoma Besch.

B. carinata (Mitt.) Jaeg.

B. chilensis (Lor.) Par.

B. chrysura (C. Muell.) Broth.

B. comosa (Mitt.) Mitt.

B. comosa Besch.

B. cuspidatissima (C. Muell.) Kindb. (Kindberg, 1889).

Syn. **Bartramia cuspidatissima** C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1894, 1904a; Brotherus, 1924a; Thériot, 1934b; Kühnemann, 1938).

B. dumosa Mitt. (Mitten, 1860).

Syn. **Bartramia dumosa** (Mitt.) Mitt. (Mitten, 1869).

Bartramia chilensis Lor. (Lorentz, 1866) fide Matteri, 1973a.

Breutelia chilensis (Lor.) Par. (Paris, 1904a) fide Matteri, 1973a.

Breutelia brachycoma Besch. (Bescherelle, 1885a) fide Matteri, 1973a.

Breutelia comosa Besch. (Bescherelle, 1885a) nom. nud.

Breutelia longipes Schimp. (Schimper in Paris, 1904a) nom. nud. pro parte.

Distr. Southern South America (Wilson and Hooker, 1847 (as **Bartramia pendula** vars. 1 and 2 fide Matteri, 1973a); Mueller, 1849 (as **Bartramia pendula** fide Matteri, 1973a), 1885 (and as **Bartramia comosa** fide Matteri, 1973a), 1901a (as **Bartramia comiramea** fide Matteri, 1973a); Sullivant, 1859; Mitten 1860, 1869; Lorentz, 1866; Jaeger, 1875 (and as **B. pendula** fide Matteri, 1973a); Bescherelle, 1885a, 1889; Dusén, 1903a (and as **B. elongata** fide Matteri, 1973a); Paris, 1904a; Cardot, 1908a (and as **B. elongata** and **B. pendula** fide Matteri, 1973a); Cardot and Brotherus, 1923; Herzog, 1923, 1939, 1954; Brotherus, 1924a; Rovainen, 1934; Kühnemann, 1938; Matteri, 1973a; Seki, 1974).

B. elongata (Hook. f. et Wils.) Mitt.

see **Breutelia aureola**, **B. dumosa** and **B. plicata**

B. glabrifolia Dus.

B. graminicola Broth.

B. hariotiana (C. Muell.) Besch.

B. hariotii Besch. ex Kindb.

B. integrifolia (Tayl.) Jaeg. (Jaeger, 1875).

Syn. **Bartramia integrifolia** Tayl. (Taylor, 1846).

Bartramia chrysura C. Muell. (Mueller, 1884 [1883]).

Breutelia chrysura (C. Muell.) Broth. (Brotherus, 1904).

Breutelia glabrifolia Dus. (Dusén, 1905b) fide Matteri, 1973a.

Bartramia graminicola C. Muell. (Mueller, 1884 [1883]) hom. illeg. fide Matteri, 1973a.

Breutelia graminicola Broth. (Brotherus, 1904) nom. illeg. fide Matteri, 1973a.

Philonotis kerguelensis Par. (Paris, 1897) fide Matteri, 1973a.

Breutelia kerguelensis (Par.) Wijk et Marg. (van der Wijk and Margadant, 1958) fide Matteri, 1973a.

Breutelia skottsbergii Card. (Cardot, 1905a) fide Matteri, 1973a.

Breutelia skottsbergii Card. var. **horrida** Card. et Broth. (Cardot and Brotherus, 1923) fide Matteri, 1973a.

Breutelia integrifolia (Tayl.) Jaeg. var. **patagonica** Card. et Broth. (Cardot and Brotherus, 1923) fide Matteri, 1973a.

Anacolia auricolor Dus. (Dusén in Paris, 1906) nom. nud. fide Matteri, 1973a.

Breutelia auricolor Herz. (Herzog, 1909) nom. nud. fide Matteri 1973a.

Breutelia thysanomitrioides Dus. (Dusén in Paris, 1904a) nom. nud. fide Matteri, 1973a.

Breutelia dumosa

Breutelia plicata

Breutelia dumosa

Breutelia integrifolia

see **Breutelia dumosa**

Breutelia dumosa

Distr. Southern South America (Sullivant, 1859; Neger, 1899; Dusén, 1903a, 1905b; Paris, 1904a, 1906; Cardot, 1905a, 1908a; Campo, 1921; Cardot and Brotherus, 1923; Brotherus, 1924a, 1925; Thériot, 1924, 1925a; Rovainen, 1934; Kühnemann, 1938; Herzog, 1940, 1954, 1957; Bartram, 1952; Matteri, 1973a; Seki, 1974). Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938; Matteri, 1973a). South Georgia (Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Steere, 1961b; Matteri, 1973a; Newton, 1974a).

var. **patagonica** Card. et Broth.

Breutelia integrifolia

B. kerguelensis (Par.) Wijk et Marg.

Breutelia integrifolia

B. longipes Schimp.

Breutelia dumosa, **Breutelia plicata** and **Breutelia subplicata**

B. lorentzii (C. Muell.) Kindb. (Kindberg, 1889).

Syn. **Bartramia lorentzii** C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. masafuerae Broth. (Brotherus, 1924b).

Syn. **Breutelia subplicata** Broth. var. **masafuerae** (Broth.) Matteri (Matteri, 1973a) fide Robinson, 1975.

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Matteri, 1973a; Robinson, 1975).

B. pendula (Sm.) Mitt.

see **Breutelia dumosa**

B. persquarrosa Dus.

Breutelia subplicata

B. perruncinata Dus.

Breutelia subplicata

B. plicata Mitt. (Mitten, 1860).

Syn. **Bartramia plicata** (Mitt.) Mitt. (Mitten, 1869).

Bartramia hariotiana Besch. ex C. Muell. (Bescherelle ex Mueller, 1885).

Breutelia hariotiana (C. Muell.) Besch. (Bescherelle, 1885a).

Breutelia hariotii Besch. ex Kindb. (Bescherelle ex Kindberg, 1888) hom. illeg.

Bartramia carinata Mitt. (Mitten, 1869) fide Matteri, 1973a.

Bartramia carinata (Mitt.) Jaeg. (Jaeger, 1875).

Bartramia rupestris Mitt. (Mitten, 1869) fide Matteri, 1973a.

Breutelia rupestris (Mitt.) Jaeg. (Jaeger, 1875) fide Matteri, 1973a.

Breutelia subelongata Broth. (Brotherus in Dusén, 1905b) fide Matteri, 1973a.

Breutelia longipes Schimp. pro parte (Schimper in Paris, 1904a) nom. nud. fide Matteri, 1973a.

Breutelia speciosa Card. (Cardot in Herzog, 1954) nom. nud. fide Seki, 1974.

Distr. Southern South America (Mitten, 1860, 1869; Lorentz, 1866; Jaeger, 1875; Mueller, 1885; Bescherelle, 1885a, 1889; Neger, 1899; Dusén, 1903a (and as **B. elongata** fide Matteri, 1973a), 1905b; Paris, 1904a; Cardot, 1905a, 1908a (and as **B. elongata** fide Matteri, 1973a); Cardot and Brotherus, 1923; Brotherus, 1924a; Reimers, 1926; Kühnemann, 1938; Herzog, 1939, 1954; Matteri, 1973a; Seki, 1974).

B. robusta (Hook. f. et Wils.) Jaeg.

Bartramia robusta

B. rupestris (Mitt.) Jaeg.

Breutelia plicata

var. **gracilior** Card.

Breutelia aureola

B. scabrifolia Dus. (Dusén in Paris, 1904a) nom. nud.

Distr. Southern South America (Paris, 1904a).

B. skottsbergii Card.

Breutelia integrifolia

var. **horrida** Card. et Broth.

Breutelia integrifolia

B. speciosa Card.

Breutelia plicata

B. subelongata Broth.

Breutelia plicata

B. subplicata Broth. (Brotherus in Thériot, 1911).

Syn. **Breutelia persquarrosa** Dus. (Dusén in Paris, 1904a) nom. nud.

Breutelia perruncinata Dus. (*Dusén in Paris*, 1904a) *nom. nud. fide* Matteri, 1973a.
Breutelia longipes Schimp. *pro parte* (*Schimper in Paris*, 1904a) *nom. nud. fide* Matteri, 1973a.

Distr. Juan Fernandez (Matteri, 1973a; Robinson, 1975).

Southern South America (Paris, 1904a; Thériot, 1911, 1934b, 1935a; Brotherus, 1924a; Roivainen, 1934; Herzog, 1938, 1939, 1940, 1954, 1960; Bartram, 1952; Matteri, 1973a; Seki, 1974).

var. masafuerae (Broth.) Matteri

B. thysanomitrioides Dus.

B. ulicina Herz. (*Herzog*, 1927).

Distr. Southern South America (*Herzog*, 1927).

Bruchia

B. brevipes Harv. ex Hook. (*Harvey ex Hooker*, 1840).

Syn. *Phascum brevipes* (Hook.) Schwaegr. (*Schwaegrichen*, 1842).

Distr. Southern South America (*Montagne*, 1850).

B. hampeana C. Muell. (*Mueller*, 1848a).

Syn. *Sporledera hampeana* (C. Muell.) Mitt. (*Mitten*, 1869).

Bruchia schwaegrichenii Jaeg. (*Jaeger*, 1869) *nom. illeg.*

Sporledera schwaegrichenii Hamp. (*Hampe in Jaeger*, 1869) *nom. nud.*

Distr. Southern South America (*Mueller*, 1848a; *Jaeger*, 1869, 1873; *Mitten*, 1869; *Paris*, 1904a; *Roth*, 1911; *Brotherus*, 1924a).

B. schwaegrichenii Jaeg.

Breutelia masafuerae

Breutelia integrifolia

Bruchia hampeana

B. uleana C. Muell. ex Roth (*Mueller ex Roth*, 1911).

Distr. Southern South America (*Hosseus*, 1940).

B. uruguensis C. Muell. (*Mueller*, 1879a).

Distr. Southern South America (*Mueller*, 1879a; *Jaeger and Sauerbeck*, 1879b; *Paris*, 1904a; *Roth*, 1911; *Brotherus*, 1924a; *Herter*, 1933a; *Kühnemann*, 1938; *Hosseus*, 1939).

Bryodusenia

B. genuflexa (C. Muell.) H. Robinson

Ancistrodes genuflexa

Bryoerythrophyllum

B. campylocarpum (C. Muell.) Crum (Crum, 1957).

Syn. *Trichostomum campylocarpum* C. Muell. (*Mueller*, 1851a).

Didymodon calymeridictyon Broth. (*Brotherus*, 1924b).

Didymodon linearis Broth. (*Brotherus*, 1924b) *hom. illeg.*

Distr. Juan Fernandez (*Brotherus*, 1924a, b; *Bartram*, 1959; *Robinson*, 1975).

B. recurvirostre (Hedw.) Chen (Chen, 1941).

Syn. *Weisia recurvirostris* Hedw. (*Hedwig*, 1801).

Bryum lacustre Brid. (*Bridel*, 1803).

Dermatodon rubellus Genth (Genth, 1835).

Didymodon rubellus B.S.G. (*Bruch and others*, 1846) *nom. illeg.*

Distr. Southern South America (*Bruch and others*, 1839; *Wilson and Hooker*, 1847; *Cardot*, 1905a, 1908a; *Kühnemann*, 1938).

var. antarctica Savicz et Z. Smirn. (*Savich-Ljubitskaya and Smirnova*, 1963b).

Distr. Antarctica, continental region (*Savich-Ljubitskaya and Smirnova*, 1963b; *Greene*, 1968a; *Savich-Ljubitskaya*, 1978).

Bryum

B. aequabile C. Muell. (*Mueller*, 1882).

Distr. Southern South America (*Mueller*, 1882; *Paris*, 1904a; *Brotherus*, 1924a; *Hosseus*, 1936, 1937, 1938b, d; *Kühnemann*, 1938).

B. aerugo Dus. (*Dusén*, 1903a) *nom. nud.*

Distr. Southern South America (*Dusén*, 1903a; *Paris*, 1904a).

B. albicans Roehl.

B. albidum (Hedw.) P. Beauv.

B. algens Card. (*Cardot*, 1907a).

Syn. *Bryum atgens* Card. (*Cardot in Holzinger*, 1911) *err.*

Distr. Antarctic, peninsula region (*Cardot*, 1911b, d, 1913a; *Dixon*, 1920; *Steere*, 1961a). Antarctic, continental region (*Cardot*, 1907a, 1908a; *Brotherus*, 1924a; *Steere*, 1961a; *Savich-Ljubitskaya and Smirnova*, 1972; *Savich-Ljubitskaya*, 1978).

B. alticaule C. Muell.

var. robustius C. Muell.

Pohlia wahlenbergii

Octoblepharum albidum

Pohlia wahlenbergii var. *glacialis*

Mniobryum alticaule var. *robustius*

B. amblyodon C. Muell. (*Mueller*, 1879a).

Distr. Southern South America (*Mueller*, 1879a; *Paris*, 1904a; *Kühnemann*, 1938).

*B. amblyolepis** Card. (*Cardot*, 1900).

Distr. Southern South America (*Costes*, 1921). Antarctic, peninsula region (*Cardot*, 1900, 1901, 1908a; *Paris*, 1904a; *Brotherus*, 1924a; *Steere*, 1961a). Antarctic, continental region (*Cardot*, 1907a, 1908a; *Steere*, 1961a).

B. amplirete C. Muell.

Pohlia wahlenbergii var. *glacialis*

B. andicola Hook. (*Hooker in Kunth*, 1822) see *Ochi*, 1972.

Syn. *Rhodobryum andicola* (Hook.) Par. (*Paris*, 1897).

Bryum lechleri C. Muell. (*Mueller*, 1856) see *Ochi*, 1972.

Rhodobryum lechleri (C. Muell.) Par. (*Paris*, 1897).

Distr. Juan Fernandez (*Brotherus*, 1924b; *Robinson*, 1975).

Southern South America (*Mueller*, 1856; *Lorentz*, 1866; *Mitten*, 1869; *Jaeger*, 1875; *Paris*, 1897, 1904a; *Thériot*, 1915, 1918, 1935a; *Campo*, 1921; *Cardot* and *Brotherus*, 1923; *Herzog*, 1923, 1938, 1939; *Brotherus*, 1924a, c; *Reimers*, 1926; *Kühnemann*, 1938; *Espinosa B.*, 1941; *Skottsberg*, 1950). Falkland Islands (*Cardot* and *Brotherus*, 1923).

B. angulosum Menz.

Conostomum pentastichum

B. anisodontacum Dus. (*Dusén*, 1903a) *nom. nud.*

Distr. Southern South America (*Dusén*, 1903a; *Paris*, 1904a).

B. anomobryoides Card. et Broth. (*Cardot* and *Brotherus*, 1923).

Distr. Southern South America (*Cardot* and *Brotherus*, 1923; *Brotherus*, 1924a; *Kühnemann*, 1938).

B. antarcticum Hook. f. et Wils. (*Hooker and Wilson in Wilson and Hooker*, 1847).

Syn. *Webera antarctica* (Hook. f. et Wils.) Jaeg. (*Jaeger*, 1875).

Distr. Antarctic, peninsula region (*Wilson and Hooker*, 1847; *Mueller*, 1851a; *Jaeger*, 1875; *Cardot*, 1906a, 1908a; *Paris*, 1906; *Darbshire*, 1923; *Brotherus*, 1924a; *Bartram*, 1957; *Steere*, 1961a; *Filson*, 1966; *Robinson*, 1972).

Antarctic, continental region (*Cardot*, 1907a, 1908a, 1910; *Dixon* and *Watts*, 1918; *Brotherus*, 1924a; *Bartram*, 1938, 1957; *Rayner*, 1940; *Clifford*, 1957; *Steere*, 1961a (as *B. algens* see *Clifford*, 1957); *Filson*, 1966; *Robinson*, 1972; *Savich-Ljubitskaya* and *Smirnova*, 1972).

* = *Bryum argenteum* Hedw. *fide* *Ochi* (1979).

Bryopoteria

B. chilensis Thér.

Camptodontium cryptodon

B. apiculatum Schwaegr. (Schwaegrichen, 1816).

Distr. Southern South America (Hornschorch, 1840; Mueller, 1848a; Mitten, 1869; Gibert, 1873; Jaeger, 1875; Paris, 1904a; Felippone, 1912; Herter, 1933a).

B. apocarpum (Hedw.) L. ex With.

var. *ciliatum* (Hedw.) With.

B. arechavaletae Broth. (Brotherus in Krieger, 1904) *nom. nud.*

Distr. Southern South America (Krieger, 1904; Herter, 1939b).

B. arenae C. Muell.***B. argenteum*** Hedw. (Hedwig, 1801).

Syn. *Bryum argyreum* C. Muell. (Mueller, 1882).

Bryum atro-sanguineum C. Muell. (Mueller, 1882).

Bryum capillisetum C. Muell. (Mueller, 1882).

Bryum compactulum C. Muell. (Mueller, 1882).

Bryum arenae C. Muell. (Mueller, 1885).

Bryum griseum Dus. ex Thér. (Dusén ex Thériot, 1917b) *fide* Ochi, 1969.

Bryum siplei Bartr. (Bartram, 1938) *fide* Greene, 1967.

Distr. Juan Fernandez (Robinson, 1975).

Southern South America (Montagne, 1850; Lorentz, 1866; Ångström, 1872; Mueller, 1882, 1885; Dusén, 1903a; Paris, 1904a; Cardot, 1908a; Felippone, 1912; Thériot, 1917b; Campo, 1921; Costes, 1921; Cardot and Brotherus, 1923; Brotherus, 1924a; Herter, 1933a, 1939b; Hosseus, 1935a, b, 1937, 1940; Herzog, 1938, 1939, 1957; Kühnemann, 1938; Bartram, 1952; Ochi, 1969; Seki, 1974).

Falkland Islands (Wilson and Hooker, 1847; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938).

Antarctic, peninsula region (Wilson and Hooker, 1847; Paris, 1904a; Cardot, 1906a, 1908a, 1911b, d, 1913a; Steere, 1961a; Robinson, 1972).

Antarctic, continental region (Gepp, 1902; Cardot, 1907a, 1908a, 1910; Bartram, 1938, 1957; Horikawa, 1961; Horikawa and Ando, 1961, 1967; Steere, 1961a; Savich-Ljubitskaya and Smirnova, 1964b; Greene, 1967; Robinson, 1972).

var. lanatum (P. Beauv.) Hamp. (Hampe, 1839).

Syn. *Mnium lanatum* P. Beauv. (Palisot de Beauvois, 1805a).

Distr. Southern South America (Hosseus, 1938a, b, c, d).

var. obtusifolium Broth. (Brotherus, 1906a).

Distr. Antarctic, continental region (Savich-Ljubitskaya, 1978).

var. subamblyolepis Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

B. argentinense Thér. (Thériot in Herzog, 1957) *nom. nud.*

Distr. Southern South America (Herzog, 1957).

B. argyreum C. Muell.***Bryum argenteum******B. aspillagae*** Thér. (Thériot, 1917b).

Distr. Southern South America (Thériot, 1917b; Costes, 1921; Brotherus, 1924a).

B. atgens Card. in Holz.***Bryum algens******B. atro-sanguineum*** C. Muell.***Bryum argenteum******B. aubertii*** (Schwaegr.) Brid.***Rhodobryum aubertii******B. austro-affine*** Broth. (Brotherus, 1916).

Distr. Southern South America (Felippone, 1928; Herter, 1933a).

B. austro-albicans C. Muell.***Pohlia wahlenbergii******B. austro-chilense*** Bartr. (Bartram, 1946).

Distr. South South America (Rovainen, 1934; Bartram, 1946).

B. austro-polare* Card. (Cardot, 1900).

Distr. Southern South America (Looser, 1932; Thériot, 1934a).

Antarctic, peninsula region (Cardot, 1900, 1901, 1906b, 1907c, 1908a; Paris, 1904a; Brotherus, 1924a; Steere, 1961a (as *B. antarcticum*); ? Filson, 1966 (as *B. antarcticum*); Ochi, 1970 (as *B. pseudotriquetrum* for Racovitza, 234b)).

B. austro-turbinatum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935a, 1937; Kühnemann, 1938).

B. badium (Brid.) Schimp. (Schimper, 1876).

Syn. *Bryum caespiticium* L. ex Hedw. var. *badium* Bruch ex Brid. (Bridel, 1827a).

Bryum caespiticium L. ex Hedw. ssp. *badium* (Brid.) Lindb. (Lindberg, 1879).

Bryum caespiticium L. ex Hedw. var. *gracilescens* B.S.G. (Bruch and others, 1839) *nom. illeg.*

Distr. Falklands Islands (Wilson and Hooker, 1847; Kühnemann, 1938).

B. bartramiospis C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. beyrichiae Hornsch. ex Kindb.***Rhodobryum beyrichianum******B. beyrichianum*** (Hornsch.) C. Muell.***Rhodobryum beyrichianum******B. beyrichii*** Aongstr.***Rhodobryum beyrichianum******B. bicolor*** Dicks. (Dickson, 1801).

Syn. *Bryum dichotomum* auct. non Hedwig *fide* Ochi, 1972.

Bryum bulbilosum Mont. (Montagne, 1841).

Bryum ferriviae C. Muell. (Mueller in Ule, 1899) *nom. nud.*

Distr. Southern South America (Mitten, 1869; Gibert, 1873; Jaeger, 1875; Paris, 1904a; Brotherus, 1924a; Herter, 1928, 1933a; Seki, 1974).

B. billardieri Schwaegr. (Schwaegrichen, 1816).

Distr. Falkland Islands (Wilson and Hooker, 1847; Kühnemann, 1938).

B. bimum (Scherb.) Turn.***Bryum pseudotriquetrum* ssp. *bimum***

ssp. *pseudotriquetrum* (Hedw.) Dix.

Bryum pseudotriquetrum***B. brachycarpum*** Bartr. (Bartram, 1946) *hom. illeg.*

Distr. Southern South America (Bartram, 1946).

B. brachychaete Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a).

B. brachymeniopsis C. Muell.***Anomobryum brachymeniopsis******B. brevigemmatum*** Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1904a; Seki, 1974).

B. bulbillinum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. bulbillosum Mont.***Bryum bicolor******B. caespiticium*** L. ex Hedw. (Hedwig, 1801).

Syn. *Mnium caespiticium* (Hedw.) With. (Withering, 1801).

Hypnum caespiticium (Hedw.) Web. et Mohr (Weber and Mohr, 1803).

* = *Bryum pseudotriquetrum* (Hedw.) Gaertn., Meyer et Scherb. *fide* Ochi (1970, 1979).

Distr. Juan Fernandez (Montagne, 1835, 1845b).
 Southern South America (Wilson and Hooker, 1847; Kühnemann, 1938; Seki, 1974).
 Falkland Islands (Wilson and Hooker, 1847).

ssp. badium (Brid.) Lindb.

var. badium Bruch ex Brid.

var. gracilescens B.S.G.

B. caespitosum (Hopp. et Hornsch.) Brid.

B. calopyxis C. Muell.

B. campoanum Thér. (Thériot, 1926).

Distr. Southern South America (Campo, 1921; Thériot, 1926).

B. campylothecium Tayl. (Taylor, 1846).

Syn. *Bryum jaffuelii* Thér. (Thériot, 1926) *fide* Ochi, 1977.

Distr. Southern South America (Ochi, 1977).

B. campylotrichum C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

B. canariense Brid. (Bridel, 1817).

Syn. *Bryum jaffuelii* Thér. (Thériot, 1926).

Distr. Southern South America (Montagne, 1850; Thériot, 1926).

B. candicans Tayl. (Taylor *in* Mitten, 1869).

Distr. Southern South America (Thériot, 1918; Campo, 1921).

B. canescens Schimp. (Schimper *in* Paris, 1904a) *nom. nud.*

Distr. Southern South America (Lechler, 1857; Paris, 1904a).

B. capillaceum (Hedw.) Dicks. *ex* With.

B. capillare Hedw. (Hedwig, 1801).

Syn. *Bryum philippianum* C. Muell. (Mueller, 1844b [1845]) *fide* Ochi, 1969.

Bryum philippii C. Muell. *ex* Kindb. (Mueller *ex* Kindberg, 1888) *nom. illeg.*

Bryum chilense Reichdt. (Reichardt, 1868) *fide* Ochi, 1969.

Bryum nanocoma C. Muell. (Mueller, 1882) *fide* Ochi, 1969.

Distr. Southern South America (Mueller, 1844b, 1848a, 1882; Montagne, 1850; Lorentz, 1866; Reichardt, 1868, 1870; Mitten, 1869; Jaeger, 1875; Neger, 1899; Paris, 1904a; Herzog, 1923, 1938, 1954; Brotherus, 1924a; Kühnemann, 1938; Ochi, 1969; Seki, 1974).

ssp. torquescens (B.S.G.) Kindb. (Kindberg, 1897).

Syn. *Bryum torquescens* B.S.G. (Bruch and others, 1839).

Distr. Southern South America (Bruch and others, 1839; Mueller, 1848a; Montagne, 1850; Mitten, 1869; Gibert, 1873; Paris, 1904a; Felippone, 1912; Thériot, 1928; Herter, 1928, 1933a, 1939b).

var. cymbifolium Hook. f. *et* Wils.

var. flaccidum (Brid.) B.S.G. (Bruch and others, 1839).

Syn. *Bryum flaccidum* Brid. (Bridel, 1826).

Distr. Southern South America (Bruch and others, 1839).

B. capillisetum C. Muell.

B. capituliforme Warnst. (Warnstorf, 1915).

Distr. Southern South America (Warnstorf, 1915).

B. cavum C. Muell. (Mueller, 1844a).

Bryum badium

Bryum badium

Bryum badium

Pohlia nutans

Pohlia calopyxis

Syn. *Bryum capillare* Hedw. var. *cymbifolium* Hook. f. *et* Wils. (Hooker, J. D. and Wilson, 1844).

Distr. Southern South America (Mitten, 1869; Gibert, 1873; Paris, 1904a; Herter, 1928, 1933a, b).

B. cephalozoides Card. (Cardot, 1906a).

Distr. Antarctic, peninsula region (Cardot, 1906a, 1908a; Steere, 1961a).

B. chilense Reichdt.

Bryum capillare

B. chiloense Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926).

B. chorizodontum Card. *et* Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a).

B. chryseum Mitt. (Mitten, 1869).

Distr. Juan Fernandez (Bartram, 1959; Robinson, 1975).

B. ciliatum (Hedw.) Dicks.

Hedwigia ciliata

B. cirrhatum Hopp. *et* Hornsch. var. *australe* Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a).

B. clavatum (Schimp.) C. Muell. (Mueller, 1848a).

Syn. *Pohlia clavata* Schimp. (Schimper, 1836).

Distr. Southern South America (Schimper, 1836; Mueller, 1848a; Montagne, 1850; Mitten, 1869; Paris, 1904a; Brotherus, 1924a).

B. climacodontium Par.

Anomobryum cygnicollum

B. cochlearifolium Card. *et* Broth. (Cardot and Brotherus, 1923) hom. illeg.

Distr. Southern South America (Donat, 1936a; Herzog, 1960; Seki, 1974).

South Georgia (Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b).

B. coelophyllum Eaton (Eaton, 1892).

Distr. Southern South America (Eaton, 1892).

B. commutatum (Schimp.) Boul.

Pohlia drummondii

B. compactulum C. Muell.

Bryum argenteum

B. concavifolium Thér. (Thériot, 1926).

Distr. Southern South America (Costes, 1921 (as *B. concarifolium*); Thériot, 1926; Herzog, 1939).

B. conicum Hornsch.

Anomobryum conicum

B. controversum (Hedw.) P. Beauv.

Weisia controversa

B. coronatum Schwaegr. (Schwaegrichen, 1816).

Distr. Southern South America (Mueller, 1848a; Montagne, 1850; Jaeger, 1875; Seki, 1974).

B. corralense Herz. *et* Thér. (Herzog and Thériot *in* Herzog, 1938).

Distr. Southern South America (Herzog, 1938).

B. crassicaule Dus.

Bryum laevigatum

B. crassinervium Lor. (Lorentz, 1866).

Distr. Southern South America (Lorentz, 1866; Jaeger, 1875; Paris, 1904a; Costes, 1921; Brotherus, 1924a).

B. crateris* Dix. (Dixon, 1920).

Distr. Antarctic, peninsula region (Dixon, 1920; Brotherus, 1924a; Steere, 1961a).

B. crispatum Dicks. ex With.

B. crudum (Hedw.) Turn.

B. cruegeri Hamp. (Hampe in Mueller, 1848a).

Syn. *Pohlia cruegeri* (Hamp.) Andrews (Andrews, 1935).

Distr. Southern South America (Kühnemann, 1938).

B. cygnicollum C. Muell.***Rhabdoweisia crispata******Pohlia cruda******Anomobryum cygnicollum******B. decurrentinervium*** C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1918, 1924a; Kühnemann, 1938).

B. delitescens Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. demissum Dus. (Dusén, 1903a) nom. nud.

Distr. Southern South America (Dusén, 1903a).

B. densifolium Brid. (Bridel, 1827a).

Syn. *Rhodobryum gracilescens* C. Muell. (Mueller in Hampe, 1874).

Distr. Southern South America (Herter, 1933a; Hosseus, 1940; Herzog, 1952a).

B. denticulatinervium Ochi (Ochi, 1977).

Distr. Southern South America (Ochi, 1977).

B. diaphanum C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a, 1901a; Paris, 1904a).

B. dicarpum Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

B. dichotomum auct. non Hedwig***Bryum bicolor******B. dissolutinerve*** C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Kühnemann, 1938).

B. donati Herz. et Thér.***Bryum redboonii******B. donatii*** Thér. (Thériot, 1934b [1935]).

Distr. Southern South America (Thériot, 1934b; Herzog, 1957).

B. drummondii C. Muell.***Pohlia drummondii******B. elegantulum*** Lor. (Lorentz, 1864).

Distr. Southern South America (Lorentz, 1864, 1866; Jaeger, 1875; Paris, 1904a; Brotherus, 1924a, c; Espinosa B., 1941).

B. ellipsifolium C. Muell. (Mueller, 1890b).

Syn. *Bryum myurella* Dus. (Dusén, 1905b) hom. illeg. fide Ochi, 1972.

Distr. Southern South America (Dusén, 1903a, 1905b; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Ochi, 1972; Seki, 1974).

B. ellipticum Mitt.***Brachymenium meyenianum* var. *ellipticum******B. emergens*** C. Muell.***Mielichhoferia emergens***

* = *Bryum pseudotriquetrum* (Hedw.) Gaertn., Meyer et Scherb. fide Ochi (1979).

B. encalyptaceum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. erubescens C. Muell.***Brachymenium erubescens******B. erythrocarpoides*** C. Muell. et Hamp. (Mueller and Hampe, 1855).

Syn. *Bryum reichertii* Herz. (Herzog, 1926) fide Seki, 1974.

Distr. Southern South America (Herzog, 1926; Seki, 1974).

B. fabronia C. Muell. (Mueller, 1882).

Syn. *Bryum fabroniae* C. Muell. ex Kindb. (Mueller ex Kindberg, 1889) nom. inval.

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. fabroniae C. Muell. ex Kindb.***Bryum fabronia******B. fabronioides*** C. Muell.***Brachymenium fabronioides******B. felipponei*** Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933a).

B. fernandezianum Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

B. ferriviae C. Muell.***Bryum bicolor******B. filicaule*** Broth. (Brotherus, 1906a).

Distr. Antarctic, continental region (Cardot, 1908a; Brotherus, 1924a; Steere, 1961a (as *B. antarcticum*); Filson, 1966 (as *B. antarcticum*)).

B. filiforme Dicks.***Anomobryum filiforme******B. flaccidum*** Brid.***Bryum capillare* var. *flaccidum******B. flagellans*** C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

B. flagellicoma C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Kühnemann, 1938).

B. flavo-pallidum Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a).

B. flexisetum Mitt. (Mitten, 1869).

Distr. Southern South America (Bartram, 1942, 1943).

B. fontanum (Hedw.) Gaertn., Meyer et Scherb.***Philonotis fontana******B. fuegianum*** Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

B. fusco-mucronatum C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1901a; Brotherus, 1924a; Herter, 1933a; Kühnemann, 1938).

B. ganophyllum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Brotherus, 1904, 1924a; Kühnemann, 1938).

B. gaudichaudii (Schwaegr.) Spreng.***Leptotheca gaudichaudii***

B. gayanum Mont. (Montagne in Mueller, 1848a).

Syn. *Bryum gayi* Mont. ex Kindb. (Montagne ex Kindberg, 1888) nom. illeg.

Distr. Southern South America (Mueller, 1848a; Montagne, 1850, 1856; Mitten, 1869; Jaeger, 1875; Paris, 1904a; Brotherus, 1924a).

B. gayi Mont. ex Kindb.***Bryum gayanum******B. gemmaceum*** Besch. (Bescherelle in Mueller, 1901a [1900] nom. nud.).

Distr. Southern South America (Mueller, 1901a).

B. gemmatum C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Paris, 1904a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).

B. gerlachei* Card. ex Broth. (Cardot ex Brotherus, 1904).

Syn. *Webera gerlachei** Card. (Cardot, 1900).

Distr. Antarctic, peninsula region (Cardot, 1900, 1901, 1906a, b, 1907c, 1908a, 1911d, 1913a; Paris, 1906; Brotherus, 1924a; Steere, 1961a (as *B. antarcticum*): ? Filson, 1966 (as *B. antarcticum*); Ochi, 1970 (as *B. pseudotriquetrum* for Racovitzia 229a)).

B. gibertii C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

B. gilliesii Hook. (Hooker, 1829).

Syn. *Pohlia gilliesii* (Hook.) Mont. (Montagne, 1839b).
Brachymenium gilliesii (Hook.) Jaeg. (Jaeger, 1875).

Distr. Southern South America (Hooker, 1829; Montagne, 1839b; Mueller, 1848a, 1879a; Mitten, 1869; Jaeger, 1875; Kurtz, 1904; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935a, 1937; Kühnemann, 1938; Herzog, 1957).

B. glaciale Brid.***Pohlia wahlenbergii* var. *glacialis******B. glauco-viride*** C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. gracilisetum Hornsch. (Hornschuch, 1840).

Distr. Southern South America (Krieger, 1904; Herter, 1943).

B. gracili-torquescens C. Muell. (Mueller in Kindberg, 1891) nom. nud.

Distr. Southern South America (Paris, 1900a; Thériot, 1917b, 1918, 1921a; Campo, 1921; Costes, 1921).

B. gracillimum Broth. (Brotherus in Felippone, 1909).

Distr. Southern South America (Felippone, 1909; Brotherus, 1924a; Herter, 1933a).

B. griffithianum Dicks.***Oedipodium griffithianum******B. griseum*** Dus. ex Thér.***Bryum argenteum******B. hallei*** Card.***Bryum schleicheri* var. *patagonicum******B. hamatum**** Dus. (Dusén, 1903a) nom. nud.***Bryum perlimbatum* pro parte*****B. hamipilum*** C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1918; Kühnemann, 1938).

B. hatcheri Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Kühnemann, 1938).

* = *Bryum pseudotriquetrum* (Hedw.) Gaertn., Meyer et Scherb. fide Ochi (1979).

* = *Bryum viridescens* Welw. et Dub. fide Ochi (1977).

B. hauthalli C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. hedwigii Brotero***Hedwigia ciliata******B. heimii*** (Hedw.) Dicks. ex With.***Pottia heimii******B. heteroblepharum*** Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a).

B. heteromallum (Hedw.) P. Beauv.***Ditrichum heteromallum******B. heterophyllum*** Card. et Broth. (Cardot and Brotherus, 1923) hom. illeg.

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

B. heterostichum (Hedw.) Dicks.***Racomitrium heterostichum******B. hickenii*** Herz.***Bryum pseudotriquetrum******B. hieronymi*** C. Muell.***Rhodobryum hieronymi******B. humile*** Mont.***Pohlia humilis******B. hypnoides*** L. ex With.***Racomitrium lanuginosum******B. imperfectum*** Card. (Cardot, 1900).

Distr. Southern South America (Herzog, 1957).
Antarctic, peninsula region (Cardot, 1900, 1901, 1908a, 1911b, d, 1913a; Paris, 1904a; Brotherus, 1924a; Steere, 1961a; Robinson, 1972).

B. inclinatum (Brid.) Bland. (Blandow, 1809) hom. illeg.

Syn. *Pohlia inclinata* Sw. ex Brid. (Swartz ex Bridel, 1803).
Leskeia inclinata (Brid.) Web. et Mohr (Weber and Mohr, 1803).

Distr. Southern South America (Montagne, 1850).

var. magellanicum Card. (Cardot, 1900).

Distr. Southern South America (Paris, 1904a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Herzog, 1923; Brotherus, 1924a; Kühnemann, 1938).
Antarctic, peninsula region (Cardot, 1900, 1901; Bartram, 1957; Steere, 1961a).

B. inconnexum* Card. (Cardot, 1900).

Distr. Antarctic, peninsula region (Cardot, 1900, 1901, 1906b, 1907c, 1908a; Paris, 1904a; Brotherus, 1924a; Steere, 1961a (as *B. antarcticum*); ? Filson, 1966 (as *B. antarcticum*); Robinson, 1972).

Antarctic, continental region (Greene, 1968a; Kuc, 1969; Robinson, 1972; Ochi, 1976).

var. fragile Hor. et Ando (Horikawa and Ando, 1961).

Distr. Antarctic, continental region (Horikawa and Ando, 1961).

var. tomentosum Card. (Cardot, 1906b).

Distr. Antarctic, peninsula region (Cardot, 1906b, 1907c, 1908a; Greene, 1968a).

B. indicum Doz. et Molk.***Brachymenium indicum******B. inflexum*** C. Muell.***Pohlia inflexa******B. jaffuelii*** Thér.

see *Bryum canariense* and *Bryum campylothecium*

B. julaceum Schrad. ex Gaertn., Meyer et Scherb.***Anomobryum filiforme******B. kermesinum*** Herz. et Thér. (Herzog and Thériot in Herzog, 1939).

Distr. Southern South America (Herzog, 1939).

B. klinggraeffii Schimp. (Schimper in Klinggraeff, 1858).

Distr. Southern South America (Crundwell and Nyholm, 1964).

* = *Bryum pseudotriquetrum* (Hedw.) Gaertn., Meyer et Scherb. fide Ochi (1979).

B. korothkevieziae Sav. et Smirn. (Savich-Ljubitskaya and Smirnova, 1959).

Distr. Antarctic, continental region (Savich-Ljubitskaya and Smirnova, 1959; Savich-Ljubitskaya, 1978).

. var. hollerbachii Sav. et Smirn. (Savich-Ljubitskaya and Smirnova, 1960).

Distr. Antarctic, continental region (Savich-Ljubitskaya and Smirnova, 1960; Greene, 1968a).

B. lacustre Brid.***Bryoerythrophyllum recurvirostre******B. laetevirens*** Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1954; Seki, 1974).

B. laevigatum Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Syn. *Bryum levigatum* Hook. f. et Wils. (Hooker and Wilson in Mitten, 1860) err.
Bryum crassicaule Dus. (Dusén in Paris, 1904a) nom. nud.

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Sullivant, 1859; Jaeger, 1875; Bescherelle, 1889; Dusén, 1903a; Paris, 1904a; Cardot, 1905a, 1908a; Campo, 1921; Cardot and Brotherus, 1923; Brotherus, 1924a; Thériot, 1934b, 1935a; Kühnemann, 1938; Herzog, 1938, 1939, 1954; Ochi, 1970; Seki, 1974).

Falkland Islands (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a; Dusén, 1903a; Paris, 1904a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. lamprocarpum C. Muell. (Mueller, 1890a).

Syn. *Zieria lamprocarpa* (C. Muell.) Kindb. (Kindberg, 1891).
Plagiobryum lamprocarpum (C. Muell.) Par. (Paris, 1897).

Distr. Falkland Islands (Kühnemann, 1938).
South Georgia (Mueller, 1890a; Paris, 1897, 1905b; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b).

B. lamprochaete Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1957).

B. lamprocumum C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. lanuginosum (Hedw.) Brotero***Racomitrium lanuginosum******B. laticeps*** C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904a; Brotherus, 1924a; Hosseus, 1937; Kühnemann, 1938).

B. lechleri C. Muell.***Bryum andicola******B. leptotrichum*** C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a).

B. leuco-aristatum Dus. (Dusén, 1903a) nom. nud.

Distr. Southern South America (Dusén, 1903a).

B. levigatum Hook. f. et Wils.***Bryum laevigatum******B. liliiputanum*** Dus. (Dusén, 1903a) nom. nud.

Distr. Southern South America (Dusén, 1903a).

B. linearifolium C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a).

B. litoris Card. (Cardot in Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. lonchochaete Dus. (Dusén, 1903a) nom. nud.

Distr. Southern South America (Dusén, 1903a).

B. longidens Thér. (Thériot, 1926).

Distr. Southern South America (Campo, 1921; Thériot, 1926).

B. longirostre Brid.***Mnium rostratum******B. lorentzianum*** C. Muell.***Rhodobryum lorentzianum******B. maceratum*** C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1918, 1924a; Kühnemann, 1938).

B. macro-carneum C. Muell. (Mueller, 1901a [1900]) nom. nud.

Distr. Southern South America (Mueller, 1901a).

B. macrochaete Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Donat, 1936a; Kühnemann, 1938; Herzog, 1954).

B. macropelma C. Muell. (Mueller, 1848a).

Distr. Southern South America (Mueller, 1848a; Montagne, 1850; Mitten, 1869; Jaeger, 1875; Paris, 1904a; Brotherus, 1924a).

B. macrophyllum Card. et Broth. (Cardot and Brotherus, 1923).

Syn. *Bryum macrophyllum* Card. et Broth. var. *latifolium* Broth. (Brotherus in Cardot and Brotherus, 1923) fide Ochi, 1967.

Bryum megalophyllum Card. (Cardot in Cardot and Brotherus, 1923) nom. nud.

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Ochi, 1967).

var. *latifolium* Broth.***Bryum macrophyllum******B. macropoma*** C. Muell.***Acidodontium macropoma******B. macrosporum*** Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

B. magellanicum (Sull.) C. Muell.***Brachymenium indicum******B. malachiticum*** C. Muell.***Brachymenium malachiticum******B. malacophyllum*** Broth. (Brotherus in Herzog, 1916a).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938; Herzog, 1954).

B. mancum C. Muell.***Mielichhoferia manca******B. megalocarpum*** Hook.***Acidodontium megalocarpum******B. megalophyllum*** Card.***Bryum macrophyllum******B. megalothecium*** Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926; Bartram, 1942, 1943; Herzog, 1957).

B. meyenianum (Hamp.) C. Muell.***Brachymenium meyenianum******B. microglossum*** C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. micron C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

B. micro-pendulum C. Muell.***Bryum nano-pendulum******B. microphyllum*** Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).

B. minusculum C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Dusén, 1903a, 1905d; Paris, 1904a; Cardot, 1908a; Brotherus, 1924a).

B. miserum Card. (Cardot, 1905a).

Distr. Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1939).

B. montevicense Broth. (Brotherus in Felippone, 1928 [1929]).

Distr. Southern South America (Felippone, 1928; Herter, 1933a).

B. montevicense Thér.***Bryum subpercurrente******B. mucronatum*** C. Muell. (Mueller, 1901a [1900]) nom. nud.

Distr. Southern South America (Mueller, 1901a).

B. murale (Hedw.) L. ex With.***Tortula muralis******B. myurella*** Dus.***Bryum ellipsifolium******B. nanocoma*** C. Muell.***Bryum capillare******B. nano-pendulum*** C. Muell. (Mueller, 1882).

Syn. *Bryum micro-pendulum* C. Muell. (Mueller, 1879b) hom. illeg.

Distr. Southern South America (Mueller, 1879a, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. nevadense C. Muell.***Pohlia nevadensis******B. nivale*** C. Muell. (Mueller, 1848a).

Distr. Southern South America (Mueller, 1848a; Montagne, 1850; Sullivant, 1859; Mitten, 1869; Jaeger, 1875; Paris, 1904a; Brotherus, 1924a).

B. nutans (Hedw.) Turn.***Pohlia nutans***

var. *minus* Hook.

Mielichhoferia longiseta***B. obliquatum*** Dus. (Dusén in Paris, 1904a) nom. nud.

Distr. Southern South America (Paris, 1904a).

B. obliquum C. Muell. (Mueller, 1890a).

Distr. South Georgia (Mueller, 1890a; Paris, 1904a; Cardot, 1908a; Steere, 1961b).

B. obscurum Card. et Broth. (Cardot and Brotherus, 1923) hom. illeg.

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. oediloma C. Muell. (Mueller in Brotherus, 1895).

Distr. Southern South America (Bartram, 1965).

B. ongiense* Hor. et Ando (Horikawa and Ando, 1961).

Distr. Antarctic, continental region (Horikawa and Ando, 1961; Savich-Ljubitskaya and Z. Smirnova, 1972).

B. orbiculatifolium Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1954; Seki, 1974).

B. orthodontoides C. Muell. (Mueller, 1848a).

Distr. Southern South America (Herter, 1943).

B. orthothecium Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a).

B. pachyneuron Dus. (Dusén in Paris, 1904a) nom. nud.

Distr. Southern South America (Paris, 1904a).

B. pallescens Schleich. ex Schwaegr. (Schleicher ex Schwaegrichen, 1816).

Syn. *Bryum pseudotriquetrum* (Hedw.) Gaertn., Meyer et Scherb. ssp. *pallescens* (Schwaegr.) Dix. (Dixon, 1924b).

Distr. Southern South America (Wilson and Hooker, 1847).

Falkland Islands (Wilson and Hooker, 1847; Kühnemann, 1938).

B. pallidipes C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. pallido-viride Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Kühnemann, 1938; Herzog, 1957).

B. palustre (Hedw.) Gaertn., Meyer et Scherb.***Aulacomnium palustre******B. papillosum*** C. Muell.***Pohlia papillosa******B. parvirete*** Dus. (Dusén in Paris, 1904a) nom. nud.

Distr. Southern South America (Paris, 1904a).

B. parvulum Card. (Cardot, 1906a) hom. illeg.

Distr. Southern South America (Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).
South Georgia (Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b).

B. parvulum Broth. (Brotherus in Felippone, 1928 [1929]) hom. illeg.

Distr. Southern South America (Felippone, 1928; Herter, 1933a).

B. patens Hook. f. et Wils.***Bryum rigidum******B. pauperculum*** Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

B. pellucens Hook.***Orthodontium pellucens******B. perangustidens**** Card. (Cardot, 1911b, d).

Distr. Antarctic, peninsula region (Cardot, 1911b, d, 1913a; Brotherus, 1924a; Steere, 1961a).

B. perlimbatum Card. (Cardot, 1905a).

Syn. *Bryum hamatum* Dus. (Dusén, 1903a) nom. nud. pro parte fide Seki, 1974.

Distr. Southern South America (Dusén, 1903a; Paris, 1906; Cardot and Brotherus, 1923; Herzog, 1923, 1954; Seki, 1974).

Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Ochi, 1970, 1972).

* = *Bryum pseudotriquetrum* (Hedw.) Gaertn., Meyer et Scherb. fide Ochi (1979).

* = *Bryum pseudotriquetrum* (Hedw.) Gaertn., Meyer et Scherb. fide Ochi (1979).

B. pertenerum† Thér. (Thériot, 1926).

Distr. Southern South America (Campo, 1921; Thériot, 1926).

B. pertomentosum Dus. (Dusén in Paris, 1904a) nom. nud.

Distr. Southern South America (Paris, 1904a).

B. pertriste C. Muell. (Mueller, 1882).

Syn. *Bryum triste* C. Muell. (Mueller, 1879a) hom. illeg.

Distr. Southern South America (Mueller, 1879a, 1882; Paris, 1904a; Brotherus, 1924a).

B. phallus C. Muell. (Mueller, 1875).

Distr. Southern South America (Mueller, 1875; Jaeger and Sauerbeck, 1879b; Paris, 1904a).

B. philippianum C. Muell.***Bryum capillare*****B. philippii** C. Muell. ex Kindb.***Bryum capillare*****B. philonoteum** C. Muell.***Mniobryum philonoteum*****B. philonotooides** C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

B. philonotula C. Muell. (Mueller, 1901a [1900]) nom. nud.

Distr. Southern South America (Mueller, 1901a).

B. piriforme Par.***Leptobryum pyriforme*****B. platense** C. Muell.***Rhodobryum platense*****B. platyphylloides** C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

B. platyphyllum (Schwaegr.) C. Muell. (Mueller, 1848a).

Syn. *Pohlia platyphylla* Schwaegr. (Schwaegrichen, 1842).

Distr. Southern South America (Schwaegrichen, 1842; Mueller, 1848a; Montagne, 1850; Mitten, 1869; Jaeger, 1875; Paris, 1904a; Brotherus, 1924a).

B. pluricolor Dus. ex Thér. (Dusén ex Thériot, 1917b).

Distr. Southern South America (Thériot, 1917b).

B. polycarpum Mitt.***Pohlia polycarpa*****B. pomiforme** (Hedw.) With.***Bartramia pomiformis*****B. porteri** Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926).

B. posthumum C. Muell. (Mueller, 1897a).

Syn. *Bryum pseudogemmatum* Card. (Cardot in Cardot and Brotherus, 1923) nom. nud.

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1924a; Kühnemann, 1938).

Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

B. pseudo-acidodontium C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. pseudo-fallax Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. pseudogemmatum Card.***Bryum posthumum***

† = *Bryum funkii* Schwaegr. fide Ochi (1977).

B. pseudo-micron C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Hosseus, 1935b, 1937; Kühnemann, 1938).

B. pseudothyridium Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

B. pseudotriquetrum (Hedw.) Gaertn., Meyer et Scherb. (Gärtner and others, 1802).

Syn. *Mnium pseudo-triquetrum* Hedw. (Hedwig, 1801).
Hypnum pseudotriquetrum (Hedw.) Web. et Mohr (Weber and Mohr, 1807).
Bryum ventricosum Dicks. ex Relh. (Dickson ex Relhan, 1802) nom. illeg.
Bryum ventricosum Relh. var. *pseudotriquetrum* (Hedw.) Hamp. (Hampe, 1837a) nom. illeg.

Hypnum ventricosum Web. et Mohr (Weber and Mohr, 1803) nom. illeg.
Bryum bimum (Scherb.) Turn. ssp. *pseudotriquetrum* (Hedw.) Dix. (Dixon, 1896) hom. illeg.
Bryum yendangaianum Card. (Cardot in Cardot and Brotherus, 1923) nom. nud.
Bryum hickenii Herz. (Herzog, 1926) fide Seki, 1974.

Distr. Southern South America (Cardot and Brotherus, 1923; Herzog, 1926; Hosseus, 1938c, d; Kühnemann, 1938; Bartram, 1952; Ochi, 1970; Seki, 1974).

ssp. *bimum* (Schreb.) Hartm. (Hartman, 1849).

Syn. *Mnium bimum* Schreb. (Schrebius, 1802).
Bryum bimum (Schreb.) Turn. (Turner, 1804a).
Hypnum bimum (Schreb.) Web. et Mohr (Weber and Mohr, 1803).
Bryum ventricosum Relh. var. *bimum* (Schreb.) Hamp. (Hampe, 1837a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Kühnemann, 1938).

ssp. *pallescens* (Schwaegr.) Dix.***Bryum pallescens*****B. puconense** Herz. et Thér. (Herzog and Thériot in Herzog, 1938).

Distr. Southern South America (Herzog, 1938).

B. pulchellum Hedw.***Mniobryum pulchellum*****B. pulvinatum** C. Muell.***Pohlia nutans*****B. pumilum** C. Muell. (Mueller in Brotherus, 1918) nom. nud.

Distr. Southern South America (Brotherus, 1918; Kühnemann, 1938).

B. purpuratum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. pycnodictyum C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

B. pycnotylum Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a).

B. pyriforme (Hedw.) Gaertn., Meyer et Scherb.***Leptobryum pyriforme***

var. *antarcticum* C. Muell.

Leptobryum pyriforme* var. *antarcticum

var. *fuegianum* C. Muell.

Leptobryum pyriforme* var. *fuegianum**B. pyrrhothrix** C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. raminerive C. Muell. (Mueller, 1901a [1900]) nom. nud.

Distr. Southern South America (Mueller, 1901a).

B. redboonii Donat (Donat, 1936a).

Syn. *Bryum donati* Herz. et Thér. (Herzog and Thériot *in* Donat, 1936b) *hom. illeg.*

Distr. Southern South America (Donat, 1936a, b; Kühnemann, 1938).

B. reichertii Herz.***Bryum erythrocarpoides******B. revolutum*** C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

B. rhizoblastum Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. rigidulum (Hedw.) Dicks.***Didymodon rigidulus******B. rigidum*** (Hornschr.) C. Muell. (Mueller, 1848a) *hom. illeg.*

Syn. *Mnium rigidum* Hornsch. (Hornschuch, 1840).

Rhodobryum rigidum (Hornsch.) Par. (Paris, 1897).

Bryum patens Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844) *hom. illeg.*

Distr. Southern South America (Hornschuch, 1840; Mueller, 1848a; Mitten, 1869; Gibert, 1873; Jaeger, 1875; Hampe, 1879; Paris, 1897, 1904a; Herter, 1928, 1933a).

B. rigochaete Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. rimicola Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. rivale C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a, 1882; Paris, 1904a; Kühnemann, 1938).

B. roivainenii Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

B. roseolum C. Muell. (Mueller, 1879a).

Syn. *Rhodobryum roseolum* (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1879a; Paris, 1897, 1904a; Brotherus, 1924a; Kühnemann, 1938).

B. rostratum (Schrad.) Sm.***Mnium rostratum******B. rosulans*** C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1924a).

B. rubellicaulis C. Muell. or Dus. (Mueller or Dusén *in* Neger, 1899) *nom. nud.*

Distr. Southern South America (Neger, 1899).

B. rubinerve Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. rurale (Hedw.) L. ex With.***Tortula ruralis******B. sabuletorum*** Broth. et Card. (Brotherus and Cardot *in* Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. schilleri Herz. (Herzog, 1916b).

Distr. Southern South America (Herzog, 1916b).

B. schleicheri Cand. (de Candolle, 1815).

Syn. *Mnium schleicheri* (Cand.) Brid. (Bridel, 1817).

Bryum turbinatum (Hedw.) Turn. ssp. *schleicheri* (Cand.) Kindb. (Kindberg, 1883).

Bryum turbinatum (Hedw.) Turn. var. *schleicheri* (Cand.) Fuernr. (Fürmrohr, 1829).

Distr. Southern South America (Amann, 1924; Thériot, 1926).

var. *patagonicum* Broth. (Brotherus *in* Cardot and Brotherus, 1923).

Syn. *Bryum hallei* Card. (Cardot *in* Cardot and Brotherus, 1923) *nom. nud.*

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

B. schnyderi C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Kühnemann, 1938).

B. semi-reticulatum C. Muell.

Anomobryum semireticulatum

B. senopyxis C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Brotherus, 1924a (as *B. stenopyxis*); Kühnemann, 1938).

B. senopyxoides C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

B. siplei Bartr.

Bryum argenteum

B. skottsbergii Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. spegazzinii C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Dusén, 1903a; Paris, 1904a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938). Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

B. sphagnadelphus C. Muell.

Pohlia nutans

B. splachnobryoides C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

B. steffenii Dus. ex Herz. (Dusén ex Herzog, 1926).

Distr. Southern South America (Paris, 1904a; Herzog, 1926; Seki, 1974).

B. subclavatum Thér. (Thériot, 1926).

Distr. Southern South America (Campo, 1921; Thériot, 1926).

B. subgracillimum Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926).

B. sublamprocarpum Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a).

B. submicrolaeavigatum Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. subpercurrente Thér. (Thériot *in* Felippone, 1930).

Syn. *Bryum montevidense* Thér. (Thériot *in* Felippone, 1928 [1929]) *hom. illeg.*

Distr. Southern South America (Felippone, 1928, 1930; Herter, 1933a).

B. synoico-crudum C. Muell.

Pohlia cruda

B. tenuicaule Mont. (Montagne, 1845a).

Distr. Southern South America (Montagne, 1845a, 1846, 1850, 1856; Mueller, 1848a; Mitten, 1869; Jaeger, 1875; Paris, 1904a; Costes, 1921; Brotherus, 1924a).

B. tenuirete Dus. ex Card. (Dusén ex Cardot, 1905a).

Distr. Southern South America (Dusén, 1903a; Paris, 1904a; Cardot, 1905a, 1908a; Brotherus, 1924a).

B. terminale C. Muell.**Anomobryum terminale****B. torquescens** B.S.G.**Bryum capillare** ssp. **torquescens****B. tortuosum** (Hedw.) L. ex With.**Tortella tortuosa****B. triste** C. Muell.**Bryum pertriste****B. truncatum** L. ex With.**Pottia truncata****B. truncatum** (Hedw.) Brotero**Pottia truncata****B. truncorum** (Brid.) Brid. (Bridel, 1819).

Syn. *Mnium truncorum* Brid. (Bridel, 1817).

Distr. Southern South America (Kühnemann, 1938).
Falkland Islands (Wilson and Hooker, 1847).

B. turbinatum (Hedw.) Turn. (Turner, 1804a).

Syn. *Mnium turbinatum* Hedw. (Hedwig, 1801).

Hypnum turbinatum (Hedw.) Web. et Mohr (Weber and Mohr, 1803).

Mnium bimum Schreb. var. *turbinatum* (Hedw.) Wahlenb. (Wahlenberg, 1826).

Distr. Southern South America (Herzog, 1923; Seki, 1974).

ssp. *schleicheri* (Cand.) Kindb.

Bryum schleicheri

var. *patagonicum* Broth. (Brotherus in Herzog, 1923) *nom. nud.*

Distr. Southern South America (Herzog, 1923).

var. *schleicheri* (Cand.) Fuernr.

Bryum schleicheri**B. uliginosum** (Brid.) B.S.G. (Bruch and others, 1839).

Syn. *Cladodium uliginosum* Brid. (Bridel, 1827a).

Distr. Southern South America (Ochi, 1967).

B. unguiculatum (Hedw.) With.**Barbula unguiculata****B. uruguense** C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

B. uspallatense Herz. (Herzog, 1916b).

Distr. Southern South America (Herzog, 1916b).

var. *turgidum* Herz. (Herzog, 1916b).

Distr. Southern South America (Herzog, 1916b).

B. uvidum Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

B. vagans Hook. f. et Wils.**Philonotis vagans****B. valdiviae** Lor. (Lorentz, 1864).

Distr. Southern South America (Lorentz, 1864, 1866; Jaeger, 1875; Brotherus, 1924a).

B. valparaisense Thér. (Thériot, 1917b).

Distr. Southern South America (Thériot, 1917b; Costes, 1921; Brotherus, 1924a; Herzog, 1938).

- B. ventricosum Dicks. ex Relh. *Bryum pseudotriquetrum*
 var. *bimum* (Schreb.) Hamp. *Bryum pseudotriquetrum* ssp. *bimum*
 var. *pseudotriquetrum* (Hedw.) Hamp. *Bryum pseudotriquetrum*
- B. vernicosum Dus. (Dusén, 1903a) *hom. illeg.*
Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Kühnemann, 1938; Herzog, 1957).
- B. violaceum** Crundw. et Nyh. (Crundwell and Nyholm, 1963).
Distr. Southern South America (Crundwell and Nyholm, 1963, 1964).
- B. viridatum** C. Muell. *Pohlia cruda*
- B. vulcanicum** Dus. (Dusén in Paris, 1904a) *nom. nud.*
Distr. Southern South America (Paris, 1904a).
- B. wahlenbergii** (Web. et Mohr) Schwaegr. *Pohlia wahlenbergii*
- B. weberoides** Card. et Broth. (Cardot and Brotherus, 1923).
Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).
- B. weisii** Dicks. ex With. *Ditrichum heteromallum*
B. wilsonii Mitt. *Leptobryum wilsonii*
- B. yendangaianum** Card. *Bryum pseudotriquetrum*
- B. zeballosicum** Card. et Broth. (Cardot and Brotherus, 1923).
Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).
- Bryum** sp.
 Syn. *Hypnum aquicola* C. Muell. (Mueller, 1879a).
Sematophyllum aquicola (C. Muell.) Kindb. (Kindberg, 1888).
Distr. Southern South America (Mueller, 1879a; Paris, 1900a, 1904a; Kurtz, 1904; Hosseus, 1935b; Kühnemann, 1938).
- Bucklandia** *hom. illeg.* *Racomitrium bartramii*
B. bartramii Roiv.
- Bucklandiella** *Racomitrium bartramii*
B. bartramii Roiv.
- Callicostella*** *Hookeriopsis incurva*
C. incurva (Hornschr.) Aongstr.
- C. scabriuscula** (C. Muell.) Jaeg. (Jaeger, 1877).
 Syn. *Hookeria scabriuscula* C. Muell. (Mueller, 1848b).
Distr. Southern South America (Mueller, 1848b, 1851a; Mitten, 1869; Jaeger, 1877; Paris, 1904b; Brotherus, 1925).
- Calliergidium**
C. austro-stramineum (C. Muell.) Bartr. (Bartram, 1946).
 Syn. *Hypnum austro-stramineum* C. Muell. (Mueller, 1890a).
Drepanocladus austro-stramineus (C. Muell.) Broth. ex Par. (Brotherus ex Paris, 1909).
Distr. Southern South America (Bartram, 1946; Seki, 1974).
 South Georgia (Mueller, 1890a; Paris, 1904c; Cardot, 1908a; Brotherus, 1925; Dixon, 1932; Steere, 1961b).

* According to Crosby (1975) the generic name should be *Schizomitrium* B.S.G.

Antarctic, peninsula region (Cardot, 1900, 1906b; Robinson, 1972).

var. *gracillimum* (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1965).

Syn. *Hypnum austro-stramineum* C. Muell. var. *gracillimum* C. Muell. (Mueller, 1890a).
Drepanocladus austro-stramineus (C. Muell.) Broth. ex Par. var. *gracillimus* (C. Muell.) Card. et Broth. (Cardot and Brotherus, 1923).

Distr. South Georgia (Mueller, 1890a; Cardot, 1908a; Dixon, 1932; Steere, 1961b).
 Antarctic, peninsula region (Cardot, 1901, 1908a; Steere, 1961a).

var. *minus* (Card.) Wijk et Marg. (van der Wijk and Margadant, 1965).

Syn. *Hypnum austro-stramineum* C. Muell. var. *minus* Card. (Cardot, 1906b).
Drepanocladus austro-stramineus (C. Muell.) Broth- ex Par. var. *minus* (Card.) Steere (Steere, 1961a).

Distr. Antarctic, peninsula region (Cardot, 1906b, 1907c, 1908a; Steere, 1961a).

var. *subfluitans* (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1965).

Syn. *Hypnum austro-stramineum* C. Muell. var. *subfluitans* C. Muell. (Mueller, 1890a).
Drepanocladus austro-stramineus (C. Muell.) Broth. ex Par. var. *subfluitans* (C. Muell.) Steere (Steere, 1961b).

Distr. South Georgia (Mueller, 1890a; Cardot, 1906a, 1908a; Steere, 1961b).

Calliergon

C. cuspidatum (Hedw.) Kindb.

Calliergonella cuspidata

C. sarmentosum (Wahlenb.) Kindb. (Kindberg, 1894).

Syn. *Hypnum sarmentosum* Wahlenb. (Wahlenberg, 1812).
Amblystegium sarmentosum (Wahlenb.) De Not. (De Notaris, 1867).
Acroceratium sarmentosum (Wahlenb.) Mitt. (Mitten in Reid and Ridley, 1888).
Acrocladium sarmentosum (Wahlenb.) Richs. et Wall. (Richards and Wallace, 1950).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Bartram, 1946).
 South Georgia (Cardot, 1906a, 1908a; Brotherus, 1925; Dixon, 1932; Steere, 1961b; Newton, 1979b).
 Antarctic, peninsula region (Cardot, 1906a, 1908a; Brotherus, 1925; Steere, 1961a; Robinson, 1972).

C. stramineum (Brid.) Kindb. (Kindberg, 1894).

Syn. *Hypnum stramineum* Dicks. ex Brid. (Dickson ex Bridel, 1801b).
Amblystegium stramineum (Brid.) De Not. (De Notaris, 1867).
Acrocladium stramineum (Brid.) Richs. et Wall. (Richards and Wallace, 1950).

Distr. Southern South America (Donat, 1936a; Seki, 1974).

Calliergonella

C. cuspidata (Hedw.) Loesk. (Loeske, 1911).

Syn. *Hypnum cuspidatum* Hedw. (Hedwig, 1801).
Acrocladium cuspidatum (Hedw.) Lindb. (Lindberg, 1879).
Calliergon cuspidatum (Hedw.) Kindb. (Kindberg, 1894).
Cuspidaria fulvo-acuta C. Muell. (Mueller, 1897a).
Hypnum fulvo-acutum (C. Muell.) Par. (Paris, 1900a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904c; Brotherus, 1925; Kühnemann, 1938).

Calliergonella hom. illeg.

C. complanata Card. et Broth.

Catagonium politum

C. nitida (Hook. f. et Wils.) Card.

Catagonium nitidum

Calymperes

C. fasciculatum (Hook. et Grev.) Mitt.

Thyridium fasciculatum

C. glaziovii Hamp. (Hampe, 1878).

Distr. Southern South America (Hosseus, 1940).

Calyptopogon

C. crispatus (C. Muell.) Broth.

Calyptopogon mnioides

C. mnioides (Schwaegr.) Broth. (Brotherus, 1902).

Syn. *Barbula mnioides* Schwaegr. (Schwaegrichen, 1842).

Tortula mnioides (Schwaegr.) Mont. (Montagne, 1850).

Streptopogon mnioides (Schwaegr.) Mitt. (Mitten, 1860).

Syntichia crispata C. Muell. (Mueller, 1897a).

Streptopogon crispatus (C. Muell.) Par. (Paris, 1898).

Barbula crispata C. Muell. ex Par. (Mueller ex Paris, 1900a) nom. inval.

Calyptopogon crispatus (C. Muell.) Broth. (Brotherus, 1902).

Platylomidium crispata C. Muell. (Mueller in Salmon, 1903) nom. nud.

Distr. Southern South America (Schwaegrichen, 1842; Montagne, 1850; Mitten, 1860, 1869; Lorentz, 1866; Jaeger, 1873; Mueller, 1897a; Paris, 1898, 1904a; Brotherus, 1902, 1924a; Salmon, 1903; Dusén, 1906; Cardot, 1908a; Thériot, 1921a; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1954, 1960; Reimers, 1926; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

var. *prostratus* (Mont.) Par.

Tortula prostrata

Camptochaete

C. arbusculans Broth. ex Bartr.

Camptochaete orbiculata

C. gracilis (Hook. f. et Wils.) Par. (Paris, 1894).

Syn. *Hypnum gracile* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844) hom. illeg.

Isothecium gracile Hook. f. et Wils. (Hooker and Wilson in Wilson, 1854).

Stereodon gracilis (Hook. f. et Wils.) Mitt. (Mitten, 1860).

Microthamnium gracile (Hook. f. et Wils.) Jaeg. (Jaeger, 1878).

Porotrichum gracile (Hook. f. et Wils.) Mitt. (Mitten, 1883).

Hypnum gracilescens C. Muell. (Mueller, 1851a) hom. illeg.

Hypnum micans Wils. var. *laxum* Hook. f. (Wilson and Hooker, 1847) fide Mueller, 1851a.

Distr. Southern South America (Wilson and Hooker, 1847; Paris, 1904a; Brotherus, 1907, 1925).

C. orbiculata (Thér.) Robins. (Robinson, 1970b).

Syn. *Weymouthia orbiculata* Thér. (Thériot, 1934b [1935]).

Camptochaete arbusculans Broth. ex Bartr. (Brotherus ex Bartram, 1946) fide Robinson, 1970b.

Distr. Southern South America (Paris, 1904a; Brotherus, 1925; Roivainen, 1934; Thériot, 1934b; Herzog, 1938, 1940, 1957, 1960; Bartram, 1946; Robinson, 1970b).

Camptodontium

C. brotheri Dus.

Camptodontium cryptodon

var. *grandirete* Dus. (Dusén, 1907).

Distr. Southern South America (Dusén, 1907; Cardot, 1908a).

C. cryptodon (Mont.) Reim. (Reimers, 1936).

Syn. *Weisia cryptodon* Mont. (Montagne, 1845a).

Dicranum cryptodon (Mont.) Mitt. (Mitten, 1869).

Dicranoweisia cryptodon (Mont.) Par. (Paris, 1895).

Grimmia cryptodon (Mont.) Hamp. (Hampe in Reimers, 1936) nom. inval.

Dicranoweisia tenella Dus. (Dusén, 1903a) nom. nud.

Camptodontium brotheri Dus. (Dusén, 1905b).

Bryopoteria chilensis Thér. (Thériot, 1933).

Distr. Southern South America (Montagne, 1845a, 1850; Mueller, 1849; Mitten, 1869; Jaeger,

1870; Paris, 1895, 1904a, b; Neger, 1899; Dusén, 1903a, 1905b, 1907; Cardot, 1908a; Campo, 1921; Brotherus, 1924a; Reimers, 1926, 1936; Thériot, 1933; Röiväinen, 1934; Herzog, 1938, 1954, 1957, 1960; Seki, 1974).

Campylium

C. polygamum (B.S.G.) C. Jens. (Jensen in Lange and Jensen, 1887).

Syn. *Amblystegium polygamum* B.S.G. (Bruch and others, 1853).

Hypnum polygamum (B.S.G.) Wils. (Wilson, 1855).

Chrysohypnum polygamum (B.S.G.) Loesk. (Loeske, 1903).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938; Seki, 1974).

C. radicale (P. Beauv.) Grout (Grout, 1929).

Syn. *Hypnum radicale* P. Beauv. (Palisot de Beauvois, 1805a).

Amblystegium radicale (P. Beauv.) Mitt. (Mitten, 1869) *hom. illeg.*

Distr. Southern South America (Bizot and Piovano, 1953; Piovano, 1954).

C. squarroso-byssoides C. Muell. (Mueller, 1897a).

Syn. *Hypnum squarroso-byssoides* (C. Muell.) Par. (Paris, 1900a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1925; Kühnemann, 1938).

Campylospodium

C. flaccidum Herz. (Herzog, 1926).

Distr. Southern South America (Herzog, 1926; Seki, 1974).

Campylopus

C. aberrans Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Frahm, 1975a, 1976a).

var. viridis Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b; Frahm, 1975a).

C. acuminatus Mitt. (Mitten, 1869).

Syn. *Dicranum acuminatum* (Mitt.) C. Muell. (Mueller, 1885).

Distr. Southern South America (Mitten, 1869; Jaeger, 1872; Mueller, 1885; Paris, 1904a; Cardot, 1908a; Frahm, 1975a, 1976a).

C. alopecurus (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Dicranum alopecurus* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938; Frahm, 1975a).

C. amabile (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Dicranum amabile* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Hosseus, 1938c, d; Kühnemann, 1938; Frahm, 1975a).

C. arctocarpus (Hornsch.) Mitt. (Mitten, 1869).

Syn. *Dicranum arctocarpum* Hornsch. (Hornschuch, 1840).

Thysanomitrion scabrisetum Hamp. (Hampe, 1844).

Distr. Southern South America (Hornschuch, 1840; Hampe, 1844; Mueller, 1848a; Mitten, 1869; Jaeger, 1872; Gibert, 1873; Paris, 1904a; Brotherus, 1924a; Herter, 1933a; Frahm, 1975a).

- C. arenicola* (C. Muell.) Mitt. *Campylopus surinamensis*
- C. areodictyon* (C. Muell.) Mitt. (Mitten, 1869).
- Syn. *Dicranum areodictyon* C. Muell. (Mueller, 1848a).
- Distr. Juan Fernandez (Brotherus, 1924b; Espinosa, B., 1941; Frahm, 1975a, 1976a).
- C. asperifolius* Mitt. *Dicranodontium asperifolium*
- C. atratus* Schimp. *Campylopus patagonicus*
- C. aurificus* (C. Muell.) Kindb. *Campylopus introflexus*
- var. *flavescens*** (C. Muell.) Par. (Paris, 1894).
- Syn. *Dicranum auricum* C. Muell. var. *flavescens* C. Muell. (Mueller, 1882).
- Distr. Southern South America (Mueller, 1882; Paris, 1894).
- C. austro-alpinus* (C. Muell.) Kindb. (Kindberg, 1889).
- Syn. *Dicranum austro-alpinum* C. Muell. (Mueller, 1882).
- Distr. Southern South America (Mueller, 1882; Paris, 1904a; Kühnemann, 1938; Frahm, 1975a).
- C. berteroanus* Dub. *Campylopus introflexus*
- C. berteroii* Dub. ex Kindb. *Campylopus introflexus*
- C. birgeri* Card. (Cardot, 1905a).
- Distr. Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Frahm, 1975a).
- C. blindiaceus* C. Muell. (Mueller, 1901a [1900]) *nom. nud.*
- Distr. Southern South America (Mueller, 1901a).
- C. blindioides* Broth. (Brotherus, 1924b).
- Distr. Juan Fernandez (Brotherus, 1924b, 1925; Frahm, 1975a, 1976a; Robinson, 1975).
- C. brachythysanos* (C. Muell.) Par. (Paris, 1900a).
- Syn. *Dicranum brachythysanos* C. Muell. (Mueller, 1897a).
- Distr. Southern South America (Mueller, 1897a; Paris, 1900a; Frahm, 1975a).
- C. cacti* (C. Muell.) Kindb. (Kindberg, 1889).
- Syn. *Dicranum cacti* C. Muell. (Mueller, 1882).
- Distr. Southern South America (Mueller, 1882; Paris, 1904a; Kühnemann, 1938; Frahm, 1975a).
- C. campanus* Thér. (Thériot, 1918).
- Distr. Southern South America (Thériot, 1918; Campo, 1921; Brotherus, 1924a; Herzog, 1938, 1939; Frahm, 1975a, 1976a).
- C. canescens* (C. Muell.) Schimp. (Schimper in Jaeger, 1872).
- Syn. *Dicranum canescens* C. Muell. (Mueller, 1859).
- Dicranum perincanum* C. Muell. (Mueller, 1885) *fide* Frahm, 1975a.
- Campylopus perincanus* (C. Muell.) Kindb. (Kindberg, 1888) *fide* Frahm, 1975a.
- Distr. Southern South America (Mueller, 1885; Paris, 1904a; Cardot, 1905a, 1908a; Kühnemann, 1938; Frahm, 1975a, 1976a).
- Falkland Islands (Mueller, 1859; Jaeger, 1872; Paris, 1904a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938; Frahm, 1975a, 1976a).
- C. carboniculus* Thér. (Thériot, 1918).
- Distr. Southern South America (Thériot, 1918; Campo, 1921; Frahm, 1975a, 1976a).

C. carinatus C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a; Brotherus, 1924a; Kühnemann, 1938; Frahm, 1975a).

C. chalarobasis C. Muell. ex Thér. (Mueller ex Thériot, 1934b [1935]).

Distr. Southern South America (Thériot, 1934b; Frahm, 1975a, 1976a).

C. chilensis De Not. (De Notaris, 1859).

Distr. Southern South America (De Notaris, 1859; Mitten, 1869; Jaeger, 1872; Paris, 1904a; Brotherus, 1924a; Frahm, 1975a).

C. chrismarii (C. Muell.) Mitt. (Mitten, 1869).

Syn. *Dicranum chrismarii* C. Muell. (Mueller, 1855).

Distr. Juan Fernandez (Frahm, 1975a).

C. clavatus (R. Brown) Wils. (Wilson, 1854).

Syn. *Dicranum clavatum* R. Brown (R. Brown in Schwaegrichen, 1829).

Campylopus leptodus Mont. (Montagne, 1845a).

Trichostomum leptodum (Mont.) Mitt. (Mitten in Hooker, 1867).

Pilopogon leptodus (Mont.) Broth. (Brotherus, 1901).

Thysanomitrion leptodus (Mont.) Broth. (Brotherus in Skottsberg, 1914).

Campylopus corralensis Broth. (Brotherus in Paris, 1904a) *nom. nud.*

Campylopus montagnei De Not. (De Notaris, 1859) *fide* Frahm, 1975a.

Distr. Juan Fernandez (Mitten, 1885; Paris, 1900a, 1905a; Skottsberg, 1914; Brotherus, 1924b; Bartram, 1965; Frahm, 1975a, 1976a; Robinson, 1975).
Southern South America (Montagne, 1845a, 1850, 1856; De Notaris, 1859; Mitten, 1869; Jaeger, 1872; Brotherus, 1901, 1924a; Paris, 1904a, 1905a; Dusén, 1905a; Cardot, 1908a; Herzog, 1923, 1938; Thériot, 1925a, 1935a; Bartram, 1952; Frahm, 1975a, 1976a, b).

C. concolor (Hook.) Brid. (Bridel, 1826).

Syn. *Dicranum concolor* Hook. (Hooker, 1816).

Distr. Southern South America (Bartram, 1965; Frahm, 1975a).

C. corralensis Broth.

Campylopus clavatus

C. crassissimus Besch. (Bescherelle, 1885a).

Syn. *Campylopus sordido-nigritus* Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Bescherelle, 1885a, 1889; Dusén, 1903a, 1905c; Paris, 1904a; Cardot, 1908a; Brotherus, 1924a; Roivainen, 1934; Frahm, 1975a, 1976a).

C. cummingii Dub. (Duby, 1870).

Distr. Southern South America (Duby, 1870; Jaeger and Sauerbeck, 1879b; Paris, 1904a).

C. curvatifolius Card. (Cardot, 1905a).

Distr. Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Frahm, 1975a).

C. fernandezianus Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Southern South America (Skottsberg, 1914).

C. fernandezianus Bartr. (Bartram, 1959).

Distr. Juan Fernandez (Bartram, 1959; Frahm, 1975a, 1976a).

C. fibrobasis Dus.

Campylopus torquatus

C. flavissimus (C. Muell.) Besch. (Bescherelle, 1885a).

Syn. *Dicranum flavissimum* C. Muell. (Mueller, 1885).

Distr. Southern South America (Bescherelle, 1885a, 1889; Mueller, 1885; Cardot, 1901, 1908a; Paris, 1904a; Dusén, 1905c; Brotherus, 1924a; Frahm, 1975a, 1976a).

C. flavo-nigritus Dus. (Dusén, 1905c).

Distr. Southern South America (Dusén, 1903a, 1905c; Paris, 1904a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Frahm, 1975a, 1976a).

C. flavo-viridis Dus. (Dusén, 1905c).

Distr. Southern South America (Paris, 1904a; Dusén, 1905c; Cardot, 1908a; Seki, 1974; Frahm, 1975a, 1976a).

C. flexuosus (Hedw.) Brid. (Bridel, 1819).

Syn. *Dicranum flexuosum* Hedw. (Hedwig, 1801).

Dicranum flexuosum Hedw. var. *minus* Schwaegr. (Schwaegrichen in Gaudichaud, 1826 [1828]) *nom. nud.*

Distr. Southern South America (Montagne, 1850).

Falkland Islands (Dumont D'Urville, 1825; Gaudichaud, 1825, 1826; Bridel, 1827b).

C. fuegianus Dus. (Dusén, 1905c).

Distr. Southern South America (Dusén, 1905c; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Frahm, 1975a, 1976a).

C. fuscatus Besch. (Bescherelle, 1891).

Distr. Southern South America (Bescherelle, 1891; Paris, 1904a; Felippone, 1912; Herter, 1933a; Frahm, 1975a).

C. fusco-luteus Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

C. gaudichaudii Besch. (Bescherelle, 1891).

Distr. Southern South America (Bescherelle, 1891; Paris, 1904a; Frahm, 1975a).

C. grimmiooides (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Dicranum grimmiooides* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938; Frahm, 1975a).

C. griseus (Hornschr.) Jaeg. (Jaeger, 1872).

Syn. *Thysanomitrion griseum* Hornsch. (Hornschuch, 1840).

Dicranum griseum (Hornsch.) C. Muell. (Mueller, 1848a).

Dicranum recurvipilum C. Muell. (Mueller, 1900).

Campylopus recurvipilus (C. Muell.) Par. (Paris, 1904a).

Distr. Southern South America (Hornschuch, 1840; Mueller, 1848a; Jaeger, 1872; Paris, 1904a; Felippone, 1912; Herter, 1933a, 1943; Frahm, 1975a).

C. guaitecae Dus. (Dusén, 1905c).

Distr. Juan Fernandez (Frahm, 1976a).

Southern South America (Paris, 1904a; Dusén, 1905c; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Thériot, 1934b; Roivainen and Bartram, 1937; Kühnemann, 1938; Frahm, 1975a, 1976a).

C. hamatus Bartr. (Bartram in Roivainen and Bartram, 1937).

Distr. Southern South America (Roivainen and Bartram, 1937; Frahm, 1975a, 1976a).

C. humifugus (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Dicranum humifugum* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938; Frahm, 1975a).

C. incrassatus C. Muell. (Mueller, 1844b [1845]).

Syn. *Dicranum incrassatum* (C. Muell.) C. Muell. (Mueller, 1848a).

Distr. Southern South America (Mueller, 1844b, 1847c, 1848a; Montagne, 1850; Mitten, 1869; Jaeger, 1872; Paris, 1904a; Dusén, 1905c; Thériot, 1917b; Costes, 1921; Brotherus, 1924a; Espinosa B., 1924; Herzog, 1938, 1960; Frahm, 1975a, 1976a).

C. introflexus (Hedw.) Brid. (Bridel, 1819).

Syn. *Dicranum introflexum* Hedw. (Hedwig, 1801).

Thysanomitrion introflexum (Hedw.) Arnott (Arnott, 1827).

Campylopus introflexus (Hedw.) Brid. var. *genuinus* Giac. (Giacomini, 1955) hom. illeg.

Campylopus xanthophyllum Mont. (Montagne, 1845a).

Dicranum xanthophyllum Mont. ex Par. (Montagne ex Paris, 1895).

Dicranum aurificum C. Muell. (Mueller, 1882) fide Frahm, 1975a.

Campylopus aurificus (C. Muell.) Kindb. (Kindberg, 1889) fide Frahm, 1975a.

Campylopus lechleri Schimp. (Schimper in Lorentz, 1866) nom. nud. fide Frahm, 1975a.

Campylopus berteroanus Dub. (Duby, 1875b).

Campylopus berteroii Dub. ex Kindb. (Duby ex Kindberg, 1888) hom. illeg.

Campylopus rectipilus C. Muell. (Mueller in Kindberg, 1891) nom. nud.

Campylopus nodicaulis C. Muell. (Mueller in Frahm, 1975a) nom. nud.

Campylopus latipilis C. Muell. (Mueller in Kindberg, 1891) nom. nud. fide Frahm, 1975a.

Distr. Juan Fernandez (Montagne, 1835; Duby, 1875a, b; Jaeger and Sauerbeck, 1879b; Johow, 1896 (as *C. leptodus* fide Brotherus, 1924b); Paris, 1904a; Skottsberg, 1914; Brotherus, 1924b; Thériot, 1927b; Espinosa B., 1941; Bartram, 1959; Frahm, 1974, 1975a; Robinson, 1975).

Southern South America (Hornschuch, 1840; Mueller, 1844b; Montagne, 1845a, 1846, 1850, 1856; Lorentz, 1866; Mitten, 1869; Jaeger, 1872; Gibert, 1873; Paris, 1895, 1904a; Dusén, 1905c; Cardot 1908a; Campo, 1915, 1921; Thériot, 1918; Brotherus, 1924a, c; Reimers, 1926; Herter, 1928, 1933a, b, 1939b; Roivainen and Bartram, 1937; Hosseus, 1938c, d; Herzog, 1938, 1939, 1954, 1960; Espinosa B., 1941; Skottsberg, 1950; Bartram, 1952; Giacomini, 1955; Bizot, 1967; Seki, 1974; Frahm, 1974, 1975a, 1976a, b).

Falkland Islands (Wilson and Hooker, 1847; Mueller, 1848a; Jaeger, 1872; Cardot, 1905a; 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938; Giacomini, 1955; Frahm, 1974, 1975a).

var. *brachycarpus* Giac. (Giacomini, 1955).

Distr. Southern South America (Giacomini, 1955; Frahm, 1975a).

var. *cordobaensis* Thér.

Campylopus pilifer

var. *genuinus* Giac.

Campylopus introflexus* var. *introflexus

C. jamesonii (Hook.) Jaeg. (Jaeger, 1872).

Syn. *Dicranum jamesonii* Hook. (Hooker, 1837).

Distr. Southern South America (Frahm, 1975a).

C. jugorum Herz. (Herzog, 1909).

Distr. Southern South America (Hosseus, 1938c, d, 1939; Frahm, 1975a).

C. kunkelli Bartr. (Bartram, 1959).

Distr. Juan Fernandez (Bartram, 1959; Frahm, 1975a, 1976a; Robinson, 1975).

C. laetivirens (Broth.) Frahm (Frahm, 1975a) hom. illeg.

Syn. *Thysanomitrion laetevirens* Broth. (Brotherus, 1924c) fide Frahm, 1975a.

Distr. Southern South America (Frahm, 1975a, b, 1976a).

C. laniger (C. Muell.) Besch.

Chorisodontium lanigerum

C. latipilus C. Muell.

Campylopus introflexus

C. laxiretis Bartr.

Campylopus masafuerae

<i>C. lechleri</i> Schimp.	<i>Campylopus introflexus</i>
<i>C. leptodus</i> Mont.	<i>Campylopus clavatus</i>
<i>C. lorentzii</i> C. Muell. ex Par.	<i>Conostomum lorentzii</i>
<i>C. maulensis</i> Broth. (Brotherus <i>in Espinosa</i> B., 1941) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Espinosa B., 1941).	
<i>C. masafuerae</i> Frahm (Frahm, 1975a).	
<i>Syn.</i> <i>Campylopus laxiretis</i> Bartr. (Bartram, 1959) <i>hom. illeg. fide</i> Frahm, 1975a.	
<i>Distr.</i> Juan Fernandez (Bartram, 1959; Frahm, 1975a, 1976a).	
<i>C. modestus</i> Card. (Cardot, 1905a).	
<i>Distr.</i> Southern South America (Herzog, 1938, 1939; Frahm, 1975a, 1976a).	
Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Frahm, 1975a, 1976a).	
var. <i>deciduus</i> Card. et Broth. (Cardot and Brotherus, 1923).	
<i>Distr.</i> Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938; Frahm, 1975a).	
<i>C. montagnei</i> De Not.	<i>Campylopus clavatus</i>
<i>C. morenoi</i> (C. Muell.) Par. (Paris, 1900a).	
<i>Syn.</i> <i>Dicranum morenoi</i> C. Muell. (Mueller, 1897a).	
<i>Distr.</i> Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Cardot, 1908a; Frahm, 1975a).	
<i>C. mucorinus</i> C. Muell. (Mueller, 1901a [1900]) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Mueller, 1901a).	
<i>C. muricatus</i> Dix.	<i>Chorisodontium aciphyllum</i>
<i>C. nanocaudatus</i> C. Muell.	<i>Chorisodontium aciphyllum</i>
<i>C. nanotrichus</i> C. Muell. (Mueller, 1901a [1900]) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Mueller, 1901a).	
<i>C. nitidus</i> Thér.	<i>Campylopus urugensis</i>
<i>C. nodicaulis</i> C. Muell.	<i>Campylopus introflexus</i>
<i>C. occultoviridis</i> Dus. (Dusén, 1903a) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Dusén, 1903a).	
<i>C. orthocomus</i> (C. Muell.) Besch.	<i>Chorisodontium aciphyllum</i>
<i>C. patagonicus</i> Broth. ex Dus. (Brotherus <i>in Dusén</i>, 1905c).	
<i>Syn.</i> <i>Campylopus atratus</i> Schimp. (Schimper <i>in Mueller</i> , 1901a [1900]) <i>nom. nud. fide</i> Frahm, 1975a.	
<i>Distr.</i> Southern South America (Dusén, 1903a, 1905c; Paris, 1904a; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974; Frahm, 1975a, 1976a).	
Falkland Islands (Mueller, 1901a).	
<i>C. perhorridus</i> Dus. (Dusén, 1905c).	
<i>Syn.</i> <i>Dicranum perhorridum</i> (Dus.) Card. (Cardot, 1908a).	
<i>Distr.</i> Southern South America (Dusén, 1905c; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Herzog, 1938; Kühnemann, 1938; Frahm, 1975a, 1976a).	
<i>C. perincanus</i> (C. Muell.) Kindb.	<i>Campylopus canescens</i>
<i>C. permollis</i> Thér. (Thériot, 1926).	
<i>Distr.</i> Southern South America (Thériot, 1926, 1934b; Frahm, 1975a, 1976a).	

C. pilifer Brid. (Bridel, 1819).

Syn. *Campylopus polytrichoides* De Not. (De Notaris, 1838b) *fide* Gradstein and Sipman, 1978.

Campylopus introflexus (Hedw.) Brid. var. *cordobaensis* Thér. (Thériot *in* Bauer, 1934) *fide* Gradstein and Sipman, 1978.

Distr. Juan Fernandez (Brotherus, 1924b).

Southern South America (Bauer, 1934; Hosseus, 1936, 1937, 1938c, d, 1939; Kühnemann, 1938; Espinosa B., 1941; Frahm, 1975a; Gradstein and Sipman, 1978).

C. pilopogon C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

C. polytrichoides De Not.***Campylopus pilifer*****C. purpureocaulis** Dus. (Dusén, 1905c).

Distr. Southern South America (Dusén, 1903a, 1905c; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1954; Seki, 1974; Frahm, 1975a, 1976a).

C. rectipilus C. Muell.***Campylopus introflexus*****C. recurvifolius** Dus. (Dusén, 1905c).

Distr. Southern South America (Dusén, 1905c; Brotherus, 1924a; Herzog, 1938, 1960; Frahm, 1975a, 1976a).

C. recurvipilus (C. Muell.) Par.***Campylopus griseus*****C. richardii** Brid. (Bridel, 1819).

Syn. *Thysanomitrion richardii* (Brid.) Schwaegr. (Schwaegrichen, 1823).

Weisia richardii (Brid.) Schwaegr. (Schwaegrichen *in* Bridel, 1826) *nom. inval.*

Grimmia richardii (Brid.) Spreng. (Sprengel, 1827).

Dicranum richardii (Brid.) C. Muell. (Mueller, 1848a).

Pilopogon richardii (Brid.) Broth. (Brotherus, 1901).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Frahm, 1975a, 1976a).

Southern South America (Mueller, 1848a; Paris, 1905a).

C. rigescens Kindb. (Kindberg, 1889) *nom. nud.*

Distr. No locality given; see *Index muscorum*, 1, p. 453.

C. saddleanus (C. Muell.) Besch.***Campylopus vesticaulis*****C. sancti-caroli** Besch. (Bescherelle, 1891).

Distr. Southern South America (Bescherelle, 1891; Paris, 1904a; Frahm, 1975a).

C. scaberimus Broth.***Chorisodontium aciphyllum*****C. scabrophylloides** (C. Muell.) Par. (Paris, 1900a).

Syn. *Dicranum scabrophylloides* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Felippone, 1912; Herter, 1933a; Frahm, 1975a).

C. senilis Lor. (Lorentz *in* Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

C. sordido-nigritus Dus.***Campylopus crassissimus*****C. spegazzinii** (C. Muell.) Kindb.***Chorisodontium spegazzinii*****C. spiralis** Dus. (Dusén, 1905c).

Distr. Southern South America (Paris, 1904a; Dusén, 1905c; Cardot, 1908a; Thériot, 1934b, 1935a; Herzog, 1939; Frahm, 1975a, 1976a).

C. stenocarpus (Hamp.) Jaeg. (Jaeger, 1872).

Syn. *Thysanomitrion stenocarpum* Hamp. (Hampe, 1844).

Dicranum stenocarpum (Hamp.) C. Muell. (Mueller, 1851a).

Dicranum stenopus C. Muell. (Mueller, 1848a) *nom. illeg.*

Distr. Southern South America (Hampe, 1844; Mueller, 1848a; Jaeger, 1872; Paris, 1904a; Felippone, 1912; Herter, 1933a; Frahm, 1975a).

C. subareodictyon Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Frahm, 1975a, 1976a).

C. subexalaris C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

C. substygius C. Muell. (Mueller in Kindberg, 1889) *nom. nud.*

Distr. No locality given, see *Index muscorum*, 1, 458.

C. sulfureo-nigritus Dus. (Dusén, 1905c).

Distr. Southern South America (Dusén, 1903a, 1905c; Paris, 1904a; Cardot, 1908a; Brotherus, 1924a; Frahm, 1975a, 1976a).

C. surinamensis C. Muell. (Mueller, 1848b).

Syn. *Dicranum arenicola* C. Muell. (Mueller, 1855).

Campylopus arenicola (C. Muell.) Mitt. (Mitten, 1869).

Distr. Southern South America (Herter, 1939b; Frahm, 1975a).

C. torquatus Mitt. (Mitten in Wilson, 1859).

Syn. *Campylopus fibrobasis* Dus. (Dusén, 1905c) *fide* Frahm, 1975a.

Distr. Southern South America (Paris, 1904a; Dusén, 1905c; Cardot, 1908a; Brotherus, 1924a; Thériot, 1934b; Herzog, 1938, 1954; Kühnemann, 1938; Frahm, 1975a, 1976a).

C. truncatus C. Muell. (Mueller, 1844b [1845]).

Syn. *Dicranum truncatum* (C. Muell.) C. Muell. (Mueller, 1848a).

Distr. Juan Fernandez (Skottsberg, 1914; Thériot, 1921b; Brotherus, 1924b; Frahm, 1976a). Southern South America (Mueller, 1844b, 1848a; Jaeger, 1872; Paris, 1904a; Reimers, 1926; Thériot, 1934b; Herzog, 1938; Frahm, 1975a, 1976a).

C. urugensis Frahm (Frahm, 1975a).

Syn. *Campylopus nitidus* Thér. (Thériot in Felippone, 1929 [1930]) *hom. illeg. fide* Frahm, 1975a.

Distr. Southern South America (Felippone, 1929; Herter, 1933a; Frahm, 1975a).

C. ventanus C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

C. verticillatus (Hamp.) Kindb. (Kindberg, 1889).

Syn. *Dicranum verticillatum* Hamp. (Hampe, 1879).

Distr. Southern South America (Hampe, 1879; Paris, 1894, 1904a; Frahm, 1975a).

C. vesticaulis Mitt. (Mitten, 1875).

Syn. *Dicranum saddleanum* Besch. ex C. Muell. (Mueller, 1885).

Campylopus saddleanus (C. Muell.) Besch. (Bescherelle, 1885a).

Distr. Southern South America (Bescherelle, 1885a, 1889; Mueller, 1885; Paris, 1904a; Cardot, 1908a; Brotherus, 1924a; Roivainen and Bartram, 1937; Frahm, 1975a, 1976a).

Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Frahm, 1975a, 1976a).

C. xanthophyllus Mont. (Montagne, 1845a).

Campylopus introflexus

Catagoniopsis

C. berteroana (Mont.) Broth. (Brotherus, 1909).

<i>Catharinea nom. rejic.</i>	
<i>C. antarctica</i> C. Muell.	<i>Psilopilum trichodon</i>
<i>C. canaliculata</i> (Hook. et Arnott) C. Muell.	<i>Oligotrichum canaliculatum</i>
var. <i>major</i> Lor.	<i>Oligotrichum canaliculatum</i> var. <i>majus</i>
<i>C. compressa</i> (Hook. f. et Wils.) C. Muell.	<i>Atrichopsis compressa</i>
<i>C. dendroides</i> (Hedw.) Hamp.	<i>Dendroligotrichum dendroides</i>
<i>C. gymnostoma</i> C. Muell. ex Par.	<i>Psilopilum gymnostomulum</i>
<i>C. gymnostomula</i> C. Muell.	<i>Psilopilum gymnostomulum</i>
<i>C. magellanica</i> (Hedw.) Brid.	<i>Polytrichadelphus magellanicus</i>
<i>C. molinae</i> (Mont.) C. Muell.	<i>Oligotrichum canaliculatum</i> var. <i>molinae</i>
<i>C. planifolia</i> C. Muell.	<i>Atrichum polycarpum</i>
<i>C. polycarpa</i> C. Muell.	<i>Atrichum polycarpum</i>
<i>C. rigida</i> (Lor.) C. Muell.	<i>Oligotrichum rigidum</i>
<i>C. simplex</i> C. Muell. (Mueller, 1901a [1900]) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Mueller, 1901a).	
<i>C. squamosa</i> (Hook. f. et Wils.) C. Muell.	<i>Dendroligotrichum squamosum</i>
<i>C. tapes</i> C. Muell.	<i>Psilopilum tapes</i>
<i>Cecalyphum nom. illeg.</i>	
<i>C. dichotomum</i> P. Beauv.	<i>Dicranoloma billardieri</i>
<i>C. semi-completum</i> (Hedw.) P. Beauv.	<i>Fissidens semicompletus</i>
<i>Cephalotrichum hom. illeg.</i>	
<i>C. oligodus</i> B.S.G.	<i>Pogonatum oligodus</i>
<i>Ceratodon</i>	
<i>C. amblyocalyx</i> C. Muell. ex Ångstr.	<i>Ceratodon purpureus</i> var. <i>amblyocalyx</i>
<i>C. antarcticus</i> Card. (Cardot, 1900).	
<i>Distr.</i> Antarctic, peninsula region (Cardot, 1900, 1901, 1908a; Paris, 1904a; Dixon, 1935; Horikawa and Ando, 1961, 1963; Steere, 1961a).	
<i>C. chilensis</i> (Mont.) Par.	<i>Cheilotrichia chilensis</i>
<i>C. convolutus</i> Reichdt.	<i>Ceratodon purpureus</i>
<i>C. corralensis</i> (Lor.) Jaeg.	<i>Zygodon corralensis</i>
<i>C. crassinervis</i> Lor. (Lorentz, 1866).	
<i>Distr.</i> Southern South America (Lorentz, 1866; Paris, 1904a; Thériot, 1918, 1921a, 1935a; Campo, 1921; Reimers, 1926; Herzog, 1938).	
<i>C. dubius</i> Dus.	<i>Cheilotrichia chilensis</i>
<i>C. grossiretis</i> Card. (Cardot, 1906a).	
<i>Distr.</i> Southern South America (Thériot, 1926). South Georgia (Dixon, 1920; Steere, 1961b; Horikawa and Ando, 1963). Antarctic, peninsula region (Cardot, 1906a, 1908a, 1911b, d, 1913a; Horikawa and Ando, 1961, 1963; Steere, 1961a; Robinson, 1972). Antarctic, continental region (Bartram, 1957; Steere, 1961a; Horikawa and Ando, 1963).	
var. <i>validus</i> Card.	<i>Ceratodon validus</i>
<i>C. minutifolius</i> Card. (Cardot, 1911b, d).	
<i>Distr.</i> Antarctic, peninsula region (Cardot, 1911b, d, 1913a; Steere, 1961a; Horikawa and Ando, 1963).	

C. purpureus (Hedw.) Brid. (Bridel, 1826).

Syn. *Dicranum purpureum* Hedw. (Hedwig, 1801).

Ceratodon purpureus (Hedw.) Brid. var. *xanthopus* Sull. (Sullivant in Sullivant and Lesquereux, 1865).

Ceratodon purpureus (Hedw.) Brid. var. *fastigiatus* Warnst. (Warnstorff, 1904).

Ceratodon convolutus Reichdt. (Reichardt, 1868).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Robinson, 1975).

Southern South America (Montagne, 1845b, 1850; Wilson and Hooker, 1847; Lorentz, 1866; Mitten, 1869; Cardot, 1900, 1905a, 1908a; Krieger, 1904; Dusén, 1905a; Thériot, 1918, 1935a; Campo, 1921; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1954, 1957; Roivainen, 1934; Roivainen and Bartram 1937; Hosseus, 1938c, d; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

Falkland Islands (Dumont d'Urville, 1825; Gaudichaud, 1825, 1826; Bridel, 1827a; Wilson and Hooker, 1847; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938).

Antarctic, peninsula region (Cardot, 1900, 1901, 1906a, 1908a, 1911b, d, 1913a; Paris, 1904a; Dixon and Watts, 1918; Bartram, 1957; Steere, 1961a; Robinson, 1972).

Antarctic, continental region (Cardot, 1907a, 1908a; Steere, 1961a; Horikawa, 1961; Horikawa and Ando, 1961, 1967; Kuc, 1969; Robinson, 1972).

var. *amblyocalyx* (Aongstr.) C. Muell. (Mueller, 1885).

Syn. *Ceratodon amblyocalyx* C. Muell. ex Aongstr. (Mueller ex Ångström, 1876a).

Distr. Juan Fernandez (Skottsberg, 1914).

Southern South America (Ångström, 1872 (as *C. purpureus*), 1876a; Jaeger and Sauerbeck, 1879b; Mueller, 1885; Bescherelle, 1889; Cardot, 1900, 1901, 1905a, 1908a; Paris, 1904a; Dusén, 1905a, d; Kühnemann, 1938).

Falkland Islands (Cardot, 1905a, 1908a; Kühnemann, 1938).

var. *falklandicus* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

var. *fastigiatus* Warnst.

Ceratodon purpureus

var. *xanthopus* Sull.

Ceratodon purpureus

C. valdiviae C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

C. validus (Card.) Hor. et Ando (Horikawa and Ando, 1963).

Syn. *Ceratodon grossiretis* Card. var. *validus* Card. (Cardot, 1906a) *fide* Horikawa and Ando, 1963.

Distr. Antarctic, peninsula region (Cardot, 1906a, 1908a; Horikawa and Ando, 1961, 1963; Greene, 1968a).

Chaetephora hom. illeg.**C. cristata** (Hedw.) Brid.

Eriopus cristatus

C. incurva Hornsch.

Hookeriopsis incurva

C. orthotrichoides (Raddi) Brid.

Macrocoma orthotrichoides

Cheilotrichia**C. chilensis** (Mont.) Broth. (Brotherus, 1901).

Syn. *Trichostomum chilense* Mont. (Montagne, 1845a).

Ceratodon chilensis (Mont.) Par. (Paris, 1904a).

Chrysoblastella chilensis (Mont.) Reim. (Reimers, 1926).

Ceratodon dubius Dus. (Dusén in Paris, 1904a) *nom. nud.*

Cheilotrichia dubia Dus. (Dusén, 1905a) *fide* Newton, 1977a.

Chrysoblastella dubia (Dus.) Reim. (Reimers, 1926).

Chrysoblastella revoluta Herz. (Herzog, 1916a) *fide* Newton, 1977a.

Cheilotrichia revoluta (Herz.) Thér. (Thériot, 1934b [1935]).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Jaeger, 1873; Brotherus, 1901, 1924a; Paris, 1904a; Dusén, 1905a; Reimers, 1926; Hilpert, 1933;

Thériot, 1934b; Herzog, 1938, 1954; Newton, 1974b; Seki, 1974).
South Georgia (Newton, 1977a).

C. dubia Dus.

Cheilotrichia chilensis

C. revoluta (Herz.) Thér.

Cheilotrichia chilensis

Chorisodontium

C. aciphyllum (Hook. f. et Wils.) Broth. (Brotherus, 1924a).

Syn. *Dicranum aciphyllum* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Dicranum rigens Besch. (Bescherelle, 1885a).

Dicranoloma rigens (Besch.) Par. (Paris, 1904b).

Dicranum leucopterum C. Muell. (Mueller, 1885).

Chorisodontium leucopterum (C. Muell.) Roiv. (Rovainen in Rovainen and Bartram, 1937).

Dicranum orthocomus Besch. ex C. Muell. (Bescherelle ex Mueller, 1885).

Campylopus orthocomum (C. Muell.) Besch. (Bescherelle, 1885a).

Chorisodontium orthocomum (C. Muell.) Roiv. (Rovainen in Rovainen and Bartram, 1937).

Dicranum tenuicuspitatum C. Muell. (Mueller, 1890a).

Campylopus nanocaudatus C. Muell. (Mueller, 1901a [1900]) nom. nud. fide Bell, 1977a.

Dicranum scaberrimum Dus. (Dusén, 1903a).

Campylopus scaberrimus Broth. (Brotherus in Dusén, 1903a) nom. nud.

Dicranum nordenskioeldii Card. (Cardot, 1906a) fide Bell, 1973a.

Chorisodontium nordenskioeldii (Card.) Roiv. (Rovainen in Rovainen and Bartram, 1937) fide Bell, 1973a.

Campylopus muricatus Dix. (Dixon, 1932) fide Dixon, 1934 and Bell, 1977a.

Dicranum atro-viride Card. (Cardot in Rovainen and Bartram, 1937).

Dicranum hatscheri Dus. (Dusén in Rovainen and Bartram, 1937) nom. nud.

Distr. Juan Fernandez (Bartram, 1959; Robinson, 1975).

Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Sullivant, 1859; Mitten, 1869; Jaeger, 1872; Bescherelle, 1885a, 1889; Cardot, 1900, 1901, 1905a, 1908a, 1913a; Dusén, 1903a, 1905a, d; Paris, 1904a, b; Spagazzini, 1922; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

Falkland Islands (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Paris, 1904b; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

South Georgia (Mueller, 1890a, 1901a; Paris, 1904b; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Dixon, 1932; Steere, 1961b; Bell, 1973b; Frahm, 1975a).

Antarctic, peninsula region (Wright, 1905 (as *Campylopus introflexus* fide Cardot, 1908a); Cardot, 1906a, 1908a, 1911b, c, d, 1912b, 1913a; Brotherus, 1924a; Dixon, 1935; Steere, 1961a; Greene, 1968a; Robinson, 1972 (as *C. nordenskioeldii*)).

C. dicranellatum (Dus.) Roiv. (Rovainen in Rovainen and Bartram, 1937).

Syn. *Dicranum dicranellatum* Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a, 1905a (and as *Campylopus perhorridus* pro parte fide Rovainen and Bartram, 1937); Cardot, 1908a; Cardot and Brotherus, 1923; Rovainen and Bartram, 1937; Kühnemann, 1938; Seki, 1974).

C. lanigerum (C. Muell.) Broth. (Brotherus, 1924a).

Syn. *Dicranum lanigerum* C. Muell. (Mueller, 1885).

Campylopus laniger (C. Muell.) Besch. (Bescherelle, 1885a).

Distr. Southern South America (Bescherelle, 1885a, 1889; Mueller, 1885; Dusén, 1903a, 1905a; Paris, 1904a; Cardot, 1908a, 1912a; Cardot and Brotherus, 1923; Brotherus, 1924a; Rovainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

C. leucopterum (C. Muell.) Roiv.

Chorisodontium aciphyllum

C. luridum Bartr. (Bartram in Rovainen and Bartram, 1937).

Distr. Southern South America (Rovainen and Bartram, 1937).

C. macropus (C. Muell.) Broth. (Brotherus, 1924a).

Syn. *Dicranum macropus* Kunz. ex C. Muell. (Kunze ex Mueller, 1844b [1845]).

Distr. Southern South America (Mueller, 1844b, 1848a; Montagne, 1850; Mitten, 1869; Jaeger, 1872; Paris, 1904b; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Seki, 1974).

C. magellanicum (Card.) Bartr. (Bartram in Roivainen and Bartram, 1937).

Syn. *Dicranum magellanicum* Card. (Cardot, 1901).

Dicranum percompactum Dus. (Dusén, 1903a) nom. nud.

Distr. Southern South America (Cardot, 1901, 1905a, 1908a; Dusén, 1903a, 1905a; Cardot and Brotherus, 1923; Roivainen and Bartram, 1937; Kühnemann, 1938).

C. microodus (Lor.) Broth. (Brotherus, 1924a).

Syn. *Dicranum microodus* Lor. (Lorentz, 1866).

Distr. Southern South America (Lorentz, 1866; Jaeger, 1872; Paris, 1904b; Brotherus, 1924a).

C. nordenskioeldii (Card.) Roiv.

Chorisodontium aciphyllum

C. orthocomum (C. Muell.) Roiv.

Chorisodontium aciphyllum

C. sericeum Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

C. spegazzinii (C. Muell.) Roiv. (Roivainen in Roivainen and Bartram, 1937).

Syn. *Dicranum spegazzinii* C. Muell. (Mueller, 1885).

Campylopus spegazzinii (C. Muell.) Kindb. (Kindberg, 1888).

Dicranum blindiaceum C. Muell. (Mueller in Kindberg, 1889) nom. nud.

Distr. Southern South America (Mueller, 1885; Paris, 1894, 1904a; Cardot, 1900, 1901, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938).

C. sphagneticola Roiv. (Roivainen in Roivainen and Bartram, 1937).

Distr. Southern South America (Roivainen and Bartram, 1937; Kühnemann, 1938).

Chrysoblastella

C. chilensis (Mont.) Reim.

Cheilotrichia chilensis

C. dubia (Dus.) Reim.

Cheilotrichia chilensis

C. revoluta (Herz.) Thér.

Cheilotrichia chilensis

***Chrysohypnum* hom. illeg.**

C. polygamum (B.S.G.) Loesk.

Campylium polygamum

Cinclidium

C. stygium Sw. (Swartz, 1803).

Syn. *Meesia stygia* (Sw.) Brid. (Bridel, 1803).

Mnium stygium (Sw.) B.S.G. (Bruch and others, 1838).

Mnium subpunctatum Card. et Broth. (Cardot and Brotherus, 1923) fide Koponen, 1969.

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938; Koponen, 1969).

Cladodium

C. uliginosum Brid.

Bryum uliginosum

Cladomnion

C. crenato-obtusum Dus.

Cladomniopsis crenato-obtusa

C. gracile (Hamp.) Mitt.

Glyphothecium gracile

<i>C. membranosum</i> Dus.	<i>Juratzkaea membranosa</i>
<i>C. montevidense</i> C. Muell.	<i>Felipponea montevidensis</i>
<i>C. poeppigianum</i> C. Muell.	<i>Glyphothecium gracile</i>

C. valdiviae C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1904a).

Cladomniopsis

C. crenato-obtusa Fleisch. (Fleischer, 1906 [1908]).

Syn. *Cladomnion crenato-obtusum* Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1904a; Fleischer, 1906; Brotherus, 1925; Matteri, 1973d; Seki, 1974; Kühnemann and Gonçalves, 1975).

Climacodontium *nom. nud.*

C. cygnicollum (C. Muell.) Hamp.

Anomobryum cygnicollum

Codonoblepharon

C. menziesii Schwaegr.

Zygodon menziesii

C. papillatum (Mont.) Jaeg.

Zygodon papillatus

C. uncinatum (Mitt.) Jaeg.

Zygodon pentastichus

Coelidium *hom. illeg.*

C. auriculatum (Mont.) Jaeg.

Acrocladium auriculatum

C. cochlearifolium (Schwaegr.) Reichdt.

Weymouthia cochlearifolia

Conomitrium *nom. illeg.*

C. atratum C. Muell.

Fissidens atratus

C. berterii Mont.

Fissidens fontanus

C. berteroii Mont. ex Kindb.

Fissidens fontanus

C. dillenii (Pyl.) Mont.

Fissidens fontanus

C. hedwigii Mont.

Fissidens semicompletus

C. julianum (Cand.) Mont.

Fissidens fontanus

C. lorentziae C. Muell.

Fissidens lorentziae

C. molle C. Muell.

Fissidens permollis

C. nigrillum C. Muell.

Fissidens nigrillum

C. nigritum C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

C. osmundoides (Hedw.) C. Muell.

Fissidens osmundoides

C. smaragdinum Lor. et C. Muell.

Fissidens smaragdinus

Conostomum

C. australe Sw.

Conostomum pentastichum

var. *microphyllum* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).
Falkland Islands (Cardot and Brotherus, 1923).

var. *piliferum* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

C. lorentzii C. Muell. (Mueller, 1882).

Syn. *Campylopus lorentzii* C. Muell. ex Par. (Mueller ex Paris, 1894) *err.*

Distr. Southern South America (Mueller, 1882; Paris, 1894, 1900a, 1904a; Brotherus, 1924a; Kühnemann, 1938).

C. magellanicum Sull. (Sullivant, 1850).

Syn. *Bartramia magellanica* (Sull.) Mitt. (Mitten, 1869).

Conostomum perangulatum Card. (Cardot, 1905a) *fide* Bell, 1973a.

Distr. Southern South America (Sullivant, 1850, 1859; Mueller, 1851a, 1885; Mitten, 1869; Jaeger, 1875; Paris, 1904a; Cardot, 1905a, 1908a; Brotherus, 1924a; Kühnemann, 1938).

South Georgia (Bell, 1973b).

Antarctic, peninsula region (Bell, 1973a).

C. pentastichum (Brid.) Lindb. (Lindberg, 1863).

Syn. *Bartramia pentasticha* Brid.: (Bridel, 1803).

Weisia pentasticha (Brid.) Hedw. (Hedwig *in* Bridel, 1826) *nom. inval.*

Conostomum australe Sw. (Swartz, 1806a) *nom. illeg.*

Philonotis australis Mitt. (Mitten, 1860).

Bartramia australis Mitt. (Mitten, 1869) *hom. illeg.*

Bryum angulosum Menz. (Menzies *in* Bridel, 1826) *nom. nud.*

Conostomum rhynchostegium C. Muell. (Mueller, 1890a).

Distr. Southern South America (Bridel, 1803, 1819, 1826; Swartz, 1806a; Schwaegrichen, 1811, 1824; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Montagne, 1850; Sullivant, 1859; Mitten, 1869; Jaeger, 1875; Bescherelle, 1889; Paris, 1904a; Cardot, 1905a, 1908a, 1913a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

Falkland Islands (Wilson and Hooker, 1847; Jaeger, 1875; Paris, 1904a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Paris, 1904a; Cardot, 1906a, 1908a; Dixon, 1920, 1934; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b; Bell, 1973b, 1977b).

Antarctic, peninsula region (Bell, 1973a).

C. perangulatum Card.

Conostomum magellanicum

var. *majus* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

C. perpusillum Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923).

C. rhynchostegium C. Muell.

Conostomum pentastichum

Costesia

C. spongiosa Thér. (Thériot, 1917b).

Distr. Southern South America (Thériot, 1917b, 1921a, 1930; Costes, 1921; Brotherus, 1924a).

Cratoneuron

C. filicinum (Hedw.) Spruc. (Spruce, 1867).

Syn. *Hypnum filicinum* Hedw. (Hedwig, 1801).

Distr. Southern South America (Robinson, 1964).

C. mendozense Herz. (Herzog, 1916b).

Distr. Southern South America (Herzog, 1916b).

C. perplicatum (Dus.) Broth. (Brotherus, 1909).

Syn. *Hypnum perplicatum* Dus. (Dusén, 1903a).

Drepanocladus perplicatus (Dus.) Roth (Roth, 1908).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Roth, 1908; Brotherus, 1909, 1925; Kühnemann, 1938).

Crossidium*C. roseae* (Williams) Bartr.*Aloina roseae****Crossomitrium****C. patrisiae* (Brid.) C. Muell. (Mueller, 1874).Syn. *Hypnum patrisiae* Brid. (Bridel, 1827b).*Hookeria patrisiae* (Brid.) Hamp. (Hampe, 1844).*Lepidopilum patrisiae* (Brid.) Mitt. (Mitten, 1869).

Distr. Southern South America (Hampe, 1844; Mueller, 1851a; Mitten, 1869; Gibert, 1873; Herter, 1933a; Welch, 1969).

Cryphaea*C. aurantiorum* C. Muell.*Cryphidium leucocoleum**C. caldensis* Aongstr. (Ångström, 1876a).

Distr. Southern South America (Herzog, 1952a).

C. consimilis Mont.*Cryphaea tenella**C. cuspidata* Sull.*Dendrocryphaea cuspidata**C. furcinervis* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1925; Kühnemann, 1938).

C. gorveana Mont.*Dendrocryphaea gorveana**C. julacea* Mitt.*Acrocryphaea julacea**C. lechleri* C. Muell.*Dendrocryphaea lechleri**C. lorentziana* C. Muell. (Mueller, 1882).Syn. *Cryphaea lorentzii* C. Muell. ex Kindb. (Mueller ex Kindberg, 1891) nom. illeg.

Distr. Southern South America (Mueller, 1882; Paris, 1904a; Brotherus, 1925; Kühnemann, 1938).

C. lorentzii C. Muell. ex Kindb.*Cryphaea lorentziana**C. mollis* Dus.*Cryphaeophilum molle**C. orbifolia* Besch.*Cryphidium leucocoleum**C. pulchella* Par. et Schimp.*Dendrocryphaea ramosissima**C. ramosissima* Schimp.*Dendrocryphaea lechleri**C. ramosissima* Par.*Dendrocryphaea ramosissima**C. rhacomitrioides* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a, 1882; Paris, 1904a; Brotherus, 1925; Kühnemann, 1938; Herzog, 1952a).

var. *gracilis* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Kühnemann, 1938).

C. tenella (Schwaegr.) Hornsch. ex C. Muell. (Hornschuch ex Mueller, 1844b [1845]).Syn. *Neckera tenella* Schwaegr. (Schwaegrichen, 1827a).*Pilotrichum tenellum* (Schwaegr.) C. Muell. (Mueller, 1851a).*Cryphaea consimilis* Mont. (Montagne, 1845a).*Pilotrichum consimile* (Mont.) C. Muell. (Mueller, 1851a).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1851a; Mitten, 1869; Jaeger, 1876; Paris, 1904a; Cardot, 1908a; Herzog, 1923, 1938; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938; Seki, 1974).

C. tenella Mitt.*Dendrocryphaea ramosissima*

Cryphaeophilum***C. molle*** (Dus.) Fleisch. (Fleischer, 1914).Syn. *Cryphaea mollis* Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1903a, 1905b; Paris, 1904a; Cardot, 1908a; Fleischer, 1914; Brotherus, 1925; Kühnemann, 1938; Herzog, 1954; Seki, 1974; Kühnemann and Gonçalves, 1976).

var. ***enerve*** Herz. (Herzog, 1954).

Distr. Southern South America (Herzog, 1954; Seki, 1974).

Cryphidium***C. aurantiacum*** (C. Muell.) Broth.***Cryphidium leucocoleum******C. leucocoleum*** (Mitt.) Jaeg. (Jaeger, 1876).Syn. *Neckera leucocolea* Mitt. (Mitten, 1869).*Cryphaea aurantiacum* C. Muell. (Mueller, 1882).*Cryphidium aurantiacum* (C. Muell.) Broth. (Brotherus, 1905b).*Cryphaea orbifolia* Besch. (Bescherelle, 1891).*Cryphidium orbifolium* (Besch.) Broth. (Brotherus, 1905b).

Distr. Southern South America (Mitten, 1869; Gibert, 1873; Jaeger, 1877; Mueller, 1882; Bescherelle, 1891; Paris, 1904a, 1905a; Brotherus, 1905b, 1925; Herter, 1928, 1933a, b; Kühnemann, 1938; Herzog, 1952a).

C. orbifolium (Besch.) Broth.***Cryphidium leucocoleum***

Cupressina hom. illeg.

C. cupressiformis (Hedw.) C. Muell.***Hypnum cupressiforme******C. pallido-nitida*** C. Muell.***Rhaphidorrhynchium pallido-nitidum***

Cuspidaria hom. illeg.

C. fulvo-acuta C. Muell.***Calliergonella cuspidata******C. morenoi*** C. Muell.***Pleurozium schreberi******C. pseudo-pura*** C. Muell.***Pseudoscleropodium pseudopurum******Cyathophorum******C. splendidissimum*** (Mont.) Hamp. et Lor.***Schimperobryum splendidissimum******Cyclodictyon******C. albicans*** (Hedw.) O. Kuntze (Kuntze, 1891).Syn. *Leskea albicans* Hedw. (Hedwig, 1801).

Distr. Southern South America (Herzog, 1952a).

C. sublimbatum (C. Muell.) O. Kuntze (Kuntze, 1891).Syn. *Hookeria sublimbata* C. Muell. (Mueller, 1851a).*Pterygophyllum sublimbatum* (C. Muell.) Mitt. (Mitten, 1885).

Distr. Southern South America (Mueller, 1851a; Mitten, 1869, 1885; Jaeger, 1877; Paris, 1904b, 1905b; Brotherus, 1925; Welch, 1969).

Cylindrothecium***C. flavo-virens*** (C. Muell.) Par.***Entodon flavo-virens***var. ***flagellaceum*** (C. Muell.) Par.***Entodon flavo-virens* var. *flagellaceus******C. platygynoides*** (C. Muell.) Par.***Erythrodontium platygynoides******C. suberythropus*** (C. Muell.) Par.***Entodon suberythropus***var. ***complanatum*** (C. Muell.) Par.***Entodon suberythropus* var. *complanatum***

var. <i>ellipticum</i> (C. Muell.) Par.	<i>Entodon suberythropus</i> var. <i>ellipticus</i>
var. <i>flaccidum</i> (C. Muell.) Par.	<i>Entodon suberythropus</i> var. <i>flaccidus</i>
var. <i>gracillimum</i> (C. Muell.) Par.	<i>Entodon suberythropus</i> var. <i>gracillimus</i>
var. <i>lutescens</i> (C. Muell.) Par.	<i>Entodon suberythropus</i> var. <i>lutescens</i>
var. <i>microtheca</i> (C. Muell.) Par.	<i>Entodon suberythropus</i> var. <i>microtheca</i>
var. <i>myosuroides</i> (C. Muell.) Par.	<i>Entodon suberythropus</i> var. <i>myosuroides</i>
var. <i>rufulum</i> (C. Muell.) Par.	<i>Entodon suberythropus</i> var. <i>rufulus</i>
<i>C. truncatum</i> (C. Muell.) Par.	<i>Entodon truncatus</i>

Cynodontium

<i>C. capillaceum</i> (Hedw.) Brid.	<i>Distichium capillaceum</i>
<i>C. conicum</i> (Mont.) Mitt.	<i>Ditrichum conicum</i>
<i>C. fuegianum</i> (Card.) Card.	<i>Oncophorus fuegianus</i>
<i>C. hookeri</i> (C. Muell.) Mitt.	<i>Ditrichum hookeri</i>
<i>C. leiocarpum</i> Lor.	<i>Aongstroemia leiocarpa</i>

Cynontodium nom. rejic.

<i>C. affine</i> (C. Muell.) Mitt.	<i>Ditrichum difficile</i>
<i>C. australe</i> Mitt.	<i>Ditrichum strictum</i>
<i>C. capillaceum</i> Hedw.	<i>Distichium capillaceum</i>
<i>C. elongatum</i> (Hook. f. et Wils.) Mitt.	<i>Ditrichum cylindricarpum</i>
<i>C. hookeri</i> (C. Muell.) Mitt.	<i>Ditrichum hookeri</i>
<i>C. hyalinum</i> (Mitt.) Mitt.	<i>Ditrichum hyalinum</i>

Cyptodon

<i>C. crassinervis</i> Broth. (Brotherus, 1924b).	
<i>Distr.</i> Juan Fernandez (Brotherus, 1924b, 1925; Robinson, 1975).	

Cyrt-Hypnum

<i>C. filarium</i> (Mitt.) Hamp.	<i>Thuidiopsis furfurosa</i>
----------------------------------	------------------------------

Cyrtopus

<i>C. curtipendulus</i> (Hedw.) Spruc.	<i>Antitrichia curtipendula</i>
--	---------------------------------

Daltonia hom. illeg.

<i>D. ariquensis</i> Schimp.	<i>Daltonia gracilis</i>
------------------------------	--------------------------

D. aristata Geh. et Hamp. (Geheebe and Hampe *in Hampe*, 1879).

Syn. *Daltonia uleana* C. Muell. (Mueller, 1898c).

Distr. Southern South America (Reimers, 1926).

D. gracilis Mitt. (Mitten, 1869).

Syn. *Daltonia krauseana* C. Muell. ex Dus. (Mueller ex Dusén, 1905b).

Daltonia ariquensis Schimp. (Schimper *in Bartram*, 1931) *nom. nud.*

Distr. Juan Fernandez (Robinson, 1975).

Southern South America (Dusén, 1905b; Brotherus, 1925; Bartram, 1931, 1952; Herzog, 1954; Welch, 1969; Seki, 1974; Matteri, 1975b).

D. krauseana C. Muell. ex Dus.

Daltonia gracilis

D. patagonica Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a).

- D. polytrichoides* (Hedw.) Arnott *Lepidopilum polytrichoides*
D. trachydonta Mitt. ex Bartr. *Daltonia trachydonta*
- D. trachydonta* Mitt. (Mitten, 1869).**
- Syn. *Daltonia trachydonta* Mitt. ex Bartr. (Mitten ex Bartram, 1931) err.
Daltonia valdiviae Herz. (Herzog, 1923).
- Distr. Southern South America (Herzog, 1923; Brotherus, 1925; Bartram, 1931; Welch, 1969; Seki, 1974; Matteri, 1975b).
- D. uleana* C. Muell. *Daltonia aristata*
D. valdiviae Herz. *Daltonia trachydonta*
- Decodon hom. illeg.* *Rhachithecium perpusillum*
D. demissus C. Muell.
- Dendrocryphaea***
- D. cuspidata* (Sull.) Broth. ex Par. (Brotherus ex Paris, 1909).**
- Syn. *Cryphaea cuspidata* Sull. (Sullivant, 1857 [1854]).
Braunia cuspidata (Sull.) Hamp. (Hampe, 1867a).
- Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b).
Southern South America (Sullivant, 1857, 1859; Mitten, 1869; Paris, 1904a, 1909; Brotherus, 1905b, 1925; Thériot, 1924, 1934b, 1935a; Herzog, 1954, 1960).
- D. gorveana* (Mont.) Par. et Schimp. (Paris and Schimper in Brotherus, 1905b).**
- Syn. *Cryphaea gorveana* Mont. (Montagne, 1845a).
Dendropogon gorveanus (Mont.) B.S.G. (Bruch and others, 1850).
Pilotrichum gorveanum (Mont.) C. Muell. (Mueller, 1851a).
- Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1851a; Lorentz, 1866; Mitten, 1869; Jaeger, 1876; Neger, 1899; Paris, 1904a; Brotherus, 1905b; Cardot, 1908a; Herzog, 1923; Kühnemann, 1938; Seki, 1974).
- var. *lechleri* Fleisch. *Dendrocryphaea lechleri*
- D. lechleri* (Fleisch.) Par. et Schimp. ex Thér. (Paris and Schimper ex Thériot, 1934b [1935]).**
- Syn. *Cryphaea lechleri* C. Muell. (Mueller in Lorentz, 1866) nom. nud.
Dendropogon lechleri C. Muell. (Mueller in Fleischer, 1914) nom. nud.
Dendrocryphaea gorveana (Mont.) Par. et Schimp. var. *lechleri* Fleisch. (Fleischer, 1914).
Cryphaea ramosissima Schimp. (Schimper in Bescherelle, 1876) nom. nud.
Dendrocryphaea ramosissima Par. et Schimp. (Paris and Schimper in Paris, 1894) nom. nud.
- Distr. Southern South America (Lorentz, 1866; Paris, 1894, 1904a; Fleischer, 1914; Thériot, 1934b).
- D. pulchella* (Par. et Schimp.) Fleisch. *Dendrocryphaea ramosissima*
D. ramosissima Par. et Schimp. *Dendrocryphaea lechleri*
- D. ramosissima* (Par.) Wijk et Marg. (van der Wijk and Margadant, 1959).**
- Syn. *Cryphaea ramosissima* Par. (Paris in Jaeger, 1876).
Cryphaea tenella Mitt. (Mitten, 1869) hom. illeg.
Dendrocryphaea tenella Thér. (Thériot, 1934b [1935]) nom. illeg.
Cryphaea pulchella Par. et Schimp. (Paris and Schimper in Paris, 1894).
Dendrocryphaea pulchella (Par. et Schimp.) Fleisch. (Fleischer, 1914).
- Distr. Southern South America (Mitten, 1869; Jaeger, 1876; Paris, 1894, 1904a; Cardot, 1908a; Brotherus, 1925; Thériot, 1934b; Kühnemann, 1938).
- D. tenella* Thér. *Dendrocryphaea ramosissima*
- Dendro-Hypnum***
- D. fasciculatum* (Hedw.) Hamp. *Porothamnium fasciculatum*

Dendroligotrichum

D. dendroides (Hedw.) Broth. (Brotherus, 1905a).

Syn. *Polytrichum dendroides* Brid. ex Hedw. (Bridel ex Hedwig, 1801).
Pogonatum dendroides (Hedw.) Brid. (Bridel, 1827b).
Catharinea dendroides (Hedw.) Hamp. (Hampe, 1847).
Polytrichadelphus dendroides (Hedw.) Mitt. (Mitten, 1869).
Oligotrichum dendroides (Hedw.) Kindb. (Kindberg, 1888).
Gymnopilum dendroides (Hedw.) Schimp. (Schimper in Paris, 1895) nom. inval.

Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).
Southern South America (Hedwig, 1801; Bridel, 1803, 1806, 1827b; Palisot de Beauvois, 1805a; Schwaegrichen, 1816, 1826; Bridel, 1827b; Montagne, 1845b, 1850; Wilson and Hooker, 1847; Mueller, 1848a, 1851a, 1885; Sullivant, 1859; Lorentz, 1866; Mitten, 1869; Jaeger, 1875; Bescherelle, 1889; Hariot, 1891; Neger, 1899; Cardot, 1900, 1901, 1908a, 1913a; Paris, 1905b; Thériot, 1918; Campo, 1921; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1940, 1954, 1957, 1960; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

D. squamosum (Hook. f. et Wils.) Broth. (Cardot, 1908a).

Syn. *Polytrichum squamosum* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Catharinea squamosa (Hook. f. et Wils.) C. Muell. (Mueller, 1848a).
Polytrichadelphus squamosus (Hook. f. et Wils.) Mitt. (Mitten, 1869).
Oligotrichum squamosum (Hook. f. et Wils.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Sullivant, 1859; Mitten, 1869; Ångström, 1872; Jaeger, 1875; Cardot, 1905a, 1908a, 1913a; Paris, 1905b; Cardot and Brotherus, 1923; Brotherus, 1925; Bartram, 1946, 1952; Herzog, 1954; Seki, 1974).
Falkland Islands (Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925).
South Georgia (Greene, 1973).

Dendropogon

D. gorveanus (Mont.) B.S.G.

Dendrocryphaea gorveana

D. lechleri C. Muell.

Dendrocryphaea lechleri

Dermatodon

D. rubellus Genth

Bryoerythrophyllum recurvirostre

Desmatodon

D. ambyophyllum Mont.

Tortula platyphylla

D. argentinus Broth. (Brotherus, 1918).

Distr. Southern South America (Brotherus, 1918, 1924a; Kühnemann, 1938).

D. atro-virens (Sm.) Jur.

Desmatodon convolutus

D. cavifolius Mitt.

Pterygoneurum ovatum

D. convolutus (Brid.) Grout (Grout, 1939).

Syn. *Trichostomum convolutum* Brid. (Bridel, 1806).

Grimmia atro-virens Sm. (Smith, 1809).

Tortula atro-virens (Sm.) Lindb. (Lindberg, 1864a).

Barbula atrovirens (Sm.) Schimp. (Schimper, 1876).

Desmatodon atro-virens (Sm.) Jur. (Juratzka, 1882).

Distr. Southern South America (Mitten, 1869; Thériot, 1921a, b; Brotherus, 1924a; Bartram, 1952; Herzog, 1954).

D. heimii (Hedw.) Mitt.

Pottia heimii

D. rigidulus Hassk.

Didymodon rigidulus

D. ruber (Mitt.) Hamp.

Tortula robusta

<i>D. schimperi</i> (Mont.) Jaeg.	<i>Didymodon schimperi</i>
<i>D. serrulatus</i> (Hook. et Grev.) Mitt.	<i>Tortula robusta</i>
<i>D. subtophaceus</i> (Williams) Williams	<i>Trichostomopsis australasiae</i>
<i>Dichodontium</i>	
<i>D. brotheri</i> Dix.	<i>Oncophorus fuegianus</i>
<i>D. capituligerum</i> (C. Muell.) Par.	<i>Anisothecium capituligerum</i>
<i>D. dicranelloides</i> Card.	<i>Dicranella cardotii</i>
var. <i>falklandicum</i> Card.	<i>Anisothecium persquarrosum</i> var. <i>falklandicum</i>
<i>D. elegans</i> (Dub.) Jaeg.	<i>Anisothecium elegans</i>
<i>D. jamesonii</i> (Mitt.) Jaeg.	<i>Anisothecium vaginatum</i>
<i>D. krausei</i> (Lor.) Jaeg.	<i>Anisothecium krausei</i>
<i>D. lorentzii</i> (C. Muell.) Par.	<i>Anisothecium lorentzii</i>
<i>D. paludella</i> Besch.	<i>Anisothecium paludella</i>
<i>D. patagonicum</i> (C. Muell.) Par.	<i>Dicranella patagonica</i>
<i>D. persquarrosum</i> (Dus.) Card.	<i>Dicranella cardotii</i>
var. <i>falklandicum</i> (Card.) Card.	<i>Anisothecium persquarrosum</i> var. <i>falklandicum</i>
<i>D. rufipes</i> (C. Muell.) Par.	<i>Anisothecium rufipes</i>
<i>D. subclathratum</i> (Lor.) Par.	<i>Dicranella hookeri</i>
<i>D. vaginatum</i> (Hook.) Jaeg.	<i>Anisothecium vaginatum</i>
<i>Dicranella</i>	
<i>D. argentinica</i> (C. Muell.) Kindb. (Kindberg, 1889).	
Syn. <i>Aongstroemia argentinica</i> C. Muell. (Mueller, 1879a).	
Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Hosseus, 1935b; Kühnemann, 1938).	
<i>D. aulacocarpa</i> (Mont.) Mitt. (Mitten, 1869).	
Syn. <i>Dicranum aulacocarpum</i> Mont. (Montagne, 1845a).	
<i>Aongstroemia aulacocarpa</i> (Mont.) C. Muell. (Mueller, 1848a).	
Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Mitten, 1869; Jaeger, 1872; Paris, 1904b).	
<i>D. capituligera</i> (C. Muell.) Kindb.	<i>Anisothecium capituligerum</i>
<i>D. cardotii</i> (R. Brown ter.) Dix. (Dixon, 1923).	
Syn. <i>Dicranum cardotii</i> R. Brown ter. (Brown, 1902 [1903]).	
<i>Aongstroemia persquarrosa</i> Dus. (Dusén, 1905a) fide Newton, 1977a.	
<i>Dichodontium persquarrosum</i> (Dus.) Card. (Cardot, 1908a).	
<i>Anisothecium persquarrosum</i> (Dus.) Broth. (Brotherus, 1924a).	
<i>Dichodontium dicranelloides</i> Card. (Cardot, 1905a).	
Distr. Southern South America (Dusén, 1903a, 1905a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen, 1934; Roivainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952; Herzog, 1954; Seki, 1974). Falkland Islands (Cardot and Brotherus, 1923). South Georgia (Newton, 1977a).	
<i>D. circinata</i> Herz. (Herzog, 1954).	
Distr. Southern South America (Herzog, 1954).	
<i>D. costata</i> Broth. (Brotherus, 1924b).	
Distr. Juan Fernandez (Brotherus, 1924a, b; Robinson, 1975).	
<i>D. euchlora</i> (Mont.) Lor.	<i>Microdus euchlorus</i>

D. flexipes Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).

D. flexiseta Dus. (Dusén *in Paris*, 1904b) *nom. nud.*

Distr. Southern South America (*Paris*, 1904b).

D. fuegiana Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

D. guilleminiana (Mont.) Mitt. (Mitten, 1869).

Syn. *Dicranum guilleminianum* Mont. (Montagne, 1841).

Distr. Southern South America (Krieger, 1904).

D. hilariana (Mont.) Mitt. (Mitten, 1869).

Syn. *Dicranum hilarianum* Mont. (Montagne, 1839a).

Aongstroemia hilariana (Mont.) C. Muell. (Mueller, 1848a).

Microdus hilarianus (Mont.) Besch. (Bescherelle *in Paris*, 1896).

Dicranum tenuirostre Kunz. ex Schwaegr. (Kunze ex Schwaegrichen, 1842).

Aongstroemia tenuirostris (Schwaegr.) C. Muell. (Mueller, 1848a).

Dicranella tenuirostris (Schwaegr.) Mitt. (Mitten, 1869).

Microdus tenuirostris (Schwaegr.) Besch. (Bescherelle *in Paris*, 1896).

Dicranum sellowii Hornsch. (Hornschuch, 1840).

Distr. Southern South America (Hornschuch, 1840; Mueller, 1844b, 1848a; Montagne, 1850, 1856; Jaeger, 1872; Gibert, 1873; Paris, 1896, 1905a; Herter, 1933a).

D. hookeri (C. Muell.) Card. (Cardot, 1906a) *fide* Newton, 1977a.

Syn. *Aongstromia hookeri* C. Muell. (Mueller, 1851a).

Anisothecium hookeri (C. Muell.) Broth. (Brotherus, 1924a).

Dicranum vaginatum Hook. f. et Wils. (Hooker and Wilson *in Mueller*, 1851a) *nom. nud.*

Dicranella subclathrata Lor. (Lorentz, 1864).

Dichodontium subclathratum (Lor.) Par. (*Paris*, 1894).

Aongstroemia subclathrata (Lor.) C. Muell. (Mueller, 1901a [1900]).

Anisothecium perpusillum Dus. (Dusén, 1903a) *fide* Newton, 1977a.

Dicranella perpusilla (Dus.) Card. (Cardot, 1908a).

Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1851a, 1885, 1901a; Lorentz, 1864, 1866; Jaeger, 1872; Paris, 1894, 1904b; Dusén, 1903a; Cardot, 1908a; Cardot and Brotherus, 1923; Thériot, 1923b, 1924; Brotherus, 1924a; Roivainen and Bartram, 1937; Herzog, 1938, 1954; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

South Georgia (Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b; Newton, 1977a).

Antarctic, continental region (Cardot, 1910; Brotherus, 1924a; Greene, 1968a; Newton, 1977b).

var. *flexisubulata* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Seki, 1974).

D. jamesonii (Mitt.) Broth.

Anisothecium vaginatum

D. krausei Lor.

Anisothecium krausei

D. kunzeana (C. Muell.) Mitt.

Microdus kunzeanus

D. lorentzii (C. Muell.) Kindb.

Anisothecium lorentzii

D. paludella (Besch.) Dus.

Anisothecium paludella

D. patagonica (C. Muell.) Broth. (Brotherus, 1901).

Syn. *Aongstroemia patagonica* C. Muell. (Mueller, 1897a).

Dichodontium patagonicum (C. Muell.) Par. (*Paris*, 1900a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1901; Cardot, 1908a; Herzog, 1923).

<i>D. perpusilla</i> (Dus.) Card.	<i>Dicranella hookeri</i>
<i>D. pseudorufescens</i> Card. et Broth.	<i>Anisothecium pseudorufescens</i>
var. <i>leptoclada</i> Card. et Broth.	<i>Anisothecium pseudorufescens</i> var. <i>leptocladus</i>
<i>D. rufipes</i> (C. Muell.) Kindb.	<i>Anisothecium rufipes</i>
<i>D. skottsbergii</i> Card. et Broth.	<i>Anisothecium skottsbergii</i>
<i>D. subclathrata</i> Lor.	<i>Dicranella hookeri</i>
<i>D. tenuirostris</i> (Schwaegr.) Mitt.	<i>Dicranella hilariana</i>
<i>D. vaginata</i> (Hook.) Card.	<i>Anisothecium vaginatum</i>
<i>Dicranobryum nom. nud.</i>	
<i>D. fabroniellum</i> Dus.	<i>Brachymenium fabroniellum</i>
<i>D. subfabroniellum</i> Dus.	<i>Brachymenium meyenianum</i>
<i>Dicranodon nom. illeg.</i>	
<i>D. adianthoides</i> (Hedw.) Behéré	<i>Fissidens adianthoides</i>
<i>D. bryoides</i> (Hedw.) Behéré	<i>Fissidens bryoides</i>
<i>Dicranodontium</i>	
<i>D. asperifolium</i> (Mitt.) Broth. (Brotherus, 1924a).	
Syn. <i>Campylopus asperifolius</i> Mitt. (Mitten, 1869).	
Distr. Southern South America (Mitten, 1869; Jaeger, 1872; Paris, 1904a).	
<i>D. schwabei</i> Herz. et Thér. (Herzog and Thériot in Herzog, 1939).	
Distr. Southern South America (Herzog, 1939).	
<i>Dicranoloma</i>	
<i>D. australe</i> (C. Muell.) Par. (Paris, 1904b).	
Syn. <i>Dicranum australe</i> Besch. ex C. Muell. (Bescherelle ex Mueller, 1885). <i>Leucoloma australe</i> (Besch.) Broth. (Brotherus, 1901). <i>Dicranum pseudo-australe</i> C. Muell. (Mueller in Kindberg, 1891) <i>nom. nud.</i> <i>Dicranoloma pseudo-australe</i> Ren. (Renauld, 1909) <i>nom. nud.</i> <i>Leucoloma robustum</i> Hook. f. et Wils. var. <i>flexuosum</i> Dus. (Dusén, 1905b) <i>fide</i> Cardot, 1908a.	
Distr. Southern South America (Bescherelle, 1885a, 1889; Mueller, 1885, 1889; Hariot, 1891; Dusén, 1903a, 1905b (and as <i>Leucoloma setosum</i> <i>fide</i> Cardot, 1908a), 1905d; Paris, 1904b; Cardot, 1905a, 1908a; Renauld, 1909; Cardot and Brotherus, 1923; Brotherus, 1924a; Rovainen and Bartram, 1937; Seki, 1974). Falkland Islands (Cardot and Brotherus, 1923).	
var. <i>condensatum</i> Broth. (Brotherus in Cardot and Brotherus, 1923).	
Distr. Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).	
var. <i>schwabei</i> Herz. et Thér. (Herzog and Thériot in Herzog, 1939).	
Distr. Southern South America (Herzog, 1939).	
<i>D. austro-georgicum</i> Broth.	<i>Dicranoloma hariotii</i>
<i>D. billardieri</i> (Brid.) Par. (Paris, 1904b).	
Syn. <i>Dicranum billardieri</i> Brid. (Bridel, 1802). <i>Oncophorus billardieri</i> (Brid.) Brid. (Bridel, 1826). <i>Leucoloma billardieri</i> (Brid.) Broth. (Brotherus, 1901). <i>Dicranum boryanum</i> Schwaegr. (Schwaegrichen, 1823). <i>Trematodon boryanus</i> (Schwaegr.) Spreng. (Sprengel, 1827) <i>hom. illeg.</i> <i>Cecalyphum dichotomum</i> P. Beauv. (Palisot de Beauvois, 1805a). <i>Dicranum dichotomum</i> (P. Beauv.) Brid. (Bridel, 1806) <i>hom. illeg.</i> <i>Oncophorus dichotomus</i> (P. Beauv.) Brid. (Bridel, 1826). <i>Dicranoloma dichotomum</i> (P. Beauv.) Ren. (Renauld, 1901).	

Dicranum callocaulon C. Muell. (Mueller *in Kindberg*, 1889) *nom. nud.*

Dicranum helvolum C. Muell. (Mueller *in Kindberg*, 1891) *nom. nud.*

Dicranoloma fernandezianum Broth. (Brotherus, 1924b) *fide* Robinson, 1975.

Distr. Juan Fernandez (Montagne, 1835; Mitten, 1869, 1885; Jaeger, 1872; Johow, 1896; Paris, 1904b; Skottsberg, 1914; Brotherus, 1924a, b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Montagne, 1850; Mitten, 1869; Jaeger, 1872; Brotherus, 1901, 1924a; Dusén, 1903a, 1905a; Paris, 1904b; Cardot, 1905a, 1908a; Thériot, 1918, 1935a; Campo, 1921; Spegazzini, 1922; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1954, 1960; Roivainen, 1934; Roivainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952; Bizot, 1967; Seki, 1974).

Falkland Islands, (Wilson and Hooker, 1847; Mueller, 1848a; Jaeger, 1872; Brotherus, 1901, 1924a; Paris, 1904b; Cardot, 1908a; Kühnemann, 1938).

var. compactum (Card.) Card. et Broth. (Cardot and Brotherus, 1923).

Syn. *Dicranum billardieri* Brid. var. *compactum* Card. (Cardot, 1905a).

Distr. Southern South America (Cardot and Brotherus, 1923).

Falkland Islands (Cardot, 1905a, 1908a).

var. moelleri Bauer (Bauer, 1934).

Distr. Southern South America (Bauer, 1934).

D. capillare (Dus.) Broth. (Brotherus, 1909).

Syn. *Leucoloma capillare* Dus. (Dusén, 1905a).

Dicranum capillare (Dus.) Card. (Cardot, 1908a).

Leucoloma nigricaule (Aongstr.) Par. var. *gracile* Dus. (Dusén, 1905a).

Leucoloma peruncinatum Dus. var. *gracile* Dus. (Dusén *in* Roivainen and Bartram, 1937) *nom. nud.*

Distr. Southern South America (Dusén, 1905a; Cardot, 1908a; Brotherus, 1909, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1939, 1957; Seki, 1974).

D. capillifolioides Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Robinson, 1975).

D. capillifolium (Card.) Broth.

Dicranoloma menziesii

D. dichotomum (P. Beauv.) Ren.

Dicranoloma billardieri

D. dusenii (Broth.) Broth. (Brotherus, 1909).

Syn. *Leucoloma dusenii* Broth. (Brotherus *in* Dusén, 1905a).

Distr. Southern South America (Dusén, 1905a; Paris, 1905a; Brotherus, 1909, 1924a; Herzog, 1954).

D. falklandicum (Card.) Broth. (Brotherus, 1909).

Syn. *Dicranum falklandicum* Card. (Cardot, 1905a).

Distr. Southern South America (Cardot and Brotherus, 1923).

Falkland Islands (Cardot, 1905a, 1908a; Brotherus, 1909, 1924a; Kühnemann, 1938).

D. fernandezianum Broth.

Dicranoloma billardieri

D. fuegianum (Dus.) Broth.

Dicranoloma harriotii

D. harriotii (C. Muell.) Par. (Paris, 1904b).

Syn. *Dicranum harriotii* C. Muell. (Mueller, 1885).

Leucoloma harriotii (C. Muell.) Broth. (Brotherus, 1901).

Leucoloma fuegianum Dus. (Dusén, 1905a).

Dicranum fuegianum (Dus.) Card. (Cardot, 1908a).

Dicranoloma fuegianum (Dus.) Broth. (Brotherus, 1909).

Leucoloma fuegianum Dus. var. *laxum* Dus. (Dusén, 1905a).

Dicranum fuegianum (Dus.) Card. var. *laxum* (Dus.) Card. (Cardot, 1908a).

Dicranoloma harriotii (C. Muell.) Par. var. *laxum* (Dus.) Card. et Broth. (Cardot and Brotherus, 1923).

Dicranum austro-georgicum C. Muell. (Mueller, 1890a).

Dicranoloma austro-georgicum Broth. (*Brotherus in Roivainen and Bartram, 1937*)
nom. nud.

Leucoloma alboalare Dus. (*Dusén in Cardot, 1908a*) *nom. nud.*

Distr. Southern South America (Bescherelle, 1885a, 1889; Mueller, 1885; Cardot, 1900, 1901, 1908a; Brotherus, 1901, 1909, 1924a; Dusén, 1904, 1905a; Paris, 1904b; Cardot and Brotherus, 1923; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1954; Seki, 1974).

Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Paris, 1904b; Cardot, 1906a, 1908a; Dixon, 1920; Steere, 1961b; Bell, 1973b, 1977b).

var. laxum (Dus.) Card. et Broth.

Dicranoloma hariotii

D. hollermayeri Thér. (*Thériot, 1935a*).

Distr. Southern South America (*Thériot, 1935a*).

D. imponens (Mont.) Ren. (*Renauld, 1909*).

Syn. *Dicranum imponens* Mont. (*Montagne, 1843*).

Leucoloma imponens (Mont.) Dus. (*Dusén, 1905a*).

Dicranum involutifolium Sull. (*Sullivant, 1850*).

Leucoloma robustum (Hook. f. et Wils.) Broth. var. *lagunicola* Dus. (*Dusén, 1905a*).

Dicranum robustum Hook. f. et Wils. var. *lagunicola* (Dus.) Card. (*Cardot, 1908a*).

Distr. Southern South America (*Montagne, 1843, 1845b, 1850, 1856; Wilson and Hooker, 1847; Mueller, 1848a, 1851a, 1885; Sullivant, 1850, 1859; De Notaris, 1859; Mitten, 1869; Jaeger, 1872; Bescherelle, 1889; Eaton, 1892; Paris, 1904b; Cardot, 1905a, 1908a; Dusén, 1905a; Renauld, 1909; Thériot, 1921a, 1934b; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen, 1934; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1939; Seki, 1974*).

D. kunkelii H. Robins. (*Robinson, 1974*).

Distr. Juan Fernandez (*Robinson, 1974, 1975*).

D. menziesii (Hook. f. et Wils.) Par. (*Paris, 1904b*).

Syn. *Dicranum menziesii* Hook. f. et Wils. (*Hooker, J. D. and Wilson, 1844*).

Leucoloma menziesii (Hook. f. et Wils.) Broth. (*Brotherus, 1901*).

Dicranum suberectum Hamp. (*Hampe, 1860*).

Leucoloma suberectum (Hamp.) Broth. (*Brotherus, 1901*).

Dicranoloma suberectum (Hamp.) Par. (*Paris, 1904b*).

Leucoloma capillifolium Broth. (*Brotherus in Dusén, 1905a*) *hom. illeg. fide* Robinson, 1975.

Dicranum capillifolium Card. (*Cardot, 1908a*).

Dicranoloma capillifolium (Card.) Broth. (*Brotherus, 1909*).

Distr. Juan Fernandez (*Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975*).

Southern South America (*Dusén, 1903a, 1905a; Paris, 1904b; Cardot, 1908a; Brotherus, 1909, 1924a; Campo, 1915, 1921; Thériot, 1918; Cardot and Brotherus, 1923; Herzog, 1938, 1954, 1960; Kühnemann, 1938*).

var. chilense Thér. (*Thériot, 1924*).

Distr. Southern South America (*Thériot, 1924*).

var. fernandezianum (Thér.) Broth. (*Brotherus, 1924b*).

Syn. *Dicranum menziesii* Hook f. et Wils. var. *fernandezianum* Card. ex Thér. (*Cardot ex Thériot, 1921b*).

Distr. Juan Fernandez (*Skottsberg, 1914; Thériot, 1921b; Brotherus, 1924b*).

D. muelleri Dus.

Dicranoloma robustum* var. *setosum

D. muelleri (Dus.) Roiv. (*Roivainen in Roivainen and Bartram, 1937*).

Syn. *Leucoloma muelleri* Dus. (*Dusén, 1905a*).

Distr. Southern South America (*Dusén, 1905a; Paris, 1905a; Cardot and Brotherus, 1923* (as *D. capillifolium* *fide* Roivainen and Bartram, 1937); *Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1954; Seki, 1974*).

var. *strictifolium* (Dus.) Herz. *Dicranoloma robustum var. strictifolium*

D. nigricaulis (Aongstr.) Par. (Paris, 1904b). *Dicranoloma robustum var. strictifolium*

Syn. *Dicranum nigricaulis* Aongstr. (Ångström, 1872).
Leucoloma nigricaulis (Aongstr.) Broth. (Brotherus, 1901).
Dicranum racovitzae Card. (Cardot, 1900).
Dicranum chilense De Not. (De Notaris, 1859).
Leucoloma peruncinatum Dus. (Dusén, 1905a).
Dicranum peruncinatum (Dus.) Card. (Cardot, 1908a).
Dicranoloma peruncinatum (Dus.) Broth. (Brotherus, 1909).

Distr. Juan Fernandez (Brotherus, 1924b; Robinson, 1975).
Southern South America (De Notaris, 1859; Ångström, 1872; Jaeger and Sauerbeck, 1879b; Mueller, 1885; Bescherelle, 1889; Cardot, 1900, 1901, 1905a, 1908a, 1913a; Brotherus, 1901, 1909, 1924a; Dusén, 1903a, 1905a; Paris, 1904b; Cardot and Brotherus, 1923; Roivainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952; Seki, 1974).
Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

var. *flexuosulum* (Dus.) Bartr. (Bartram in Roivainen and Bartram, 1937).

Syn. *Leucoloma nigricaulis* (Aongstr.) Broth. var. *flexuosulum* Dus. (Dusén, 1905a).
Dicranum nigricaulis Aongstr. var. *flexuosulum* (Dus.) Card. (Cardot, 1908a).

Distr. Southern South America (Dusén, 1905a; Cardot, 1908a; Roivainen and Bartram, 1937).

D. perremotifolium (Dus.) Broth. (Brotherus, 1924a).

Syn. *Leucoloma perremotifolium* Dus. (Dusén, 1905a).
Dicranum perremotifolium (Dus.) Card. (Cardot, 1908a).

Distr. Southern South America (Paris, 1904a, 1905a; Dusén, 1905a; Cardot, 1908a; Brotherus, 1924a; Roivainen, 1934; Kühnemann, 1938; Herzog, 1954, 1960; Seki, 1974).

var. *fragile* (Dus.) Thér. (Thériot, 1935a).

Syn. *Leucoloma perremotifolium* Dus. var. *fragile* Dus. (Dusén, 1905a).

Distr. Southern South America (Dusén, 1905a; Thériot, 1935a).

D. peruncinatum (Dus.) Broth.

Dicranoloma nigricaulis

D. pseudo-australe Ren.

Dicranoloma australe

D. pungens (Hook. f. et Wils.) Par.

Dicranoloma robustum

D. rigens (Besch.) Par.

Chorisodontium aciphyllum

D. robustum (Hook. f. et Wils.) Par. (Paris, 1904b).

Syn. *Dicranum robustum* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Leucoloma robustum (Hook. f. et Wils.) Broth. (Brotherus, 1901).
Dicranum grandialare Dus. (Dusén, 1903a) nom. nud.
Leucoloma grandialare Dus. (Dusén, 1905a).
Dicranum pungens Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Leucoloma pungens (Hook. f. et Wils.) Broth. (Brotherus, 1901).
Dicranoloma pungens (Hook. f. et Wils.) Par. (Paris, 1904b).
Dicranum robustum Hook. f. et Wils. var. *pungens* (Hook. f. et Wils.) Hook. f. (Hooker, 1867).
Dicranum hookeri Schimp. (Schimper in Paris, 1895) nom. nud.

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Sullivant, 1859; Mitten, 1869; Ångström, 1872; Jaeger 1872; Eaton, 1892; Cardot, 1900, 1908a, 1913a; Brotherus, 1901, 1924a; Dusén, 1903a, 1905a, d; Paris, 1904b; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1957; Reimers, 1926; Thériot, 1934b, 1935a; Kühnemann, 1938; Bartram, 1952; Seki, 1974).
Falkland Islands (Mueller, 1848a, 1885; Jaeger, 1872; Cardot, 1908a).

var. *longisetaceum* Herz. (Herzog, 1938).

Distr. Southern South America (Herzog, 1938).

var. setosum (Hook. f. et Wils.) Sains. (Sainsbury, 1931 [1932]).

- Syn. *Dicranum setosum* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Leucoloma setosum (Hook. f. et Wils.) Broth. (Brotherus, 1901).
Dicranoloma setosum (Hook. f. et Wils.) Par. (Paris, 1904b).
Dicranoloma muelleri Dus. (Dusén in Paris, 1904b) *nom. nud.*

Distr. Southern South America (Ångström, 1872; Paris, 1904b; Dusén, 1905a; Cardot, 1908a; Herzog, 1923, 1954; Brotherus, 1924a; Roivainen, 1934; Kühnemann, 1938; Bartram, 1952; Bizot, 1967; Seki, 1974).

var. strictifolium (Dus.) Wijk et Marg. (van der Wijk and Margadant, 1960a).

- Syn. *Leucoloma muelleri* Dus. var. *strictifolium* Dus. (Dusén, 1905a).
Dicranoloma muelleri (Dus.) Roiv. var. *strictifolium* (Dus.) Herz. (Herzog, 1940).
Dicranum setosum Hook. f. et Wils. var. *strictifolium* (Dus.) Card. (Cardot, 1908a).
Dicranoloma setosum (Hook. f. et Wils.) Par. var. *strictifolium* (Dus.) Kühnemann (Kühnemann, 1938).

Distr. Southern South America (Dusén, 1905a; Cardot, 1908a; Kühnemann, 1938; Herzog, 1940).

D. setosum (Hook. f. et Wils.) Par.

Dicranoloma robustum* var. *setosum

var. *strictifolium* (Dus.) Kühnemann

Dicranoloma robustum* var. *strictifolium

D. skottsbergii (Card.) Broth. (Brotherus, 1909).

- Syn. *Dicranum skottsbergii* Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Brotherus, 1909, 1924a; Kühnemann, 1938).

D. suberectum (Hamp.) Par.

Dicranoloma menziesii

D. subimponens (Card.) Broth. (Brotherus, 1909).

- Syn. *Dicranum subimponens* Card. (Cardot, 1905a).

Dicranum turgidissimum Dus. (Dusén in Roivainen and Bartram, 1937) *nom. nud.*

Distr. Southern South America (Cardot, 1905a, 1908a; Brotherus, 1909, 1924a; Cardot and Brotherus, 1923; Roivainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952; Seki, 1974).
South Georgia (Bell, 1973b).

D. subnigraule Dus. (Dusén in Paris, 1904b) *nom. nud.*

Distr. Southern South America (Paris, 1904b).

D. truncorum (C. Muell.) Par. (Paris, 1904b).

- Syn. *Dicranum truncorum* Schimp. ex C. Muell. (Schimper ex Mueller, 1859).

Distr. Southern South America (Lechler, 1857; Mueller, 1859; Lorentz, 1866; Jaeger, 1872; Paris, 1904b).

Dicranoweisia

D. antarctica (C. Muell.) Kindb. (Kindberg, 1888).

- Syn. *Blindia antarctica* C. Muell. (Mueller, 1848a).

Dicranum antarcticum (C. Muell.) Mitt. (Mitten, 1869).

Weisia antarctica (C. Muell.) Wils. (Wilson in Jaeger, 1870).

Distr. Southern South America (Wilson and Hooker, 1847 (as *Weissia crispula* *fide* Mueller, 1848a); Mueller, 1848a, 1879a, 1885; Mitten, 1869; Jaeger, 1870; Paris, 1904b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1939, 1954; Bartram, 1952; Seki, 1974).
Antarctic peninsula region (Bell, 1976).

var. nigricans Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

D. auriculata (C. Muell.) Broth.

Holodontium inerme

D. austro-crispula (C. Muell.) Par. (Paris, 1895).

- Syn.* *Blindia austro-crispula* C. Muell. (Mueller, 1885).
Dicranoweisia perpulvinata Dus. (Dusén, 1903a).
Blindia sulphurea Dus. (Dusén, 1903a) *nom. nud.*
Dicranoweisia sulfurea Dus. (Dusén in Paris, 1904b) *nom. nud.*
Hymenoloma nordenskjoeldii Dus. (Dusén, 1905a).
Dicranoweisia nordenskjoeldii (Dus.) Reim. (Reimers, 1936) *hom. illeg.*
- Distr.* Southern South America (Mueller, 1885; Paris, 1895, 1904b; Dusén, 1903a, 1905a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Donat, 1936a; Roivainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

D. brevipes (C. Muell.) Card. (Cardot, 1905a).

- Syn.* *Blindia brevipes* C. Muell. (Mueller, 1890a).
Distr. South Georgia (Mueller, 1890a; Paris, 1904a; Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b).

D. breviseta Card. (Cardot, 1905a).

- Syn.* *Dicranoweisia breviseta* Card. var. *atrata* Card. et Broth. (Cardot and Brotherus, 1923).
Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Roivainen and Bartram, 1937; Kühnemann, 1938; Seki, 1974).

var. *atrata* Card. et Broth.**Dicranoweisia breviseta****D. cincta** (Hook. f. et Wils.) Par.**Blindia cincta****D. cryptodon** (Mont.) Par.**Camptodontium cryptodon****D. funiculipes** Card. et Broth. (Cardot and Brotherus, 1923).

- Distr.* Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

D. grimmiae (C. Muell.) Broth. (Brotherus, 1901).

- Syn.* *Blindia grimmiae* C. Muell. (Mueller, 1890a).
Distr. South Georgia (Mueller, 1890a; Brotherus, 1901, 1924a; Paris, 1904b; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Dixon, 1934; Steere, 1961b).
Antarctic, peninsula region (Cardot, 1906a, 1908a, 1911c, 1912b; Brotherus, 1924a; Bartram, 1957; Steere, 1961a; Greene, 1968a; Robinson, 1972; Bell, 1976).

D. humilis (C. Muell.) Broth.**Holodontium pumilum****D. inermis** (Mitt.) Par.**Holodontium inerme****D. jugellifera** Dus. (Dusén, 1905a).

- Distr.* Southern South America (Dusén, 1905a; Cardot, 1908a).

D. nordenstjernae (Dus.) Reim.**Dicranoweisia austro-crispula****D. pallidifolia** (C. Muell.) Par.**Dicranoweisia subinclinata****D. perpulvinata** Dus.**Dicranoweisia austro-crispula****D. subinclinata** (C. Muell.) Broth. (Brotherus, 1901).

- Syn.* *Blindia subinclinata* C. Muell. (Mueller, 1890a).
Blindia pallidifolia C. Muell. (Mueller, 1890a).
Dicranoweisia pallidifolia (C. Muell.) Par. (Paris, 1895).

- Distr.* Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).
Falkland Islands (Cardot and Brotherus, 1923).
South Georgia (Mueller, 1890a; Paris, 1895, 1904b; Brotherus, 1901, 1924a; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Dixon, 1934; Steere, 1961b).
Antarctic, peninsula region (Cardot, 1911b, d, 1913a; Steere, 1961a; Robinson, 1972).

D. sulfurea Dus.**Dicranoweisia austro-crispula****D. tenella** Dus.**Camptodontium cryptodon**

Dicranum

- D. aciphyllum* Hook. f. et Wils.
D. acuminatum (Mitt.) C. Muell.
D. adianthoides (Hedw.) Web. et Mohr
D. albo-alare C. Muell. (*Mueller in Kindberg, 1889*) *nom. nud.*
Distr. No locality given; see *Index muscorum*, 2, p. 56.
D. alopecurus C. Muell.
D. amabile C. Muell.
D. antarcticum (C. Muell.) Mitt.
D. arctocarpum Hornsch.
D. arenicola C. Muell.
D. areodictyon C. Muell.
D. asplenioides (Hedw.) Web. et Mohr
D. atro-viride Card.
D. aulacocarpum Mont.
D. aurificum C. Muell.
var. *flavescens* C. Muell.
D. australe Besch. ex C. Muell.
D. austro-alpinum C. Muell.
D. austro-georgicum C. Muell.
D. berteroanum Brid.
D. billardieri Brid.
var. *compactum* Card.
var. *flavicans* Card. (*Cardot, 1908a*).
Distr. Southern South America (*Bescherelle, 1889* (as *D. imponens* for île Wellington, leg. Dr. Conraud *fide Cardot, 1908a*); *Cardot, 1908a*).
D. blindiaceum C. Muell.
D. boryanum Schwaegr.
D. brachythysanos C. Muell.
D. bryoides (Hedw.) Sw.
D. cacti C. Muell.
D. callocaulon C. Muell.
D. canescens C. Muell.
D. capillare (Dus.) Card.
D. capillifolium Card.
D. cardotii R. Brown ter.
D. chilense De Not.
D. chiloense C. Muell. (*Mueller in Kindberg, 1889*) *nom. nud.*
Distr. No locality given; see *Index muscorum*, 2, p. 63.
D. chrismarii C. Muell.
D. cirrhifolium Schimp. ex Dus.
D. clavatum R. Brown
D. concolor Hook.
D. cryptodon (Mont.) Mitt.
D. dichotomum (P. Beauv.) Brid.

- Chorisodontium aciphyllum***
Campylopus acuminatus
Fissidens adianthoides
Campylopus alopecurus
Campylopus amabilis
Dicranoweisia antarctica
Campylopus arctocarpus
Campylopus surinamensis
Campylopus areodictyon
Fissidens asplenioides
Chorisodontium aciphyllum
Dicranella aulacocarpa
Campylopus introflexus
Campylopus aurificus* var. *flavescens
Dicranoloma australe
Campylopus austro-alpinus
Dicranoloma hariotii
Syrrhopodon incompletus* var. *berteroanus
Dicranoloma billardieri
Dicranoloma billardieri* var. *compactum
Chorisodontium spegazzinii
Dicranoloma billardieri
Campylopus brachythysanos
Fissidens bryoides
Campylopus cacti
Dicranoloma billardieri
Campylopus canescens
Dicranoloma capillare
Dicranoloma menziesii
Dicranella cardotii
Dicranoloma nigricaulle
Campylopus chrismarii
Platyneurum laticostatum
Campylopus clavatus
Campylopus concolor
Camptodontium cyptodon
Dicranoloma billardieri

<i>D. dicranellatum</i> Dus.	<i>Chorisodontium dicranellatum</i>
<i>D. elegans</i> Dub.	<i>Anisothecium elegans</i>
<i>D. euchlorum</i> Mont.	<i>Microdus euchlorus</i>
<i>D. falklandicum</i> Card.	<i>Dicranoloma falklandicum</i>
<i>D. flavissimus</i> C. Muell.	<i>Campylopus flavissimus</i>
<i>D. flexifolium</i> Hook.	<i>Ditrichum difficile</i>
<i>D. flexuosum</i> Hedw.	<i>Campylopus flexuosus</i>
var. <i>minus</i> Schwaegr.	<i>Campylopus flexuosus</i>
<i>D. fuegianum</i> (Dus.) Card.	<i>Dicranoloma hariotii</i>
var. <i>laxum</i> (Dus.) Card.	<i>Dicranoloma hariotii</i>
<i>D. gayanum</i> Mont.	<i>Aongstroemia gayana</i>
<i>D. gracilescens</i> Dus.	<i>Blindia curviseta</i>
<i>D. grandialare</i> Dus.	<i>Dicranoloma robustum</i>
<i>D. grimmoides</i> C. Muell.	<i>Campylopus grimmoides</i>
<i>D. griseum</i> (Hornschr.) C. Muell.	<i>Campylopus griseus</i>
<i>D. guilleminianum</i> Mont.	<i>Dicranella guilleminiana</i>
<i>D. hariotii</i> C. Muell.	<i>Dicranoloma hariotii</i>
<i>D. hatscheri</i> Dus.	<i>Chorisodontium aciphyllum</i>
<i>D. helvolum</i> C. Muell.	<i>Dicranoloma billardieri</i>
<i>D. hilarianum</i> Mont.	<i>Dicranella hilarihana</i>
<i>D. hookeri</i> Schimp.	<i>Dicranoloma robustum</i>
<i>D. humifugum</i> C. Muell.	<i>Campylopus humifugus</i>
<i>D. imponens</i> Mont.	<i>Dicranoloma imponens</i>
<i>D. incrassatum</i> (C. Muell.) C. Muell.	<i>Campylopus incrassatus</i>
<i>D. inerme</i> Mitt.	<i>Holodontium inerme</i>
<i>D. introflexum</i> Hedw.	<i>Campylopus introflexus</i>
<i>D. involutifolium</i> Sull.	<i>Dicranoloma imponens</i>
<i>D. jamesonii</i> Hook.	<i>Campylopus jamesonii</i>
<i>D. jamesonii</i> Tayl.	<i>Anisothecium vaginatum</i>
<i>D. lanigerum</i> C. Muell.	<i>Chorisodontium lanigerum</i>
<i>D. laticostatum</i> Card.	<i>Platyneurum laticostatum</i>
 <i>D. leucobryoides</i> Besch. ex C. Muell. (Bescherelle ex Mueller, 1901a [1900]).	
Distr. Southern South America (Mueller, 1901a).	
 <i>D. leucopterum</i> C. Muell.	<i>Chorisodontium aciphyllum</i>
var. <i>oncophoropsis</i> C. Muell. (Mueller, 1885).	
Distr. Southern South America (Mueller, 1885; Paris, 1904b; Cardot, 1908a).	
 <i>D. lonchotrix</i> C. Muell. (Mueller, 1901a [1900]) <i>nom. nud.</i>	
Distr. Southern South America (Mueller, 1901a).	
 <i>D. macropus</i> Kunz. ex C. Muell.	<i>Chorisodontium macropus</i>
 <i>D. magellanicum</i> Card.	<i>Chorisodontium magellanicum</i>
 <i>D. menziesii</i> Hook. f. et Wils.	<i>Dicranoloma menziesii</i>
var. <i>fernandezianum</i> Card. ex Thér.	<i>Dicranoloma menziesii</i> var. <i>fernandezianum</i>
 <i>D. microdus</i> Lor.	<i>Chorisodontium microdus</i>
 <i>D. morenoi</i> C. Muell.	<i>Campylopus morenoi</i>

D. nano-cespitosum C. Muell. (Mueller *in Kindberg*, 1891) *nom. nud.*

Distr. No locality given; see *Index muscorum*, 2, p. 86.

D. nigricaule Aongstr.

Dicranoloma nigricaule

var. *flexuosulum* (Dus.) Card.

Dicranoloma nigricaule* var. *flexuosulum

D. nordenskioeldii Card.

Chorisodontium aciphyllum

D. oleodictyon Dix. (Dixon, 1934).

Distr. South Georgia (Dixon, 1934; Steere, 1961b; Bell, 1973b).
Antarctic, peninsula regionN (Bell, 1973a).

D. orthocomum Besch. ex C. Muell.

Chorisodontium aciphyllum

D. osmundoides (Hedw.) Turn.

Fissidens osmundoides

D. percompactum Dus.

Chorisodontium magellanicum

D. perhorridum (Dus.) Card.

Campylopus perhorridus

D. perincanum C. Muell.

Campylopus canescens

D. perremotifolium (Dus.) Card.

Dicranoloma perromotifolium

D. peruncinatum (Dus.) Card.

Dicranoloma nigricaule

D. pseudo-australe C. Muell.

Dicranoloma australe

D. pseudo-leucoloma C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Kühnemann, 1938).

D. pseudo-truncorum C. Muell. (Mueller *in Kindberg*, 1889) *nom. nud.*

Distr. No locality given; see *Index muscorum*, 2, p. 93.

D. pumilum (Mitt.) Mitt.

Holodontium pumilum

D. pungens Hook. f. et Wils.

Dicranoloma robustum

D. purpureum Hedw.

Ceratodon purpureus

D. racovitzae Card.

Dicranoloma nigricaule

D. ramulosum Mitt.

Holodontium ramulosum

D. recurvipilum C. Muell.

Campylopus griseus

D. richardii (Brid.) C. Muell.

Campylopus richardii

D. rigens Besch.

Chorisodontium aciphyllum

D. robustum Hook. f. et Wils.

Dicranoloma robustum

Card. (Cardot, 1908a).

Distr. Southern South America (Cardot, 1901 (as *D. robustum* *fide* Cardot, 1908a); Dusén, 1903a (p. 68 as *D. australe* *fide* Cardot, 1908a); Cardot, 1908a).

var. *lagunicola* (Dus.) Card.

Dicranoloma imponens

var. *pungens* (Hook. f. et Wils.) Hook. f.

Dicranoloma robustum

D. saddleanum Besch. ex C. Muell.

Campylopus vesticaulis

D. scaberrimum Dus.

Chorisodontium aciphyllum

D. scabrophyllyum C. Muell.

Campylopus scabrophyllyum

D. scoparium Hedw. var. ***reflectens*** Brid. (Bridel, 1826).

Distr. Southern South America (Montagne, 1845b, 1850; Kühnemann, 1938).

D. sellowii Hornsch.

Dicranella hilariana

D. semicompletum (Hedw.) Mohr

Fissidens semicompletus

D. setosum Hook. f. et Wils.

Dicranoloma robustum* var. *setosum

var. *strictifolium* (Dus.) Card.

Dicranoloma robustum* var. *strictifolium

D. skottsbergii Card.

Dicranoloma skottsbergii

<i>D. spegazzinii</i> C. Muell.	<i>Chorisodontium spegazzinii</i>
<i>D. starkei</i> Web. et Mohr	<i>Kiaeria starkei</i>
<i>D. stenocarpum</i> (Hamp.) C. Muell.	<i>Campylopus stenocarpus</i>
<i>D. stenopus</i> C. Muell.	<i>Campylopus stenocarpus</i>
<i>D. strictum</i> Schwaegr.	<i>Pilopogon gracilis</i>
<i>D. suberectum</i> Hamp.	<i>Dicranoloma menziesii</i>
<i>D. subimponens</i> Card.	<i>Dicranoloma subimponens</i>
 <i>D. syrrhopodontoides</i> C. Muell. (Mueller, 1897a).	
Syn. <i>Leucoloma syrrhodictyon</i> (C. Muell.) Broth. (Brotherus, 1901).	
Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1901).	
<i>D. taxifolium</i> (Hedw.) Web. et Mohr	<i>Fissidens taxifolius</i>
<i>D. tenuicupidatum</i> C. Muell.	<i>Chorisodontium aciphyllum</i>
<i>D. tenuifolium</i> Hook. f. et Wils.	<i>Blindia robusta</i>
<i>D. tenuirostre</i> Kunz. ex Schwaegr.	<i>Dicranella hilariana</i>
 <i>D. toninii</i> C. Muell. (Mueller, 1897a).	
Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Cardot, 1908a).	
<i>D. truncatum</i> (C. Muell.) C. Muell.	<i>Campylopus truncatus</i>
<i>D. truncorum</i> Schimp. ex C. Muell.	<i>Dicranoloma truncorum</i>
<i>D. turgidissimum</i> Dus.	<i>Dicranoloma subimponens</i>
<i>D. vaginatum</i> Hook.	<i>Anisothecium vaginatum</i>
<i>D. vaginatum</i> Hook. f. et Wils.	<i>Dicranella hookeri</i>
<i>D. verticillatum</i> Hamp.	<i>Campylopus verticillatus</i>
<i>D. xanthophyllum</i> Mont. ex Par.	<i>Campylopus introflexus</i>
 <i>Didymodon</i>	
 <i>D. amblyophyllum</i> (Hook.) Broth. (Brotherus, 1902).	
Syn. <i>Gymnostomum amblyophyllum</i> Hook. (Hooker, 1830).	
<i>Pottia amblyophylla</i> (Hook.) C. Muell. (Mueller, 1849).	
<i>Tortula amblyophylla</i> (Hook.) Mitt. (Mitten, 1869).	
<i>Barbula amblyophylla</i> (Hook.) Jaeg. (Jaeger, 1873).	
Distr. Southern South America (Hooker, 1830; Mueller, 1849; Mitten, 1869; Jaeger, 1873; Paris, 1904b; Warnstorf, 1916; Brotherus, 1924a; Kühnemann, 1938).	
 <i>D. ampliretis</i> Bartr. (Bartram, 1946).	
Distr. Southern South America (Bartram, 1946).	
 <i>D. andreaeoides</i> Card. et Broth. (Cardot and Brotherus, 1923).	
Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).	
<i>D. angustifolius</i> Herz. (Herzog, 1960) hom. illeg.	
Distr. Southern South America (Bartram, 1965 (as <i>D. angustatus</i> Herz.)).	
 <i>D. argentinicus</i> (Par.) Par. (Paris, 1904b).	
Syn. <i>Barbula argentinica</i> Par. (Paris, 1894).	
<i>Barbula rubiginosa</i> C. Muell. (Mueller, 1882) hom. illeg.	
<i>Didymodon rubiginosus</i> Broth. (Brotherus, 1902) hom. illeg.	
<i>Barbula subrubiginosa</i> Kindb. (Kindberg, 1889) nom. nud.	
Distr. Southern South America (Mueller, 1882; Paris, 1894, 1904b; Brotherus, 1902, 1924a; Kühnemann, 1938).	
<i>D. argentinicus</i> Warnst. ex Broth.	<i>Didymodon argentinensis</i>

D. argentiniensis Warnst. (Warnstorf, 1916).

Syn. *Didymodon argentinicus* Warnst. ex Broth. (Warnstorf ex Brotherus, 1924a) *hom. illeg.*

Distr. Southern South America (Warnstorf, 1916; Brotherus, 1924a; Kühnemann, 1938).

D. berthoanus Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926).

D. brachydontius (Bruch) Wils.

Trichostomum brachydontium

D. brunneus (C. Muell.) Warnst. (Warnstorf, 1916).

Syn. *Trichostomum brunneum* C. Muell. (Mueller, 1879a).

Anacalypta brunnea (C. Muell.) Kindb. (Kindberg, 1889).

Pottia brunnea (C. Muell.) Par. (Paris, 1897).

Barbula brunnea (C. Muell.) Broth. (Brotherus, 1902) *hom. illeg.*

Trichostomum nudum C. Muell. (Mueller, 1882).

Anacalypta nuda (C. Muell.) Kindb. (Kindberg, 1889).

Pottia nuda (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1879a, 1882; Paris, 1897, 1905b; Brotherus, 1902, 1924a; Warnstorf, 1916; Hosseus, 1935b, 1937; Kühnemann, 1938).

D. calympericidyon Broth.

Bryoerythrophyllum campylocarpum

D. capillaceus (Hedw.) Web. et Mohr

Distichium capillaceum

var. *compactus* Hueb.

Distichium capillaceum var. *compactum*

D. crassinervis Thér. (Thériot in Costes, 1921) *nom. nud.*

Distr. Southern South America (Costes, 1921 (as *Didymodon*)).

D. cyathicarpus (Mont.) Mitt.

Amphidium tortuosum

D. dusenii Card. (Cardot in Cardot and Brotherus, 1923) *nom. nud.*

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

D. gelidus Card. (Cardot, 1907a).

Distr. Antarctic, peninsula region (Dixon, 1920).

Antarctic, continental region (Cardot, 1907a, 1908a; Brotherus, 1924a; Steere, 1961a).

D. glacialis Hook. f. et Wils.

Sarconeurum glaciale

D. glauco-viridis (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Trichostomum glauco-viride* C. Muell. (Mueller, 1882).

Anacalypta glauco-viridis C. Muell. (Mueller, 1882) *nom. nud.*

Pottia glauco-viridis (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1882; Paris, 1897, 1904b; Brotherus, 1902, 1924a; Warnstorf, 1916; Hosseus, 1938c; Kühnemann, 1938).

D. gracilescens Bartr. (Bartram, 1936) *hom. illeg.*

Distr. Southern South America (Bartram, 1936; Hosseus, 1938c, d, 1939).

D. gracilis Hook.

Pilopogon gracilis

var. *robustum* Hook.

Pilopogon gracilis

D. gymnus (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Trichostomum gymnum* C. Muell. (Mueller, 1879a).

Anacalypta gymna (C. Muell.) Kindb. (Kindberg, 1889).

Pottia gymna (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1879a; Paris, 1897, 1904b; Warnstorf, 1916; Brotherus, 1902, 1918, 1924a; Kühnemann, 1938).

D. heimii (Hedw.) Kindb.

Pottia heimii

D. heteromallus (Hedw.) Hook. et Tayl.

Ditrichum heteromallum

D. homomallus Hedw.

Ditrichum heteromallum

D. inundatus (Mitt.) Broth. (Brotherus, 1902).

Syn. *Tortula inundata* Mitt. (Mitten, 1869).
Barbula inundata (Mitt.) Jaeg. (Jaeger, 1873).

Distr. Southern South America (Mitten, 1869; Gibert, 1873; Jaeger, 1873; Brotherus, 1902, 1924a; Paris, 1904b; Herter, 1928, 1933a; Hosseus, 1938d).

D. linearis Broth.

Bryoerythrophyllum campylocarpum

D. longifolius (Brid.) Hook. f. et Wils.

Leucoloma longifolium

var. *curvifolius* Hook. f. et Wils. (Hooker and Wilson *in* Wilson and Hooker, 1847).

Distr. Southern South America (Wilson and Hooker, 1847; Kühnemann, 1938).

var. *penicillatus* Hook. f. et Wils.

Ditrichum strictum

var. *tenuifolius* Hook. f. et Wils.

Ditrichum hyalinum

D. lorentzianus (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Pottia lorentziana* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Kurtz, 1904; Paris, 1904b; Warnstorf, 1916; Hosseus, 1935b, 1937; Kühnemann, 1938).

D. microthecius (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Trichostomum microthecium* C. Muell. (Mueller, 1882).

Anacalypta microthecium C. Muell. (Mueller, 1882) *nom. nud.*

Pottia microthecia (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1882; Paris, 1897, 1904b; Brotherus, 1902, 1924a; Warnstorf, 1916; Kühnemann, 1938).

D. montevidensis Broth. (Brotherus *in* Felippone, 1917).

Distr. Southern South America (Felippone, 1917; Brotherus, 1924a; herter, 1933a).

D. oligodontus Card. (Cardot *in* Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b).

D. orbignyanus (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Pottia orbigniana* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a, c; Paris, 1904b; Warnstorf, 1916; Espinosa B., 1941).

D. patagonicus (Mitt.) Broth. (Brotherus, 1924a).

Syn. *Tortula patagonica* Mitt. (Mitten, 1869).

Barbula patagonica (Mitt.) Jaeg. (Jaeger, 1873).

Distr. Southern South America (Mitten, 1869; Jaeger, 1873; Paris, 1904a; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

D. pichinchensis Tayl.

Zygodon pichinchensis

D. polyccephalus Mont. (Montagne, 1845a).

Distr. Southern South America (Montagne, 1845a, 1856; Mueller, 1848a; Jaeger, 1873).

D. pungens Mitt.

Leptodontium pungens

D. reflexus Thér. (Thériot, 1924).

Distr. Southern South America (Thériot, 1924; Brotherus, 1925; Hosseus, 1938d).

D. reticulatus Gillies ex Grev. (Gillies ex Greville, 1830).

Distr. Southern South America (Greville, 1830).

D. rigidulus Hedw. (Hedwig, 1801).

- Syn.* *Bryum rigidulum* (Hedw.) Dicks. (Dickson, 1801).
Trichostomum rigidulum (Hedw.) Turn. (Turner, 1804a).
Desmatodon rigidulus Hassk. (Hasskarl, 1851) *err.*
Barbula rigidula (Hedw.) Mitt. (Mitten, 1864).
Tortula rigidula (Hedw.) Lindb. (Lindberg, 1864a).
- Distr.* Southern South America (Rovainen, 1934).
Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).
- D. rubellus* B.S.G. *Bryoerythrophyllum recurvirostre*
D. rubiginosus Broth. *Didymodon argentinicus*
- D. schilleri* Herz. et Thér.** (Herzog and Thériot *in Herzog and others*, 1933).
Distr. Southern South America (Herzog and others, 1933; Herzog, 1938, 1957; Kühnemann, 1938).
- D. schimperi* (Mont.) Broth.** (Brotherus, 1902).
Syn. *Trichostomum schimperi* Mont. (Montagne, 1845a).
Barbula schimperi (Mont.) Mitt. (Mitten, 1859).
Desmatodon schimperi (Mont.) Jaeg. (Jaeger, 1873).
Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Jaeger, 1873; Brotherus, 1902, 1924a; Paris, 1904b; Williams, 1930).
- D. spathulato-linearis* (C. Muell.) Broth.** (Brotherus, 1902).
Syn. *Trichostomum spathulato-lineare* C. Muell. (Mueller, 1879a).
Anacalypta spathulato-linearis (C. Muell.) Kindb. (Kindberg, 1889).
Pottia spathulato-linearis (C. Muell.) Par. (Paris, 1897).
Distr. Southern South America (Mueller, 1879a; Paris, 1897, 1904b; Brotherus, 1902; Warnstorff, 1916; Kühnemann, 1938).
- D. subtophaceus* Williams *Trichostomopsis australasiae*
- Dimerodontium***
- D. acuminatum* C. Muell.** (Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a; Brotherus, 1925; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939; Herzog, 1952a).
- D. aurescens* C. Muell.** (Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1925; Kühnemann, 1938; Hosseus, 1939).
- D. chlorophyllum* C. Muell.** (Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1925; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939; Herzog, 1952a).
- D. mendozense* Mitt.** (Mitten, 1869).
Distr. Southern South America (Mitten, 1869; Gibert, 1873; Jaeger, 1878; Mueller, 1879a; Paris, 1904b; Krieger, 1904; Kurtz, 1904; Brotherus, 1925; Herter, 1928, 1933a, 1939a, b; Hosseus, 1935a, b, 1936, 1937, 1938a, b, c, d; Kühnemann, 1938; Herzog, 1952a).
- D. ovatifolium* Herz.** (Herzog, 1916b).
Distr. Southern South America (Herzog, 1916b).
- D. pellucidum* (Schwaegr.) Mitt.** (Mitten, 1869).
Syn. *Sclerodontium pellucidum* Schwaegr. (Schwaegrichen, 1830).
Leskeia pellucida Hook. (Hooker *in Schwaegrichen*, 1830) *nom. nud.*
Hypnum sclerodontium C. Muell. (Mueller, 1851a).
Fabriona longidens Dub. (Duby, 1868).
Distr. Southern South America (Duby, 1868; Mitten, 1869; Jaeger, 1878; Paris, 1904b; Thériot, 1919; Brotherus, 1925; Kühnemann, 1938).

D. rivulare C. Muell.

Amblystegium rivulare

D. schnyderi C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1925; Kühnemann, 1938).

Diplostichum

D. longirostre (Brid.) Mont. (Montagne, 1845a).

Syn. *Pterigynandrum longirostre* Brid. (Bridel, 1819).
Eustichia longirostris (Brid.) Brid. (Bridel, 1827b).

Distr. Juan Fernandez (Robinson, 1975).

Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Lorentz, 1866; Jaeger, 1873; Paris, 1904b; Seki, 1974).

D. lorentzii C. Muell. (Mueller, 1897a).

Syn. *Eustichia lorentzii* (C. Muell.) Par. (Paris, 1900a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a; Kurtz, 1904; Brotherus, 1924a; Hosseus, 1936, 1937; Kühnemann, 1938).

D. poeppigii C. Muell. ex Card. ex Thér. (Mueller ex Cardot ex Thériot, 1921a).

Syn. *Eustichia poeppigii* Par. (Paris, 1900a) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959).

Southern South America (Paris, 1900a; Cardot, 1908a; Campo, 1921; Thériot, 1921a; Cardot and Brotherus, 1923; Brotherus, 1924a; Reimers, 1926; Roivainen, 1934; Kühnemann, 1938; Herzog, 1939, 1940, 1954, 1960; Bartram, 1952).

var. *laetevirens* Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

Discophyllum

D. dicksonii (Hook. et Grev.) Mitt.

Distichophyllum dicksonii

Dissodon

D. dubyanus Besch.

Tayloria dubyi

D. magellanicus (Brid.) Hamp.

Tayloria magellanica

D. mirabilis Card.

Tayloria mirabilis

D. plagiopus (Mont.) C. Muell.

Tayloria octoblepharum

Distichium

D. austro-georgicum C. Muell. (Mueller, 1890a).

Distr. South Georgia (Mueller, 1890a; Paris, 1904b; Cardot, 1908a; Dixon, 1932; Steere, 1961b).

D. austro-inclinatum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

D. capillaceum (Hedw.) B.S.G. (Bruch and others, 1846).

Syn. *Cynodontium capillaceum* Hedw. (Hedwig, 1801).

Bryum capillaceum (Hedw.) Dicks. ex With. (Dickson ex Withering, 1801).

Didymodon capillaceus (Hedw.) Web. et Mohr (Weber and Mohr, 1803).

Trichostomum capillaceum (Hedw.) Sm. (Smith, 1803).

Cynodontium capillaceum (Hedw.) Brid. (Bridel, 1806).

Leptotrichum capillaceum (Hedw.) Mitt. (Mitten, 1860).

Ditrichum capillaceum (Hedw.) Watts et Whitel. (Watts and Whitelegge, 1902).

Distr. Southern South America (Wilson and Hooker, 1847; Dusén, 1903a, 1905a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Bartram, 1952).
 South Georgia (Brotherus, 1924a; Newton, 1977a).
 Antarctic, peninsula region (Cardot, 1900, 1901, 1908a, 1911b, d, 1913a; Paris, 1904b; Steere, 1961a (all these references as *D. capillaceum* var. *brevifolium* *fide* Newton, 1977b); Newton 1977b).
 Antarctic, continental region (the record for Cardot, 1907a given by Steere, 1961a and mentioned by Newton, 1977b is erroneous as is the Cardot, 1908a record for a Skottsberg specimen from Moss Island, see original Cardot papers).

var. *brevifolium* B.S.G.

Distichium capillaceum* var. *compactum

var. *compactum* (Hueb.) Torre et Sarnth. (Dalla Torre and Sarnthein, 1904).

Syn. *Didymodon capillaceus* (Hedw.) Web. et Mohr var. *compactus* Hueb. (Hübener, 1833).

Distichium capillaceum (Hedw.) B.S.G. var. *brevifolium* B.S.G. (Bruch and others, 1846) *nom. illeg.*

Distr. Southern South America (Roivainen and Bartram, 1937).
 Falkland Islands (Cardot and Brotherus, 1923).

***D. lorentzii* C. Muell. (Mueller, 1882).**

Distr. Southern South America (Mueller, 1882; Krieger, 1904; Paris, 1904b; Kühnemann, 1938).

Distichophyllum

***D. apiculatum* (Hook. f. et Wils.) Mitt.**

Eriopus apiculatus

***D. assimile* Broth. (Brotherus, 1924b).**

Syn. *Mniadelphus assimilis* Broth. (Brotherus *in* Fleischer, 1922) *nom. nud.*

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Welch, 1969; Matteri, 1975b; Robinson, 1975).

***D. cavifolium* (Card.) Card.**

Distichophyllum eremitae

var. *montanum* Card.

Distichophyllum eremitae

***D. crispatissimum* Dus.**

Distichophyllum ellipticum

***D. dicksonii* (Hook. et Grev.) Mitt. (Mitten, 1869).**

Syn. *Hookeria dicksonii* Hook. et Grev. (Hooker and Greville, 1825a).

Mniadelphus dicksonii (Hook. et Grev.) Hamp. (Hampe *in* Mueller, 1850).

Discophyllum dicksonii (Hook. et Grev.) Mitt. (Mitten, 1868).

Distr. Southern South America (Hooker and Greville, 1825a; Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1850, 1885; Sullivant, 1859; Mitten, 1869; Jaeger, 1877; Bescherelle, 1889; Paris, 1904b; Cardot, 1908a; Brotherus, 1925; Herzog, 1938, 1954; Kühnemann, 1938; Welch, 1969; Seki, 1974; Matteri, 1975a, b).

Falkland Islands (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Paris, 1904b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Matteri, 1975b).

***D. ellipticum* Herz. (Herzog, 1954).**

Syn. *Distichophyllum crispatissimum* Dus. (Dusén, 1903a) *nom. nud. fide* Matteri, 1975a.

Distr. Southern South America (Dusén, 1903a; Paris, 1904b; Herzog, 1939, 1954; Welch, 1969; Seki, 1974; Matteri, 1975a, b).

***D. eremitae* (Jaeg.) Par. (Paris, 1895).**

Syn. *Mniadelphus eremitae* Jaeg. (Jaeger, 1877).

Distichophyllum procumbens Mitt. (Mitten, 1869) *hom. illeg.*

Mniadelphus procumbens C. Muell. (Mueller, 1885) *hom. illeg.*

Mniadelphus cavifolius Card. (Cardot, 1905a) *fide* Matteri, 1975a.

Distichophyllum cavifolium (Card.) Card. (Cardot, 1908a) *fide* Matteri, 1975a.

Distichophyllum cavifolium (Card.) Card. var. *montanum* Card. (Cardot *in* Cardot and Brotherus, 1923) *nom. nud. fide* Matteri, 1975a.

Distr. Southern South America (Mitten, 1869; Jaeger, 1877; Mueller, 1885, 1889; Paris, 1895, 1904b; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Welch, 1969; Seki, 1974; Matteri, 1975a, b).
Falkland Islands (Matteri, 1975b).

D. fasciculatum Mitt. (Mitten, 1876).

Distr. Southern South America (Herzog, 1960).

D. fernandezianum Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924a, b; Welch, 1969; Matteri, 1975b; Robinson, 1975 (as synonym of *D. assimile*)).

D. flaccidum (Hook. f. et Wils.) Mitt. (Mitten, 1869).

Syn. *Hookeria flaccida* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Mniadelphus flaccidus (Hook. f. et Wils.) Hamp. (Hampe in Mueller, 1850).

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1850, 1885; Mitten, 1869; Jaeger, 1877; Paris, 1905a; Cardot, 1908a; Brotherus, 1925; Welch, 1969; Matteri, 1975a, b).

D. hahnianum Fleisch.

Eriopus apiculatus

D. fuegianum Matteri (Matteri, 1975a).

Distr. Southern South America (Matteri, 1975a, b).

D. grandifolium Matteri (Matteri, 1975b).

Distr. Southern South America (Matteri, 1975b).

D. kraussei (Lor.) Mitt. (Mitten, 1885).

Syn. *Mniadelphus kraussei* Lor. (Lorentz, 1864).
Mniadelphus krausei Lor. (Lorentz in Jaeger, 1877) err.
Mniadelphus nigricans C. Muell. (Mueller in Bescherelle, 1889) nom. nud. fide Matteri, 1975a.
Distichophyllum nigricans Besch. (Bescherelle, 1889) fide Matteri, 1975a.
Distichophyllum latilimbatum Card. (Cardot in Matteri, 1975a) nom. nud. fide Matteri, 1975a.

Distr. Southern South America (Lorentz, 1864, 1866; Jaeger, 1877; Bescherelle, 1889; Neger, 1899; Paris, 1904b, 1905a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938; Welch, 1969; Seki, 1974; Matteri, 1975a, b).

D. latilimbatum Card.

Distichophyllum kraussei

D. molle Besch.

Eriopus bescherellei

D. nanospathulatum Herz. (Herzog, 1954).

Distr. Southern South America (Herzog, 1954; Welch, 1969; Matteri, 1975b).

D. nigricans Besch.

Distichophyllum kraussei

D. patagonicum Besch. (Bescherelle, 1889).

Distr. Southern South America (Bescherelle, 1889; Paris, 1904b; Cardot, 1908a; Brotherus, 1925; Welch, 1969).

D. procumbens Mitt.

Distichophyllum eremitae

D. rotundifolium (Hook. f. et Wils.) C. Muell. et Broth. (Mueller and Brotherus, 1900).

Syn. *Hookeria rotundifolia* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Mniadelphus rotundifolius (Hook. f. et Wils.) C. Muell. (Mueller, 1851a).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938; Herzog, 1960; Welch, 1969; Seki, 1974; Matteri, 1975b).

D. subelimbatum Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Welch, 1969; Matteri, 1975b; Robinson, 1975).

D. submersum Matteri (Matteri, 1975a).

Distr. Southern South America (Matteri, 1975a, b).

D. theriotianum Matteri (Matteri, 1975b).

Distr. Southern South America (Matteri, 1975b).

Ditrichum

D. affine (C. Muell.) Hamp.

Ditrichum difficile

D. australe (Mitt.) Mitt.

Ditrichum strictum

D. capillaceum (Hedw.) Watts et Whitel.

Distichium capillaceum

D. conicum* (Mont.) Mitt. (Mitten, 1879).

Syn. *Aschistodon conicus* Mont. (Montagne, 1845a).

Cynontodium conicum (Mont.) Mitt. (Mitten, 1869).

Leptotrichum montagnei C. Muell. (Mueller, 1848a) *nom. illeg.*

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Lorentz, 1866; Mitten, 1869; Jaeger, 1873; Paris, 1904b; Dusén, 1905a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Thériot, 1935a; Roivainen and Bartram, 1937; Herzog, 1938, 1939, 1957; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

var. *glaciale* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Roivainen and Bartram, 1937).

D. cylindricarpum (C. Muell.) Hamp. (Hampe, 1880c).

Syn. *Leptotrichum cylindricarpum* C. Muell. (Mueller, 1851b).

Trichostomum cylindricarpum (C. Muell.) Wils. (Wilson *in Wilson*, 1859).

Trichostomum elongatum Hook. f. et Wils. (Hooker and Wilson *in Wilson*, 1854).

Cynontodium elongatum (Hook. f. et Wils.) Mitt. (Mitten, 1869).

Leptotrichum elongatum (Hook. f. et Wils.) Jaeg. (Jaeger, 1873).

Ditrichum elongatum (Hook. f. et Wils.) Mitt. (Mitten, 1883).

Distr. Southern South America (Mitten, 1869; Jaeger, 1873; Paris, 1904b; Dusén, 1905a; Cardot, 1908a; Herzog, 1923, 1957; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).

D. difficile (Dub.) Fleisch. (Fleischer, 1904).

Syn. *Trichostomum difficile* Dub. (Duby *in Moritzi*, 1846).

Leptotrichum difficile (Dub.) C. Muell. (Mueller, 1901a [1900]).

Leptotrichum affine C. Muell. (Mueller, 1847c).

Trichostomum affine (C. Muell.) Mont. (Montagne, 1850) *hom. illeg.*

Ditrichum affine (C. Muell.) Hamp. (Hampe, 1867b).

Cynontodium affine (C. Muell.) Mitt. (Mitten, 1869).

Dicranum flexifolium Hook. (Hooker, 1819) *hom. illeg.*

Leptotrichum flexifolium Mitt. (Mitten, 1856).

Ditrichum flexifolium (Mitt.) Hamp. (Hampe, 1867b).

Trichostomum laxifolium Hook. f. et Wils. (Hooker and Wilson *in Wilson*, 1854).

Ditrichum laxifolium (Hook. f. et Wils.) Mitt. (Mitten, 1883).

Leptotrichum laxifolium (Hook. f. et Wils.) C. Muell. (Mueller, 1901a [1900]).

Distr. Juan Fernandez (Mitten, 1869; Jaeger, 1873; Paris, 1904b; Skottsberg, 1914; Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Mueller, 1847c, 1848a; Montagne, 1850; Lorentz, 1866; Jaeger, 1873; Paris, 1904b; Dusén, 1905a; Cardot, 1908a; Thériot, 1915, 1921a; Herzog, 1923, 1938, 1954; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).

D. elongatum (Hook. f. et Wils.) Mitt.

Ditrichum cylindricarpum

D. flexifolium (Mitt.) Hamp.

Ditrichum difficile

D. fontanum Herz. (Herzog, 1954).

Distr. Southern South America (Herzog, 1954).

*See also footnote to entries for *Pseudodistichium atlanticum* Dix. and *P. taitaoense* Froehl.

- D. glaciale* (C. Muell.) O. Kuntze *Sarconeurum glaciale*
- D. hallei* Card. et Broth. (Cardot and Brotherus, 1923).
Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen, 1934; Roivainen and Bartram, 1937; Kühnemann, 1938).
- D. heteromallum* (Hedw.) Britt. (Britton, 1913).
Syn. *Weisia heteromalla* Hedw. (Hedwig, 1801).
Grimmia heteromalla (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Bryum heteromallum (Hedw.) P. Beauv. (Palisot de Beauvois, 1822) *hom. illeg.*
Didymodon heteromallus (Hedw.) Hook. et Tayl. (Hooker and Taylor, 1818).
Didymodon homomallus Hedw. (Hedwig, 1801).
Grimmia homomalla (Hedw.) Sm. (Smith, 1804b).
Leptotrichum homomallum (Hedw.) Hamp. (Hampe *in* Mueller, 1848a).
Ditrichum homomallum (Hedw.) Hamp. (Hampe, 1867b).
Bryum weisii Dicks. ex With. (Dickson ex Withering, 1801).
Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).
- D. homomallum* (Hedw.) Hamp. *Ditrichum heteromallum*
- var. *leptocephalum*** Card. et Broth. (Cardot and Brotherus, 1923).
Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).
- D. hookeri* (C. Muell.) Hamp. (Hampe, 1867b).
Syn. *Leptotrichum hookeri* C. Muell. (Mueller, 1848a).
Cynontodium hookeri (C. Muell.) Mitt. (Mitten, 1869).
Cynodontium hookeri (C. Muell.) Mitt. (Mitten, 1876).
Blindia leptotrichocarpa C. Muell. (Mueller, 1885).
Blindia leptotrichacea C. Muell. ex Par. (Mueller *ex* Paris, 1894) *err.*
Distr. Southern South America (Mueller, 1848a, 1885; Mitten, 1869; Jaeger, 1873; Bescherelle, 1889; Paris, 1894, 1904a, b; Dusén, 1905a; Cardot, 1905a, 1908a, 1913a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1954; Seki, 1974).
Falkland Islands (Cardot and Brotherus, 1923).
var. *strictiusculum* (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1959).
Syn. *Blindia leptotrichocarpa* C. Muell. var. *strictiuscula* C. Muell. (Mueller, 1885).
Distr. Southern South America (Mueller, 1885; Paris, 1904a; Cardot, 1908a; Kühnemann, 1938).
- D. hyalinocuspitatum* Card. (Cardot, 1906a).
Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).
Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).
South Georgia (Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b).
- D. hyalinum* (Mitt.) O. Kuntze (Kuntze, 1891).
Syn. *Leptotrichum hyalinum* Mitt. (Mitten, 1860).
Cynontodium hyalinum (Mitt.) Mitt. (Mitten, 1869).
Didymodon longifolius (Brid.) Hook. f. et Wils. var. *tenuifolius* Hook. f. et Wils. (Hooker and Wilson *in* Hooker, 1847).
Distr. Southern South America (Mitten, 1860, 1869; Jaeger, 1873; Mueller, 1885; Bescherelle, 1889; Dusén, 1903a, 1905a; Paris, 1904b; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1954; Seki, 1974).
Falkland Islands (Wilson and Hooker, 1847; Mitten, 1860; Paris, 1904b; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).
- D. inundatum* Card. *Blindia magellanica* var. *inundata*
- D. laxifolium* (Hook. f. et Wils.) Mitt. *Ditrichum difficile*
- D. longisetum* (Lor.) Hamp. (Hampe, 1867b).
Syn. *Leptotrichum longisetum* Lor. (Lorentz, 1866).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Lorentz, 1866; Jaeger, 1873; Bescherelle, 1889; Paris, 1904b; Dusén, 1905a; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1939; Seki, 1974).

D. perporodictyon Dix. (Dixon, 1932).

Distr. South Georgia (Dixon, 1932; Steere, 1961b).

D. praealtum (Mitt.) O. Kuntze (Kuntze, 1891).

Syn. *Leptotrichum praealtum* Mitt. (Mitten, 1860).

Distr. Southern South America (Mitten, 1860; Jaeger and Sauerbeck, 1879b; Mueller, 1885; Paris, 1904b; Cardot, 1908a; Brotherus, 1924a).

D. roivanenii Bartr. et Roiv. (Bartram and Roivainen in Roivainen and Bartram, 1937).

Syn. Southern South America (Roivainen and Bartram, 1937).

D. rufescens (Hamp.) Hamp. (Hampe, 1867b).

Syn. *Leptotrichum rufescens* Hamp. (Hampe, 1862a).

Trichostomum strictum Web. et Mohr (Weber and Mohr, 1803).

Ditrichum strictum (Web. et Mohr) Card. (Cardot, 1905a) *hom. illeg.*

Ditrichum stenostomum Card. (Cardot, 1908a).

Distr. Southern South America (Jaeger, 1873; Mueller, 1879a; Cardot, 1905a, 1908a; Brotherus, 1924a; Roivainen and Bartram, 1937; Kühnemann, 1938).
Falkland Islands (Cardot, 1908a; Kühnemann, 1938).

D. stenostomum Card.

Ditrichum rufescens

D. strictum (Hook. f. et Wils.) Hamp. (Hampe, 1867b).

Syn. *Lophiodon strictus* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Trichostomum australe Mitt. (Mitten in Wilson, 1859).

Leptotrichum australe Mitt. (Mitten, 1860) *nom. illeg.*

Cynontodium australe Mitt. (Mitten, 1869).

Ditrichum australe (Mitt.) Mitt. (Mitten, 1879) *nom. illeg.*

Didymodon longifolius (Brid.) Hook. f. et Wils. var. *penicillatus* Hook. f. et Wils. (Hooker and Wilson in Wilson and Hooker, 1847).

Distr. Southern South America (Jaeger, 1873; Paris, 1905a; Brotherus, 1924a).

Falkland Islands (Wilson and Hooker, 1847; Mitten, 1860; Paris, 1905a; Brotherus, 1924a).

D. strictum (Web. et Mohr) Card.

Ditrichum rufescens

Dorcadion nom. illeg.

D. rupestre (Schwaegr.) Lindb.

Orthotrichum rupestre

Drepanium

D. circinale (Hook.) Roth

Hypnum circinale

D. cypressiforme (Hedw.) Roth

Hypnum cypressiforme

D. revolutum (Mitt.) C. Jens.

Hypnum revolutum

Drepanocladus

D. abbreviatus Card. et Broth.

Drepanocladus brachiatus var. ***abbreviatus***

D. aduncus (Hedw.) Warnst. (Warnstorff, 1903).

Syn. *Hypnum aduncum* Hedw. (Hedwig, 1801).

Amblystegium aduncum (Hedw.) De Not. (De Notaris, 1867).

Harpidium aduncum (Hedw.) Con. (Conard, 1945) *nom. inval.*

Distr. Southern South America (Sullivant, 1859; Cardot, 1908a).

Falkland Islands (Dumont d'Urville, 1825; Gaudichaud, 1825, 1826; Bridel, 1827b).

var. *polycarpus* (Voit) Roth (Roth, 1904).

- Syn. *Hypnum polycarpon* Bland. ex Voit (Blandow ex Voit in Sturm, 1813).
Amblystegium polycarpum (Voit) Vent. et Bott. (Venturi and Bottini, 1884).
Drepanocladus polycarpus (Voit) Warnst. (Warnstorf, 1903).
Harpidium polycarpum (Voit) Williams (Williams, 1903b).
Hypnum aduncum Hedw. var. *polycarpum* (Voit) B.S.G. (Bruch and others, 1854).
Hypnum aduncum Hedw. ssp. *polycarpum* (Voit) Meyl. (Meylan, 1905).
Hypnum fluitans Hedw. var. *polycarpon* (Voit) Fiedl. (Fiedler, 1844).
Amblystegium kneiffii B.S.G. var. *polycarpum* (Voit) Braithw. (Braithwaite, 1898).
Hypnum kneiffii (B.S.G.) Schimp. var. *polycarpus* (Voit) Fam. (Familler, 1913).
Amblystegium varium (Hedw.) Lindb. var. *patagonicum* Card. (Cardot in Cardot and Brotherus, 1923) *nom. nud.*

Distr. Southern South America (Cardot and Brotherus, 1923).

var. <i>plumosus</i> (Schimp.) Warnst.	<i>Drepanocladus uncinatus</i>
var. <i>subjulaceum</i> (B.S.G.) Warnst.	<i>Drepanocladus uncinatus</i> var. <i>subjulaceus</i>
<i>D. austro-stramineus</i> (C. Muell.) Broth. ex Par.	<i>Calliergidium austro-stramineum</i>
var. <i>gracillimus</i> (C. Muell.) Card. et Broth.	<i>Calliergidium austro-stramineum</i> var. <i>gracillimum</i>
var. <i>minus</i> (Card.) Steere	<i>Calliergidium austro-stramineum</i> var. <i>minus</i>
var. <i>subfluitans</i> (C. Muell.) Steere	<i>Calliergidium austro-stramineum</i> var. <i>subfluitans</i>

var. *tenuissimus* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. South Georgia (Cardot and Brotherus, 1923; Steere, 1961b).

D. brachiatus (Mitt.) Dix. (Dixon, 1912).

- Syn. *Hypnum brachiatum* Mitt. (Mitten in Hooker, 1867).
Amblystegium brachiatum (Mitt.) Broth. (Brotherus in Mueller and Brotherus, 1900).
Amblystegium longifolium Mitt. (Mitten, 1869).
Hypnum longifolium (Mitt.) Jaeg. (Jaeger and Sauerbeck, 1879a) *hom. illeg.*
Drepanocladus longifolius (Mitt.) Broth. ex Par. (Brotherus ex Paris, 1909).

Distr. Southern South America (Jaeger and Sauerbeck, 1879a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Dixon, 1929a, b; Roivainen, 1934; Kühnemann, 1938; Bartram, 1952; Herzog, 1957).
Falkland Islands (Mitten, 1869; Jaeger and Sauerbeck, 1879a; Paris, 1904c; Cardot, 1908a; Cardot and Brotherus, 1923; Dixon, 1929b).
South Georgia (Dixon, 1932; Steere, 1961b).
Antarctic, peninsula region (Dixon, 1929b, 1935).

var. *abbreviatus* (Card. et Broth.) Dix. (Dixon, 1929b).

Syn. *Drepanocladus abbreviatus* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Dixon, 1929b).

D. exannulatus (B.S.G.) Warnst. (Warnstorf, 1903).

- Syn. *Hypnum exannulatum* B.S.G. (Bruch and others, 1854).
Amblystegium exannulatum (B.S.G.) De Not. (De Notaris, 1869).
Amblystegium fluitans (Hedw.) De Not. var. *exannulatum* (B.S.G.) Lindb. (Lindberg and Arnell, 1890) *nom. inval.*
Amblystegium fluitans (Hedw.) De Not. ssp. *exannulatum* (B.S.G.) Arn. et C. Jens. (Arnell and Jensen, 1910).

Distr. Southern South America (Seki, 1974).

D. fluitans (Hedw.) Warnst. (Warnstorf, 1903).

- Syn. *Hypnum fluitans* Hedw. (Hedwig, 1801).
Harpidium fluitans (Hedw.) Spruc. (Spruce, 1867).
Amblystegium fluitans (Hedw.) De Not. (De Notaris, 1867).
Hypnum fluitans Hedw. var. *falcifolium* Ren. (Renauld in Husnot, 1894).
Amblystegium fluitans (Hedw.) De Not. var. *falcifolium* (Ren.) Arn. et C. Jens. (Arnell and Jensen, 1910) *nom. inval.*
Drepanocladus fluitans (Hedw.) Warnst. var. *falcifolium* (Ren.) Thér. (Thériot, 1924).

Distr. Southern South America (Montagne, 1845b, 1850; Wilson and Hooker, 1847; Sullivant, 1859; Eaton, 1892; Paris, 1904c; Cardot, 1908a; Thériot, 1924; Bartram, 1952; Seki, 1974).

var. *australe* (Card.) Broth. (*Brotherus*, 1908).

Syn. *Hypnum fluitans* Hedw. var. *australe* Card. (*Cardot*, 1905a).

Distr. Southern South America (Cardot, 1905a; *Brotherus*, 1908, 1925).

Falkland Islands (Cardot, 1905a, 1908a; *Brotherus*, 1908, 1925; Kühnemann, 1938).

var. *falcifolius* (Ren.) Thér.

Drepanocladus fluitans

D. fuegianus (Mitt.) Broth. ex Par. (*Brotherus* ex *Paris*, 1909).

Syn. *Amblystegium fuegianum* Mitt. (Mitten, 1869).

Hypnum fuegianum (Mitt.) C. Muell. (Mueller, 1885).

Distr. Southern South America (Mitten, 1869; Jaeger and Sauerbeck, 1879a; Mueller, 1885; Paris, 1904c; Cardot, 1908a; Cardot and Brotherus, 1923; *Brotherus*, 1925; Kühnemann, 1938).

var. *stenophyllum* Card. et Broth. (*Cardot* and *Brotherus*, 1923).

Distr. Falkland Islands (Cardot and *Brotherus*, 1923; Kühnemann, 1938).

D. kneiffii (B.S.G.) Warnst. **var. *chilense*** Thér. (*Thériot*, 1924).

Distr. Southern South America (*Thériot*, 1924).

D. laculosus (C. Muell.) Broth. (*Brotherus* in *Cardot* and *Brotherus*, 1923).

Syn. *Hypnum laculosum* C. Muell. (Mueller, 1885).

Hypnum fluitans Hedw. var. *laculosum* (C. Muell.) Par. (*Paris*, 1896).

Distr. Southern South America (Mueller, 1885; Paris, 1896, 1904c; Cardot, 1908a; Cardot and *Brotherus*, 1923; *Brotherus*, 1925; Kühnemann, 1938).

D. longifolius (Mitt.) Broth. ex Par.

Drepanocladus brachiatus

D. orthothecoides (Lindb.) Roth

Drepanocladus uncinatus* var. *subjulaceus

D. perfoliatus C. Muell. or Dus. (Mueller or Dusén in *Neger*, 1899) nom. nud.

Distr. Southern South America (*Neger*, 1899).

D. perplicatus (Dus.) Roth

Cratoneuron perplicatum

D. plicatus Dix. (Dixon, 1934).

Distr. South Georgia (Dixon, 1934; Steere, 1961b).

D. polycarpus (Voit) Warnst.

Drepanocladus aduncus* var. *polycarpus

D. revolvens (Sw.) Warnst. (*Warnstorff*, 1903).

Syn. *Hypnum revolvens* Sw. (Swartz, 1801).

Amblystegium revolvens (Sw.) De Not. (De Notaris, 1867).

Harpidium revolvens (Sw.) C. Jens. (Jensen in Lange and Jensen, 1887).

Limprechtia revolvens (Sw.) Loesk. (Loeske, 1907).

Hypnum aduncum Hedw. var. *revolvens* (Sw.) Bland. (Blandow, 1809).

Hypnum intermedium Lindb. var. *revolvens* (Sw.) San. (Sanione, 1883) nom. illeg.

Amblystegium intermedium Lindb. var. *revolvens* (Sw.) Vent. et Bott. (Venturi and Bottini, 1884) nom. illeg.

Distr. Southern South America (Wilson and Hooker, 1847; Cardot and *Brotherus*, 1923; Kühnemann, 1938).

D. secundifolius (C. Muell.) Dix. (Dixon, 1921).

Syn. *Hypnum secundifolium* C. Muell. (Mueller, 1851a).

Sematophyllum secundifolium (C. Muell.) Mitt. (Mitten, 1869).

Rapidostegium secundifolium (C. Muell.) Jaeg. (Jaeger, 1878).

Hypnum lithophilum Hook. f. et Wils. (*Hooker and Wilson* in *Wilson and Hooker*, 1847) hom. illeg. fide Dixon, 1921.

Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1851a, 1885; Mitten, 1869; Jaeger, 1878; Paris, 1905b; Cardot, 1908a; Dixon, 1921; *Brotherus*, 1925).

- D. subjulaceus* (B.S.G.) Roiv. *Drepanocladus uncinatus* var. *subjulaceus*
- D. symmetricus* (Ren. et Card.) Card. *Drepanocladus uncinatus* var. *symmetricus*
- D. uncinatus* (Hedw.) Warnst. (Warnstorf, 1903).**
- Syn. *Hypnum uncinatum* Hedw. (Hedwig, 1801).
Amblystegium uncinatum (Hedw.) De Not. (De Notaris, 1867).
Harpidium uncinatum (Hedw.) C. Jens. (Jensen in Lange and Jensen, 1887).
Sanionia uncinata (Hedw.) Loesk. (Loeske, 1907).
Hypnum georgico-uncinatum C. Muell. (Mueller, 1890a).
Hypnum uncinatum Hedw. var. *plumosum* Schimp. (Schimper, 1860).
Drepanocladus uncinatus (Hedw.) Warnst. var. *plumosus* (Schimp.) Warnst. (Warnstorf, 1903).
Amblystegium uncinatum (Hedw.) De Not. var. *plumosum* (Schimp.) Moell. (Möller, 1907).
Amblystegium aduncum (Hedw.) De Not. var. *plumosum* (Schimp.) Hult (Hult, 1886).
Drepanocladus aduncus Warnst. var. *plumosus* (Schimp.) Warnst. (Warnstorf, 1906).
- Distr. Southern South America (Wilson and Hooker, 1847; Ångström, 1872; Jaeger and Sauerbeck, 1879a; Dusén, 1903a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Roivainen, 1934; Kühnemann, 1938; Herzog, 1939, 1940, 1954, 1957; Seki, 1974).
Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938).
South Georgia (Mueller, 1890a; Paris, 1904c; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Dixon, 1932; Steere, 1961b).
Antarctic, peninsula region (Cardot, 1900, 1901, 1906a, b, 1907c, 1908a, 1911b, c, d, 1912b, 1913a; Paris, 1904c; Wright, 1905; Brotherus, 1925; Dixon, 1935; Bartram, 1957; Steere, 1961a; Greene, 1968a; Robinson, 1972).
- var. *orthotheciooides* (Lindb.) Broth. *Drepanocladus uncinatus* var. *subjulaceus*
- var. *plumosus* (Schimp.) Warnst. *Drepanocladus uncinatus*
- var. *stenocarpus*** Card. et Broth. (Cardot and Brotherus, 1923).
- Distr. Southern South America (Cardot and Brotherus, 1923).
- var. *subjulaceus* (B.S.G.) Warnst. (Warnstorf, 1903).**
- Syn. *Hypnum uncinatum* Hedw. var. *subjulaceum* B.S.G. (Bruch and others, 1854).
Amblystegium uncinatum (Hedw.) De Not. var. *subjulaceum* (B.S.G.) C. Jens. (Jensen, 1898 [1897]).
Drepanocladus subjulaceus (B.S.G.) Roiv. (Roivainen, 1954).
Drepanocladus aduncus Warnst. var. *subjulaceum* (B.S.G.) Warnst. (Warnstorf, 1906).
Hypnum uncinatum Hedw. var. *orthotheciooides* (Lindb.) Zett. (Zetterstedt, 1876).
Harpidium uncinatum (Hedw.) C. Jens. var. *orthotheciooides* (Lindb.) C. Jens. (Jensen in Lange and Jensen, 1887).
Amblystegium uncinatum (Hedw.) De Not. var. *orthotheciooides* (Lindb.) C. Jens. (Jensen, 1898 [1897]).
Drepanocladus uncinatus (Hedw.) Warnst. var. *orthotheciooides* (Lindb.) Broth. (Brotherus, 1910b).
Amblystegium aduncum (Hedw.) De Not. ssp. *orthotheciooides* (Lindb.) Lindb. (Lindberg, 1879).
Amblystegium orthotheciooides (Lindb.) Lindb. ex Par. (Lindberg ex Paris, 1894).
Hypnum orthotheciooides (Lindb.) Par. (Paris, 1904c).
Sanionia orthotheciooides (Lindb.) Loesk. (Loeske, 1907).
Drepanocladus orthotheciooides (Lindb.) Roth (Roth, 1904).
- Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).
South Georgia (Cardot and Brotherus, 1923; Steere, 1961b).
Antarctic, peninsula region (Cardot, 1913a).
- var. *symmetricus* (Ren. et Card.) Grout (Grout, 1929).**
- Syn. *Hypnum uncinatum* Hedw. ssp. *symmetricum* Ren. et Card. (Renauld and Cardot, 1888).
Amblystegium uncinatum (Hedw.) De Not. var. *symmetricum* (Ren. et Card.) Dus. (Dusén, 1903a).
Hypnum symmetricum (Ren. et Card.) Ren. et Card. (Renauld and Cardot, 1889).

- Sanionia symmetrica* (Ren. et Card.) Wheld. (Wheldon, 1921).
Drepanocladus symmetricus (Ren. et Card.) Card. (Cardot in Cardot and Brotherus, 1923).
Hypnum uncinatum Hedw. var. *cylindropyxis* Dus. (Dusén in Paris, 1905a) nom. nud.
Distr. Southern South America (Dusén, 1903a; Paris, 1905a; Cardot, 1908a, 1913a; Cardot and Brotherus, 1923; Kühnemann, 1938).

Drummondia

D. obtusifolia C. Muell. (Mueller, 1849).

Syn. *Leiotheca obtusifolia* (C. Muell.) Hamp. (Hampe, 1867a).

Distr. Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1874; Brotherus, 1925; Herzog, 1938, 1954, 1960).

Dryptodon

D. crispulus Hook. f. et Wils.

Racomitrium crispulum

D. donnianus (Sm.) Hartm.

Grimmia donniana

D. rupestris Hook. f. et Wils.

Racomitrium crispulum* var. *rupestre

D. trichophyllum (Grev.) Brid.

Grimmia trichophylla

Dusenia nom. illeg.

D. coronata (Mont.) C. Muell.

Forsstroemia coronata

Duseniella

D. genuflexa (C. Muell.) Broth.

Ancistrodes genuflexa

Ectropothecium

E. berberidis Dus.

Rhaphidorrhynchium berberidis

E. cygnisetum C. Muell.

Ptychomnion cygnisetum

E. pallido-nitidum (C. Muell.) Par.

Rhaphidorrhynchium pallido-nitidum

E. rutilans (Brid.) Mitt.

Vesicularia vesicularis

E. spirifolium Dus.

Vesiculariopsis spirifolium

E. sublimatum (Lor.) Par.

Hypnum chrysogaster

Encalypta

E. armata Broth. ex Dus. (Brotherus ex Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Brotherus, 1924a; Kühnemann, 1938).

E. asperifolia Mitt. (Mitten, 1869).

Syn. *Encalypta emersa* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

E. australis Mitt.

Encalypta vulgaris

E. austrociliata Broth. ex Dus. (Brotherus ex Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1957).

E. berthoana Thér. (Thériot, 1923b).

Distr. Southern South America (Thériot, 1923b; Brotherus, 1925).

E. crispata Hedw.

Ptychomitrium crispatum

E. emersa C. Muell.

Encalypta asperifolia

E. obtusata Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

E. patagonica Broth. (*Brotherus in Dusén*, 1905b).

Distr. Southern South America (Dusén, 1903a (as *Eucalypta*), 1905b, 1906; Paris, 1904b; Cardot, 1905a, 1908a; Brotherus, 1924a; Kühnemann, 1938).
Antarctic, peninsula region (Newton, 1974c).

E. procera Bruch (Bruch, 1832).

Distr. South Georgia (Newton, 1974c).
Antarctic, peninsula region (Newton, 1974c).

E. vulgaris Hedw. (Hedwig, 1801).

Syn. *Encalypta australis* Mitt. (Mitten in Wilson, 1859).
Distr. Southern South America (Mitten, 1869; Jaeger, 1873; Paris, 1904b).

Entodon***E. argentinicus*** C. Muell.***Erythrodontium argentinicum******E. beyrichii*** (Schwaegr.) C. Muell. (Mueller, 1844b [1845]).

Syn. *Neckera beyrichii* Schwaegr. (Schwaegrichen, 1827a).
Distr. Southern South America (Hosseus, 1940; Herzog, 1952a).

E. flavo-virens C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium flavo-virens* (C. Muell.) Par. (Paris, 1894).
Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Brotherus, 1925; Kühnemann, 1938).

var. *flagellaceus* C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium flavo-virens* (C. Muell.) Par. var. *flagellaceum* (C. Muell.) Par. (Paris, 1894).
Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Kühnemann, 1938).

E. laxo-alaris (C. Muell.) Broth. ex Par. (*Brotherus ex Paris*, 1909).

Syn. *Aptychus laxo-alaris* C. Muell. (Mueller, 1897a).
Rhaphidostegium laxo-alare (C. Muell.) Par. (Paris, 1897).
Distr. Southern South America (Mueller, 1897a; Paris, 1897, 1905b).

E. micans Herz. (Herzog, 1916a).

Distr. Southern South America (Hosseus, 1940).

E. platygyrioides C. Muell.***Erythrodontium platygyrioides******E. schreberi*** (Brid.) Moenk.***Pleurozium schreberi******E. suberythropus*** C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium suberythropus* (C. Muell.) Par. (Paris, 1894).
Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Kurtz, 1904; Brotherus, 1918, 1925; Hosseus, 1938b, c, d; Kühnemann, 1938; Herzog, 1952a).

var. *complanatulus* C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium suberythropus* (C. Muell.) Par. var. *complanatum* (C. Muell.) Par. (Paris, 1894).
Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Kühnemann, 1938).

var. *ellipticus* C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium suberythropus* (C. Muell.) Par. var. *ellipticum* (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Hosseus, 1935a, b, 1937, 1938b; Kühnemann, 1938).

var. *flaccidus* C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium suberythropus* (C. Muell.) Par. var. *flaccidum* (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Kühnemann, 1938).

var. *gracillimus* C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium suberythropus* (C. Muell.) Par. var. *gracillimum* (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Kühnemann, 1938).

var. *lutescens* C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium suberythropus* (C. Muell.) Par. var. *lutescens* (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Kühnemann, 1938).

var. *microtheca* C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium suberythropus* (C. Muell.) Par. var. *microtheca* (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Hosseus, 1935a, 1937, 1938d; Kühnemann, 1938).

var. *myosuroides* C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium suberythropus* (C. Muell.) Par. var. *myosuroides* (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Kühnemann, 1938).

var. *rufulus* C. Muell. (Mueller, 1879a).

Syn. *Cylindrothecium suberythropus* (C. Muell.) Par. var. *rufulum* (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Kühnemann, 1938).

***E. truncatus* (C. Muell.) Mitt. (Mitten, 1869).**

Syn. *Neckera truncata* C. Muell. (Mueller, 1850).

Cylindrothecium truncatum (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1850; Mitten, 1869; Jaeger, 1878; Paris, 1894, 1904b; Brotherus, 1925).

Entopogon nom. nud.

E. cylindricus C. Muell.

Eriodon conostomus

Entosthodon

E. acutifolius Hamp.

Funaria acutifolia

E. balansae Besch.

Funaria balansae

E. bonplandii (Hook.) Mitt.

Funaria bonplandii

E. brachyodus Mont.

Funaria integra

E. chiloënsis Mitt.

Funaria chiloënsis

E. clavellatus Mitt.

Funaria clavellata

E. flexisetus C. Muell.

Funaria flexiseta

E. mathewsi Hook. f. var. *integer* C. Muell.

Funaria integra

var. *integrifolius* Par.

Funaria integra

E. papillosum C. Muell.

Funaria papillosa

E. plagiothecius C. Muell.

Funaria plagiothecia

E. rhizomaticus C. Muell.

Funaria rhizomatrica

Ephemerum***E. conicum*** C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a, 1882; Roth, 1911; Kühnemann, 1938).

E. uleanum C. Muell. (Mueller, 1898c).

Distr. Southern South America (Hosseus, 1940; Herzog, 1952a).

*Eremodon nom. illeg.****E. magellanicus*** (Brid.) Brid.***Tayloria magellanica******E. spathulatus*** Hook. f. et Wils.***Tayloria spathulata******Eriodon******E. conostomus*** Mont. (Montagne, 1845a).Syn. *Hypnum conostomum* (Mont.) Mitt. (Mitten, 1869) *nom. illeg.**Neckera eriodon* C. Muell. (Mueller, 1850) *nom. illeg.**Hookeria eriodon* C. Muell. ex Card. (Mueller ex Cardot, 1908a) *err.**Entopogon cylindricus* C. Muell. (Mueller in Paris, 1895) *nom. nud.*

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1850; Lorentz, 1866; Mitten, 1869; Jaeger, 1878; Neger, 1899; Paris, 1904b; Cardot, 1908a; Thériot, 1918; Campo, 1921; Herzog, 1923, 1954; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938; Bartram, 1952; Gunckel, 1964; Seki, 1974).

Eriopus***E. apiculatus*** (Hook. f. et Wils.) Mitt. (Mitten, 1869).Syn. *Hookeria apiculata* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).*Pterygophyllum apiculatum* (Hook. f. et Wils.) Jaeg. (Jaeger, 1877).*Distichophyllum apiculatum* (Hook. f. et Wils.) Mitt. (Mitten, 1883).*Mniadelphus apiculatus* (Hook. f. et Wils.) Mitt. (Mitten in Paris, 1896).*Mniadelphus hahnii* C. Muell. (Mueller in Kindberg, 1889) *nom. nud. fide* Matteri, 1975a.*Mniadelphus hahnianus* C. Muell. (Mueller in Fleischer, 1922) *nom. nud. fide* Matteri, 1975a.*Distichophyllum hahnianum* Fleisch. (Fleischer, 1922) *nom nud. fide* Matteri, 1975a.*Eriopus apiculatus* (Hook. f. et Wils.) Mitt. var. *platyloma* Card. et Broth. (Cardot and Brotherus, 1923) *fide* Matteri, 1975a.

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1851a, 1885; Mitten, 1869; Jaeger, 1877; Paris, 1904b; Cardot, 1908a; Fleischer, 1922; Brotherus, 1925; Bartram, 1952; Herzog, 1954; Welch, 1969; Seki, 1974; Matteri, 1975a, b).

Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938; Welch, 1969; Matteri, 1975b).

var. *platyloma* Card. et Broth.***Eriopus apiculatus***var. *spathulus* Matteri (Matteri, 1975b).

Distr. Southern South America (Matteri, 1975b).

E. bescherellei Thér. (Thériot, 1925a).Syn. *Distichophyllum molle* Besch. (Bescherelle, 1889) *nom. nud. fide* Matteri, 1975a.*Mniadelphus mollis* C. Muell. (Mueller in Bescherelle, 1889) *nom. nud. fide* Matteri, 1975a.

Distr. Southern South America (Bescherelle, 1889; Paris, 1904b; Cardot, 1908a; Thériot, 1917b, 1925a; Costes, 1921; Brotherus, 1925; Herzog, 1954; Seki, 1974; Matteri, 1975a, b).

E. cristatus (Hedw.) Brid. (Bridel, 1827b).Syn. *Leskea cristata* Hedw. (Hedwig, 1801).*Chaetophora cristata* (Hedw.) Brid. (Bridel, 1819).*Mnium cristatum* (Hedw.) P. Beauv. (Palisot de Beauvois, 1822).*Hookeria cristata* (Hedw.) Hook. et Grev. (Hooker and Greville, 1825a).

Hypnum cristatum Poir. (Poiret in Bridel, 1827b) *nom. nud.*
Lepidopilum cristatum (Hedw.) Hamp. (Hampe, 1865).

Distr. Southern South America (Wilson and Hooker, 1847; Welch, 1969).

E. flexicollis (Mitt.) Jaeg. (Jaeger, 1877).

Syn. *Hookeria flexicollis* Mitt. (Mitten in Hooker, 1867).

Distr. Southern South America (Cardot and Brotherus, 1923; Welch, 1969; Seki, 1974).

E. grandiretis Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Bartram, 1959; Welch, 1969; Matteri, 1975b).

E. leptoloma Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Welch, 1969; Matteri, 1975b).

E. odontoloma Dus. ex Matteri (Matteri, 1975b).

Distr. Southern South America (Dusén, 1903a; Paris, 1904b; Seki, 1974; Matteri, 1975b).

Erpodium

E. balansae C. Muell.

Erpodium glaziouii

E. beccarii C. Muell. ex Vent. (Mueller ex Venturi, 1872).

Syn. *Erpodium lorentzianum* C. Muell. (Mueller, 1879a) *fide* Crum, 1972.

Erpodium lorentzii C. Muell. ex Kindb. (Mueller ex Kindberg, 1888) *nom. illeg.*

Erpodium paraguense Besch. (Bescherelle, 1877) *fide* Crum, 1972.

Distr. Southern South America (Mueller, 1879a; Jaeger and Sauerbeck, 1879b; Paris, 1904b; Brotherus, 1918, 1925; Kühnemann, 1938; Herzog, 1952a; Crum, 1972).

E. chlorophyllum C. Muell.

Erpodium coronatum

E. coronatum (Hook. et Wils.) Mitt. (Mitten, 1869).

Syn. *Anictangium coronatum* Hook. et Wils. (Hooker, W. J. and Wilson, 1844).

Erpodium chlorophyllum C. Muell. (Mueller, 1879a) *fide* Crum, 1972.

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1925; Kühnemann, 1938; Crum, 1972).

E. glaziouii Hamp. (Hampe, 1872).

Syn. *Erpodium balansae* C. Muell. (Mueller, 1887b) *fide* Crum, 1972.

Distr. Southern South America (Brotherus, 1925; Kühnemann, 1938; Hosseus, 1940; Herzog, 1952a).

E. lorentzinum C. Muell.

Erpodium beccarii

E. lorentzii C. Muell. ex Kindb.

Erpodium beccarii

E. paraguense Besch.

Erpodium beccarii

Erythrodontium

E. argentinicum (C. Muell.) Par. (Paris, 1904b).

Syn. *Entodon argentinicus* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1918, 1925; Kühnemann, 1938).

E. bicolor (Aongstr.) Broth.

Erythrodontium squarrosum

E. platygynoides (C. Muell.) Fleisch. (Fleischer, 1922).

Syn. *Entodon platygynoides* C. Muell. (Mueller, 1879a).

Cylindrothecium platygynoides (C. Muell.) Par. (Paris, 1894).

Distr. Southern South America (Mueller, 1879a; Paris, 1894, 1904b; Fleischer, 1922; Brotherus, 1925; Kühnemann, 1938).

E. squarrosum (Hamp.) Par. (Paris, 1904b).

Syn. *Leptohymenium squarrosum* Hamp. (Hampe, 1862b).

Erythrodontium bicolor (Aongstr.) Broth. (Brotherus in Ule, 1899).

Pterigynandrum bicolor Lindb. ex Aongstr. (Lindberg ex Ångström, 1876a).

Distr. Southern South America (Krieger, 1904; Hosseus, 1940; Herzog, 1952a).

Erythrophyllopsis

E. boliviensis Broth.

Erythrophyllopsis fuscula

E. fuscula (C. Muell.) Hilp. (Hilpert, 1933).

Syn. *Barbula fuscula* C. Muell. (Mueller, 1879a).

Erythrophyllopsis boliviensis Broth. (Brotherus in Herzog, 1916a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904a; Brotherus, 1924a; Hilpert, 1933; Hosseus, 1935b, 1937; Kühnemann, 1938; Bartram, 1965).

Eucamptodon

E. perichaetialis (Mont.) Mont. (Montagne, 1845a).

Syn. *Weisia perichaetialis* Mont. (Montagne, 1845a).

Distr. Juan Fernandez (Brotherus, 1924b; Robinson, 1975).

Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Mitten, 1869; Jaeger, 1872; Paris, 1904b; Dusén, 1905c; Cardot, 1908a; Herzog, 1923, 1938; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).

Eucatagonium nom. illeg.

E. politum (Hook. f. et Wils.) Broth.

Catagonium politum

var. *filescens* (Card. et Broth.) Herz.

Catagonium politum var. *filescens*

Eucladium

E. calcareum (Nees et Hornsch.) C. Jens.

Gymnostomum aeruginosum

E. recurvirostre (Hedw.) C. Jens.

Hymenostylium recurvirostre

E. verticillatum (Brid.) B.S.G. (Bruch and others, 1846).

Syn. *Weisia verticillata* Brid. (Bridel, 1801a).

Distr. Southern South America (Rovainen, 1934).

Euglossophyllum nom. inval.

E. brevipes (C. Muell.) Hamp.

Stereophyllum brevipes

Eugrimmia nom. nud.

E. trichophylla (Grev.) Hamp.

Grimmia trichophylla

Eurhynchiella

E. acanthophylla (Mont.) Fleisch. (Fleischer, 1923).

Syn. *Hypnum acanthophyllum* Mont. (Montagne, 1845a).

Rhynchostegium acanthophyllum (Mont.) Jaeg. (Jaeger, 1878) nom. inval.

Rhynchostegiella acanthophylla (Mont.) Broth. (Brotherus, 1909).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1851a; Mitten, 1869; Jaeger, 1878; Paris, 1905b; Thériot, 1915, 1918; Campo, 1921; Costes, 1921; Herzog, 1923, 1938, 1939, 1954, 1957, 1960; Brotherus, 1924c, 1925; Espinosa B., 1941; Seki, 1974).

E. fuegiana (Card.) Fleisch.

Oxyrrhynchium fuegianum

E. tenuinervis Herz. (Herzog, 1952a).

Distr. Southern South America (Herzog, 1952a).

Eurhynchium

<i>E. acanthophyllum</i> (Mont.) Dus. ex Thér.	<i>Oxyrrhynchium corralense</i>
<i>E. confusum</i> Thér.	<i>Oxyrrhynchium corralense</i>
<i>E. corralense</i> (Lor.) Jaeg.	<i>Oxyrrhynchium corralense</i>
<i>E. cuestarum</i> C. Muell. (Mueller, 1897a).	
Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b).	
<i>E. cygnisetum</i> C. Muell.	<i>Ptychomnion cygnisetum</i>
<i>E. dives</i> C. Muell. (Mueller, 1897a).	
Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b).	
<i>E. encalypratum</i> Schimp.	<i>Acrocladium auriculatum</i>
<i>E. eucalypratum</i> Schimp. ex Par.	<i>Acrocladium auriculatum</i>
<i>E. fuegianum</i> Card.	<i>Oxyrrhynchium fuegianum</i>
<i>E. pendulum</i> (Brid.) Jaeg.	<i>Helicodontium pendulum</i>
<i>E. pulchellum</i> (Hedw.) Jenn. (Jennings, 1913).	
Syn. <i>Hypnum strigosum</i> Hoffm. ex Web. et Mohr (Hoffman ex Weber and Mohr, 1803).	
Distr. Southern South America (Mueller, 1851a).	
<i>E. remotifolium</i> (Grev.) Jaeg.	<i>Oxyrrhynchium clinocarpum</i>
<i>E. riparioides</i> (Hedw.) Richs.	<i>Platyhypnidium riparioides</i>
<i>E. rusciforme</i> Mild.	<i>Platyhypnidium riparioides</i>
<i>E. striatellum</i> Schimp.	<i>Rigodium toxarium</i>
<i>Eustichia hom. illeg.</i>	
<i>E. longirostris</i> (Brid.) Brid.	<i>Diplostichum longirostre</i>
<i>E. lorentzii</i> (C. Muell.) Par.	<i>Diplostichum lorentzii</i>
<i>E. poeppigii</i> Par.	<i>Diplostichum poeppigii</i>

Exodokidium

<i>E. donatii</i> Herz. (Herzog, 1937).	
Distr. Southern South America (Herzog, 1937).	
<i>E. subsymmetricum</i> (Card.) Card.	<i>Bartramia subsymmetrica</i>

Fabronia

<i>F. affinis</i> Thér. (Thériot in Felippone, 1929 [1930]).	
Distr. Southern South America (Felippone, 1929; Herter, 1933a; Herzog, 1957).	
<i>F. andina</i> Mitt. (Mitten, 1869).	
Distr. Southern South America (Mitten, 1869; Jaeger, 1878; Paris, 1904a; Felippone, 1909; Brotherus, 1925; Herter, 1933a; Herzog, 1938, 1954).	
var. <i>jaffuelii</i> Thér. (Thériot, 1924).	
Distr. Southern South America (Thériot, 1924).	
<i>F. apophysatula</i> C. Muell. (Mueller, 1897a).	
Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Kühnemann, 1938; Herzog, 1960).	
<i>F. argentinica</i> C. Muell. (Mueller, 1879a).	
Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Hosseus, 1934, 1935b, 1937, 1939; Kühnemann, 1938).	

F. balansae Besch. (Bescherelle, 1891).

Syn. *Fabronia balansaeana* Besch. (Bescherelle, 1885b) *nom. nud.*

Distr. Southern South America (Herzog, 1952a).

F. balansaeana Besch.

Fabronia balansae

F. basilaris C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Fleischer, 1920; Brotherus, 1925; Kühnemann, 1938).

F. contermina Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933a).

F. enervis Herz. (Herzog, 1960 [1961]) *hom. illeg.*

Distr. Southern South America (Herzog, 1960).

F. felipponei Broth. (Brotherus in Felippone, 1917).

Distr. Southern South America (Felippone, 1917; Brotherus, 1925; Herter, 1933a, 1939a).

F. filamentosa C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1925; Kühnemann, 1938).

F. imbricata Hamp. (Hampe, 1878).

Distr. Southern South America (Thériot, 1936; Hosseus, 1938a, d, 1939; Herter, 1939b).

F. latidens Dub. (Duby, 1870).

Distr. Southern South America (Duby, 1870; Jaeger, 1878; Paris, 1904a; Brotherus, 1925; Herzog, 1960).

F. latifolia C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Herter, 1933a; Hosseus, 1934, 1939; Kühnemann, 1938).

F. longidens Dub.

Dimerodontium pellucidum

F. lorentziae C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1925; Hosseus, 1934; Kühnemann, 1938; Herzog, 1954).

F. lorentzii C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Fleischer, 1920; Brotherus, 1925; Williams, 1930; Hosseus, 1935a, b, 1937; Kühnemann, 1938).

F. minutissima Dub. (Duby, 1880a).

Distr. Southern South America (Duby, 1880a, b; Paris, 1904b; Brotherus, 1925).

F. obtusatula C. Muell. (Mueller, 1897a).

Syn. *Helicondontium obtusatulum* (C. Muell.) Fleisch. (Fleischer, 1920).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Fleischer, 1920; Brotherus, 1925; Herter, 1933a; Hosseus, 1934, 1939; Kühnemann, 1938).

F. palmicola C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1925; Herter, 1933a, 1939b; Hosseus, 1934, 1939; Kühnemann, 1938).

F. physcomitriocarpa C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1925; Kühnemann, 1938).

***F. podocarpi* C. Muell. (Mueller, 1879a).**

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1925; Kühnemann, 1938).

***F. polycarpa* Hook. (Hooker, 1818).**

Distr. Southern South America (Mueller, 1869; Gibert, 1873; Paris, 1904b; Herter, 1928, 1933a; Herzog, 1952a).

***F. spegazzinii* C. Muell. (Mueller, 1897a).**

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1925; Hosseus, 1938c, d; Kühnemann, 1938).

***F. tucumanensis* C. Muell. (Mueller, 1897a).**

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1925; Hosseus, 1934, 1939; Kühnemann, 1938).

***F. uruguensis* C. Muell. (Mueller, 1879a).**

Distr. Southern South America (Mueller, 1879a; Brotherus, 1925; Herter, 1933a; Hosseus, 1934, 1939; Kühnemann, 1938).

Fabroniella***F. krausei* Lor. ex C. Muell.*****Juratzkaea krausei******Felipponea******F. hollermayeri* Thér. (Thériot, 1935a).**

Distr. Southern South America (Thériot, 1935a).

***F. montevidensis* (C. Muell.) Broth. (Brotherus in Felippone, 1912).**

Syn. *Cladomnion montevidense* C. Muell. (Mueller, 1897a).
Pterogoniadelphus montevidensis (C. Muell.) Fleisch. (Fleischer, 1917).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Felippone, 1912; Fleischer, 1917; Brotherus, 1925; Herter, 1933a, 1939b).

Fissidens***F. acacioides* Schrad.*****Fissidens osmundoides******F. adianthoides* Hedw. (Hedwig, 1801).**

Syn. *Hypnum adianthoides* (Hedw.) L. ex With. (Linnaeus ex Withering, 1801).
Dicranum adianthoides (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Skitophyllum adianthoides (Hedw.) Pyl. (De la Pylaie, 1814 [1815]).
Dicranodon adianthoides (Hedw.) Béhére (Béhére, 1826).
Schistophyllum adianthoides (Hedw.) Lindb. (Lindberg, 1879).
Fissidens taxifolius Hedw. var. *adianthoides* (Hedw.) Wahlenb. ex Hueb. (Hübener, 1833) nom. inval.

Distr. Southern South America (Dusén, 1906; Cardot, 1908a).

F. anacamptodes* Besch.**Fissidens hausthalii* var. *obtusatulus******F. angustelimbatus* Mitt. (Mitten, 1869).**

Distr. Southern South America (Gibert, 1873; Herter, 1928, 1933a).

***F. aspleniooides* Hedw. (Hedwig, 1801).**

Syn. *Hypnum aspleniooides* (Hedw.) Dicks. ex With. (Dickson ex Withering, 1801).
Dicranum aspleniooides (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Skitophyllum aspleniooides (Hedw.) Pyl. (De la Pylaie, 1814 [1815]).
Schistophyllum aspleniooides (Hedw.) Brid. (Bridel, 1827b) nom. inval.
Fissidens pycnotylus Broth. (Brotherus, 1924b) fide Robinson, 1975.

Distr. Juan Fernandez (Mitten, 1869, 1885; Johow, 1896; Paris, 1904b; Brotherus, 1924b, 1925; Robinson, 1975).

Southern South America (Hosseus, 1938c, d, 1939, 1940; Herzog, 1938, 1939, 1954; Bartram, 1952, 1965).

var. *australis* Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

F. atratus (C. Muell.) Broth. (Brotherus, 1901).

Syn. *Conomitrium atratum* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1901, 1924a; Kühnemann, 1938).

F. berterii (Mont.) C. Muell.

Fissidens fontanus

F. brachyloma Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

F. brevipes Besch. (Bescherelle, 1891).

Distr. Southern South America (Herzog, 1952a).

F. brotheri Dus.

Fissidens brotherianus

F. brotherianus Par. (Paris in Thériot, 1921a).

Syn. *Fissidens brotheri* Dus. (Dusén, 1906) *hom. illeg.*

Fissidens valdiviensis Herz. (Herzog, 1923).

Distr. Southern South America (Dusén, 1906; Costes, 1921; Thériot, 1921a, 1924, 1930; Herzog, 1923, 1938, 1954; Brotherus, 1924a; Bartram, 1952).

var. *longisetus* Thér. (Thériot, 1921a).

Distr. Southern South America (Costes, 1921; Thériot, 1921a).

F. bryoides Hedw. (Hedwig, 1801).

Syn. *Hypnum bryoides* (Hedw.) L. ex With. (Linnaeus ex Withering, 1801).

Dicranum bryoides (Hedw.) Sw. (Swartz, 1801).

Skitophyllum bryoides (Hedw.) Pyl. (De la Pylaie, 1814 [1815]).

Dicranodon bryoides (Hedw.) Béhétré (Béhétré, 1826).

Schistophyllum bryoides (Hedw.) Lindb. (Lindberg, 1879).

Fissidens viridulus (Sw.) Wahlenb. var. *bryoides* (Hedw.) Roehl. (Röhling, 1813) *nom. illeg.*

Distr. Juan Fernandez (Montagne, 1835).

Southern South America (Montagne, 1850).

ssp. *incurvus* (Roehl.) Bertsch (Bertsch, 1949).

Syn. *F. incurvus* Stark ex Roehl. (Röhling, 1813).

Distr. Southern South America (Montagne, 1850).

var. *osmundoides* (Hedw.) Mathieu

Fissidens osmundoides

F. campylopus Mont. (Montagne, 1845a).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Mitten, 1869; Jaeger, 1876; Paris, 1904b; Dusén, 1906; Brotherus, 1924a).

F. chilensis Dus. (Dusén, 1906).

Distr. Southern South America (Paris, 1904b; Dusén, 1906; Thériot, 1921a; Brotherus, 1924a; Herzog, 1938, 1954).

F. colonialis C. Muell.

Fissidens haustallii* var. *minor

F. compactus Mitt. (Mitten in Gibert, 1873) *nom. nud.*

Distr. Southern South America (Gibert, 1873; Herter, 1928, 1933a).

F. costesii Thér. (Thériot, 1921a).

Distr. Southern South America (Costes, 1921; Thériot, 1921a; Brotherus, 1924a).

F. crassicuspis Broth. (brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Robinson, 1975).

F. crispus Mont. (Montagne, 1838).

Distr. Southern South America (Montagne, 1838, 1856; Jaeger, 1876; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. curvinervis Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

F. decursivus C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

F. dietrichiae C. Muell.

Fissidens fontanus

F. dillenii (Pyl.) Hamp.

Fissidens fontanus

F. distichellus Besch. (Bescherelle, 1891).

Distr. Southern South America (Bescherelle, 1891; Paris, 1904b; Brotherus, 1924a; Herter, 1933a).

F. diversiformis Broth. (Brotherus, 1924c).

Distr. Southern South America (Brotherus, 1924c, 1925; Espinosa B., 1941; Skottsberg, 1950).

F. echinellus Herz. (Herzog, 1954).

Distr. Southern South America (Herzog, 1954).

F. felipponei Broth. (Brotherus in Felippone, 1909).

Distr. Southern South America (Felippone, 1909; Brotherus, 1924a; Herter, 1933a, 1939a, 1943).

F. fernandezianus Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

F. fernandezianus Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Bartram, 1959; Robinson, 1975).

F. fontanus (Pyl.) Steud. (Steudel, 1824).

Syn. *Skitophyllum fontanum* Pyl. (De la Pylaie, 1814 [1815]).

Octodiceras fontanum (Pyl.) Lindb. (Lindberg, 1863).

Skitophyllum dillenii Pyl. (De la Pylaie, 1814 [1815]).

Octodiceras dillenii (Pyl.) Brid. (Bridel, 1827b).

Conomitrium dillenii (Pyl.) Mont. (Montagne, 1837).

Fissidens dillenii (Pyl.) Hamp. (Hampe, 1839).

Fontinalis juliana Savi ex Cand. (Savi ex de Candolle, 1815).

Skitophyllum julianum (Cand.) Brid. (Bridel, 1827b) *nom. inval.*

Octodiceras julianum (Cand.) Brid. (Bridel, 1827b).

Conomitrium julianum (Cand.) Mont. (Montagne, 1837).

Fissidens julianus (Cand.) Schimp. (Schimper, 1838).

Schistophyllum julianum (Cand.) Lindb. (Lindberg, 1879).

Conomitrium berterii Mont. (Montagne, 1837).

Fissidens berterii (Mont.) C. Muell. (Mueller, 1848a).

Octodiceras berterii (Mont.) Jaeg. (Jaeger, 1876).

Conomitrium berteroii Mont. ex Kindb. (Montagne ex Kindberg, 1888) *err.*

Fissidens dietrichae C. Muell. (Mueller, 1872).

Fissidens parvus Mitt. (Mitten, 1869) *hom. illeg.*

Distr. Southern South America (De la Pylaie, 1814; Bridel, 1827b; Montagne, 1837, 1839c, 1850, 1856; Schwaegrichen, 1842; Mueller, 1848a; Mitten, 1869; Jaeger, 1876; Paris, 1904a; Dusén, 1906; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1954).

F. fossicola C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. geheebei C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. glossophyllus C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Hosseus, 1937; Kühnemann, 1938).

var. *brevifolius* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Hosseus, 1935a; Kühnemann, 1938).

F. hauthalii C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Felippone, 1912; Brotherus, 1924a; Herter, 1933a; Kühnemann, 1938).

var. *angustissimus* C. Muell. (Mueller, 1897a).

Syn. *Fissidens stoloniferus* C. Muell. (Mueller, 1897a) *nom. nud.*

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1924a; Herter, 1933a (as *F. stolonifer* C. Muell.).

var. *gracillimus* C. Muell. (Mueller, 1897a).

Syn. *Fissidens semilaminatus* C. Muell. (Mueller, 1897a) *nom. nud.*
Fissidens semi-lamellosus C. Muell. (Mueller, 1897a) *nom. nud.*

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b).

var. *minor* C. Muell. (Mueller, 1897a).

Syn. *Fissidens colonialis* C. Muell. (Mueller, 1897a) *nom. nud.*

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b).

var. *obtusatulus* C. Muell. (Mueller, 1897a).

Syn. *Fissidens anacamptodes* Besch. (Bescherelle in Mueller, 1897a) *nom. nud.*

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b).

F. hedwigii C. Muell.

Fissidens semicompletus

F. hornschuchii Mont. (Montagne, 1840a).

Distr. Southern South America (Montagne, 1850).

F. inclinatus C. Muell.

Fissidens platensis

F. incompletus Hedw. ex Dix.

Fissidens semicompletus

F. incurvus Stark ex Roehl.

Fissidens bryoides* ssp. *incurvus

F. indistinctus C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

var. *major* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b).

F. jaffueli Thér. (Thériot in Costes, 1921) *nom. nud.*

Distr. Southern South America (Costes, 1921).

F. jujuiensis Broth. (Brotherus, 1918).

Distr. Southern South America (Brotherus, 1918, 1924a; Kühnemann, 1938).

F. julaceus (Hedw.) Lindb.

Leucodon julaceus

- F. julianus* (Cand.) Schimp. *Fissidens fontanus*
- F. jungermanniopsis* C. Muell. (Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a; Paris, 1900a, 1904b; Kurtz, 1904; Brotherus, 1924a; Hosseus, 1937; Kühnemann, 1938).
- F. lechleri* Hamp. ex C. Muell. *Fissidens rigidulus*
- F. leptocaulis* C. Muell. (Mueller, 1897a).
Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Felippone, 1912; Brotherus, 1924a; Herter, 1933a).
- F. leptochaete* Dus. (Dusén, 1906).
Distr. Juan Fernandez (Brotherus, 1924b; Robinson, 1975).
Southern South America (Dusén, 1906; Brotherus, 1924a).
- F. leucodictyus* C. Muell. (Mueller, 1897a).
Distr. Southern South America (Mueller, 1897a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).
- F. lonchothecius* C. Muell. (Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Hosseus, 1938a, d, 1939; Kühnemann, 1938).
- F. lorentziae* (C. Muell.) Broth. (Brotherus, 1901).
Syn. *Conomitrium lorentziae* C. Muell. (Mueller, 1897a).
Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1901, 1924a; Felippone, 1928; Herter, 1933a; Hosseus, 1936, 1937, 1938c, d, 1939; Kühnemann, 1938).
- F. macro-bryoides* C. Muell. (Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).
- F. maschalanthus* Mont. (Montagne, 1845a).
Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).
Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Lorentz, 1866; Mitten, 1869; Jaeger, 1876; Paris, 1904b; Dusén, 1906; Cardot, 1908a; Campo, 1921; Thériot, 1921a; Brotherus, 1924a; Reimers, 1926; Herzog, 1938, 1939, 1954, 1960; Kühnemann, 1938; Seki, 1974).
- var. minor* Dus. ex Thér. (Dusén ex Thériot, 1918).
Distr. Juan Fernandez (Skottsberg, 1914).
Southern South America (Paris, 1906; Thériot, 1918; Campo, 1921).
- F. mollis* (C. Muell.) Broth. *Fissidens permollis*
- F. nigritellus* (C. Muell.) Broth. (Brotherus, 1901).
Syn. *Conomitrium nigritellum* C. Muell. (Mueller, 1897a).
Distr. Southern South America (Mueller, 1897a; Brotherus, 1901, 1924a; Paris, 1904a; Kühnemann, 1938).
- F. obliquifolius* C. Muell. (Mueller, 1897a).
Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1924a; Kühnemann, 1938).
- F. odontoloma* C. Muell. (Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).
- F. oeditrachelos* C. Muell. *Fissidens vitreo-limbatus*

F. ornaticostatus Robins. (Robinson, 1974).

Distr. Juan Fernandez (Robinson, 1974, 1975).

F. osmundoides Hedw. (Hedwig, 1801).

Syn. *Dicranum osmundoides* (Hedw.) Turn. (Turner, 1804a).

Skitophyllum osmundoides (Hedw.) Pyl. (De la Pylaie, 1814 [1815]).

Conomitrium osmundoides (Hedw.) C. Muell. (Mueller, 1851a).

Schistophyllum osmundoides (Hedw.) Lindb. (Lindberg, 1879).

Fissidens bryoides Hedw. var. *osmundoides* (Hedw.) Mathieu (Mathieu, 1853).

Fissidens osmundoides Hedw. var. *microcarpus* B.S.G. (Bruch and others, 1843).

Schistophyllum osmundoides (Hedw.) Lindb. var. *microcarpum* (B.S.G.) Kindb. (Kindberg, 1883).

Fissidens acacioides Schrad. (Schrader, 1803b).

Hypnum acacioides (Schrad.) L. ex Turn. (Linnaeus ex Turner, 1804b).

Skitophyllum acacioides (Schrad.) Pyl. (De la Pylaie, 1814 [1815]).

Schistophyllum acacioides (Schrad.) Lindb. (Lindberg, 1883).

Distr. Southern South America (Bridel, 1803, 1806, 1827b; Mueller, 1848a, 1851a; Jaeger, 1876; Lindberg, 1883; Paris, 1904b).

var. *microcarpus* B.S.G.

Fissidens osmundoides

F. pachylomatos C. Muell.

Fissidens vitreo-limbatus

F. pachylomulus C. Muell.

Fissidens vitreo-limbatus

F. palmatus Hedw. (Hedwig, 1801).

Distr. Southern South America (Montagne, 1850).

F. parvus Mitt.

Fissidens fontanus

F. permollis Thér. (Thériot, 1931b [1932]).

Syn. *Conomitrium molle* C. Muell. (Mueller, 1879a).

Fissidens mollis (C. Muell.) Broth. (Brotherus, 1900) *hom. illeg.*

Distr. Southern South America (Mueller, 1879a; Paris, 1904a; Brotherus, 1924a; Herter, 1933a, 1939b; Kühnemann, 1938).

F. plagiotheciooides C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. platensis Par. (Paris, 1900a).

Syn. *Fissidens inclinatus* C. Muell. (Mueller, 1897a) *hom. illeg.*

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. prionocheilos C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. pulvinatus Timm ex Hedw.

Grimmia pulvinata

F. pycnoglossus C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Brotherus, 1924a; Hosseus, 1937; Kühnemann, 1938).

F. pycnotylus Broth.

Fissidens asplenoides

F. repandus Wils. (Wilson in Mitten, 1851).

Distr. Southern South America (Gibert, 1873; Herter, 1928, 1933a; Bartram, 1965).

F. rigidulus Hook. f. et Wils. (Hooker and Wilson in Wilson, 1854).

Syn. *Fissidens lechleri* Hamp. ex C. Muell. (Hampe ex Mueller, 1864).

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Robinson, 1975).
 Southern South America (Lechler, 1857; Mueller, 1864; Mitten, 1869; Jaeger, 1876;
 Neger, 1899; Paris, 1904b; Dusén, 1906; Cardot, 1908a; Campo, 1921; Thériot,
 1921a, 1925a; Brotherus, 1924a; Reimers, 1926; Roivainen, 1934; Roivainen and
 Bartram, 1937; Kühnemann, 1938; Herzog, 1939, 1954; Espinosa B., 1941; Seki,
 1974).

F. scalaris Mitt. (Mitten, 1869).

Distr. Southern South America (Mitten, 1869; Gibert, 1873; Jaeger, 1876; Paris, 1904b;
 Campo, 1921; Thériot, 1921a; Brotherus, 1924c; Herter, 1928, 1933a; Herzog, 1938,
 1954, 1960; Kühnemann, 1938; Espinosa B., 1941; Bartram, 1952).

F. semicompletus Hedw. (Hedwig, 1801).

Syn. *Cecalyphum semi-completum* (Hedw.) P. Beauv. (Palisot de Beauvois, 1805a).
Dicranum semicompletum (Hedw.) Mohr (Mohr, 1806).
Skitophyllum semi-completum (Hedw.) Pyl. (De la Pylaie, 1814 [1815]).
Schistophyllum semicompletum (Hedw.) Lindb. (Lindberg, 1883).
Fissidens incompletus Hedw. ex Dix. (Dixon, 1933 [1934]) *err.*
Conomitrium hedwigii Mont. (Montagne, 1837) *nom. illeg.*
Fissidens hedwigii C. Muell. (Mueller, 1848a) *nom. illeg.*
Octodiceras hedwigii Lindb. (Lindberg, 1863) *nom. illeg.*
Octodiceras fissidentoides Brid. (Bridel, 1827b) *nom. illeg.*

Distr. Southern South America (Hedwig, 1801; Bridel, 1803, 1806; Schwaegrichen, 1816,
 1842; Montagne, 1837, 1850; Mueller, 1848a; Mitten, 1869; Jaeger, 1876; Paris,
 1904a; Brotherus, 1924a).

F. semi-lamellosus C. Muell.

Fissidens hauthalii* var. *gracillimus

F. semilaminatus C. Muell.

Fissidens hauthalii* var. *gracillimus

F. smaragdinus (Lor. et C. Muell.) Broth. (Brotherus, 1901).

Syn. *Conomitrium smaragdinum* Lor. et C. Muell. (Lorentz and Mueller in Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1901, 1924a; Kurtz, 1904; Paris,
 1904a; Hosseus, 1935b, 1937; Kühnemann, 1938; Bruggeman-Nannenga, 1973).

F. stolonaceus C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Brotherus, 1924a;
 Hosseus, 1935b, 1937; Kühnemann, 1938).

F. stoloniferus C. Muell.

Fissidens hauthalii* var. *angustissimus

F. subaloma Dus. (Dusén, 1906).

Distr. Southern South America (Paris, 1904b; Dusén, 1906; Costes, 1921; Thériot, 1921a,
 1924; Brotherus, 1924a, c; Espinosa B., 1941).

F. synoicus C. Muell. (Mueller, 1879a) *hom. illeg.*

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann,
 1938).

F. taxifolius Hedw. (Hedwig, 1801).

Syn. *Hypnum taxifolium* (Hedw.) L. ex With. (Linnaeus ex Withering, 1801).
Dicranum taxifolium (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Skitophyllum taxifolium (Hedw.) Pyl. (De la Pylaie, 1814 [1815]).
Schistophyllum taxifolium (Hedw.) Lindb. (Lindberg, 1879).

Distr. Southern South America (Brotherus, 1924c; Espinosa B., 1941).

var. *adianthoides* (Hedw.) Wahlenb. ex Hueb.

Fissidens adianthoides

F. tener Dus. (Dusén, 1906).

Distr. Southern South America (Dusén, 1906).

F. terebrifolius C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904a; Brotherus, 1924a;
 Kühnemann, 1938).

F. valdiviensis Herz.

Fissidens brotherianus

F. ventanae C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. viridulus (Sw.) Wahlenb. var. *bryoides* (Hedw.) Roehl.

Fissidens bryoides

F. vitreo-limbatus C. Muell. (Mueller, 1897a).

Syn. *Fissidens oeditrachelos* C. Muell. (Mueller, 1897a) *nom. nud.*

Fissidens pachylomatos C. Muell. (Mueller in Paris, 1900a) *nom. nud.*

Fissidens pachylomulus C. Muell. (Mueller, 1897a) *nom. nud.*

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Felippone, 1917; Brotherus, 1924a; Herter, 1933a; Kühnemann, 1938).

Floribundaria

F. cardotii (Broth.) Broth. (Brotherus, 1906b).

Syn. *Papillaria cardotii* Broth. (Brotherus, 1895).

Distr. Southern South America (Herzog, 1952a).

F. usneoides (Broth.) Broth. (Brotherus, 1906b).

Syn. *Papillaria usneoides* Broth. (Brotherus, 1893).

Distr. Southern South America (Hosseus, 1940).

Florschuetzia hom. illeg.

F. pilmaiquen Crosby

Muscoflorschuetzia pilmaiquen

Fontinalis

F. juliana (Savi) Savi ex Cand.

Fissidens fontanus

Forsstroemia

F. coronata (Mont.) Par. (Paris, 1895).

Syn. *Leptodon coronatus* Mont. (Montagne, 1838).

Neckera coronata (Mont.) C. Muell. (Mueller, 1850).

Leucodon coronatus (Mont.) Mitt. (Mitten, 1869).

Lasia coronata (Mont.) C. Muell. (Mueller, 1879a).

Dusenia coronata (Mont.) C. Muell. (Mueller in Fleischer, 1917) *nom. inval.*

Distr. Southern South America (Montagne, 1838, 1839c; Mueller, 1850, 1879a, 1882; Mitten, 1869; Jaeger, 1877; Felippone, 1917; Brotherus, 1918; Herter, 1933a; Kühnemann, 1938; Herzog, 1952a).

F. felipponei Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933a).

F. subcoronata (Besch.) Par. (Paris, 1895).

Syn. *Lasia subcoronata* Besch. (Bescherelle, 1877).

Distr. Southern South America (Herzog, 1952a).

Funaria

F. acutifolia (Hamp.) Broth. (Brotherus, 1903).

Syn. *Entosthodon acutifolius* Hamp. (Hampe in Mueller, 1859).

Distr. Southern South America (Thériot, 1927a).

F. aristatula C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

F. balansae (Besch.) Broth. (Brotherus, 1900).

Syn. *Entosthodon balansae* Besch. (Bescherelle, 1877).

Distr. Southern South America (Herter, 1933a, 1939a; Herzog, 1939).

F. berteroana Hamp. ex C. Muell. (Hampe ex Mueller, 1862).

Syn. *Funaria fontanesii* B.S.G. non Schwaegr. (Bruch and others, 1841) *pro parte fide Paris, 1904b.*

Funaria berteroii Hamp. ex Kindb. (Hampe ex Kindberg, 1888) *nom. illeg.*

Distr. Southern South America (Mueller, 1844b, 1851a, 1862; Jaeger, 1874; Paris, 1904b; Costes, 1921; Herzog, 1923; Brotherus, 1924a; Thériot, 1927a).

var. *inclinata* Thér. (Thériot, 1927a).

Distr. Southern South America (Thériot, 1927a).

F. berteroii Hamp. ex Kindb.

Funaria berteroana

F. bonplandii (Hook.) Broth. (Brotherus, 1903).

Syn. *Gymnostomum bonplandii* Hook. (Hooker, 1816).

Physcomitrium bonplandii (Hook.) De Not. (De Notaris, 1838b).

Entosthodon bonplandii (Hook.) Mitt. (Mitten, 1869).

Amphoritheca bonplandii (Hook.) Jaeg. (Jaeger, 1874).

Distr. Southern South America (Mueller, 1844b, 1848a; Montagne, 1850; Jaeger, 1874; Brotherus, 1903, 1924a; Paris, 1904b; Thériot, 1927a).

F. calvescens Schwaegr.

Funaria hygrometrica var. **calvescens**

var. *crispa* C. Muell.

Funaria hygrometrica var. **crispa**

F. cavifolia Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923).

F. chilensis (Thér.) Thér. (Thériot, 1926).

Syn. *Funaria clavellata* (Mitt.) Broth. var. *chilensis* Thér. (Thériot, 1924).

Distr. Southern South America (Thériot, 1924, 1926, 1927a).

F. chiloënsis (Mitt.) Broth. (Brotherus, 1903).

Syn. *Entosthodon chiloënsis* Mitt. (Mitten, 1869).

Distr. Southern South America (Mitten, 1869; Jaeger, 1874; Brotherus, 1903, 1924a; Krieger, 1904; Paris, 1904b; Thériot, 1927a).

F. clavellata (Mitt.) Broth. (Brotherus, 1903).

Syn. *Entosthodon clavellatus* Mitt. (Mitten, 1869).

Amphoritheca clavellata (Mitt.) Jaeg. (Jaeger, 1874).

Distr. Southern South America (Mitten, 1869; Gibert, 1873; Jaeger, 1874; Brotherus, 1903, 1924a; Paris, 1904b; Felippone, 1909; Herter, 1928, 1933a).

var. *chilensis* Thér.

Funaria chilensis

F. commixta Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926, 1927a; Herzog, 1954).

F. costesii Thér. (Thériot, 1921a).

Distr. Southern South America (Thériot, 1921a, 1927a; Costes, 1921; Brotherus, 1924a).

F. erectiuscula Mitt. (Mitten, 1869).

Distr. Southern South America (Mitten, 1869; Jaeger, 1874; Paris, 1904b; Brotherus, 1924a; Thériot, 1927a).

F. flexisetosa (C. Muell.) Broth. (Brotherus, 1903).

Syn. *Entosthodon flexisetus* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1903, 1924a; Paris, 1904b; Kühnemann, 1938).

F. fontanesii B.S.G. *non* Schwaegr. see *Funaria berteroana* and *F. integra*
F. fuegiana C. Muell. *Funaria hygrometrica* var. *fuegiana*

F. hosseusii Bartr. (Bartram, 1936).

Distr. Southern South America (Bartram, 1936; Hosseus, 1938c, d, 1939).

F. hygrometrica Hedw. (Hedwig, 1801).

Distr. Juan Fernandez (Mitten, 1885; Johow, 1896; Skottsberg, 1914; Brotherus, 1924b; Espinosa B., 1941; Robinson, 1975).
 Southern South America (Schwaegrichen, 1816; Montagne, 1850; Lorentz, 1866; Ångström, 1872; Gibert, 1873; Bescherelle, 1889; Dusén, 1903a; Cardot, 1905a, 1908a; Felippone, 1912; Williams, 1915, 1930; Thériot, 1915, 1917b, 1918, 1921a, 1927a; Campo, 1921; Costes, 1921; Cardot and Brotherus, 1923; Brotherus, 1924c; Espinosa B., 1924, 1941; Reimers, 1926; Herter, 1928, 1933a, 1939a, b; Roivainen, 1934; Hosseus, 1938c, d; Kühnemann, 1938; Herzog, 1939, 1940, 1954, 1957; Bartram, 1952; Bizot and Piovano, 1953; Piovano, 1954; Seki, 1974).
 Falkland Islands (Wilson and Hooker, 1847; Kühnemann, 1938).

var. *calvescens* (Schwaegr.) Mont. (Montagne, 1839a).

Syn. *Funaria calvescens* Schwaegr. (Schwaegrichen, 1816).

Distr. Southern South America (Montagne, 1846, 1850; Mitten, 1869; Mueller, 1879a; Krieger, 1904; Felippone, 1912; Thériot, 1927a; Herter, 1933a; Hosseus, 1938a, c, d; Kühnemann, 1938; Herzog, 1954).

var. *crispa* (C. Muell.) Par. (Paris, 1895).

Syn. *Funaria calvescens* Schwaegr. var. *crispa* C. Muell. (Mueller, 1848a).

Distr. Southern South America (Mueller, 1848a; Paris, 1895, 1904b; Thériot, 1927a).

var. *fuegiana* (C. Muell.) Besch. (Bescherelle, 1889).

Syn. *Funaria fuegiana* C. Muell. (Mueller, 1885).

Funaria magellanica C. Muell. (Mueller in Bescherelle, 1889) *nom. nud.*

Distr. Southern South America (Mueller, 1885; Bescherelle, 1889; Paris, 1904b; Cardot, 1908a; Brotherus, 1924a; Thériot, 1927a; Kühnemann, 1938).

F. incompleta C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. integra (C. Muell.) Broth. (Brotherus, 1903).

Syn. *Entosthodon mathewsi* Hook. f. var. *integer* C. Muell. (Mueller, 1848a).

Entosthodon mathewsi Hook. f. var. *integifolius* Par. (Paris, 1895) *nom. inval.*

Entosthodon brachyodus Mont. (Montagne in Mueller, 1851a) *nom. nud.*

Funaria fontanesii B.S.G. *non* Schwaegr. (Bruch and others, 1841) *pro parte fide* Paris, 1904b.

Distr. Southern South America (Mueller, 1844b, 1848a, 1851a; Montagne, 1850; Jaeger, 1874; P-aris, 1895, 1904b; Brotherus, 1903, 1924a, c; Thériot, 1927a; Skottsberg, 1950; Herzog, 1954).

var. *serrulata* Broth. (Brotherus, 1924c).

Distr. Southern South America (Brotherus, 1924c; Espinosa B., 1941).

F. jujuiensis C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. linearidens C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. lorentzii C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

F. magellanica C. Muell.

Funaria hygrometrica var. *fuegiana*

F. meeseacea C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. papillosa (C. Muell.) Wijk. et Marg. (van der Wijk and Margadant, 1960a).

Syn. *Entosthodon papillosum* C. Muell. (Mueller, 1882).

Amphoritheca papillosa (C. Muell.) Kindb. (Kindberg, 1889).

Funaria tucumanica Broth. (Brotherus, 1903) *nom. illeg.*

Distr. Southern South America (Mueller, 1882; Brotherus, 1903, 1924a; Paris, 1904b; Kühnemann, 1938).

F. plagiothecia (C. Muell.) Broth. (Brotherus, 1903).

Syn. *Entosthodon plagiothecius* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1903, 1924a; Paris, 1904b; Kühnemann, 1938).

F. porteri Thér. (Thériot, 1928 [1929]).

Distr. Southern South America (Thériot, 1928).

F. pulchricolor C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1924a; Hosseus, 1938a, d; Kühnemann, 1938).

F. pungens Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

F. rhizomatica (C. Muell.) Broth. (Brotherus, 1903).

Syn. *Entosthodon rhizomaticus* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1903, 1924a; Paris, 1904b; Kühnemann, 1938).

F. schnyderi C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

F. suberecta Mitt. (Mitten, 1869).

Distr. Southern South America (Williams, 1915; Thériot, 1927a).

F. tenella C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Brotherus, 1924a; Hosseus, 1935b, 1937, 1938a, d; Kühnemann, 1938).

F. tucumanica Broth.

Funaria papillosa

F. valdiviae C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

Gigaspermum

G. imbricatum (Mitt.) Kindb.

Lorentziella imbricata

Glossophyllum hom. illeg.

G. brevipes (C. Muell.) Hamp.

Stereophyllum brevipes

Glyphocarpa*G. laevisphaera* Tayl.**Anacolia laevisphaera***Glyphocarpus nom. illeg.**G. cygneus* (Mont.) Jaeg.**Bartramidula cygnea***G. exiguus* (Sull.) Jaeg.**Philonotis scabrifolia***G. krausei* (C. Muell.) Jaeg.**Philonotis krausei***G. laevisphaerus* (Tayl.) Jaeg.**Anacolia laevisphaera****Glyhomitrium***G. balansae* (Besch.) Broth.**Ptychomitrium balansae***G. brevifolium* (C. Muell.) Broth.**Ptychomitrium balansae***G. cummingii* (Dub.) Broth.**Ptychomitrium fernandesianum***G. emersum* (C. Muell.) Broth.**Ptychomitrium emersum***G. fernandesianum* Mitt.**Ptychomitrium fernandesianum***G. hieronymi* (Besch.) Broth.**Ptychomitrium hieronymi***G. kurtzii* Broth. (Brotherus *in Kurtz*, 1904) *nom. nud.**Distr.* Southern South America (Kurtz, 1904; Hosseus, 1937; Kühnemann, 1938).*G. ligulatum* Mitt.**Ptychomitrium ligulatum***G. sellowianum* (C. Muell.) Mitt.**Ptychomitrium sellowianum***G. uruguense* (C. Muell.) Broth.**Ptychomitrium balansae***G. vernicosum* (C. Muell.) Broth.**Ptychomitrium vernicosum****Glyphothecium*****G. gracile*** (Hamp.) Broth. (Brotherus *in Fleischer*, 1906 [1908]).Syn. *Leucodon gracilis* Hamp. (Hampe, 1844).*Cladomnion gracile* (Hamp.) Mitt. (Mitten, 1869).*Glyptothecium gracile* (Hamp.) Broth. (Brotherus, 1905b).*Neckera poeppigiana* C. Muell. (Mueller, 1850).*Cladomnion poeppigianum* C. Muell. (Mueller *in Fleischer*, 1917) *nom. nud.**Leskea distans* Mont. (Montagne, 1845a).*Distr.* Southern South America (Hampe, 1844; Montagne, 1845a, 1850, 1856; Mueller, 1850; Lorentz, 1866; Mitten, 1869; Jaeger, 1877; Neger, 1899; Paris, 1904a; Brotherus, 1905a, 1925; Fleischer, 1906, 1917; Cardot, 1908a; Thériot, 1915; Reimers, 1926; Herzog, 1938, 1954, 1960; Kühnemann, 1938; Bizot, 1967; Seki, 1974; Kühnemann and Gonçalves, 1975).**var. *nanum*** Herz. (Herzog, 1923).*Distr.* Southern South America (Herzog, 1923).*Glyptothecium nom. illeg.**G. gracile* (Hamp.) Broth.***Glyptothecium gracile*****Goniobryum***G. reticulatum* Lindb.***Philonotis vagans*****var. *evanidum*** Herz. (Herzog *in Donat*, 1936a).*Distr.* Southern South America (Donat, 1936a; Kühnemann, 1938).***G. subbasilare*** (Hook.) Lindb. (Lindberg, 1864b [1865]).Syn. *Hypnum subbasilare* Hook. (Hooker, 1818).*Rhizogonium subbasilare* (Hook.) Schimp. (Schimper *in Mueller*, 1847b).*Mnium subbasilare* (Hook.) C. Muell. (Mueller, 1848a).*Trachyloma subbasilare* (Hook.) Mitt. (Mitten, 1860).*Photinophyllum subbasilare* (Hook.) Mitt. (Mitten, 1868).*Webera hariotiana* Besch. (Bescherelle *in Mueller*, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Hooker, 1818; Bridel, 1827b; Schwaegrichen, 1829; Wilson and Hooker, 1847; Mueller, 1848a, 1851a, 1885, 1901a; Mitten, 1869; Ångström, 1872; Jaeger, 1875; Bescherelle, 1889; Dusén, 1903a; Paris, 1904b; Cardot, 1908a, 1913a; Spegazzini, 1922; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

Grimmia

G. abscondita Card. (Cardot, 1908a).

Distr. Southern South America (Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

G. alpicola sw. ex Hedw.

Schistidium alpicola

var. *maritima* (Turn.) Wahlenb.

Schistidium maritimum

var. *rivularis* (Brid.) Wahlenb.

Schistidium apocarpum

G. amblyophylla C. Muell. (Mueller, 1849).

Syn. **Schistidium amblyophyllum** (C. Muell.) Herz. (Herzog, 1960 [1961]).

Distr. Southern South America (Wilson and Hooker, 1847 (as *G. apocarpa* var. *pro parte fide* Mueller, 1849); Mueller, 1849, 1885; Mitten, 1869; Jaeger, 1874; Paris, 1904b; Dusén, 1907; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Bartram, 1952; Herzog, 1960).

G. andreaeacea C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Kühnemann, 1938).

G. antarctici Card.

Schistidium antarctici

var. *percompacta* Bartr.

Schistidium antarctici

var. *pilifera* Bartr.

Schistidium antarctici

G. apocarpa Hedw.

Schistidium apocarpum

ssp. *alpicola* (Hedw.) C. Hartm.

Schistidium alpicola

ssp. *conferta* (Funck) Boul.

Schistidium apocarpum ssp. *confertum*

var. *conferta* (Funck) Spreng.

Schistidium apocarpum ssp. *confertum*

var. *rivularis* (Brid.) Roehl.

Schistidium apocarpum

G. aquatica (Schrad.) C. Muell.

Racomitrium aquaticum

G. arachnoidea Dus.

Grimmia orbicularis var. *patagonica*

G. atro-virens Sm.

Desmatodon convolutus

G. austro-georgica C. Muell. ex Par.

Racomitrium austro-georgicum

G. austro-leucophaea Besch.

Willia austro-leucophaea

G. austro-patens C. Muell.

Racomitrium austro-georgicum

G. calotricha C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

G. celata Card. (Cardot, 1906a).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).
South Georgia (Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b).

G. chilensis Thér. (Thériot in Herzog, 1957).

Distr. Southern South America (Herzog, 1957; Seki, 1974).

G. chubutensis Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

G. conferta Funck

Schistidium apocarpum ssp. *confertum*

G. consobrina Kunz. ex C. Muell. (Kunze ex Mueller, 1849).

Distr. Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1874; Paris, 1904b; Dusén, 1907; Thériot, 1917b, 1921a, 1935a; Campo, 1921; Costes, 1921; Cardot and Brotherus, 1923; Brotherus, 1924a; Reimers, 1926; Röiväinen, 1934; Herzog, 1938, 1939, 1954, 1957; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

G. controversa (Hedw.) Web. et Mohr***Weisia controversa******G. convoluta*** (Mont.) C. Muell.***Racomitrium crispulum******G. crassiretis*** Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

G. crispula (Hook. f. et Wils.) C. Muell.***Racomitrium crispulum******G. cryptodon*** (Mont.) Hamp.***Camptodontium cryptodon******G. decurrentipilis*** Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933a).

G. depressa C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Bescherelle, 1889; Paris, 1904b; Cardot, 1908a; Kühnemann, 1938).

var. *terrestris* C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Paris, 1904b; Cardot, 1908a).

G. dicksonii (Dus.) Dus.***Grimmia flexicaulis* var. *dicksonii******G. didyma*** Mont.***Racomitrium didymum******G. donatii*** Herz. et Thér. (Herzog and Thériot in Herzog, 1937).

Distr. Southern South America (Herzog, 1937).

G. doniana Sm. ex Spruc.***Grimmia donniana******G. donii*** Sm. ex Lindb.***Grimmia donniana******G. donniana*** Sm. (Smith, 1804a).

Syn. *Weisia donniana* (Sm.) Poir. (Poiret in Bridel, 1826) nom. inval.

Dryptodon donnianus (Sm.) Hartm. (Hartman, 1838).

Grimmia obtusa Schwaegr. var. *donniana* (Sw.) Hartm. (Hartman, 1849) nom. illeg.

Grimmia donnii Gray (Gray, 1821) nom. illeg.

Grimmia doniana Sm. ex Spruc. (Smith ex Spruce, 1847) nom. illeg.

Grimmia donii Sm. ex Lindb. (Smith ex Lindberg, 1879) nom. illeg.

Distr. Antarctic, peninsula region (Cardot, 1901, 1908a, 1911b, d, 1913a; Paris, 1904b;

Brotherus, 1924a; Bartram, 1957, Steere, 1961a; Robinson, 1972).

Antarctic, continental region (Kuc, 1969; Savich-Ljubitskaya and Smirnova, 1969; Savich-Ljubitskaya, 1978).

G. donii Gray***Grimmia donniana******G. fallax*** Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a, 1907; Cardot, 1908a; Brotherus, 1924a; Thériot, 1924; Kühnemann, 1938).

G. fasciculata Dus. (Dusén, 1905b) hom. illeg.

Distr. Southern South America (Paris, 1904b; Dusén, 1905b, 1907; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

G. fastigiata Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

Antarctic, continental region (Dixon and Watts, 1918; Steere, 1961a).

G. flexicaulis C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

var. dicksonii Dus. (Dusén, 1905b).

Syn. *Grimmia dicksonii* (Dus.) Dus. (Dusén, 1907) hom. illeg.
Grimmia pycnophylla Dus. (Dusén, 1903a) nom. nud.
Grimmia pachyphylla Dus. (Dusén, 1905b) hom. illeg.

Distr. Southern South America (Dusén, 1903a, 1905b, 1907; Paris, 1904b; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).

G. flexiseta Dus.***Grimmia reflexidens*****G. gibertii** Mitt. (Mitten, 1869).

Distr. Southern South America (Mitten, 1869; Gibert, 1873; Jaeger, 1874; Krieger, 1904; Paris, 1904b; Felippone, 1909; Brotherus, 1924a; Herter, 1928, 1933a, 1939a, b, 1943).

G. glacialis C. Muell.***Racomitrium lanuginosum*****G. grisea** Card. (Cardot, 1906a).

Distr. Southern South America (Cardot and Brotherus, 1923; Thériot, 1934b; Kühnemann, 1938).
 South Georgia (Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Dixon, 1932).

G. gusindei Froehl. (Froehlich, 1953).

Distr. Southern South America (Froehlich, 1953).

G. heteromalla (Hedw.) Web. et Mohr***Ditrichum heteromallum*****G. heterosticha** (Hedw.) C. Muell.***Racomitrium heterostichum*****G. homomalla** (Hedw.) Sm.***Ditrichum heteromallum*****G. humilis** Mitt. (Mitten, 1869).

Syn. *Grimmia macropulvinata* Dus. (Dusén, 1903a) fide Cardot, 1908a.

Distr. Southern South America (Mitten, 1869; Jaeger, 1874; Mueller, 1885; Dusén, 1903a, 1907; Paris, 1904b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Espinosa B., 1924; Kühnemann, 1938; Bartram, 1952; Herzog, 1957; Seki, 1974).

G. hyalino-cuspidata C. Muell. (Mueller, 1890a).

Syn. *Racomitrium hyalino-cuspidatum* (C. Muell.) Kindb. (Kindberg, 1891).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Paris, 1905b; Cardot, 1908a; Brotherus, 1924a; Steere, 1961b).

var. *mutica* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. South Georgia (Cardot and Brotherus, 1923; Steere, 1961b).

G. imberbis Kunz. ex C. Muell. (Kunze ex Mueller, 1849).

Distr. Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1874; Paris, 1904b; Dusén, 1907; Brotherus, 1924a; Herzog, 1938, 1957, 1960).

G. immerso-leucophaea (C. Muell.) Kindb. (Kindberg, 1891).

Syn. *Guembelia immerso-leucophaea* C. Muell. (Mueller, 1890a).

Distr. South Georgia (Mueller, 1890a; Paris, 1904b; Cardot, 1908a; Brotherus, 1924a; Steere, 1961b).

G. integrifolia C. Muell.***Grimmia subovata*****G. julacea** (Hedw.) Web. et Mohr***Leucodon julaceus*****G. julacea** Hornsch.***Acrocryphaea julacea*****G. laevigata** Mitt.***Racomitrium laevigatum***

- G. lamprocarpa* C. Muell. *Racomitrium lamprocarpum*
G. lanuginosa (Hedw.) C. Muell. *Racomitrium lanuginosum*
- G. lawiana*** J. H. Willis (*Willis in Filson*, 1966).
Distr. Antarctic, continental region (*Filson*, 1966; *Greene*, 1968a).
- G. leucophaeola*** C. Muell. (*Mueller*, 1882).
Distr. Southern South America (*Mueller*, 1882; *Paris*, 1904b; *Brotherus*, 1924a; *Kühnemann*, 1938).
- G. lorentzii* C. Muell. *ex Kindb.* ***Grimmia lorentziana***
- G. lorentziana*** (C. Muell.) Par. (*Paris*, 1895).
Syn. *Guembelia lorentziana* C. Muell. (*Mueller*, 1882).
Grimmia lorentzii C. Muell. *ex Kindb.* (*Mueller ex Kindberg*, 1889) *nom. illeg.*
Distr. Southern South America (*Mueller*, 1882; *Paris*, 1895, 1904b; *Brotherus*, 1924a; *Hosseus*, 1936; *Kühnemann*, 1938).
- G. macropulvinata* Dus. ***Grimmia humilis***
G. macrothylla Card. et Broth. ***Grimmia macrotyla***
- G. macrotyla*** Card. et Broth. (*Cardot and Brotherus*, 1923).
Syn. *Grimmia macrothylla* Card. et Broth. *in Broth.* (*Cardot and Brotherus in Brotherus*, 1924a) *err.*
Distr. Southern South America (*Cardot and Brotherus*, 1923; *Brotherus*, 1924a; *Kühnemann*, 1938).
- G. maritima* Turn. ***Schistidium maritimum***
- G. montevidensis*** Thér. (*Thériot in Felippone*, 1929 [1930]).
Distr. Southern South America (*Felippone*, 1929; *Herter*, 1933a).
- G. murina* C. Muell. ***Grimmia reflexidens***
G. nigrita C. Muell. ***Racomitrium crispulum***
- G. nivea*** Dus. (*Dusén*, 1905b).
Distr. Southern South America (*Dusén*, 1905b, 1907; *Cardot*, 1908a; *Brotherus*, 1924a; *Kühnemann*, 1938; *Seki*, 1974).
- G. nordenskioldii*** Card. (*Cardot*, 1906a).
Distr. Southern South America (*Cardot and Brotherus*, 1923).
 South Georgia (*Cardot*, 1906a, 1908a; *Cardot and Brotherus*, 1923; *Brotherus*, 1924a; *Steere*, 1961b).
- G. obtusa* Schwaegr. var. *donniana* (Sm.) Hartm. ***Grimmia donniana***
- G. occulta*** C. Muell. (*Mueller*, 1890a).
Distr. Southern South America (*Dusén*, 1907; *Cardot*, 1908a; *Cardot and Brotherus*, 1923; *Kühnemann*, 1938).
 South Georgia (*Mueller*, 1890a; *Paris*, 1904b; *Cardot*, 1906a, 1908a; *Cardot and Brotherus*, 1923; *Brotherus*, 1924a; *Dixon*, 1935; *Steere*, 1961b).
- G. orbicularis*** Bruch (*Bruch in Wilson*, 1844).
Syn. *Guembelia orbicularis* (Bruch) Hamp. (*Hampe in Mueller*, 1849).
Distr. Southern South America (*Cardot and Brotherus*, 1923; *Brotherus*, 1924a; *Kühnemann*, 1938 (as *G. orvicularis*); *Seki*, 1974).
- var. patagonica*** Card. (*Cardot*, 1908a).
Syn. *Grimmia arachnoidea* Dus. (*Dusén*, 1903a) *fide Cardot*, 1908a, p. 284.
Distr. Southern South America (*Dusén*, 1903a, 1905b (as *G. orbicularis* *fide Cardot*, 1908a), 1907 (as *G. orbicularis* *fide Cardot*, 1908a); *Cardot*, 1908a (and as *G. orbicularis*)).

<i>G. pachyloma</i> (Mont.) Mitt.	<i>Sciaromium pachyloma</i>
<i>G. pachyphylla</i> C. Muell.	<i>Andreaea pachyphylla</i>
<i>G. pachyphylla</i> Dus.	<i>Grimmia flexicaulis</i> var. <i>dicksonii</i>
<i>G. patagonica</i> Mitt.	<i>Scouleria patagonica</i>
<i>G. perplexa</i> Thér. (Thériot, 1923a).	
Distr. Southern South America (Thériot, 1923a, 1934b; Brotherus, 1925).	
<i>G. phyllorhizans</i> Broth. (Brotherus, 1924b).	
Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Robinson, 1975).	
<i>G. plagiopodia</i> Hedw. (Hedwig, 1801).	
Syn. <i>Weisia plagiopodi</i> (Hedw.) Poir. (Poiret, 1808).	
<i>Schistidium plagiopodium</i> (Hedw.) Loesk. (Loeske, 1913).	
Distr. Antarctic, peninsula region (Bartram, 1957; Steere, 1961a).	
Antarctic, continental region (Bartram, 1957; Steere, 1961a).	
var. <i>antarctica</i> Bartr. (Bartram, 1957).	
Distr. Antarctic, continental region (Bartram, 1957; Steere, 1961a; Savich-Ljubitskaya and Smirnova, 1970; Savich-Ljubitskaya, 1978).	
<i>G. praemorsa</i> (C. Muell.) Kindb. (Kindberg, 1889).	
Syn. <i>Guembelia praemorsa</i> C. Muell. (Mueller, 1882).	
<i>Schistidium praemorsum</i> (C. Muell.) Herz. (Herzog, 1916a).	
Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Hosseus, 1938c, d; Kühnemann, 1938).	
<i>G. protensum</i> (Braun) Mitt.	<i>Racomitrium aquaticum</i>
<i>G. ptychophylla</i> Mitt.	<i>Racomitrium ptychophyllum</i>
<i>G. pulvinata</i> (Hedw.) Sm. (Smith, 1807).	
Syn. <i>Fissidens pulvinatus</i> Timm ex Hedw. (Hedwig, 1801).	
Distr. Southern South America (Montagne, 1850).	
<i>G. pycnophylla</i> Dus.	<i>Grimmia flexicaulis</i> var. <i>dicksonii</i>
<i>G. quadricruris</i> C. Muell. (Mueller, 1882).	
Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938 (as <i>quaticruris</i> C. Muell.); Bartram, 1965).	
<i>G. raphidostega</i> C. Muell. ex Kindb.	<i>Grimmia raphidostega</i>
<i>G. reflexidens</i> C. Muell. (Mueller, 1849).	
Syn. <i>Grimmia murina</i> C. Muell. (Mueller, 1897a).	
<i>Grimmia flexiseta</i> Dus. (Dusén, 1903a) nom. nud.	
Distr. Southern South America (Mueller, 1849, 1897a; Montagne, 1850; Mitten, 1869; Jaeger, 1874; Paris, 1900a, 1904b; Dusén, 1903a, 1905b, 1907; Brotherus, 1924a; Kühnemann, 1938).	
<i>G. raphidostega</i> C. Muell. (Mueller, 1882).	
Syn. <i>Grimmia raphidostega</i> C. Muell. ex Kindb. (Mueller ex Kindberg, 1889) err.	
Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).	
<i>G. richardii</i> (Brid.) Spreng.	<i>Campylopus richardii</i>
<i>G. rivularis</i> Brid.	<i>Schistidium apocarpum</i>
<i>G. rupestris</i> (Hook. f. et Wils.) C. Muell.	<i>Racomitrium crispulum</i> var. <i>rupestre</i>

- G. scabripes** Bartr. (Bartram, 1946).
Distr. Southern South America (Rovainen, 1934; Bartram, 1946).
- G. senilis** Besch. **Willia austro-leucophaea**
- G. serripila** Dus. (Dusén *in Paris*, 1904b) *nom. nud.*
Distr. Southern South America (Paris, 1904b; Thériot, 1923b).
- G. stolonifera** C. Muell. (Mueller, 1884 [1883]).
Distr. Antarctic, continental region (Dixon and Watts, 1918; Steere, 1961a).
- G. striata** Schrad. **Rhabdoweisia fugax**
- G. subcrispipila** C. Muell. **Racomitrium subcrispipilum**
- G. sublamprocarpa** C. Muell. **Racomitrium sublamprocarpum**
- G. subnigrita** C. Muell. **Racomitrium subnigritum**
- G. subovata** Schimp. (*Schimper in Mueller*, 1882).
Syn. *Grimmia integrifolia* C. Muell. (Mueller, 1882).
Distr. Southern South America (Mueller, 1882; Kurtz, 1904; Paris, 1904b; Brotherus, 1924a; Hosseus, 1936, 1937; Kühnemann, 1938).
- G. symphyodon** Mitt. **Racomitrium symphyodon**
- G. symphyodontata** C. Muell. **Racomitrium symphyodontatum**
- G. syntrichiacea** C. Muell. (Mueller, 1890a).
Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).
 South Georgia (Mueller, 1890a; Paris, 1904b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b).
- G. tenuicaulis** Hamp. (*Hampe in Jaeger*, 1874) *nom. nud.*
Distr. Southern South America (Jaeger, 1874).
- G. tortuosa** Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Distr. Falkland Islands (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1849; Jaeger, 1874; Paris, 1904b; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).
- G. trichophylla** Grev. (Greville, 1824).
Syn. *Dryptodon trichophyllum* (Grev.) Brid. (Bridel, 1827a).
Eugrimmia trichophylla (Grev.) Hamp. (*Hampe, 1880c*) *comb. inval.*
Distr. Southern South America (Montagne, 1850).
- G. unicurvis** C. Muell. (Mueller, 1882).
Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1957).
- G. urnulacea** C. Muell. (Mueller, 1890a).
Distr. South Georgia (Mueller, 1890a; Paris, 1904b; Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b).
- G. vernicosula** C. Muell. (Mueller, 1882).
Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).
- G. willii** C. Muell. **Racomitrium willii**
- Groutiella**
- G. apiculata** (Hook.) Crum et Steere (Crum and Steere, 1950).
Syn. *Orthotrichum apiculatum* Hook. (Hooker, 1818).
Macromitrium apiculatum (Hook.) Schwaeggr. (Schwaegrichen, 1826).

Distr. Southern South America (Krieger, 1904).

G. tomentosa (Hornschr.) Wijk et Marg. (van der Wijk and Margadant, 1960a).

Syn. *Macromitrium tomentosum* Hornsch. (Hornschuch, 1840).

Micromitrium tomentosum (Hornsch.) Par. (Paris, 1905a).

Distr. Southern South America (Hornschuch, 1840; Mueller, 1849; Gibert, 1873; Jaeger, 1874; Paris, 1905a; Brotherus, 1925; Herter, 1933a).

Guembelia

G. austro-leucophaea C. Muell.

Willia austro-leucophaea

G. immerso-leucophaea C. Muell.

Grimmia immerso-leucophaea

G. lorentziana C. Muell.

Grimmia lorentziana

G. orbicularis (Bruch) Hamp.

Grimmia orbicularis

G. praemorsa C. Muell.

Grimmia praemorsa

Gymnocybe nom. rejic.

G. palustris (Hedw.) Fries

Aulacomnium palustre

Gymnopilum nom. nud.

G. dendroides (Hedw.) Schimp.

Dendroligotrichum dendroides

G. magellanicum (Hedw.) Schimp.

Polytrichadelphus magellanicus

Gymnostomum

G. aeruginosum Sm. (Smith, 1804b).

Syn. *Gymnostomum calcareum* Nees et Hornsch. (Nees and Hornschuch in Nees von Esenbeck and others, 1823) *fide* Zander, 1977.

Weisia calcarea (Nees et Hornsch.) C. Muell. (Mueller, 1849) *hom. illeg. fide* Zander, 1977.

Hymenostylium calcareum (Nees et Hornsch.) Mitt. (Mitten, 1859) *fide* Zander, 1977.
Trichostomum calcareum (Nees et Hornsch.) Lindb. (Lindberg, 1864a) *fide* Zander, 1977.

Eucladium calcareum (Nees et Hornsch.) C. Jens. (Jensen, 1923).

Gymnostomum calcareum Nees et Hornsch. var. *tenellum* B.S.G. (Bruch and others, 1846).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Robinson, 1975).

Southern South America (Montagne, 1850; Mitten, 1869; Paris, 1904b; Dusén, 1906; Thériot, 1921a; Hosseus, 1938c, d; Herzog, 1938, 1940, 1954; Bizot and Piovano, 1953; Zander, 1977).

G. aestivum Hedw. (Hedwig, 1801).

Syn. *Gymnostomum euchloron* Schwaegr. (Schwaegrichen, 1827a) *fide* Zander, 1977.
Anoectangium euchloron (Schwaegr.) Mitt. (Mitten, 1869) *fide* Zander, 1977.

Distr. Southern South America (Thériot, 1936; Hosseus, 1938c, d, 1939; Bartram, 1965; Zander, 1977).

G. amblyophyllum Hook.

Didymodon amblyophyllum

G. antarcticum Aongstr.

Pottia antarctica

G. bonplandii Hook.

Funaria bonplandii

G. calcareum Nees et Hornsch.

Gymnostomum aeruginosum

var. *tenellum* B.S.G.

Gymnostomum aeruginosum

G. curvirostre Hedw. ex Brid.

Hymenostylium recurvirostre

G. euchloron Schwaegr.

Gymnostomum aestivum

G. griffithianum (Dicks.) Sm.

Oedipodium griffithianum

G. heimii Hedw.

Pottia heimii

G. imberbe Sm.

Hedwigidium integrifolium

<i>G. kunzeanum</i> (C. Muell.) Par.	<i>Hymenostylium kunzeanum</i>
<i>G. laxifolium</i> Kunz. ex Par.	<i>Gymnostomum tenerimum</i>
<i>G. longirostre</i> Kunz. ex C. Muell.	<i>Hymenostylium kunzeanum</i>
<i>G. magellanicum</i> Aongstr. ex Par.	<i>Pottia antarctica</i>
<i>G. marginatum</i> Kunz. ex Mont.	<i>Hymenostomum kunzeanum</i>
<i>G. menziesii</i> (R. Brown) Hook.	<i>Leptostomum menziesii</i>
<i>G. orbignyanum</i> Mont.	<i>Physcomitrium orbignyanum</i>
<i>G. ovatum</i> Hedw.	<i>Pterygoneurum ovatum</i>
<i>G. pachyloma</i> Mont.	<i>Sciaromium pachyloma</i>
<i>G. pyriforme</i> Hedw.	<i>Physcomitrium pyriforme</i>
<i>G. recurvirostre</i> Hedw.	<i>Hymenostylium recurvirostre</i>
<i>G. subsessile</i> Brid.	<i>Pterygoneurum subsessile</i>

G. tenerimum (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1959).

Syn. *Anoectangium tenerimum* C. Muell. (Mueller, 1847c).
Zygodon tenerimus (C. Muell.) C. Muell. (Mueller, 1849).
Anoectangium laxifolium Par. (Paris, 1894) *hom. illeg.*
Gymnostomum laxifolium Kunz. ex Par. (Kunze ex Paris, 1904b) *nom. illeg.*
Distr. Southern South America (Mueller, 1847c, 1849; Lorentz, 1866; Mitten, 1869; Jaeger, 1870; Paris, 1894, 1904b; Brotherus, 1924a).

<i>G. tortula</i> Schwaegr.	<i>Hyophila tortula</i>
<i>G. truncatum</i> Hedw.	<i>Pottia truncata</i>

Gyroweisia

G. detersifolia (Hamp.) Broth. (*In* Piovano, 1954) *nom. nud.*
Distr. Southern South America (Piovano, 1954).

Haplocladium

<i>H. amblyostomum</i> (C. Muell.) Broth.	<i>Haplocladium microphyllum</i>
<i>H. austro-serpens</i> (C. Muell.) Broth.	<i>Haplocladium microphyllum</i>
<i>H. laplatae</i> (C. Muell.) Broth.	<i>Haplocladium microphyllum</i>
<i>H. laterculi</i> (C. Muell.) Broth.	<i>Haplocladium microphyllum</i>
<i>H. malachiticum</i> C. Muell. (Mueller <i>in</i> Reimers, 1937) <i>nom. nud.</i>	

Distr. Southern South America (Reimers, 1937).

H. microphyllum (Hedw.) Broth. (Brotherus, 1907).

Syn. *Hypnum microphyllum* Hedw. (Hedwig, 1801).
Leskea microphylla (Hedw.) Sw. (Swartz *in* Mitten, 1859).
Thuidium microphyllum (Hedw.) Jaeg. (Jaeger, 1878).
Pseudoleskea microphylla (Hedw.) Sauerb. (Sauerbeck *in* Jaeger and Sauerbeck, 1879b).
Hypnum austro-serpens C. Muell. (Mueller, 1879a).
Amblystegium austro-serpens (C. Muell.) Kindb. (Kindberg, 1891).
Thuidium austro-serpens (C. Muell.) Kindb. (Kindberg, 1891).
Haplocladium austro-serpens (C. Muell.) Broth. (Brotherus, 1907).
Hypnum laterculi C. Muell. (Mueller, 1879a).
Thuidium laterculi (C. Muell.) Kindb. (Kindberg, 1891).
Thuidium laterale Par. (Paris, 1898) *err.*
Haplocladium laterculi (C. Muell.) Broth. (Brotherus, 1907).
Hypnum attenuatifolium C. Muell. (Mueller, 1882) *nom. nud.*
Hypnum paludicola C. Muell. (Mueller, 1882) *nom. nud.*
Hypnum amblyostomum C. Muell. (Mueller, 1882).
Thuidium amblyostomum (C. Muell.) Kindb. (Kindberg, 1891).
Haplocladium amblyostomum (C. Muell.) Broth. (Brotherus, 1907).

- Hypnum amblyostomum* C. Muell. var. *attenuatifolium* C. Muell. (Mueller, 1882).
Thuidium amblyostomum (C. Muell.) Kindb. var. *attenuatifolium* (C. Muell.) Par. (Paris, 1898).
Hypnum amblyostomum C. Muell. var. *paludicola* C. Muell. (Mueller, 1882).
Thuidium amblyostomum (C. Muell.) Kindb. var. *paludicola* (C. Muell.) Par. (Paris, 1898).
Haplocladium pseudo-gracile C. Muell. (Mueller, 1897a).
Pseudoleskea pseudogracilis (C. Muell.) Par. (Paris, 1896).
Thuidium pseudo-gracile (C. Muell.) Par. (Paris, 1906) *hom. illeg.*
Haplodontium pseudo-gracile C. Muell. ex Par. (Mueller ex Paris, 1906) *err.*
Pseudoleskea uruguensis C. Muell. (Mueller, 1897a).
Haplocladium uruguense (C. Muell.) Broth. (Brotherus, 1907).
Thuidium semi-lunare C. Muell. (Mueller, 1897a).
Haplocladium semilunare (C. Muell.) Broth. (Brotherus, 1907).
Hypnum semilunare C. Muell. (Mueller in Reimers, 1937) *nom. nud.*
Thuidium torskii Kiaer ex C. Muell. (Kiaer ex Mueller, 1897a).
Haplocladium torskii (C. Muell.) Broth. (Brotherus, 1907).
Pseudoleskea laplatae C. Muell. (Mueller, 1897a).
Haplocladium laplatae (C. Muell.) Broth. (Brotherus, 1907).

Distr. Southern South America (Montagne, 1839c; Mueller, 1879a, 1882, 1897a; Paris, 1896, 1898, 1905b, 1906; Brotherus, 1907, 1918, 1925; Herter, 1933a, 1943; Kühnemann, 1938; Hosseus, 1939, 1940; Herzog, 1952a).

H. persistens (C. Muell.) Broth. (Brotherus, 1907).

- Syn.* *Hypnum persistens* C. Muell. (Mueller, 1882).
Thuidium persistens (C. Muell.) Kindb. (Kindberg, 1891).

Distr. Southern South America (Mueller, 1882; Paris, 1906; Brotherus, 1907, 1925; Kühnemann, 1938).

H. pseudo-gracile C. Muell.

Haplocladium microphyllum

H. semilunare (C. Muell.) Broth.

Haplocladium microphyllum

H. torskii (C. Muell.) Broth.

Haplocladium microphyllum

H. uruguense (C. Muell.) Broth.

Haplocladium microphyllum

Haplodontium

H. brachycladum Broth. (Brotherus in Felippone, 1909).

- Syn.* *Haplodontium brachycladulum* Broth. (Brotherus, 1924a) *err.*

Distr. Southern South America (Felippone, 1909; Brotherus, 1924a; Herter, 1933a).

H. brachycladulum Broth.

Haplodontium brachycladum

H. humipetens C. Muell. (Mueller, 1879a).

- Syn.* *Mielichhoferia humipetens* (C. Muell.) Par. (Paris, 1896).

Distr. Southern South America (Mueller, 1879a; Paris, 1896, 1904b; Brotherus, 1924a; Kühnemann, 1938).

H. pernanum C. Muell.

Haplodontium sanguinolentum

var. robustior C. Muell. (Mueller, 1879a).

- Syn.* *Mielichhoferia pernana* (C. Muell.) Par. var. *robustior* (C. Muell.) Par. (Paris, 1896).

Distr. Southern South America (Mueller, 1879a; Paris, 1896, 1904b; Kühnemann, 1938).

H. pseudo-gracile C. Muell. ex Par.

Haplocladium microphyllum

H. sanguinolentum C. Muell. (Mueller, 1879a).

- Syn.* *Mielichhoferia sanguinolenta* (C. Muell.) Par. (Paris, 1896).

Haplodontium pernanum C. Muell. (Mueller, 1879a).

Mielichhoferia pernana (C. Muell.) Par. (Paris, 1896).

Distr. Southern South America (Mueller, 1879a; Paris, 1896, 1904b; Brotherus, 1924a; Hosseus, 1936, 1937; Kühnemann, 1938; Bartram, 1965).

*Haptymenium nom. illeg.**H. curtipendulum* (Hedw.) Fuernr.*Antitrichia curtipendula**Harpidium hom. illeg.**H. aduncum* (Hedw.) Con.*Drepanocladus aduncus**H. fluitans* (Hedw.) Spruc.*Drepanocladus fluitans**H. polycarpum* (Voit) Williams*Drepanocladus aduncus* var. *polycarpus**H. revolvens* (Sw.) C. Jens.*Drepanocladus revolvens**H. uncinatum* (Hedw.) C. Jens.*Drepanocladus uncinatus*var. *orthotheciooides* (Lindb.) C. Jens.*Drepanocladus uncinatus* var. *subjulaceus**Harrisonia hom. illeg.**H. australis* (Hook. f. et Wils.) Hamp.*Rhacocarpus purpurascens**H. crasso-limbata* C. Muell.*Rhacocarpus purpurascens**H. humboldtii* (Hook.) Spreng.*Rhacocarpus purpurascens*var. *argentinica* C. Muell.*Rhacocarpus purpurascens* var. *argentinicus**H. orbiculata* (Mitt.) Jaeg.*Rhacocarpus orbiculatus**H. purpurascens* (Brid.) C. Muell.*Rhacocarpus purpurascens**H. rhabdocarpa* Hamp.*Hedwigidium integrifolium**H. secunda* (Hook.) Spreng. var. *cirrhifolia* Mitt.*Braunia cirrhifolia**Hedwigia**H. albicans* Lindb.*Hedwigia ciliata**H. argentinica* C. Muell.*Hedwigia ciliata**H. ciliata* (Hedw.) P. Beauv. (Palisot de Beauvois, 1805a).Syn. *Anictangium ciliatum* Hedw. (Hedwig, 1801).*Bryum ciliatum* (Hedw.) Dicks. (Dickson, 1801).*Anoectangium ciliatum* (Hedw.) Roehl. (Röhling, 1809).*Perisiphorus ciliatus* (Hedw.) P. Beauv. (Palisot de Beauvois, 1822).*Hedwigidium ciliatum* (Hedw.) Hartm. (Hartman, 1849).*Pilotrichum ciliatum* (Hedw.) C. Muell. (Mueller, 1851a).*Bryum apocarpum* (Hedw.) L. ex With. var. *ciliatum* (Hedw.) With. (Withering, 1801).*Bryum hedwigii* Brotero (Brotero, 1804).*Hedwigia albicans* Lindb. (Lindberg, 1864b) *nom. illeg.**Hedwigia argentinica* C. Muell. (Mueller in Fleischer, 1920) *nom. nud.*

Distr. Southern South America (Jaeger, 1876; Fleischer, 1920).

H. cirrhifolia Mitt.*Braunia cirrhifolia**H. humboldtii* (Hook.) Hook.*Rhacocarpus purpurascens*var. *australis* (Hook. f. et Wils.) Wils.*Rhacocarpus purpurascens**H. imberbis* (Sm.) Spruc.*Hedwigidium integrifolium**H. integrifolia* P. Beauv.*Hedwigidium integrifolium**H. orbiculata* Mitt.*Rhacocarpus orbiculatus**H. patagonica* C. Muell. (Mueller in Kindberg, 1891) *nom. nud.*Distr. No locality given, see *Index muscorum*, 5, p. 674.*Hedwigidium**H. chilense* Fleisch. (Fleischer, 1920) *nom. nud.*Syn. *Braunia chilensis* Broth. (*Brotherus* in Fleischer, 1920) *nom. nud.*

Distr. Southern South America (Fleischer, 1920).

H. ciliatum (Hedw.) Hartm.*Hedwigia ciliata*

H. gibertii Fleisch. (Fleischer, 1920) *nom. nud.*

Syn. *Braunia gibertii* Besch. (Bescherelle in Fleischer, 1920) *nom. nud.*

Distr. Southern South America (Fleischer, 1920).

H. imberbe (Sm.) B.S.G.

Hedwigidium integrifolium

H. integrifolium (P. Beauv.) Dix. (Dixon in Jensen, 1939).

Syn. *Hedwigia integrifolia* P. Beauv. (Palisot de Beauvois, 1805a).

Anoectangium integrifolium (P. Beauv.) Schwaegr. (Schwaegrichen, 1811).

Gymnostomum imberbe Sm. (Smith, 1811).

Anictangium imberbe (Sm.) Hook. et Tayl. (Hooker and Taylor, 1818).

Schistidium imberbe (Sm.) Nees et Hornsch. (Nees von Esenbeck and others, 1823).

Hedwigidium imberbe (Sm.) B.S.G. (Bruch and others, 1846).

Hedwigia imberbis (Sm.) Spruc. (Spruce, 1847).

Neckera imberbis (Sm.) C. Muell. (Mueller, 1850).

Schistidium ciliatum (Hedw.) Brid. var. *imberbe* (Sm.) Hueb. (Hübener, 1833).

Harrisonia rhabdocarpa Hamp. (Hampe, 1863).

Braunia rhabdocarpa (Hamp.) C. Muell. (Mueller, 1879a).

Hedwigidium rhabdocarpum (Hamp.) Jaeg. (Jaeger in Jaeger and Sauerbeck, 1879b).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1905a, 1925; Reimers, 1926; Herzog, 1938; Kühnemann, 1938; Bartram, 1952).

H. patagonicum Fleisch. (Fleischer, 1920) *nom. nud.*

Syn. *Braunia patagonica* C. Muell. (Mueller in Fleischer, 1920) *nom. nud.*

Distr. Southern South America (Fleischer, 1920).

H. rhabdocarpum (Hamp.) Jaeg.

Hedwigidium integrifolium

Helicodontium

H. acuminatum C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1925; Kühnemann, 1938).

H. chloronema C. Muell. (Mueller, 1897a).

Syn. *Helicodontium montevidense* C. Muell. (Mueller in Fleischer, 1920) *nom. nud.*

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Fleischer, 1920; Brotherus, 1925; Hosseus, 1934, 1939; Kühnemann, 1938).

H. clarazii (Dub.) Par. (Paris, 1895).

Syn. *Hypnum clarazii* Dub. (Duby, 1868).

Rudia clarazii (Dub.) Jaeg. (Jaeger, 1878) *comb. inval.*

Amblystegium clarazii (Dub.) Jaeg. (Jaeger in Jaeger and Sauerbeck, 1879a) *nom. inval.*

Distr. Southern South America (Duby, 1868; Jaeger, 1878; Paris, 1895, 1904b).

H. complanatum Broth. (Brotherus, 1895).

Distr. Southern South America (Herzog, 1952a).

H. montevidense C. Muell.

Helicodontium chloronema

H. obtusatulum (C. Muell.) Fleisch.

Fabronia obtusatula

H. pendulum (Brid.) Lindb. (Lindberg, 1883).

Syn. *Isothecium pendulum* Brid. (Bridel, 1827b).

Hypnum pendulum (Brid.) C. Muell. (Mueller, 1851a) *hom. illeg.*

Eurhynchium pendulum (Brid.) Jaeg. (Jaeger, 1878).

Rhynchostegium pendulum (Brid.) Jaeg. (Jaeger, 1878).

Distr. Southern South America (Bridel, 1827b; Mueller, 1851a; Mitten, 1869; Gibert, 1873; Jaeger, 1878; Lindberg, 1883; Paris, 1904b; Herter, 1928, 1933b).

H. pervirens (C. Muell.) Broth. (Brotherus, 1895).

Syn. *Hypnum pervirens* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1904b; Brotherus, 1925; Herter, 1933a, 1943; Hosseus, 1938; Kühnemann, 1938; Herzog, 1952a).

H. pseudo-limnobium (C. Muell.) Kindb. (Kindberg, 1891).

Syn. *Hypnum pseudo-limnobium* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1925; Kühnemann, 1938; Herzog, 1952a).

H. rhyparobolax C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1925; Hosseus, 1934, 1939; Kühnemann, 1938).

H. seminerve (Schwaegr.) Schwaegr.

Juratzaea seminervis

H. siambonense C. Muell. (Mueller, 1897a).

Syn. *Helicodontium siambonicum* C. Muell. (Mueller in Fleischer, 1920) nom. nud.

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Fleischer, 1920; Brotherus, 1925; Kühnemann, 1938).

H. siambonicum C. Muell.

Helicodontium siambonense

H. tenuifolium (Hedw.) C. Muell.

Rhynchostegium tenuifolium

H. tenuirostre Schwaegr. (Schwaegrichen, 1830).

Syn. *Hypnum tenuirostre* (Schwaegr.) C. Muell. (Mueller, 1851a) hom. illeg.

Distr. Southern South America (Jaeger, 1878; Paris, 1904b; Brotherus, 1925).

Hepaticina nom. inval.

H. lamellata Fleisch.

Pterygophyllum lamellatum

H. peruncinata Dus.

Pterygophyllum peruncinatum

Herpetineuron

H. toccae (Sull. et Lesq.) Card. (Cardot, 1905c).

Syn. *Anomodon toccae* Sull. et Lesq. (Sullivan and Lesquereux, 1856).

Distr. Southern South America (Hosseus, 1940; Herzog, 1952a).

Heterocladium

H. austro-alpinum (C. Muell.) Broth.

Leptopterigynandrum austro-alpinum

H. implexum (Schwaegr.) Lor.

Rigodium implexum

H. lechleri Schimp.

Rigodium arborescens and **R. toxarion**

H. prolixum Schimp.

Rigodium arborescens and **R. toxarion**

Heterophyllum

H. breviflagellosum C. Muell.

Acanthocladium breviflagellosum

Holodontium

H. inerme (Mitt.) Broth. (Brotherus, 1924a).

Syn. *Dicranum inerme* Mitt. (Mitten, 1869).

Dicranoweisia inermis (Mitt.) Par. (Paris, 1895).

Blindia auriculata C. Muell. (Mueller, 1885).

Dicranoweisia auriculata (C. Muell.) Broth. (Brotherus, 1901).

Distr. Southern South America (Mueller, 1885; Paris, 1895, 1904a; Brotherus, 1901, 1924a;

Dusén, 1905a; Cardot, 1905a, 1908a, 1913a; Cardot and Brotherus, 1923; Roivainen, 1934; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1939).

South Georgia (Bell, 1973b, p. 33).

H. pumilum (Mitt.) Broth. (Brotherus, 1924a).

Syn. *Symblepharis pumila* Mitt. (Mitten in Hooker, 1867).
Dicranum pumilum (Mitt.) Mitt. (Mitten, 1869) *hom. illeg.*
Holomitrium pumilum (Mitt.) Kindb. (Kindberg, 1888).
Blindia humilis C. Muell. (Mueller, 1885).
Dicranoweisia humilis (C. Muell.) Broth. (Brotherus, 1901).
Blindia churuccana Besch. (Bescherelle, 1885a).

Distr. Southern South America (Mitten, 1869; Mueller, 1885; Bescherelle, 1885a, 1889; Cardot, 1900, 1901, 1905a, 1908a; Brotherus, 1901, 1924a; Paris, 1904a, b; Dusén, 1905a; Cardot and Brotherus, 1923; Roivainen and Bartram, 1937; Kühnemann, 1938; Herzog, 1954; Seki, 1974).
Falkland Islands (Cardot and Brotherus, 1923).

H. ramulosum (Mitt.) Broth. (Brotherus, 1924a).

Syn. *Dicranum ramulosum* Mitt. (Mitten, 1869).

Distr. Southern South America (Mitten, 1869; Mueller, 1885; Paris, 1904b; Cardot, 1908a; Brotherus, 1924a (as *H. ramosum*)).

Holodontium sp.

Syn. *Blindia savatieriana* C. Muell. (Mueller in Bescherelle, 1889).
Blindia savatieri C. Muell. (Mueller, 1901a [1900]) *nom. illeg.*

Distr. Southern South America (Bescherelle, 1889; Mueller, 1901a; Paris, 1904a; Cardot, 1908a; Brotherus, 1924a).

Holomitrium

H. pumilum (Mitt.) Kindb.

Holodontium pumilum

Homalia

H. glabella (Hedw.) B.S.G. (Bruch and others, 1850).

Syn. *Leskea glabella* Hedw. (Hedwig, 1801).
Neckera glabella (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Neckera patagonica Brid. (Bridel, 1801b).
Pilotrichum patagonicum (Brid.) P. Beauv. (Palisot de Beauvois, 1805a).
Hypnum patagonicum Brid. (Bridel in Steudel, 1824) *nom. nud.*
Neckera dillenii Lindb. (Lindberg, 1883) *nom. illeg.*

Distr. Southern South America (Bridel, 1801b, 1827b; Schwaegrichen, 1816; Lindberg, 1883).

Hookeria

H. ancistrodes Mont.

Ancistrodes genuflexa

H. anomala (Schwaegr.) C. Muell.

Achrophyllum anomalum

H. apiculata Hook. f. et Wils.

Eriopus apiculatus

H. cirrhifolia Arnott

Sematophyllum cirrhifolium

H. concinna (Hook.) Hook. et Grev.

Lopidium concinnum

H. cristata (Hedw.) Hook. et Grev.

Eriopus cristatus

H. dentata Hook. f. et Wils.

Achrophyllum dentatum

H. denticulata Hook. f. et Wils.

Achrophyllum dentatum

H. dicksonii Hook. et Grev.

Distichophyllum dicksonii

H. eriodon C. Muell. ex Card.

Eriodon conostomus

H. erectiuscula Tayl.

Lepidopilum erectiusculum

H. flaccida Hook. f. et Wils.

Distichophyllum flaccidum

H. flexicollis Mitt.

Eriopus flexicollis

H. incurva (Hornsch.) Hook. et Grev.

Hookeriopsis incurva

H. lorentzii C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1925; Kühnemann, 1938; Welch, 1969).

H. magellanica (P. Beauv.) Arnott (Arnott, 1827).

Syn. *Hypnum magellanicum* P. Beauv. (Palisot de Beauvois, 1805a).

Distr. Southern South America (Palisot de Beauvois, 1805a; Bridel, 1817, 1827b; Wilson and Hooker, 1847; Mueller, 1850; Paris, 1904c; Kühnemann, 1938; Welch, 1969).

H. nigella Hook. f. et Wils.

Achrophyllum dentatum

H. obscura Mont.

Pterygophyllum crassirete

H. patrisiae (Brid.) Hamp.

Crossomitrium patrisiae

H. polytrichoides (Hedw.) Spreng.

Lepidopilum polytrichoides

H. rigida (Schwaegr.) Arnott

Lepidopilum polytrichoides

H. rotundifolia Hook. f. et Wils.

Distichophyllum rotundifolium

H. scabriuscula C. Muell.

Callicostella scabriuscula

H. sellovii Hornsch.

Mittenothamnium sellovii

H. speciosissima Mont. ex Schwaegr.

Schimperobryum splendidissimum

H. splendidissima Mont.

Schimperobryum splendidissimum

H. sublimbata C. Muell.

Cyclodictyon sublimbatum

H. tenella Hook. f. et Wils.

Sauloma tenella

H. uliginosa C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1925; Kühnemann, 1938; Welch, 1969).

Hookeriopsis

H. incurva (Hornsch.) Broth. (Brotherus, 1907).

Syn. *Chaetephora incurva* Hornsch. (Hornschuch, 1820a).

Hookeria incurva (Hornsch.) Hook. et Grev. (Hooker and Greville, 1825a).

Lepidopilum incurvum (Hornsch.) C. Muell. (Mueller, 1848b).

Callicostella incurva (Hornsch.) Aongstr. (Ångström, 1876a).

Distr. Southern South America (Hornschuch, 1820a, b; Hooker and Greville, 1825a; Bridel, 1827b; Schwaegrichen, 1829; Montagne, 1850; Mueller, 1851a; Jaeger, 1877; Paris, 1904b; Brotherus, 1907, 1925; Welch, 1969; Matteri, 1972).

Husnotiella

H. baueri Bartr. (Bartram, 1936).

Distr. Southern South America (Bartram, 1936; Hosseus, 1938c, d, 1939).

H. glossophylla Herz. (Herzog, 1916a).

Distr. Southern South America (Hosseus, 1936, 1937 (both as *H. glaucophylla* Herz.); Kühnemann, 1938).

Hygroamblystegium

H. antarcticum (Card.) Reim. (Reimers, 1926).

Syn. *Pseudoleskea antarctica* Card. (Cardot, 1900).

Distr. Southern South America (Reimers, 1926).

South Georgia (Cardot, 1906a, 1908a; Brotherus, 1925; Reimers, 1926; Steere, 1961b). Antarctic, peninsula region (Cardot, 1900, 1901, 1908a; Paris, 1905b; Brotherus, 1925; Steere, 1961a).

H. austro-fluviatile (C. Muell.) Broth. (Brotherus, 1908).

Syn. *Hypnum austro-fluviatile* C. Muell. (Mueller, 1890a).

Amblystegium austro-fluviatile (C. Muell.) Kindb. (Kindberg, 1891).

Distr. Southern South America (Thériot, 1924; Bartram, 1942, 1943).

South Georgia (Mueller, 1890a; Paris, 1903; Cardot, 1906a, 1908a; Brotherus, 1908, 1925; Reimers, 1926; Steere, 1961b).

***H. calochroum* (Card.) Reim. (Reimers, 1926).**

Syn. Pseudoleskea calochroa Card. (Cardot, 1906a).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

South Georgia (Cardot, 1906a, 1908a; Brotherus, 1925; Reimers, 1926; Steere, 1961b).

Antarctic, peninsula region (Cardot, 1911b, d, 1913a; Steere, 1961a).

***H. chilense* (Lor.) Reim. (Reimers, 1926).**

Syn. Amblystegium chilense Lor. (Lorentz, 1866).

Orthotheciella austro-chilense C. Muell. (Mueller in Reimers, 1926) *nom. nud.*

Distr. Southern South America (Lorentz, 1866; Jaeger and Sauerbeck, 1879a; Paris, 1903; Reimers, 1926; Herzog, 1939, 1940, 1954; Seki, 1974).

H. crassicostatum* Bartr. (Bartram, 1942) *hom. illeg.

Distr. Southern South America (Bartram, 1942, 1943).

H. excurrens* (Card. et Broth.) Reim.**Hygroamblystegium fuegianum* var. *excurrens******H. filum* (C. Muell.) Reim. (Reimers, 1926).**

Syn. Hypnum filum C. Muell. (Mueller, 1884 [1883]).

Orthothecium filum (C. Muell.) Kindb. (Kindberg, 1891).

Pseudoleskea filum (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Cardot, 1905a, 1908a; Thériot, 1918; Campo, 1921; Herzog, 1923, 1938, 1939; Brotherus, 1925; Reimers, 1926; Roivainen, 1934; Kühnemann, 1938; Bartram, 1946, 1952).

var. *compactum* Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

***H. fuegianum* (Besch.) Reim. (Reimers, 1926).**

Syn. Leskea fuegiana Besch. (Bescherelle, 1889).

Pseudoleskea fuegiana (Besch.) Broth. (Brotherus in Cardot, 1905a).

Distr. Southern South America (Bescherelle, 1889; Dusén, 1903a; Cardot, 1905a, 1908a; Paris 1905a; Thériot, 1915, 1917b; Campo, 1921; Costes, 1921; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1960; Reimers, 1926; Looser, 1932; Roivainen, 1934; Kühnemann, 1938; Bartram, 1946; Seki, 1974).

Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938).

var. *excurrens* (Card. et Broth.) Bartr. (Bartram, 1946).

Syn. Amblystegium excurrens Card. et Broth. (Cardot and Brotherus, 1923).

Hygroamblystegium excurrens (Card. et Broth.) Reim. (Reimers, 1926).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Reimers, 1926; Bartram, 1946, 1952).

var. *gracilis* (Card. et Broth.) Bartr. (Bartram, 1946).

Syn. Pseudoleskea fuegiana (Besch.) Broth. var. *gracilis* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Bartram, 1946).

var. *secundum* Bartr. (Bartram, 1946).

Distr. Southern South America (Bartram, 1946).

var. *skottsbergii* (Card.) Bartr. (Bartram in Roivainen, 1934).

Syn. Pseudoleskea fuegiana (Besch.) Broth. var. *skottsbergii* Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Roivainen, 1934; Kühnemann, 1938; Bartram, 1946).

Falkland Islands (Cardot and Brotherus, 1923).

H. luridum (Card.) Reim. (Reimers, 1926).

Syn. *Pseudoleskea lurida* Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Herzog, 1923; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938).

H. platyphyllum (Card.) Reim. (Reimers, 1926).

Syn. *Pseudoleskea platyphylla* Card. (Cardot, 1906a).

Distr. South Georgia (Cardot, 1906a, 1908a; Brotherus, 1925; Reimers, 1926; Steere, 1961b).

H. sordidoviride (Card. et Broth.) Reim. (Reimers, 1926).

Syn. *Pseudoleskea sordidoviridis* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938).

H. strictulum (Card.) Reim. (Reimers, 1926).

Syn. *Pseudoleskea strictula* Card. (Cardot, 1906a).

Distr. South Georgia (Cardot, 1906a, 1908a; Brotherus, 1925; Reimers, 1926; Steere, 1961b).

H. tenellum (Card. et Broth.) Reim. (Reimers, 1926).

Syn. *Amblystegium tenellum* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938; Herzog, 1954).

H. varium (Hedw.) Moenk.

Amblystegium varium

Hygrodicranum

H. falklandicum Card. (Cardot, 1911a).

Distr. Falkland Islands (Cardot, 1911a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

Hygrohypnum

H. aureum Herz. (Herzog, 1916a).

Distr. Southern South America (Bartram, 1965).

H. falcifolium Bartr. (Bartram, 1965).

Distr. Southern South America (Bartram, 1965).

H. peruviense Williams (Williams, 1916).

Distr. Southern South America (Bartram, 1942, 1943).

Hylocomium

H. lagurus (Hook.) Kindb.

Lepyrodon lagurus

H. schreberi (Brid.) De Not.

Pleurozium schreberi

H. splendidissimum De Not.

Ptychomnion cygnisetum

Hymenocleiston

H. magellanicum Dub.

Tayloria dubyi

var. *edenense* Besch.

Tayloria dubyi* var. *edenensis

Hymenodon

H. ovatus (Hook. f. et Wils.) C. Muell.

Leptotheca gaudichaudii

Hymenoloma

H. intermedium (Dix.) Broth.

Verrucidens intermedius

H. nordenskjoeldii Dus.

Dicranoweisia austro-crispula

H. subglobosum Herz. (Herzog, 1937).

Distr. Southern South America (Herzog, 1937).

H. turpe (Card.) Card. et Broth.

Verrucidens turpis

Hymenostomum

H. argentinicum (C. Muell.) Kindb. (Kindberg, 1891).

Syn. *Weisia argentinica* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904b; Brotherus, 1924a; Hosseus, 1935b, 1938c, d; Kühnemann, 1938).

H. balansae Besch. ex Kindb.

Hymenostomum balansaenum

H. balansaenum Besch. (Bescherelle, 1877).

Syn. *Hymenostomum balansae* Besch. ex Kindb. (Bescherelle ex Kindberg, 1888) nom. illeg.

Weisia balansaeana (Besch.) C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Felippone, 1909; Brotherus, 1924a; Herter, 1933a, 1943).

var. *densirete* Thér. (Thériot, 1935b).

Distr. Southern South America (Thériot, 1935b; Hosseus, 1938a, d).

H. griffithianum (Dicks.) Spreng.

Oedipodium griffithianum

H. kunzeanum (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Weisia kunzeana* C. Muell. (Mueller, 1849).

Gymnostomum marginatum Kunz. ex Mont. (Kunze ex Montagne, 1850) nom. illeg.

Hymenostomum marginatum Par. (Paris, 1904b) nom. illeg.

Distr. Juan Fernandez (Brotherus, 1924b; Robinson, 1975).

Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1870; Brotherus, 1902, 1924a; Paris, 1904b; Thériot, 1935a; Herzog, 1939, 1954).

H. marginatum Par.

Hymenostomum kunzeanum

H. semi-involutum (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Weisia semi-involuta* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1904b; Brotherus, 1924a; Kühnemann, 1938).

Hymenostylium

H. calcareum (Nees et Hornsch.) Mitt.

Gymnostomum aeruginosum

H. curvirostre Mitt.

Hymenostylium recurvirostre

H. kunzeanum (C. Muell.) C. Muell. (Mueller, 1901a [1900]).

Syn. *Pottia kunzeana* C. Muell. (Mueller in Montagne, 1850).

Gymnostomum kunzeanum (C. Muell.) Par. (Paris, 1895) nom. illeg.

Gymnostomum longirostris Kunz. ex C. Muell. (Mueller, 1847c) hom. illeg.

Pottia longirostris C. Muell. (Mueller, 1849) hom. illeg.

Weisia longirostris Mitt. (Mitten, 1869) hom. illeg.

Hymenostylium longirostre Broth. (Brotherus, 1902) hom. illeg.

Distr. Southern South America (Mueller, 1847c, 1849, 1901a; Montagne, 1850; Mitten, 1869; Jaeger, 1870; Paris, 1895, 1904b; Brotherus, 1902, 1924a; Warnstorf, 1916).

H. longirostre Broth.

Hymenostylium kunzeanum

H. longo-pulvinatum Dus.

Hymenostylium recurvirostre

H. recurvirostre (Hedw.) Dix. (Dixon, 1933 [1934]).

Syn. *Gymnostomum recurvirostre* Hedw. (Hedwig, 1801).

Pottia recurvirostra Ehrh. ex Hedw. (Hedwig, 1801) nom. inval.

Weisia recurvirostris (Hedw.) Dix. (Dixon, 1902) hom. illeg.

Eucladium recurvirostre (Hedw.) C. Jens. (Jensen, 1939).
Gymnostomum curvirostre Hedw. ex Brid. (Hedwig ex Bridel, 1801a) nom. illeg.
Weisia curvirostris C. Muell. (Mueller, 1849) hom. illeg.
Hymenostylium curvirostre Mitt. (Mitten, 1859) nom. illeg.
Barbula curvirostris Lindb. (Lindberg, 1879) hom. illeg.
Hymenostylium longo-pulvinatum Dus. (Dusén, 1906).
Barbula ripicola Card. (Cardot in Cardot and Brotherus, 1923) nom. nud.

Distr. Southern South America (Paris, 1904b; Dusén, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).

Hyophila

H. argentinica Thér. (Thériot, 1935b).

Distr. Southern South America (Bauer, 1934; Thériot, 1935b; Hosseus, 1936, 1937, 1938c, d, 1939; Kühnemann, 1938).

H. macrocarpa (Schimp.) Kindb.

Pottia macrocarpa

H. poeppigiana (C. Muell.) Jaeg.

Trichostomopsis australasiae

H. poeppigii C. Muell. ex Kindb.

Trichostomopsis australasiae

H. tortula (Schwaegr.) Hamp. (Hampe, 1846).

Syn. *Gymnostomum tortula* Schwaegr. (Schwaegrichen, 1826).

Distr. Southern South America (Herzog, 1952a).

Hypnodendron

H. arcuatum (Hedw.) Lindb. (Lindberg, 1861).

Syn. *Hypnum arcuatum* Hedw. (Hedwig, 1801).

Pterygophyllum arcuatum (Hedw.) Brid. (Bridel, 1827b).

Hypopterygium arcuatum (Hedw.) C. Muell. (Mueller, 1850).

Racopilum arcuatum (Hedw.) Jaeg. (Jaeger, 1876).

Hypnum spininervium Hook. (Hooker, 1818).

Rhizogonium spininervium (Hook.) Schimp. (Schimper in Mueller, 1847c).

Isothecium spininervium (Hook.) Wils. (Wilson, 1854).

Hypnodendron spininervium (Hook.) Reichdt. (Reichardt, 1870).

Isothecium spininervium (Hook.) Wils. var. *arcuatum* (Hook.) Wils. (Wilson, 1854).

Hypnodendron spininervium (Hook.) Reichdt. var. *arcuatum* (Hook.) Jaeg. (Jaeger and Sauerbeck, 1879a) nom. illeg.

Distr. Southern South America (Lorentz, 1866).

H. krausei (C. Muell.) Jaeg.

Hypnodendron microstictum

H. microstictum Mitt. (Mitten, 1869).

Syn. *Hypnum krausei* C. Muell. (Mueller, 1874).

Hypnodendron krausii (C. Muell.) Jaeg. (Jaeger and Sauerbeck, 1879a).

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Touw, 1971; Robinson, 1975).

Southern South America (Mitten, 1869; Mueller, 1874a; Jaeger and Sauerbeck, 1879a;

Neger, 1899; Paris, 1904b; Cardot, 1908a; Herzog, 1923, 1938, 1939, 1954, 1960; Brotherus, 1924a; Reimers, 1926; Kühnemann, 1938; Bizot, 1967; Touw, 1971; Seki, 1974).

H. naumannii (C. Muell.) Kindb.

Porothamnium arbusculans

H. spininervium (Hook.) Reichdt.

Hypnodendron arcuatum

var. *arcuatum* (Hedw.) Jaeg.

Hypnodendron arcuatum

Hypnodon

H. demissus C. Muell.

Rhachithecium perpusillum

H. perpusillus (Thwait. et Mitt.) C. Muell.

Rhachithecium perpusillum

Hypnopsis

H. schreberi (Brid.) Podp.

Pleurozium schreberi

Hypnum

<i>H. acacioides</i> (Schrad.) L. ex Turn.	<i>Fissidens osmundoides</i>
<i>H. acanthophyllum</i> Mont.	<i>Eurhynchiella acanthophylla</i>
<i>H. aciculare</i> Brid.	<i>Ptychomnion aciculare</i>
<i>H. adianthoides</i> (Hedw.) L. ex With.	<i>Fissidens adianthoides</i>
<i>H. aduncum</i> Hedw.	<i>Drepanocladus aduncus</i>
ssp. <i>polycarpum</i> (Voit) Meyl.	<i>Drepanocladus aduncus</i> var. <i>polycarpus</i>
var. <i>polycarpum</i> (Voit) B.S.G.	<i>Drepanocladus aduncus</i> var. <i>polycarpus</i>
var. <i>revolvens</i> (Sw.) Bland.	<i>Drepanocladus revolvens</i>
<i>H. albicans</i> Web. et Mohr	<i>Pohlia wahlenbergii</i>
<i>H. alopecurum</i> L. ex Hedw. var. <i>flagelliferum</i> Mont. (Montagne, 1835).	
Distr. Juan Fernandez (Montagne, 1835).	
<i>H. altisetum</i> (C. Muell.) Broth.	<i>Oxyrrhynchium altisetum</i>
<i>H. amblyostomum</i> C. Muell.	<i>Haplocladium microphyllum</i>
var. <i>attenuatifolium</i> C. Muell.	<i>Haplocladium microphyllum</i>
var. <i>paludicola</i> C. Muell.	<i>Haplocladium microphyllum</i>
<i>H. amoenum</i> Hedw.	<i>Rhaphidorrhynchium amoenum</i>
<i>H. aquicola</i> C. Muell.	<i>Bryum</i> sp.
<i>H. arborescens</i> C. Muell.	<i>Rigodium arborescens</i>
<i>H. arbuscula</i> P. Beauv.	<i>Hypopterygium arbuscula</i>
<i>H. arbusculans</i> C. Muell.	<i>Porothamnium arbusculans</i>
<i>H. arcuatum</i> Hedw.	<i>Hypnodendron arcuatum</i>
<i>H. argentinicum</i> C. Muell.	<i>Rigodium argentinicum</i>
<i>H. asplenoides</i> (Hedw.) Dicks. ex With.	<i>Fissidens asplenoides</i>
<i>H. attenuatifolium</i> C. Muell.	<i>Haplocladium microphyllum</i>
<i>H. auriculatum</i> Mont.	<i>Acrocladium auriculatum</i>
<i>H. austro-fluviatile</i> C. Muell.	<i>Hygroamblystegium austro-fluviatile</i>
<i>H. austro-glareosum</i> C. Muell.	<i>Brachythecium austro-glareosum</i>
<i>H. austro-salebrosum</i> C. Muell.	<i>Brachythecium austro-salebrosum</i>
<i>H. austro-serpens</i> C. Muell.	<i>Haplocladium microphyllum</i>
<i>H. austro-stramineum</i> C. Muell.	<i>Calliergidium austro-stramineum</i>
var. <i>gracillimum</i> C. Muell.	<i>Calliergidium austro-stramineum</i> var. <i>gracillimum</i>
var. <i>minus</i> Card.	<i>Calliergidium austro-stramineum</i> var. <i>minus</i>
var. <i>subfluitans</i> C. Muell.	<i>Calliergidium austro-stramineum</i> var. <i>subfluitans</i>
<i>H. berteroanum</i> Mont.	<i>Catagoniopsis berteroana</i>
<i>H. beskeanum</i> C. Muell.	<i>Rhynchosstegium beskeanum</i>
<i>H. bimum</i> (Schreb.) Web. et Mohr	<i>Bryum pseudotriquetrum</i> ssp. <i>bimum</i>
<i>H. brachiatum</i> Mitt.	<i>Drepanocladus brachiatus</i>
<i>H. brachypelma</i> C. Muell.	<i>Rigodium brachypodium</i>
<i>H. brachypodium</i> C. Muell.	<i>Rigodium brachypodium</i>
<i>H. brachypus</i> Hamp.	<i>Rhaphidorrhynchium brachypus</i>
<i>H. bracteatum</i> C. Muell.	<i>Sematophyllum bracteatum</i>
<i>H. brevipes</i> C. Muell.	<i>Stereophyllum brevipes</i>
<i>H. bryoides</i> (Hedw.) L. ex With.	<i>Fissidens bryoides</i>

<i>H. cacti</i> C. Muell.	<i>Isopterygium cacti</i>
<i>H. cacticola</i> C. Muell.	<i>Rhynchostegium cacticola</i>
<i>H. caespiticium</i> (Hedw.) Web. et Mohr	<i>Bryum caespiticium</i>
<i>H. calerae</i> C. Muell.	<i>Brachythecium calerae</i>
<i>H. callichroum</i> C. Muell.	<i>Catagonium politum</i> var. <i>phyllgonium</i>
<i>H. callidum</i> Mont.	<i>Rhaphidorrhynchium callidum</i>
<i>H. campicola</i> C. Muell.	<i>Leptodictyum campicola</i>
 <i>H. campoanum</i> (Thér.) Broth. (Brotherus, 1925).	
<i>Syn.</i> <i>Stereodon campoanus</i> Thér. (Thériot, 1918).	
<i>Distr.</i> Southern South America (Thériot, 1918; Campo, 1921; Brotherus, 1925; Herzog, 1938, 1954, 1960).	
<i>H. camptorrhynchium</i> Hamp.	<i>Mittenothamnium camptorrhynchium</i>
<i>H. chlamydophyllum</i> Hook. f. et Wils.	<i>Acrocladium auriculatum</i>
 <i>H. chrysogaster</i> C. Muell. (Mueller, 1851a).	
<i>Syn.</i> <i>Hypnum cupressiforme</i> L. ex Hedw. var. <i>chrysogaster</i> (C. Muell.) Sains. (Sainsbury, 1952).	
<i>Hypnum lechleri</i> C. Muell. (Mueller, 1856) <i>fide</i> Ando, 1972.	
<i>Stereodon lechleri</i> (C. Muell.) Mitt. (Mitten, 1869) <i>fide</i> Ando, 1972.	
<i>Hypnum lechleri</i> C. Muell. var. <i>fernandezianum</i> Card. ex Thér. (Cardot ex Thériot, 1921b) <i>fide</i> Ando, 1976.	
<i>Hypnum sublimatum</i> Lor. (Lorentz, 1866) <i>fide</i> Ando, 1976.	
<i>Ectropothecium sublimatum</i> (Lor.) Par. (Paris, 1895) <i>fide</i> Ando, 1976.	
<i>Hypnum cupressoides</i> Schimp. (Schimper in Lechler, 1857) <i>nom. nud. fide</i> Ando, 1976.	
<i>Distr.</i> Juan Fernandez (Jaeger and Sauerbeck, 1879a; Johow, 1896; Paris, 1904c; Skottsberg, 1914; Thériot, 1921b; Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975; Ando, 1976).	
Southern South America (Mueller, 1856; Lechler, 1857; Lorentz, 1866; Mitten, 1869; Jaeger and Sauerbeck, 1879a; Paris, 1895, 1904b, c; Cardot, 1908a; Campo, 1915; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1954, 1960; Brotherus, 1925; Roivainen, 1934; Kühnemann, 1938; Seki, 1974; Ando, 1976).	
 <i>H. circinale</i> Hook. (Hooker, 1819).	
<i>Syn.</i> <i>Hypnum circinnale</i> Hook. ex Macoun et Kindb. (Macoun and Kindberg, 1892).err <i>Drepanium circinale</i> (Hook.) Roth (Roth, 1904).	
<i>Distr.</i> Juan Fernandez (Montagne, 1835; Johow, 1896; Paris, 1904c). Southern South America (Montagne, 1850; Mueller, 1851a).	
<i>H. circinnale</i> Hook. ex Macoun et Kindb.	<i>Hypnum circinale</i>
<i>H. cirrifolium</i> (Arnott) Schwaegr.	<i>Sematophyllum cirrifolium</i>
<i>H. clarazii</i> Dub.	<i>Helicodontium clarazii</i>
<i>H. clinocarpum</i> Tayl.	<i>Oxyrrhynchium clinocarpum</i>
<i>H. cochlearifolium</i> Schwaegr.	<i>Weymouthia cochlearifolia</i>
<i>H. complanum</i> Mitt.	<i>Rhynchostegium complanum</i>
<i>H. confertum</i> Dicks.	<i>Rhynchostegium confertum</i>
<i>H. confluens</i> C. Muell.	<i>Sciaromium confluens</i>
<i>H. conostomum</i> Tayl.	<i>Brachythecium conostomum</i>
<i>H. conostomum</i> (Mont.) Mitt.	<i>Eriodon conostomus</i>
<i>H. conspissatum</i> Hook. f. et Wils.	<i>Sciaromium conspissatum</i>
<i>H. corralense</i> Lor.	<i>Oxyrrhynchium corralense</i>
<i>H. crassiusculum</i> Brid.	<i>Sematophyllum crassiusculum</i>
<i>H. crinitifolium</i> C. Muell.	<i>Leucomium crinitifolium</i>

<i>H. cristatum</i> Poir.	<i>Eriopus cristatus</i>
<i>H. crudum</i> (Hedw.) Web. et Mohr	<i>Pohlia cruda</i>
<i>H. cucullifolium</i> P. Beauv.	<i>Ptychomnion aciculare</i>
<i>H. cupressiforme</i> L. ex Hedw. (Hedwig, 1801).	
Syn. <i>Stereodon cupressiformis</i> (Hedw.) Brid. ex Mitt. (Bridel ex Mitten, 1859). <i>Cupressina cupressiformis</i> (Hedw.) C. Muell. (Mueller, 1896a). <i>Drepanium cupressiforme</i> (Hedw.) Roth (Roth, 1904). <i>Hypnum cupressiforme</i> L. ex Hedw. var. <i>tectorum</i> Brid. (Bridel, 1827b).	
Distr. Southern South America (Campo, 1915; Cardot and Brotherus, 1923; Brotherus, 1925; Reimers, 1926; Herzog, 1938, 1940; Kühnemann, 1938; Bartram, 1952; Seki, 1974).	
var. <i>chrysogaster</i> (C. Muell.) Sains.	<i>Hypnum chrysogaster</i>
var. <i>tectorum</i> Brid.	<i>Hypnum cupressiforme</i>
<i>H. cupressoides</i> Schimp.	<i>Hypnum chrysogaster</i>
<i>H. curtipedulum</i> (Hedw.) L. ex With.	<i>Antitrichia curtipedula</i>
<i>H. cuspidatum</i> Hedw.	<i>Calliergonella cuspidata</i>
<i>H. cygnisetum</i> C. Muell.	<i>Ptychomnion cygnisetum</i>
<i>H. delicatulum</i> Hedw.	<i>Thuidium recognitum</i> var. <i>delicatulum</i>
<i>H. densifolium</i> Brid.	<i>Ptychomnion densifolium</i>
<i>H. denticulatum</i> Hedw.	<i>Plagiothecium denticulatum</i>
var. <i>obtusifolium</i> Turn.	<i>Plagiothecium denticulatum</i> var. <i>obtusifolium</i>
<i>H. diminutivum</i> Hamp.	<i>Mittenothamnium diminutivum</i>
<i>H. dolomiticum</i> Mild.	<i>Hypnum revolutum</i> var. <i>dolomiticum</i>
<i>H. donnianum</i> Sm.	<i>Plagiothecium denticulatum</i> var. <i>obtusifolium</i>
<i>H. drepanocladoides</i> C. Muell.	<i>Rhynchosstegium drepanocladoides</i>
<i>H. encalyptatum</i> Schimp.	<i>Acrocladium auriculatum</i>
<i>H. eucalyptatum</i> Lindb. ex Card.	<i>Acrocladium auriculatum</i>
<i>H. exannulatum</i> B.S.G.	<i>Drepanocladus exannulatus</i>
<i>H. falcatum</i> Brid.	see <i>Hypnum longifolium</i>
<i>H. fasciculatum</i> Hedw.	<i>Porothamnium fasciculatum</i>
<i>H. filarium</i> (Mitt.) Hamp.	<i>Thuidiopsis furfurosa</i>
<i>H. filicinum</i> Hedw.	<i>Cratoneuron filicinum</i>
<i>H. filum</i> C. Muell.	<i>Hygroamblystegium filum</i>
<i>H. fissidens</i> C. Muell.	<i>Rhynchosstegium fissidens</i>
<i>H. flagellare</i> Hedw.	<i>Brachythecium plumosum</i>
<i>H. fluitans</i> Hedw.	<i>Drepanocladus fluitans</i>
var. <i>australe</i> Card.	<i>Drepanocladus fluitans</i> var. <i>australe</i>
var. <i>falcifolium</i> Ren.	<i>Drepanocladus fluitans</i>
var. <i>iaculosum</i> (C. Muell.) Par.	<i>Drepanocladus laculosus</i>
var. <i>polycarpon</i> (Voit) Fiedl.	<i>Drepanocladus aduncus</i> var. <i>polycarpus</i>
<i>H. fuegianum</i> (Mitt.) C. Muell.	<i>Drepanocladus fuegianus</i>
<i>H. fulvo-acutum</i> (C. Muell.) Par.	<i>Calliergonella cuspidata</i>
<i>H. furfurosum</i> Hook. f. et Wils.	<i>Thuidiopsis furfurosa</i>
<i>H. gayanum</i> (Mont.) Lor.	<i>Ptychomnion ptychocarpum</i>
<i>H. georgico-antarcticum</i> C. Muell.	<i>Plagiothecium georgico-antarcticum</i>

<i>H. georgico-glareosum</i> C. Muell.	<i>Brachythecium austro-salebrosum</i>
<i>H. georgico-uncinatum</i> C. Muell.	<i>Drepanocladus uncinatus</i>
<i>H. glauco-virescens</i> C. Muell.	<i>Rhynchosstegium glauco-virescens</i>
<i>H. globipyxis</i> C. Muell.	<i>Rhynchosstegium globipyxis</i>
<i>H. gracile</i> Hook. f et Wils.	<i>Camptochaete gracilis</i>
<i>H. gracile</i> Par. (Paris, 1904c) <i>nom. nud.</i>	
Syn. <i>Stereodon gracilis</i> Dus. (Dusén in Paris, 1904c) <i>nom. nud.</i>	
Distr. Southern South America (Paris, 1904c).	
<i>H. gracilescens</i> C. Muell.	<i>Camptochaete gracilis</i>
<i>H. harrisona</i> Poir.	<i>Neckeropsis undulata</i>
<i>H. heufleri</i> Jur.	<i>Hypnum revolutum</i>
<i>H. homaliocaulon</i> C. Muell.	<i>Rhynchosstegium homaliocaulon</i>
<i>H. hookeri</i> Lindb.	<i>Catagonium nitidum</i>
<i>H. implexum</i> (Schwaegr.) C. Muell.	<i>Rigodium implexum</i>
<i>H. intermedium</i> Lindb. var. <i>revolvens</i> (Sw.) San.	<i>Drepanocladus revolvens</i>
<i>H. julaceum</i> (Hedw.) Schleich.	<i>Leucodon julaceus</i>
<i>H. kneiffii</i> (B.S.G.) Schimp. var. <i>polycarpus</i> (Voit) Fam.	<i>Drepanocladus aduncus</i> var. <i>polycarpus</i>
<i>H. krausei</i> C. Muell.	<i>Hypnodendron microstictum</i>
<i>H. laculosum</i> C. Muell.	<i>Drepanocladus laculosus</i>
<i>H. lagurum</i> Menz.	<i>Lepyrodon lagurus</i>
<i>H. laterculi</i> C. Muell.	<i>Haplocladium microphyllum</i>
<i>H. lechleri</i> C. Muell.	<i>Hypnum chrysogaster</i>
var. <i>fernandezianum</i> Card. ex Thér.	<i>Hypnum chrysogaster</i>
<i>H. lepidopilooides</i> C. Muell.	<i>Stereophyllum lepidopilooides</i>
<i>H. leptorhynchum</i> Brid.	<i>Rhaphidorrhynchium amoenum</i>
<i>H. leucocytus</i> C. Muell.	<i>Sematophyllum leucocytus</i>
<i>H. leucophyllum</i> Hamp.	<i>Isopterygium leucophyllum</i>
<i>H. leucostegum</i> (Brid.) C. Muell.	<i>Stereophyllum leucostegum</i>
<i>H. lithophilum</i> Hook. f. et Wils.	<i>Drepanocladus secundifolius</i>
<i>H. longidens</i> C. Muell.	<i>Brachythecium longidens</i>
<i>H. longifolium</i> Wils. (Wilson, 1855) <i>hom. illeg.</i>	
Distr. Falkland Islands (Wilson and Hooker, 1847 (as <i>H. falcatum</i> Brid. <i>fide</i> Wilson, 1855); Wilson, 1855; Kühnemann, 1938 (as <i>H. falcatum</i> Brid. <i>fide</i> Wilson, 1855)).	
<i>H. longifolium</i> (Mitt.) Jaeg.	<i>Drepanocladus brachiatus</i>
<i>H. loxense</i> Hook.	<i>Sematophyllum caespitosum</i>
<i>H. lucidulum</i> Hook. f. et Wils.	<i>Plagiothecium lucidulum</i>
<i>H. lucidum</i> Hook. f. et Wils.	<i>Plagiothecium lucidum</i>
<i>H. macroconum</i> C. Muell.	<i>Brachythecium plumosum</i>
var. <i>robustius</i> C. Muell.	<i>Brachythecium macroconum</i> var. <i>robustius</i>
<i>H. magellanicum</i> P. Beauv.	<i>Hookeria magellanica</i>
<i>H. magellanicum</i> C. Muell.	<i>Isopterygium fuegianum</i>
<i>H. masafuerae</i> Bartr. (Bartram, 1959).	
Distr. Juan Fernandez (Bartram, 1959).	

<i>H. micans</i> Wils. var. <i>laxum</i> Hook. f.	<i>Campochaete gracilis</i>
<i>H. microcarpoides</i> C. Muell.	<i>Sematophyllum microcarpoides</i>
<i>H. microphyllum</i> Hedw.	<i>Haplocladium microphyllum</i>
<i>H. micropyxis</i> C. Muell.	<i>Trichosteleum</i> sp.
<i>H. mnioides</i> Hook.	<i>Rhizogonium mnioides</i>
<i>H. morenoi</i> (C. Muell.) Par.	<i>Pleurozium schreberi</i>
<i>H. nanum</i> C. Muell.	<i>Rhaphidostegium nanum</i>
<i>H. naumanii</i> C. Muell.	<i>Porothamnium arbusculans</i>
<i>H. neckera</i> Schimp.	<i>Porothamnium valdiviae</i>
<i>H. neckeraeforme</i> Hamp.	<i>Porothamnium neckeraeforme</i>
<i>H. neckeroides</i> Hook.	<i>Thamnium neckeroides</i>
<i>H. neei</i> Mohr	<i>Rigodium implexum</i>
<i>H. nematogonium</i> C. Muell.	<i>Brachythecium nematogonium</i>
<i>H. nigrescens</i> Hedw.	<i>Papillaria nigrescens</i>
<i>H. nitidum</i> (Hook. f. et Wils.) C. Muell.	<i>Catagonium nitidum</i>
<i>H. obtusifolium</i> (Turn.) Brid.	<i>Plagiothecium denticulatum</i> var. <i>obtusifolium</i>
<i>H. oligorrhizon</i> Guemb.	<i>Amblystegium serpens</i>
<i>H. ornithopodioides</i> Brid.	<i>Racopilum tomentosum</i>
<i>H. orthocarpon</i> Hornsch. ex C. Muell.	<i>Plagiothecium orthocarpum</i>
<i>H. orthothecioides</i> (Lindb.) Par.	<i>Drepanocladus uncinatus</i> var. <i>subjulaceus</i>
<i>H. pachyloma</i> (Mont.) C. Muell.	<i>Sciaromium pachyloma</i>
<i>H. pallens</i> Schimp. ex Card.	<i>Hypnum skottsbergii</i>
<i>H. paludicola</i> C. Muell.	<i>Haplocladium microphyllum</i>
<i>H. palustre</i> (Hedw.) Web. et Mohr	<i>Aulacomnium palustre</i>
<i>H. pampae</i> C. Muell.	<i>Rhynchosstegium pampae</i>
<i>H. panduraefolium</i> C. Muell.	<i>Porothamnium panduraefolium</i>
<i>H. paradoxum</i> Hook. f. et Wils. var. <i>leptocladum</i> C. Muell. (Mueller, 1889).	<i>Brachythecium paradoxum</i>
<i>Distr.</i> Southern South America (Mueller, 1889).	
<i>H. parietinum</i> Wahlenb.	<i>Pleurozium schreberi</i>
<i>H. patagonicum</i> Brid.	<i>Homalia glabella</i>
<i>H. patrisiae</i> Brid.	<i>Crossomitrium patrisiae</i>
<i>H. pendulum</i> (Brid.) C. Muell.	<i>Helicodontium pendulum</i>
<i>H. perplicatum</i> Dus.	<i>Cratoneuron perplicatum</i>
<i>H. persistens</i> C. Muell.	<i>Haplocladium persistens</i>
<i>H. pervirens</i> C. Muell.	<i>Helicodontium pervirens</i>
<i>H. phyllogonium</i> C. Muell.	<i>Catagonium politum</i> var. <i>phyllogonium</i>
<i>H. pinnicaule</i> C. Muell.	<i>Rhynchosstegium pinnicaule</i>
<i>H. piriforme</i> Web. et Mohr ex Par.	<i>Leptobryum pyriforme</i>
<i>H. plicatile</i> (Mitt.) Lesq. et Jam. ssp. <i>revolutum</i> (Mitt.) Kindb.	<i>Hypnum revolutum</i>
<i>H. plinthophilum</i> C. Muell.	<i>Mittenothamnium plinthophilum</i>
<i>H. plumosum</i> Hedw.	<i>Brachythecium plumosum</i>
<i>H. politum</i> Hook. f. et Wils.	<i>Catagonium politum</i>
<i>H. polycarpon</i> Bland. ex Voit	<i>Drepanocladus aduncus</i> var. <i>polycarpus</i>

<i>H. polygamum</i> (B.S.G.) Wils.	<i>Campylium polygamum</i>
<i>H. polytrichoides</i> Hedw.	<i>Lepidopilum polytrichoides</i>
<i>H. pseudo-limnobium</i> C. Muell.	<i>Helicodontium pseudo-limnobium</i>
<i>H. pseudo-pallens</i> Dus. (Dusén in Paris, 1905a) <i>nom. nud.</i>	
Distr. Southern South America (Paris, 1905a).	
var. <i>gracile</i> Dus. (Dusén in Paris, 1905a) <i>nom. nud.</i>	
Distr. Southern South America (Paris, 1905a).	
<i>H. pseudo-purum</i> (C. Muell.) Par.	<i>Pseudoscleropodium pseudopurum</i>
<i>H. pseudo-triquetrum</i> (Hedw.) Web. et Mohr	<i>Bryum pseudotriquetrum</i>
<i>H. ptychocarpum</i> Schwaegr.	<i>Ptychomnion ptychocarpum</i>
<i>H. purpurascens</i> Brid.	<i>Rhacocarpus purpurascens</i>
<i>H. pyriforme</i> (Hedw.) Web. et Mohr	<i>Leptobryum pyriforme</i>
<i>H. radicale</i> P. Beauv.	<i>Campylium radicale</i>
<i>H. raphidostegum</i> C. Muell.	<i>Thuidium raphidostegum</i>
<i>H. remotifolium</i> Grev.	<i>Oxyrrhynchium clinocarpum</i>
<i>H. reticulatum</i> Hook. f. et Wils.	<i>Philonotis vagans</i>
<i>H. revolutum</i> (Mitt.) Lindb. (Lindberg, 1866 [1867]).	
<i>Syn. Stereodon revolutus</i> Mitt. (Mitten, 1859).	
<i>Drepanium revolutum</i> (Mitt.) C. Jens. (Jensen in Lange and Jensen, 1887).	
<i>Hypnum plicatile</i> (Mitt.) Lesq. et Jam. ssp. <i>revolutum</i> (Mitt.) Kindb. (Kindberg, 1897) <i>nom. illeg.</i>	
<i>Hypnum heufleri</i> Jur. (Juratzka, 1861).	
Distr. Antarctic, peninsula region (Cardot, 1900, 1901, 1908a; Paris, 1905a; Steere, 1961a; Robinson, 1972; Ando, 1973).	
Antarctic, continental region (Bartram, 1957; Steere, 1961a).	
<i>var. dolomiticum</i> (Mild.) Moenk. (Mönkemeyer, 1927).	
<i>Syn. Hypnum dolomiticum</i> Mild. (Milde, 1864a).	
Distr. Antarctic, peninsula region (Ando, 1973).	
<i>H. revolvens</i> Sw.	<i>Drepanocladus revolvens</i>
<i>H. raphidostegum</i> C. Muell. ex Par.	<i>Thuidium raphidostegum</i>
<i>H. rigidum</i> Schwaegr.	<i>Lepidopilum polytrichoides</i>
<i>H. riparioides</i> Hedw.	<i>Platyhypnidium riparioides</i>
<i>H. riparium</i> L. ex Hedw. ssp. <i>kochii</i> (B.S.G.) Boul.	<i>Leptodictyum kochii</i>
<i>H. rivularioides</i> C. Muell.	<i>Brachythecium rivularioides</i>
<i>H. rostratum</i> (Schrad.) Web. et Mohr	<i>Mnium rostratum</i>
<i>H. rusciforme</i> Brid.	<i>Platyhypnidium riparioides</i>
<i>H. rutabulum</i> Hedw.	<i>Brachythecium rutabulum</i>
<i>H. sarmentosum</i> Wahlenb.	<i>Calliergon sarmentosum</i>
<i>H. scabrifolium</i> Hook. f. et Wils.	<i>Philonotis scabrifolia</i>
<i>H. schreberi</i> Brid.	<i>Pleurozium schreberi</i>
<i>H. sclerodontium</i> C. Muell.	<i>Dimerodontium pellucidum</i>
<i>H. scorpiurus</i> Mont.	<i>Rhaphidorrhynchium scorpiurus</i>
<i>H. secundifolium</i> C. Muell.	<i>Drepanocladus secundifolius</i>
<i>H. sellowianum</i> C. Muell.	<i>Rhynchostegium sellowii</i>
<i>H. sellowii</i> Hornsch.	<i>Rhynchostegium sellowii</i>
<i>H. semilunare</i> C. Muell.	<i>Haplocladium microphyllum</i>

<i>H. semi-nerve</i> (Schwaegr.) C. Muell.	<i>Juratzkaea seminervis</i>
<i>H. semi-tortulum</i> C. Muell.	<i>Rhynchosstegium semi-tortulum</i>
<i>H. sericeo-virens</i> C. Muell.	<i>Brachythecium sericeo-virens</i>
<i>H. serpens</i> Hedw.	<i>Amblystegium serpens</i>
<i>H. serrulatum</i> Hedw.	<i>Rhynchosstegium serrulatum</i>
<i>H. simorhynchum</i> Hamp.	<i>Mittenothamnium simorhynchum</i>
<i>H. simplex</i> Brid.	<i>Sematophyllum subsimplex</i>
<i>H. skottsbergii</i> Ando (Ando, 1972) <i>nom nud.</i>	
Syn. <i>Rhaphidostegium pallens</i> Dus. (Dusén, 1903a) <i>nom. nud.</i>	
<i>Stereodon pallens</i> Broth. (Brotherus, 1908) <i>nom. nud.</i>	
<i>Hypnum pallens</i> Schimp. ex Card. (Schimper ex Cardot, 1908a) <i>hom. illeg.</i>	
<i>Distr.</i> Southern South America (Mueller, 1885; Dusén, 1903a; Paris, 1904c; Cardot, 1905a, 1908a; Brotherus, 1908, 1925; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1940, 1954, 1957, 1960; Thériot, 1934b; Kühnemann, 1938; Seki, 1974).	
<i>H. smithii</i> Hedw.	<i>Leptodon smithii</i>
<i>H. sparsifolium</i> Hamp.	<i>Amblystegium sparsifolium</i>
<i>H. speciosum</i> C. Muell.	<i>Hypopterygium arbuscula</i>
<i>H. spegazzinii</i> C. Muell. (Mueller, 1885).	
Syn. <i>Brachythecium spegazzinii</i> (C. Muell.) Kindb. (Kindberg, 1888).	
<i>Stereodon spegazzinii</i> (C. Muell.) Broth. (Brotherus, 1908).	
<i>Distr.</i> Southern South America (Mueller, 1885; Paris, 1905a; Brotherus, 1908; Cardot, 1908a).	
<i>H. spiniforme</i> Hedw.	<i>Rhizogonium spiniforme</i>
<i>H. spininervium</i> Hook.	<i>Hypnodendron arcuatum</i>
<i>H. squalidissimum</i> C. Muell.	<i>Brachythecium squalidissimum</i>
<i>H. squarroso-byssoides</i> (C. Muell.) Par.	<i>Campylium squarroso-byssoides</i>
<i>H. stenopyxidium</i> C. Muell.	<i>Sematophyllum stenopyxidium</i>
<i>H. stramineum</i> Dicks. ex Brid.	<i>Calliergon stramineum</i>
<i>H. strigosum</i> Hoffm. ex Web. et Mohr	<i>Eurhynchium pulchellum</i>
<i>H. subbasilare</i> Hook.	<i>Goniobryum subbasilare</i>
<i>H. sublimatum</i> Lor.	<i>Hypnum chrysogaster</i>
<i>H. subpilosum</i> Hook. f. et Wils.	<i>Brachythecium subpilosum</i>
<i>H. subplicatum</i> Hamp.	<i>Brachythecium subplicatum</i>
<i>H. subsimplex</i> Hedw.	<i>Sematophyllum subsimplex</i>
<i>H. sulphureum</i> Geh. et Hamp.	<i>Brachythecium sulphureum</i>
<i>H. symmetricum</i> (Ren. et Card.) Ren. et Card.	<i>Drepanocladus uncinatus</i> var. <i>symmetricus</i>
<i>H. tamariscinum</i> Hedw.	<i>Thuidium tamariscinum</i>
var. <i>delicatulum</i> (Hedw.) Brid.	<i>Thuidium recognitum</i> var. <i>delicatulum</i>
<i>H. taxarion</i> Schwaegr. ex Mont.	<i>Rigodium toxarion</i>
<i>H. taxifolium</i> (Hedw.) L. ex With.	<i>Fissidens taxifolius</i>
<i>H. taylorii</i> Mitt.	<i>Brachythecium conostomum</i>
<i>H. tenerum</i> Sw.	<i>Isopterygium tenerum</i>
<i>H. tenuifolium</i> Hedw.	<i>Rhynchosstegium tenuifolium</i>
<i>H. tenuirostre</i> (Schwaegr.) C. Muell.	<i>Helicodontium tenuirostre</i>
<i>H. thelistegum</i> C. Muell.	<i>Mittenothamnium diminutivum</i>
<i>H. thieleanum</i> C. Muell.	<i>Pinnatella thieleana</i>

<i>H. thouinii</i> Schwaegr.	<i>Hypopterygium arbuscula</i>
<i>H. tomentosum</i> Hedw.	<i>Racopilum tomentosum</i>
<i>H. torquatum</i> Hedw.	<i>Schlotheimia torquata</i>
<i>H. toxarion</i> Schwaegr.	<i>Rigodium toxarion</i>
<i>H. trichopodium</i> Schultz var. <i>kochii</i> Lindb. ex Par.	<i>Leptodictyum kochii</i>
<i>H. turbinatum</i> (Hedw.) Web. et Mohr	<i>Bryum turbinatum</i>
<i>H. uncinatum</i> Hedw.	<i>Drepanocladus uncinatus</i>
ssp. <i>symmetricum</i> Ren. et Card.	<i>Drepanocladus uncinatus</i> var. <i>symmetricus</i>
var. <i>cylindropyxis</i> Dus.	<i>Drepanocladus uncinatus</i> var. <i>symmetricus</i>
var. <i>orthotheciooides</i> (Lindb.) Zett.	<i>Drepanocladus uncinatus</i> var. <i>subjulaceus</i>
var. <i>plumosum</i> Schimp.	<i>Drepanocladus uncinatus</i>
var. <i>subjulaceum</i> B.S.G.	<i>Drepanocladus uncinatus</i> var. <i>subjulaceus</i>
<i>H. unguiculatum</i> Hook. f. et Wils.	<i>Thuidiopsis furfurosa</i>
<i>H. valdiviae</i> C. Muell.	<i>Porothamnium valdiviae</i>
var. <i>pallidum</i> Lor.	<i>Porothamnium valdiviae</i> var. <i>pallidum</i>
<i>H. varium</i> (Hedw.) P. Beauv.	<i>Amblystegium varium</i>
<i>H. ventricosum</i> Web. et Mohr	<i>Bryum pseudotriquetrum</i>
<i>H. vesiculare</i> Schwaegr.	<i>Vesicularia vesicularis</i>
<i>H. wahlenbergii</i> Web. et Mohr	<i>Pohlia wahlenbergii</i>

Hypopterygium

H. arbuscula Brid. (Bridel, 1827b).

Syn. *Hypnum arbuscula* P. Beauv. (Palisot de Beauvois, 1805a) hom. illeg.
Hypnum speciosum C. Muell. (Mueller, 1844b [1845]) hom. illeg.
Hypopterygium speciosum C. Muell. (Mueller, 1844b [1845]).
Hypnum thouinii Schwaegr. (Schwagrichen, 1830).
Hypopterygium thouinii (Schwaegr.) Mont. (Montagne, 1845a).

Distr. Juan Fernandez (Brotherus, 1924a, b; Espinosa B., 1941; Bartram, 1959; Matteri, 1973b; Robinson, 1975).

Southern South America (Palisot de Beauvois, 1805a; Bridel, 1827b; Schwaegrichen, 1830; Mueller, 1844b, 1850, 1885; Montagne, 1845a, 1850, 1856; Wilson and Hooker, 1847; Lorentz, 1866; Mitten, 1869; Jaeger, 1876; Bescherelle, 1885a; Hariot, 1891; Eaton, 1892; Neger, 1899; Paris, 1905a; Cardot, 1908a; Campo, 1915, 1921; Thériot, 1915, 1918; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1940, 1954, 1957, 1960; Brotherus, 1924a; Reimers, 1926; Roivainen, 1934; Kühnemann, 1938; Bartram, 1952; Bizot, 1967; Matteri, 1973b, 1975a; Seki, 1974).

H. arcuatum (Hedw.) C. Muell.

Hypnodendron arcuatum

H. argentinicum Lor. ex C. Muell. (Lorentz ex Mueller, 1879a).

Syn. *Hypopterygium tamarisci* (Sw.) Brid. ex C. Muell. ssp. *argentinicum* (C. Muell.) Kindb. (Kindberg, 1901).

Distr. Southern South America (Mueller, 1879a; Kindberg, 1901; Paris, 1905a; Brotherus, 1925; Herter, 1933a, 1939a; Kühnemann, 1938; Herzog, 1952a).

H. concinnum (Hook.) Brid.

Lopidium concinnum

H. didictyon C. Muell. (Mueller, 1850).

Distr. Southern South America (Wilson and Hooker, 1847 (as *H. laricinum* fide Mueller, 1850); Mueller, 1850, 1885; Sullivant, 1859; Mitten, 1869; Jaeger, 1876; Bescherelle, 1889; Hariot, 1891; Neger, 1899; Kindberg, 1901; Cardot, 1905a, 1908a; Paris, 1905a; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1940, 1954, 1960; Thériot, 1924, 1934b; Brotherus, 1925; Kühnemann, 1938; Bartram, 1952; Matteri, 1973b, 1975a; Seki, 1974).

H. flavescens Hamp. (Hampe, 1847).

Distr. Southern South America (Herzog, 1952a).

H. flexisetum Hamp.

Lopidium concinnum

H. incrassato-limbatum C. Muell.

Hypopterygium laricinum ssp. *incrassato-limbatum*

H. krauseanum C. Muell.

Hypopterygium novae-seelandiae var. *chilensis*

H. laricinum (Hook.) Brid. ssp. *incrassato-limbatum* (C. Muell.) Kindb. (Kindberg, 1901).

Syn. *Hypopterygium incrassato-limbatum* C. Muell. (Mueller, 1850).

Distr. Southern South America (Krieger, 1904).

H. novae-seelandiae C. Muell. var. *chilensis* Lor. (Lorentz, 1866).

Syn. *Hypopterygium krauseanum* C. Muell. (Mueller in Kindberg, 1901).

Distr. Southern South America (Lorentz, 1866; Kindberg, 1901; Paris, 1905a).

H. pallens (Hook. f. et Wils.) Mitt.

Lopidium concinnum

ssp. *plumarium* (Mitt.) Kindb.

Lopidium concinnum

H. plumarium Mitt.

Lopidium concinnum

H. speciosum C. Muell.

Hypopterygium arbuscula

H. squarrulosum C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1925; Herter, 1933a, 1943; Kühnemann, 1938 (as *H. squamulosum* C. Muell.); Hosseus, 1939).

H. tamarisci (Sw.) Brid. ex C. Muell. ssp. *argentinicum* (C. Muell.) Kindb.

Hypopterygium argentinicum

H. thouinii (Schwaegr.) Mont.

Hypopterygium arbuscula

H. tomentosum (Hedw.) C. Muell.

Racopilum tomentosum

H. uliginosum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

H. wolffhuegelii Herz. (Herzog, 1923).

Distr. Southern South America (Herzog, 1923; Brotherus, 1925).

Isopterygium

I. adflatum (C. Muell.) Par. (Paris, 1900a).

Syn. *Taxicaulis adflatus* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1925; Kühnemann, 1938).

I. arechavaletae Broth. (Brotherus in Krieger, 1904) nom. nud.

Distr. Southern South America (Krieger, 1904).

I. byssobolax (C. Muell.) Par. (Paris, 1900a).

Syn. *Taxicaulis byssobolax* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1918, 1925; Kühnemann, 1938).

I. cacti (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum cacti* C. Muell. (Mueller, 1879a).

Plagiothecium cacti (C. Muell.) Kindb. (Kindberg, 1891).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1925; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939).

I. elegans (Brid.) Lindb. var. *nanum* (Jur.) Jaeg. (Jaeger, 1878).

Syn. *Plagiothecium nanum* Jur. (Juratzka in Milde, 1864b).

Isopterygium nanum (Jur.) Loesk. (Loeske, 1910) nom. inval.

Plagiothecium schimperi Jur. et Mild. var. *nanum* (Jur.) Walth. et Mol. (Walther and Molendo, 1868).

Plagiothecium elegans (Brid.) Sull. var. *nanum* (Jur.) Mol. (Molendo, 1875).

Distr. Falkland Islands (Cardot and Brotherus, 1923).

I. fernandezianum Broth.

Isopterygium tenerum

var. *longisetum* Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959).

I. fuegianum Besch. (Bescherelle, 1889).

Syn. *Plagiothecium fuegianum* (Besch.) Dus. (Dusén, 1903a).

Hypnum magellanicum C. Muell. (Mueller, 1885) nom. illeg.

Plagiothecium magellanicum Par. (Paris, 1896) nom. illeg.

Distr. Southern South America (Mueller, 1885; Bescherelle, 1889; Paris, 1896, 1905b; Dusén, 1903a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

I. leptoplumosum (Dus.) Card. (Cardot, 1908a).

Syn. *Plagiothecium leptoplumosum* Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Kühnemann, 1938; Seki, 1974).

I. leucophyllum (Hamp.) Mitt. (Mitten, 1869).

Syn. *Hypnum leucophyllum* Hamp. (Hampe in Mueller, 1851a).

Plagiothecium leucophyllum Hamp. (Hampe in Mueller, 1851a).

Distr. Southern South America (Mueller, 1851a; Mitten, 1869; Jaeger, 1878).

I. nanum (Jur.) Loesk.

Isopterygium elegans* var. *nanum

I. pyrrhopus (C. Muell.) Par. (Paris, 1900a).

Syn. *Taxicaulis pyrrhopus* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1925; Kühnemann, 1938).

I. saprophilum (C. Muell.) Par. (Paris, 1900a).

Syn. *Taxicaulis saprophilus* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1925; Kühnemann, 1938).

I. tenerum (Sw.) Mitt. (Mitten, 1869).

Syn. *Hypnum tenerum* Sw. (Swartz, 1806b).

Isopterygium fernandezianum Broth. (Brotherus, 1924b) fide Robinson, 1975.

Distr. Juan Fernandez (Skottsberg, 1914; Brotheus, 1924b, 1925; Bartram, 1959; Robinson, 1975).

Isothecium

I. amoenum (Hedw.) Brid.

Rhaphidorrhynchium amoenum

I. cochlearifolium (Schwaegr.) Mitt.

Weymouthia cochlearifolia

I. gracile Hook. f. et Wils.

Camptochaete gracilis

I. molle (Hedw.) Mitt.

Weymouthia mollis

I. neckeroides (Hook.) Brid.

Thamnium neckeroides

I. orthocarpum Brid.

Plagiothecium orthocarpum

I. pendulum Brid.

Helicodontium pendulum

I. pumilum Hook. f. et Wils.

Thamnobryum pumilum

I. serpens Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Neger, 1899; Dusén, 1903a; Paris, 1905a; Seki, 1974).

I. spininervium (Hook.) Wils.

Hypnodendron arcuatum

var. *arcuatum* (Hedw.) Wils.

Hypnodendron arcuatum

I. subsimplex (Hedw.) Brid.

Sematophyllum subsimplex

I. tenuifolium (Hedw.) Brid.

Rhynchostegium tenuifolium

Juratzkaea

J. krausei (C. Muell.) Fleisch. (Fleischer, 1923).

Syn. *Fabroniella krausei* Lor. ex C. Muell. (Lorentz ex Mueller, 1875).

Stereophyllum krausei (C. Muell.) Broth. (Brotherus, 1907).

Distr. Southern South America (Mueller, 1875; Jaeger, 1878; Paris, 1904b; Brotherus, 1907, 1925; Campo, 1921 (as *Atereophyllum krausei* Lor.); Fleischer, 1923; Thériot, 1935).

J. membranosa Fleisch. (Fleischer, 1917) *nom. nud.*

Syn. *Cladomnion membranosum* Dus. (Dusén in Fleischer, 1917) *nom. nud.*

Distr. Southern South America (Fleischer, 1917).

J. seminervis (Schwaegr.) Lor. (Lorentz, 1866).

Syn. *Leskea seminervis* Kunz. ex Schwaegr. (Kunze ex Schwaegrichen, 1829).

Helicodontium seminerve (Schwaegr.) Schwaegr. (Schwaegrichen, 1830) *nom. inval.*

Hypnum semi-nerve (Schwaegr.) C. Muell. (Mueller, 1851a).

Stereophyllum seminerve (Schwaegr.) Mitt. (Mitten, 1869).

Distr. Southern South America (Schwaegrichen, 1829; Mueller, 1851a; Sullivant, 1859 (as *Pylaisea seminervis* Kunz.); Lorentz, 1866; Mitten, 1869; Reichardt, 1870; Jaeger, 1878; Paris, 1904b; Thériot, 1917b, 1918, 1921b, 1928; Campo, 1921; Costes, 1921; Herzog, 1923, 1938, 1952b, 1954; Brotherus, 1924a, c; Hosseus, 1938b; Espinosa B., 1941; Bartram, 1942, 1943, 1952).

var. *arenicola* Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926).

var. *argentinica* Thér. (Thériot in Bauer, 1931).

Distr. Southern South America (Bauer, 1931; Thériot, 1935b; Hosseus, 1936, 1937, 1938b, c, d, 1939; Kühnemann, 1938; Herzog, 1952a).

var. *pilifera* Thér. (Thériot in Verdoorn, 1938).

Distr. Southern South America (Verdoorn, 1938).

Kiaeria

K. starkei (Web. et Mohr) Hag. (Hagen, 1915).

Syn. *Dicranum starkei* Web. et Mohr (Weber and Mohr, 1807).

Oncophorus starkei (Web. et Mohr) Brid. (Bridel, 1826).

Distr. Southern South America (Wilson and Hooker, 1847 (as vars 1 and 2)).

Lamprophyllum hom. illeg.

L. crudum (Hedw.) Lindb.

Pohlia cruda

L. nutans (Hedw.) Lindb.

Pohlia nutans

Lamprophyllum hom. illeg.

L. splendidissimum (Mont.) Broth.

Schimperobryum splendidissimum

Lasia hom. illeg.

L. coronata (Mont.) C. Muell.

Forsstroemia coronata

L. orthotrichoides Raddi

Macrocoma orthotrichoides

L. subcoronata Besch.

Forsstroemia subcoronata

Leiotheca nom. rejic

- L. filiformis* Brid.
L. microstoma (Hook. et Grev.) Brid.
L. obtusifolia (C. Muell.) Hamp.
L. urceolata (Hook.) Brid.
- Macrocoma orthotrichoides*
Macromitrium microstomum
Drummondia obtusifolia
Macromitrium urceolatum

Lembophyllum

- L. auriculatum* (Mont.) Par.
L. cochlearifolium (Schwaegr.) Lindb.
- Acrocladium auriculatum*
Weymouthia cochlearifolia

Lepidopilum

- L. aurescens* C. Muell. (Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938; Welch, 1969).
- L. cristatum* (Hedw.) Hamp.
- Eriopus cristatus*
- L. erectiusculum* (Tayl.) Mitt. (Mitten, 1869).
Syn. *Hookeria erectiuscula* Tayl. (Taylor, 1846).
Distr. Southern South America (Mitten, 1869; Jaeger, 1877; Paris, 1905a; Brotherus, 1925; Welch, 1969).
- L. incurvum* (Hornschr.) C. Muell.
- Hookeriopsis incurva*
- L. patrisiae* (Brid.) Mitt.
- Crossomitrium patrisiae*
- L. plebejum* C. Muell. (Mueller, 1900).
Distr. Southern South America (Welch, 1969).

- L. polytrichoides* (Hedw.) Brid. (Bridel, 1827b).
Syn. *Hypnum polytrichoides* Hedw. (Hedwig, 1801).
Orthotrichum polytrichoides (Hedw.) Brid. (Bridel, 1801a).
Neckera polytrichoides (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Pilotrichum polytrichoides (Hedw.) Brid. (Bridel, 1819).
Daltonia polytrichoides (Hedw.) Arnott (Arnott, 1827).
Hookeria polytrichoides (Hedw.) Spreng. (Sprengel, 1827).
Hypnum rigidum Schwaegr. (Schwaegrichen, 1816).
Pterygophyllum rigidum (Schwaegr.) Brid. (Bridel, 1819).
Hookeria rigida (Schwaegr.) Arnott (Arnott, 1827).

Distr. Southern South America (Paris, 1905a; Brotherus, 1925; Welch, 1969).

- L. scabrisetum* (Schwaegr.) Steere (Steere, 1948).
Syn. *Neckera scabriseta* Schwaegr. (Schwaegrichen, 1816).
Pilotrichum scabrisetum (Schwaegr.) Brid. (Bridel, 1819).
Lepidopilum subnerve Brid. (Bridel, 1827b) hom. illeg.

Distr. Southern South America (Welch, 1969).

- L. splendidissimum* (Mont.) Mitt.
- Schimperobryum splendidissimum*
- L. subnerve* Brid.
- Lepidopilum scabrisetum*
- L. uruguense* Broth. (Brotherus in Krieger, 1904) nom. nud.
- Distr.* Southern South America (Krieger, 1904).

Leptangium hom. illeg.

- L. imbricatum* Mitt.
- Lorentziella imbricata*

Leptobryum

- L. pottiaceum* Dus. (Dusén, 1903a).
Syn. *Wollnya pottiacea* (Dus.) Herz. (Herzog, 1954).

Distr. Southern South America (Dusén, 1903a, 1905d; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1954).

L. pyriforme (Hedw.) Wils. (Wilson, 1855).

Syn. *Webera pyriformis* Hedw. (Hedwig, 1801).

Bryum pyriforme (Hedw.) Gaertn., Meyer and Scherb. (Gärtner and others, 1802) *hom. illeg.*

Bryum piriforme Par. (Paris, 1894) *err.*

Hypnum pyriforme (Hedw.) Web. et Mohr (Weber and Mohr, 1803).

Hypnum piriforme Web. et Mohr ex Par. (Weber and Mohr ex Paris, 1896) *err.*

Distr. Southern South America (Bescherelle, 1889; Dusén, 1903a, 1905d; Krieger, 1904; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938 (as *L. pysiforme* Dus.); Seki, 1974).

var. antarcticum Besch. (Bescherelle, 1889) *nom. nud.*

Syn. *Bryum pyriforme* (Hedw.) Gaertn., Meyer and Scherb. var. *antarcticum* C. Muell. (Mueller, 1885) *nom. nud.*

Distr. Southern South America (Mueller, 1885; Paris, 1905a; Cardot, 1908a; Kühnemann, 1938).

var. fuegianum Par. (Paris, 1896) *nom. nud.*

Syn. *Bryum pyriforme* (Hedw.) Gaertn., Meyer and Scherb. var. *fuegianum* C. Muell. (Mueller, 1889) *nom. nud.*

Distr. Southern South America (Mueller, 1889; Paris, 1905a; Cardot, 1908a).

L. wilsonii (Mitt.) Broth. (Brotherus, 1903).

Syn. *Bryum wilsonii* Mitt. (Mitten, 1851).

Webera wilsonii (Mitt.) Jaeg. (Jaeger, 1875).

Brachymenium wilsonii (Mitt.) Jaeg. (Jaeger, 1875).

Wollnya wilsonii (Mitt.) Herz. (Herzog, 1911).

Distr. Southern South America (Bartram, 1952; Herzog, 1954).

Leptochlaena hom. illeg.

L. chilensis Mont.

Mielichhoferia chilensis

Leptodictyum

L. brachypelmatum (C. Muell.) Broth. (Brotherus, 1925).

Syn. *Amblystegium brachypelmatum* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1903; Brotherus, 1925; Hosseus, 1938c, d; Kühnemann, 1938).

L. campicola (C. Muell.) Broth. (Brotherus, 1925).

Syn. *Hypnum campicola* C. Muell. (Mueller, 1879a).

Amblystegium campicola (C. Muell.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Mueller, 1879a; Paris, 1903; Kurtz, 1904; Brotherus, 1925; Herter, 1933a; Hosseus, 1935a, 1936, 1937; Kühnemann, 1938).

L. kochii (B.S.G.) Warnst. (Warnstorff, 1906).

Syn. *Amblystegium kochii* B.S.G. (Bruch and others, 1853).

Amblystegium riparium (Hedw.) B.S.G. var. *kochii* (B.S.G.) Buyss. (Du Buysson, 1883).

Amblystegium riparium (Hedw.) B.S.G. ssp. *kochii* (B.S.G.) Ren. et Card. (Renauld and Cardot, 1893a).

Hypnum riparium L. ex Hedw. ssp. *kochii* (B.S.G.) Boul. (Boulay, 1884).

Amblystegium trichopodium (Schultz) Hartm. var. *kochii* (B.S.G.) Lindb. (Lindberg, 1872).

Amblystegium trichopodium (Schultz) Hartm. ssp. *kochii* (B.S.G.) Loeske. (Loeske, 1903).

Hypnum trichopodium Schultz var. *kochii* Lindb. ex Par. (Lindberg ex Paris, 1896) *err.*
Leptodictyum trichopodium (Schultz) Warnst. var. *kochii* (B.S.G.) Broth. (Brotherus, 1925).

Distr. Southern South America (Cardot and Brotherus, 1923).

L. riparium (Hedw.) Warnst. (Warnstorff, 1906).

var. *jaffuelii* (Thér.) Thér. (Thériot, 1934b [1935]).

Syn. *Amblystegium jaffuelii* Thér. (Thériot, 1924).

Distr. Southern South America (Thériot, 1924; Brotherus, 1925).

var. *looseri* Thér. (Thériot, 1934b [1935]).

Distr. Southern South America (Thériot, 1934b).

L. trichopodium (Schultz) Warnst. var. *kochii* (B.S.G.) Broth.

Leptodictyum kochii

Leptodon

L. coronatus Mont.

Forsstroemia coronata

L. dillenii Lindb. (Lindberg, 1883).

Distr. Southern South America (Lindberg, 1883).

L. patagonicus C. Muell.

Leptodon smithii

L. smithii (Hedw.) Web. et Mohr (Weber and Mohr, 1803).

Syn. *Hypnum smithii* Hedw. (Hedwig, 1801).

Orthotrichum smithii (Hedw.) Brid. (Bridel, 1801a).

Pilotrichum smithii (Dicks.) P. Beauv. (Palisot de Beauvois, 1805a).

Pterigynandrum smithii (Hedw.) Lam. et Cand. (Lamark and de Candolle, 1805).

Maschalocarpus smithii (Hedw.) Spreng. (Sprengel, 1827).

Neckera smithii (Hedw.) C. Muell. (Mueller, 1850).

Leptodon patagonicus C. Muell. (Mueller in Paris, 1905a) *nom. nud. fide* Thériot, 1929c.

Distr. Juan Fernandez (Johow, 1896; Paris, 1905a; Brotherus, 1924b; Bartram, 1959; Robinson, 1975).

Southern South America (Mitten, 1869; Paris, 1905a; Cardot, 1908a; Fleischer, 1917; Cardot and Brotherus, 1923; Brotherus, 1924a; Thériot, 1924; Roivainen, 1934; Herzog, 1938, 1954, 1960) Kühnemann, 1938;

var. *longirepens* Card. et Thér. (Cardot and Thériot in Thériot, 1924).

Distr. Juan Fernandez (Thériot, 1924).

Leptodontium

L. arachnoideum C. Muell.

Leptodontium capituligerum

L. braunioides C. Muell.

Leptodontium pungens

L. bullockii Thér.

Leptostomum bullockii

L. capituligerum C. Muell. (Mueller, 1879a).

Syn. *Leptodontium arachnoideum* C. Muell. (Mueller, 1879a) *fide* Zander, 1972.

Leptodontium felipponei Broth. (Brotherus in Felippone, 1917) *fide* Zander, 1972.

Leptodontium gracile C. Muell. ex Britt. var. *gemmaescens* Bartr. (Bartram, 1936) *fide* Zander, 1972.

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905a; Felippone, 1917; Brotherus, 1924a; Herter, 1933a; Hosseus, 1935b, 1937, 1938c, d, 1939; Bartram, 1936; Kühnemann, 1938; Zander, 1972; Seki, 1974).

L. chiloense C. Muell. (Mueller in Kindberg, 1889) *nom. nud.*

Distr. No locality given, see *Index muscorum*, 3, p. 233.

L. felipponei Broth.

Leptodontium capituligerum

L. fernandezianum Broth.

Leptodontium longicaule* var. *microruncinatum

L. filicaule Dus. (Dusén in Paris, 1905a) *nom. nud.*

Distr. Southern South America (Paris, 1905a).

L. filicola Herz. (Herzog, 1916a).

Distr. Southern South America (Zander, 1972).

L. gracile C. Muell. ex Britt. var. *gemmascens* Bartr.

Leptodontium capituligerum

L. gracile C. Muell. ex Britt. var. *gracilescens* Bartr. *nom. nud.*

Distr. Southern South America (Bartram, 1936; Hosseus, 1938c, 1939).

L. kurtzii Broth. (Brotherus *in Kurtz*, 1904) *nom. nud.*

Distr. Southern South America (Kurtz, 1904; Hosseus, 1936, 1937; Kühnemann, 1938).

L. longicaule Mitt. var. *microruncinatum* (Dus.) Zander (Zander, 1972).

Syn. *Barbula micro-runcinata* Dus. (Dusén, 1903a) *nom. nud. fide* Zander, 1972.

Leptodontium microruncinatum Dus. (Dusén, 1906) *fide* Zander, 1972.

Syntrichia micro-runcinata Dus. (Dusén *in Paris*, 1906) *nom. nud.*

Tortula micro-runcinata Par. (Paris, 1906) *nom. nud.*

Leptodontium fernandezianum Broth. (Brotherus, 1924b) *fide* Zander, 1972.

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Zander, 1972; Robinson, 1975).

Southern South America (Dusén, 1903a, 1906; Paris, 1906; Cardot, 1908a; Brotherus, 1924a; Thériot, 1925a; Herzog, 1938, 1954; Zander, 1972).

L. menziesii R. Brown ex Thér.

Leptostomum menziesii

L. microruncinatum Dus.

Leptodontium longicaule* var. *microruncinatum

L. orthotrichoides (C. Muell.) Par. (Paris, 1900a).

Syn. *Trichostomum orthotrichoides* C. Muell. (Mueller, 1874).

Zygodon orthotrichoides (C. Muell.) Broth. (Brotherus, 1918).

Zygodon palmarum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1906; Brotherus, 1918, 1925; Kühnemann, 1938).

L. pungens (Mitt.) Kindb. (Kindberg, 1888).

Syn. *Didymodon pungens* Mitt. (Mitten, 1863).

Trichostomum pungens (Mitt.) Par. (Paris, 1898).

Leptodontium braunioides C. Muell. (Mueller, 1879a) *fide* Zander, 1972.

Leptodontium skottsbergii Bartr. (Bartram, 1959) *fide* Zander, 1972.

Distr. Juan Fernandez (Bartram, 1959; Zander, 1972; Robinson, 1975).

Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938; Zander, 1972).

L. quennoae C. Muell.

Leptodontium viticulosoides* var. *viticulosoides

L. racomitrioides Lor. et C. Muell.

Leptodontium viticulosoides* var. *viticulosoides

L. rhacomitrioides Lor. et C. Muell.

Leptodontium viticulosoides* var. *viticulosoides

L. skottsbergii Bartr.

Leptodontium pungens

L. splachnoides (Hook. et Arnott) Thér.

Leptostomum splachnoides

L. viticulosoides (P. Beauv.) Wijk et Marg. (van der Wijk and Margadant, 1960a).var. *viticulosoides*

Syn. *Neckera viticulosoides* P. Beauv. (Palisot de Beauvois, 1805a).

Leptodontium quennoae C. Muell. (Muller, 1879a) *fide* Zander, 1972.

Leptodontium rhacomitroides Lor. et C. Muell. (Lorentz and Mueller *in* Mueller, 1879a).

Leptodontium racomitroides Lor. et C. Muell. (Lorentz and Mueller *in* Kindberg, 1889) *err.*

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938; Zander, 1972).

L. zygondontoides C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Zander, 1972).

Leptohymenium*L. squarrosum* Hamp.***Erythrodontium squarrosum******Leptopterigynandrum****L. austro-alpinum* C. Muell. (Mueller, 1879a).Syn. *Heterocladium austro-alpinum* (C. Muell.) Broth. (Brotherus, 1907).*Pterigynandrum austro-alpinum* C. Muell. (Mueller in Fleischer, 1912) nom. nud.

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1907, 1925; Fleischer, 1912, 1922; Kühnemann, 1938).

Leptostomum*L. bullockii* (Thér.) Thér. (Thériot in Andrews, 1951 [1952]).Syn. *Leptodontium bullockii* Thér. (Thériot, 1929b).

Distr. Southern South America (Thériot, 1929b, 1930; Andrews, 1951).

L. inclinans R. Brown (Brown, 1811).

Distr. Southern South America (Montagne, 1850).

L. menziesii R. Brown (Brown, 1811).Syn. *Gymnostomum menziesii* (R. Brown) Hook. (Hooker, 1820).*Leptodontium menziesii* R. Brown ex Thér. (R. Brown ex Thériot, 1929b) err.

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Brown, 1811; Schwaegrichen, 1816, 1823; Hooker, 1820; Montagne, 1845b, 1850; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Sullivant, 1859; Mitten, 1869; Jaeger, 1875; Bescherelle, 1889; Hariot, 1891; Neger, 1899; Cardot, 1900; 1901, 1908a, 1913a; Dusén, 1903a; Paris, 1905a; Spegazzini, 1922; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1954; Brotherus, 1924a; Thériot, 1929b; Roivainen, 1934; Kühnemann, 1938; Andrews, 1951; Bartram, 1952).

L. splachnoides Hook. et Arnott (Hooker and Arnott, 1832).Syn. *Leptodontium splachnoides* (Hook. et Arnott) Thér. (Thériot, 1929a) err.*Leptostomum sulcinerve* Kunz. (Kunze in Mueller, 1848a) nom. nud.

Distr. Southern South America (Hooker and Arnott, 1832; Schwaegrichen, 1842; Montagne, 1845b, 1850; Mueller, 1848a; Lorentz, 1866; Mitten, 1869; Jaeger, 1875; Paris, 1905a; Campo, 1921; Brotherus, 1924a; Thériot, 1929b, 1935a; Andrews, 1951; Herzog, 1954; Seki, 1974).

L. sulcinerve Kunz.***Leptostomum splachnoides******Leptotheca****L. gaudichaudii* Schwaegr. (Schwaegrichen, 1824).Syn. *Bryum gaudichaudii* (Schwaegr.) Spreng. (Sprengel, 1827).*Aulacomnium gaudichaudii* (Schwaegr.) Mitt. (Mitten, 1856).*Brachymenium ovatum* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).*Hymenodon ovatus* (Hook. f. et Wils.) C. Muell. (Mueller, 1851a).*Leptotheca sphaerocarpa* C. Muell. (Mueller, 1885).

Distr. Juan Fernandez (Robinson, 1975).

Southern South America (Mueller, 1848a, 1885; Jaeger, 1875; Cardot, 1905a, 1908a; Paris, 1905b; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1939; Bartram, 1952; Seki, 1974).

Falkland Islands (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1851a; Mitten, 1869; Jaeger, 1875; Cardot, 1905a, 1908a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

South Georgia (Clarke, 1973a).

L. sphaerocarpa C. Muell.***Leptotheca gaudichaudii****Leptotrichum hom. illeg.**L. affine* C. Muell.***Ditrichum difficile***

<i>L. australe</i> Mitt.	<i>Ditrichum strictum</i>
<i>L. capillaceum</i> (Hedw.) Mitt.	<i>Distichium capillaceum</i>
<i>L. cylindricarpum</i> C. Muell.	<i>Ditrichum cylindricarpum</i>
<i>L. difficile</i> (Dub.) C. Muell.	<i>Ditrichum difficile</i>
<i>L. elongatum</i> (Hook. f. et Wils.) Jaeg.	<i>Ditrichum cylindricarpum</i>
<i>L. flexifolium</i> Mitt.	<i>Ditrichum difficile</i>
<i>L. glaciale</i> C. Muell.	<i>Sarconeurum glaciale</i>
<i>L. homomallum</i> (Hedw.) Hamp.	<i>Ditrichum heteromallum</i>
<i>L. hookeri</i> C. Muell.	<i>Ditrichum hookeri</i>
<i>L. hyalinum</i> Mitt.	<i>Ditrichum hyalinum</i>
<i>L. jamesonii</i> Mitt.	<i>Anisothecium vaginatum</i>
<i>L. kunzeanum</i> C. Muell.	<i>Microdus kunzeanus</i>
<i>L. laxifolium</i> (Hook. f. et Wils.) C. Muell.	<i>Ditrichum difficile</i>
<i>L. longisetum</i> Lor.	<i>Ditrichum longisetum</i>
<i>L. lorifolium</i> C. Muell. (Mueller, 1901a [1900]) <i>nom. nud.</i>	
Syn. <i>Leptotrichum sericeum</i> Lor. (Lorentz in Mueller, 1901a [1900]) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Mueller, 1901a).	
<i>L. montagnei</i> C. Muell.	<i>Ditrichum conicum</i>
<i>L. praealtum</i> Mitt.	<i>Ditrichum praealtum</i>
<i>L. rufescens</i> Hamp.	<i>Ditrichum rufescens</i>
<i>L. sericeum</i> Lor.	<i>Leptotrichum lorifolium</i>
<i>Lepyrodon</i>	
<i>L. glaucus</i> (C. Muell.) Kindb.	<i>Lepyrodon hexastichus</i>
<i>L. gunckelii</i> Thér. (Thériot, 1934b [1935]).	
<i>Distr.</i> Southern South America (Thériot, 1934b).	
<i>L. hexastichus</i> (Mont.) Wijk et Marg. (van der Wijk and Margadant, 1960b).	
Syn. <i>Leucodon hexastichus</i> Mont. (Montagne, 1845a).	
<i>Neckera implexa</i> C. Muell. (Mueller, 1850) <i>nom. illeg.</i>	
<i>Leucodon implexus</i> Kunz. ex Hook. f. (Kunze ex Hooker, 1867) <i>nom. illeg.</i>	
<i>Lepyrodon implexus</i> Kindb. (Kindberg, 1888) <i>nom. illeg.</i>	
<i>Neckera glauca</i> C. Muell. (Mueller, 1859).	
<i>Leucodon glaucus</i> (C. Muell.) Jaeg. (Jaeger, 1877).	
<i>Lepyrodon glaucus</i> (C. Muell.) Kindb. (Kindberg, 1888).	
<i>Leucodon kunzeanus</i> C. Muell. (Mueller, 1844b [1845]).	
<i>Distr.</i> Juan Fernandez (Brotherus, 1924b, 1925; Thériot, 1927b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).	
Southern South America (Mueller, 1844b, 1850, 1859; Montagne, 1845a, 1850, 1856; Jaeger, 1877; Dusén, 1903a; Paris, 1905a; Cardot, 1908a; Fleischer, 1917; Herzog, 1923, 1938, 1939, 1940, 1954, 1957, 1960; Thériot, 1924; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938; Bartram, 1952; Seki, 1974).	
<i>L. implexus</i> Kindb.	<i>Lepyrodon hexastichus</i>
var. <i>reptans</i> Reim. (Reimers, 1926).	
<i>Distr.</i> Southern South America (Reimers, 1926).	
<i>L. lagurus</i> (Hook.) Mitt. (Mitten, 1869).	
Syn. <i>Leucodon lagurus</i> Hook. (Hooker, 1819).	
<i>Hypnum lagurum</i> Menz. (Menzies in Hooker, 1819) <i>nom. nud.</i>	
<i>Neckera lagura</i> (Hook.) C. Muell. (Mueller, 1850).	
<i>Stereodon lagurus</i> (Hook.) Mitt. (Mitten, 1860).	
<i>Hylocomium lagurus</i> (Hook.) Kindb. (Kindberg, 1885).	

Distr. Southern South America (Hooker, 1819; Schwaegrichen, 1824; Bridel, 1827b; Montagne, 1845b, 1850; Wilson and Hooker, 1847; Mueller, 1850, 1885, 1889; Sullivant, 1859; Mitten, 1869; Jaeger, 1877; Bescherelle, 1889; Neger, 1899; Cardot, 1900, 1901, 1905a, 1908a, 1913a; Dusén, 1903a, 1905d (as *Leptodon lagurus*); Paris, 1905a; Fleischer, 1917; Campo, 1921; Cardot and Brotherus, 1923; Herzog, 1923, 1939; Brotherus, 1925; Roivainen, 1934; Thériot, 1934b; Kühnemann, 1938; Bartram, 1952; Seki, 1974).
South Georgia (Cardot, 1906a, 1908a; Steere, 1961b).

var. *laxior* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923).

***L. parvulus* Mitt. (Mitten, 1869).**

Syn. *Leucodon parvulus* (Mitt.) Jaeg. (Jaeger, 1877).

Distr. Juan Fernandez (Mitten, 1885; Johow, 1896; Paris, 1905a; Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).
Southern South America (Mitten, 1869; Jaeger, 1877; Fleischer, 1917; Brotherus, 1925; Herzog, 1939, 1954).

L. suborthostichus (C. Muell.) Hamp.

Lepyrodon tomentosus

***L. tomentosus* (Hook.) Mitt. (Mitten, 1869).**

Syn. *Leucodon tomentosus* Hook. (Hooker, 1818).

Neckera tomentosa (Hook.) C. Muell. (Mueller, 1850).

Neckera suborthosticha C. Muell. (Mueller, 1850).

Leucodon suborthostichus (C. Muell.) De Not. (De Notaris, 1859).

Lepyrodon suborthostichus (C. Muell.) Hamp. (Hampe, 1862b).

Distr. Juan Fernandez (Brotherus, 1924, 1925; Bartram, 1959; Robinson, 1975).

Southern South America (Lorentz, 1866; Mitten, 1869; Jaeger, 1877; Kurtz, 1904; Paris, 1905a; Campo, 1921; Cardot and Brotherus, 1923; Brotherus, 1925; Roivainen, 1934; Thériot, 1935a; Hosseus, 1936, 1937; Herzog, 1938, 1957, 1960; Kühnemann, 1938).

var. *minor* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923).

var. *patagonicus* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Seki, 1974).

Lescureaea

***L. chilensis* Herz. (Herzog, 1954).**

Distr. Southern South America (Herzog, 1954).

Leskea

<i>L. albicans</i> Hedw.	<i>Cyclodictyon albicans</i>
<i>L. caespitosa</i> Hedw.	<i>Sematophyllum caespitosum</i>
<i>L. concinna</i> Hook.	<i>Lopidium concinnum</i>
<i>L. cristata</i> Hedw.	<i>Eriopus cristatus</i>
<i>L. distans</i> Mont.	<i>Glyphothecium gracile</i>
<i>L. fuegiana</i> Besch.	<i>Hygroamblystegium fuegianum</i>
<i>L. gayana</i> Mont.	<i>Ptychomnion ptychocarpum</i>
<i>L. glabella</i> Hedw.	<i>Homalia glabella</i>
<i>L. inclinata</i> (Brid.) Web. et Mohr	<i>Bryum inclinatum</i>
<i>L. julacea</i> Schwaegr.	<i>Myurella julacea</i>
<i>L. leucostega</i> Brid.	<i>Stereophyllum leucostegum</i>
<i>L. microphylla</i> (Hedw.) Sw.	<i>Haplocladium microphyllum</i>
<i>L. mollis</i> Hedw.	<i>Weymouthia mollis</i>

- L. nitida* Hook. f. et Wils. *Catagonium nitidum*
L. ornithopodioides (Brid.) Brid. *Racopilum tomentosum*
- L. pellicula* (C. Muell.) Broth. (Brotherus, 1907).
 Syn. *Anomodon pellicula* C. Muell. (Mueller in Kindberg, 1891) *nom. nud.*
Anomodon pellicula C. Muell. (Mueller, 1897a).
 Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1903; Brotherus, 1907, 1918,
 1925; Kühnemann, 1938).
- L. pellucida* Hook. *Dimerodontium pellucidum*
L. pseudo-leskeoides Broth. (Brotherus in Kurtz, 1904) *nom. nud.*
 Distr. Southern South America (Kurtz, 1904; Hosseus, 1936, 1937; Kühnemann, 1938).
- L. remotifolia* C. Muell. *Meteoriopsis remotifolia*
L. rutilans Brid. *Vesicularia vesicularis*
L. seminervis Kunz. ex Schwaegr. *Juratzkaea seminervis*
L. varia Hedw. *Amblystegium varium*
- Leskeadelphus**
L. angustifolius Herz. (Herzog, 1952a).
 Distr. Southern South America (Herzog, 1952a).
- L. catenulatulus* (C. Muell.) Herz. (Herzog, 1916a).
 Syn. *Pseudoleskea catenulatula* C. Muell. (Mueller, 1897b).
 Distr. Southern South America (Mueller, 1897a, b; Paris, 1905b; Brotherus, 1925;
 Kühnemann, 1938).
- Leucobryum**
L. argentinicum C. Muell. (Mueller, 1879a).
 Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Kühnemann, 1938).
- L. crispum* C. Muell. (Mueller, 1848a).
 Distr. Southern South America (Hosseus, 1940).
- L. giganteum* C. Muell. (Mueller, 1848a).
 Syn. *Leucobryum longifolium* Hamp. (Hampe, 1847) *hom. illeg.*
 Distr. Southern South America (Jaeger, 1872, 1873).
L. longifolium Hamp. *Leucobryum giganteum*
- L. sordidum* Aongstr. (Ångström, 1876a).
 Distr. Southern South America (Hosseus, 1940).
- Leucodon**
L. coronatus (Mont.) Mitt. *Forsstroemia coronata*
L. curtipendulus (Hedw.) T. Jens. *Antitrichia curtipendula*
- L. felipponei* Thér. (Thériot, 1939 [1941]).
 Distr. Southern South America (Thériot, 1939).
- L. glaucus* (C. Muell.) Jaeg. *Lepyrodon hexastichus*
L. gracilis Hamp. *Glyphothecium gracile*
L. hexastichus Mont. *Lepyrodon hexastichus*
L. implexus Kunz. ex Hook. f. *Lepyrodon hexastichus*
- L. julaceus* (Hedw.) Sull. (Sullivant, 1846).
 Syn. *Pterigynandrum julaceum* Hedw. (Hedwig, 1801).

Maschalocarpus julaceus (Hedw.) Spreng. (Sprengel, 1827).
Grimmia julacea (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Pterogonium julaceum (Hedw.) Schwaegr. (Schwaegrichen, 1811).
Hypnum julaceum (Hedw.) Schleich. (Schleicher, 1821) *hom. illeg.*
Fissidens julaceus (Hedw.) Lindb. (Lindberg, 1883).
Leucodontiella julacea (Hedw.) Nog. (Noguchi, 1947).

Distr. Southern South America (Hornschorch, 1820b; Bridel, 1827b; Montagne, 1850).

<i>L. kunzeanus</i> C. Muell.	<i>Lepyrodon hexastichus</i>
<i>L. lagurus</i> Hook.	<i>Lepyrodon lagurus</i>
<i>L. parvulus</i> (Mitt.) Jaeg.	<i>Lepyrodon parvulus</i>
<i>L. suborthostichus</i> (C. Muell.) De Not.	<i>Lepyrodon tomentosus</i>
<i>L. tomentosus</i> Hook.	<i>Lepyrodon tomentosus</i>

Leucodontella

<i>L. julacea</i> (Hedw.) Nog.	<i>Leucodon julaceus</i>
--------------------------------	---------------------------------

Leucoloma

<i>L. alboalare</i> Dus.	<i>Dicranoloma hariotii</i>
<i>L. australe</i> (Besch.) Broth.	<i>Dicranoloma australe</i>
<i>L. billardieri</i> (Brid.) Broth.	<i>Dicranoloma billardieri</i>
<i>L. capillare</i> Dus.	<i>Dicranoloma capillare</i>
<i>L. capillifolium</i> Broth.	<i>Dicranoloma menziesii</i>
<i>L. dusenii</i> Broth.	<i>Dicranoloma dusenii</i>
<i>L. fuegianum</i> Dus.	<i>Dicranoloma hariotii</i>
var. <i>laxum</i> Dus.	<i>Dicranoloma hariotii</i>
<i>L. grandialare</i> Dus.	<i>Dicranoloma robustum</i>
<i>L. hariotii</i> (C. Muell.) Broth.	<i>Dicranoloma hariotii</i>
<i>L. imponens</i> (Mont.) Dus.	<i>Dicranoloma imponens</i>

***L. longifolium* (Brid.) Wijk et Marg. (van der Wijk and Margadant, 1960b).**

Syn. *Trichostomum longifolium* Brid. (Bridel, 1806).
Didymodon longifolius (Brid.) Hook. f. et Wils. *hom. illeg.*

Distr. Juan Fernandez (Montagne, 1835, 1850; Johow, 1896).
Southern South America (Wilson and Hooker, 1847).

<i>L. menziesii</i> (Hook. f. et Wils.) Broth.	<i>Dicranoloma menziesii</i>
<i>L. muelleri</i> Dus.	<i>Dicranoloma muelleri</i>
var. <i>strictifolium</i> Dus.	<i>Dicranoloma robustum</i> var. <i>strictifolium</i>
<i>L. nigricaula</i> (Aongstr.) Broth.	<i>Dicranoloma nigricaula</i>
var. <i>flexuosulum</i> Dus.	<i>Dicranoloma nigricaula</i> var. <i>flexuosulum</i>
var. <i>gracile</i> Dus.	<i>Dicranoloma capillare</i>
<i>L. perremotifolium</i> Dus.	<i>Dicranoloma perremotifolium</i>
var. <i>fragile</i> Dus.	<i>Dicranoloma perremotifolium</i> var. <i>fragile</i>
<i>L. peruncinatum</i> Dus.	<i>Dicranoloma nigricaula</i>
var. <i>gracile</i> Dus.	<i>Dicranoloma capillare</i>
<i>L. pungens</i> (Hook. f. et Wils.) Broth.	<i>Dicranoloma robustum</i>
<i>L. robustum</i> (Hook. f. et Wils.) Broth.	<i>Dicranoloma robustum</i>
var. <i>flexuosum</i> Dus.	<i>Dicranoloma australe</i>
var. <i>lagunicola</i> Dus.	<i>Dicranoloma imponens</i>
<i>L. setosum</i> (Hook. f. et Wils.) Broth.	<i>Dicranoloma robustum</i> var. <i>setosum</i>

- L. suberectum* (Hamp.) Broth. *Dicranoloma menziesii*
- L. subnigraule* Dus. (Dusén *in Paris*, 1905a) *nom. nud.*
- Distr.* Southern South America (Paris, 1905a).
- L. syrrhodictyon* (C. Muell.) Broth. *Dicranum syrrhopodontoides*
- Leucomium**
- L. crinitifolium* (C. Muell.) Mitt. (Mitten, 1869).
- Syn.* *Hypnum crinitifolium* C. Muell. (Mueller, 1851a).
- Distr.* Southern South America (Mueller, 1851a; Mitten, 1869; Jaeger, 1878; Paris, 1905a; Brotherus, 1925).
- Limbella* *nom. nud.*
- L. confluens* (C. Muell.) C. Muell. *Sciaromium confluens*
- L. conspissata* (Hook. f. et Wils.) C. Muell. *Sciaromium conspissatum*
- L. conspissatula* C. Muell. *Sciaromium conspissatulum*
- L. drepanophyllopsis* C. Muell. *Sciaromium drepanophyllopsis*
- L. krauseana* C. Muell. *Sciaromium krauseanum*
- L. negeri* Broth. (Brotherus *in Neger*, 1899) *nom. nud.*
- Distr.* Southern South America (Neger, 1899).
- L. pachyloma* (Mont.) C. Muell. *Sciaromium pachyloma*
- L. pachylomata* C. Muell. *Sciaromium pachylomatum*
- L. platylomata* C. Muell. *Sciaromium platylomatum*
- Limnobryum**
- L. palustre* (Hedw.) Rabenh. *Aulacomnium palustre*
- Limprichtia**
- L. revolvens* (Sw.) Loesk. *Drepanocladus revolvens*
- Lindigia**
- L. lorentzii* C. Muell. (Mueller, 1879a).
- Distr.* Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938; Herzog, 1952a).
- Lophiodon* *nom. rejic.*
- L. strictus* Hook. f. et Wils. *Ditrichum strictum*
- Lopidium**
- L. concinnum* (Hook.) Wils. (Wilson, 1854).
- Syn.* *Leskea concinna* Hook. (Hooker, 1818).
- Hookeria concinna* (Hook.) Hook. et Grev. (Hooker and Greville, 1825a).
- Hypopterygium concinnum* (Hook.) Brid. (Bridel, 1827b).
- Lopidium pallens* Hook. f. et Wils. (Hooker and Wilson *in Wilson*, 1854).
- Hypopterygium pallens* (Hook. f. et Wils.) Mitt. (Mitten, 1856).
- Hypopterygium plumarium* Mitt. (Mitten, 1869).
- Lopidium plumarium* (Mitt.) Hamp. (Hampe, 1879).
- Hypopterygium pallens* (Hook. f. et Wils.) Mitt. ssp. *plumarium* (Mitt.) Kindb. (Kindberg, 1901).
- Hypopterygium flexisetum* Hamp. (Hampe *in Lorentz*, 1866) *nom. nud.*
- Lopidium flexisetum* Fleisch. (Fleischer, 1922) *nom. nud.*
- Distr.* Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Matteri, 1973b; Robinson, 1975).
- Southern South America (Montagne, 1850; Lorentz, 1866; Mitten, 1869; Jaeger, 1876; Neger, 1899; Kindberg, 1901; Paris, 1905a; Cardot, 1908a; Campo, 1915; Fleischer,

1922; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1954, 1960; Brotherus, 1925; Reimers, 1926; Thériot, 1934a; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

<i>L. flexisetum</i> Fleisch.	<i>Lopidium concinnum</i>
<i>L. pallens</i> Hook. f. et Wils.	<i>Lopidium concinnum</i>
<i>L. plumarium</i> (Mitt.) Hamp.	<i>Lopidium concinnum</i>

Lorentziella

<i>L. gibertii</i> C. Muell.	<i>Lorentziella imbricata</i>
------------------------------	-------------------------------

L. glauca C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Jaeger and Sauerbeck, 1879b; Paris, 1905a; Roth, 1911; Brotherus, 1924a; Kühnemann, 1938).

<i>L. globiceps</i> C. Muell.	<i>Lorentziella imbricata</i>
-------------------------------	-------------------------------

L. imbricata (Mitt.) Broth. (Brotherus, 1903).

Syn. *Leptangium imbricatum* Mitt. (Mitten, 1869).
Gigaspernum imbricatum (Mitt.) Kindb. (Kindberg, 1888).
Lorentziella globiceps C. Muell. (Mueller, 1879a).
Lorentziella paraguensis Besch. (Bescherelle, 1877).
Lorentziella gibertii C. Muell. (Mueller, 1888).

Distr. Southern South America (Mitten, 1869; Gibert, 1873; Jaeger and Sauerbeck, 1879b; Mueller, 1879a, 1888; Paris, 1896, 1904b, 1905a; Brotherus, 1903, 1924a; Kurtz, 1904; Roth, 1911, 1913; Felippone, 1912; Herter, 1928, 1933a, b, 1939a; Hosseus, 1935b, 1937, 1938a, d; Kühnemann, 1938).

<i>L. paraguensis</i> Besch.	<i>Lorentziella imbricata</i>
------------------------------	-------------------------------

L. uruguensis C. Muell. (Mueller in Bescherelle, 1877).

Distr. Southern South America (Jaeger and Sauerbeck, 1879b; Paris, 1905a).

Macrocoma

<i>M. filiforme</i> (Hook. et Grev.) Grout	<i>Macrocoma orthotrichoides</i>
--	----------------------------------

M. hymenostomum (Mont.) Grout (Grout, 1946).

Syn. *Macromitrium hymenostomum* Mont. (Montagne, 1845a).
Macromitrium hariotii Besch. ex C. Muell. (Bescherelle ex Mueller, 1885).

Distr. Juan Fernandez (Brotherus, 1924b, 1925).
Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849, 1885; Mitten, 1869; Jaeger, 1874; Bescherelle, 1885a, 1889; Paris, 1905a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

M. orthotrichoides (Raddi) Wijk et Marg. (van der Wijk and Margadant, 1962).

Syn. *Lasia orthotrichoides* Raddi (Raddi, 1823).
Chaetephora orthotrichoides (Raddi) Brid. (Bridel, 1827b).
Orthotrichum filiforme Hook. et Grev. (Hooker and Greville, 1824a) nom. illeg.
Macromitrium filiforme Schwaegr. (Schwaegrichen, 1826) nom. illeg.
Leiotheca filiformis Brid. (Bridel, 1826) nom. illeg.
Macrocoma filiforme (Hook. et Grev.) Grout (Grout, 1944) nom. illeg.
Macromitrium quinquefarium Hornsch. (Hornschuch, 1840).

Distr. Southern South America (Schwaegrichen, 1826; Bridel, 1827a; Montagne, 1839c, 1850; Hornschuch, 1840; Mueller, 1849; Mitten, 1869; Paris, 1905a; Felippone, 1912; Brotherus, 1925; Herter, 1933a; Kühnemann, 1938; Herzog, 1952a).

M. sullivantii (C. Muell.) Grout (Grout, 1944).

Syn. *Macromitrium sullivantii* C. Muell. (Mueller, 1862).
Macromitrium paraphysatum Mitt. (Mitten, 1869) fide Vitt, 1973.
Macromitrium paraphysatum Mitt. var. *chilense* Thér. (Thériot, 1921a) fide Vitt, 1973.

Distr. Juan Fernandez (Robinson, 1975).
 Southern South America (Mitten, 1869; Jaeger, 1874; Paris, 1905a; Thériot, 1921a;
 Brotherus, 1925).

Macromitrium

M. aciculare Brid.

see ***Macromitrium filicaule***

M. anacamptophyllum C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Kühnemann, 1938).

M. angulicaule C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1925;
 Kühnemann, 1938).

M. apiculatum (Hook.) Schwaegr.

Groutiella apiculata

M. bifasciculare C. Muell. ex Dus. (Mueller ex Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Kühnemann, 1938).

M. bifasciculatum C. Muell. (Mueller, 1898a).

Distr. Southern South America (Mueller, 1898a; Paris, 1900a, 1905a; Cardot, 1908a; Cardot
 and Brotherus, 1923; Brotherus, 1925).

M. campoanum Thér. (Thériot in Herzog, 1938).

Distr. Southern South America (Herzog, 1938).

M. coriaceum Bartr.

Macromitrium tenax

M. crassiusculum Lor. (Lorentz, 1866).

Distr. Southern South America (Lorentz, 1866; Jaeger, 1874; Paris, 1905a; Brotherus, 1925;
 Thériot, 1925a; Herzog, 1938, 1960).

M. didymodon Schwaegr.

Macromitrium richardii

M. fernandezianum Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson,
 1975).

M. filicaule C. Muell. (Mueller, 1849).

Distr. Juan Fernandez (? Montagne, 1835 (as *M. aciculare* Brid.)).
 Southern South America (Paris, 1905a).

M. filiforme Schwaegr.

Macrocoma orthotrichoides

M. fimbriatum (P. Beauv.) Schwaeg. (Schwaegrichen, 1823).

Syn. *Orthotrichum fimbriatum* P. Beauv. (Palisot de Beauvois, 1805a).

Distr. Juan Fernandez (Montagne, 1850; Johow, 1896).

M. gracillimum (Besch.) Broth. (Brotherus, 1902).

Syn. *Schlotheimia gracillima* Besch. (Bescherelle, 1885a).

Distr. Southern South America (Bescherelle, 1885a, 1889; Brotherus, 1902, 1925; Paris,
 1905b; Cardot, 1908a; Herzog, 1926, 1954; Kühnemann, 1938; Seki, 1974).

M. hariotii Besch. ex C. Muell.

Macrocoma hymenostomum

M. hymenostomum Mont.

Macrocoma hymenostomum

M. jamesonii (Arnott) C. Muell.

Schlotheimia jamesonii

M. krausei Lor. (Lorentz, 1866).

Distr. Southern South America (Lorentz, 1866; Jaeger, 1874; Mueller, 1885; Cardot, 1908a;
 Brotherus, 1925; Herzog, 1938; Bartram, 1952).

M. linearifolium C. Muell.

Schlotheimia linearifolia

M. longipes (Hook.) Schwaegr. (Schwaegrichen, 1824).

Syn. *Orthotrichum longipes* Hook. (Hooker, 1818).

Distr. Southern South America (Wilson and Hooker, 1847).

M. longirostre (Hook.) Schwaegr. (Schwaegrichen, 1823).

Syn. *Orthotrichum longirostre* Hook. (Hooker, 1818).

Distr. Southern South America (Mitten, 1869; Paris, 1905a; Brotherus, 1925).

M. macrocomoides C. Muell. (Mueller, 1898a).

Distr. Southern South America (Mueller, 1898a; Paris, 1900a, 1905a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

M. masafuerae Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Bartram, 1959; Robinson, 1975).

M. microcarpum C. Muell. (Mueller, 1849).

Distr. Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1874; Paris, 1905a; Brotherus, 1925; Thériot, 1935a; Herzog, 1939, 1960; Bartram, 1952).

M. microstomum (Hook. et Grev.) Schwaegr. (Schwaegrichen, 1827a).

Syn. *Orthotrichum microstomum* Hook. et Grev. (Hooker and Greville, 1824a).

Leiotheca microstoma (Hook. et Grev.) Brid. (Bridel, 1826).

Macromitrium saxatile Mitt. (Mitten, 1869) *fide* Robinson, 1975.

Distr. Juan Fernandez (Montagne, 1835; Mitten, 1869, 1885; Jaeger, 1874; Johow, 1896; Paris, 1905a; Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

M. ottonis (Schwaegr.) C. Muell.

Schlottheimia torquata

M. papillosum Thér. (Thériot *in* Herzog, 1938).

Distr. Southern South America (Herzog, 1938, 1954).

M. paraphysatum Mitt.

Macrocoma sullivantii

var. *chilense* Thér.

Macrocoma sullivantii

M. pertriste C. Muell. (Mueller, 1898a).

Distr. Southern South America (Mueller, 1898a; Paris, 1900a, 1905a; Thériot, 1921a, 1934b; Brotherus, 1925).

M. poeppigii Dub.

Zygodon pentastichus

M. pseudoserrulatum Bartr. (Bartram, 1965).

Distr. Southern South America (Bartram, 1965).

M. quinquefarium Hornsch.

Macrocoma orthotrichoides

M. richardii Schwaegr. (Schwaegrichen, 1826).

Syn. *Macromitrium didymodon* Schwaegr. (Schwaegrichen, 1827a).

Distr. Southern South America (Krieger, 1904).

M. saddleanum Besch. ex C. Muell. (Bescherelle ex Mueller, 1885).

Distr. Southern South America (Bescherelle, 1885a, 1889; Mueller, 1885; Paris, 1905a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Herzog, 1938, 1954; Kühnemann, 1938).

M. saxatile Mitt.

Macromitrium microstomum

M. spurio-crispulum Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1905a).

M. subpiliferum Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a).

M. sullivantii C. Muell.

Macrocoma sullivantii

M. tenax C. Muell. (Mueller, 1884 [1883]).

Syn. *Macromitrium coriaceum* Bartr. (Bartram, 1946) *fide* Seki, 1974.

Distr. Southern South America (Mueller, 1884, 1885, 1889; Dusén, 1903a; Paris, 1905a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Roivainen, 1934; Kühnemann, 1938; Bartram, 1946; Seki, 1974).

var. theriotii Card. (Cardot, 1908a).

Syn. *Macromitrium theriotii* Card. (Cardot, 1908a) *nom. inval.*

Distr. Southern South America (Cardot, 1908a).

M. theriotii Card.

Macromitrium tenax* var. *theriotii

M. tomentosum Hornsch.

Groutiella tomentosa

M. torquatum (Hedw.) C. Muell.

Schlotheimia torquata

M. trichomitrium (Schwaegr.) C. Muell.

Schlotheimia trichomitria

M. urceolatum (Hook.) Brid. (Bridel, 1826).

Syn. *Orthotrichum urceolatum* Hook. (Hooker, 1819).

Leiotheca urceolata (Hook.) Brid. (Bridel, 1826).

Distr. Juan Fernandez (Montagne, 1835; Johow, 1896).

Southern South America (Montagne, 1850).

***Macromitrium* sp.**

Syn. *Zygodon hatcheri* Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

Maschalocarpus *nom. illeg.*

M. julaceus (Hedw.) Spreng.

Leucodon julaceus

M. smithii (Hedw.) Spreng.

Leptodon smithii

Meesia

M. alpina Funck ex Bruch

Meesia uliginosa* var. *alpina

M. austro-georgica C. Muell.

Philonotis acicularis

M. hymenostoma Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

M. patagonica Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Brotherus, 1924a; Kabiersch, 1937; Kühnemann, 1938).

M. stygia (Sw.) Brid.

Cinclidium stygium

M. trichodes Spruc. var. *alpina* (Bruch) Braithw.

Meesia uliginosa* var. *alpina

M. uliginosa Hedw. var. *alpina* (Bruch) Hamp. (Hampe, 1837a).

Syn. *Meesia alpina* Funck ex Bruch (Funck ex Bruch, 1826).

Meesia trichodes Spruc. var. *alpina* (Bruch) Braithw. (Braithwaite, 1895).

Distr. Southern South America (Cardot and Brotherus, 1923; Kabiersch, 1937; Kühnemann, 1938).

M. vagans (Hook. f. et Wils.) C. Muell.

Philonotis vagans

Meteoriopsis

M. ambigua (Hornschr.) Broth. (Brotherus, 1906).

Syn. *Pilotrichum ambiguum* Hornsch. (Hornschuch, 1840).
Meteorium ambiguum (Hornsch.) Mitt. (Mitten, 1869).

Distr. Southern South America (Hampe, 1879).

M. characea (C. Muell.) Broth. (Brotherus, 1906).

Syn. *Neckera characea* C. Muell. (Mueller, 1879a).
Pilotrichella characea (C. Muell.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1906, 1925;
Kühnemann, 1938).

var. *nigrescens* (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1962).

Syn. *Neckera characea* C. Muell. var. *nigrescens* C. Muell. (Mueller, 1879a).
Pilotrichella characea (C. Muell.) Kindb. var. *nigrescens* (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1879a; Paris, 1897, 1905b).

var. *tenuis* (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1962).

Syn. *Neckera characea* C. Muell. var. *tenuis* C. Muell. (Mueller, 1879a).
Pilotrichella characea (C. Muell.) Kindb. var. *tenuis* (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1879a; Paris, 1897, 1905b).

M. decurrens (Broth.) Broth. (Brotherus, 1906).

Syn. *Meteorium decurrens* Broth. (Brotherus, 1900).

Distr. Southern South America (Herzog, 1952a).

M. implanata (Mitt.) Broth. (Brotherus, 1906b).

Syn. *Meteorium implanatum* Mitt. (Mitten, 1869).

Distr. Southern South America (Herzog, 1952a).

M. onusta (Mitt.) Broth. (Brotherus, 1906b).

Syn. *Meteorium onustum* Spruc. ex Mitt. (Spruce ex Mitten, 1869).

Distr. Southern South America (Herzog, 1952a).

M. piligera (C. Muell.) Broth. (Brotherus, 1906b).

Syn. *Meteorium piligerum* C. Muell. (Mueller, 1901b).

Distr. Southern South America (Herzog, 1952a).

M. remotifolia (C. Muell.) Broth. (Brotherus, 1906b).

Syn. *Leskea remotifolia* C. Muell. (Mueller, 1847a [1846]).

Distr. Southern South America (Hosseus, 1940).

M. straminea (C. Muell.) Broth. ex Par. (Brotherus ex Paris, 1909).

Syn. *Pilotrichum stramineum* C. Muell. (Mueller, 1850).
Meteorium stramineum (C. Muell.) Mitt. (Mitten, 1869).

Distr. Southern South America (Mueller, 1850; Mitten, 1869; Jaeger, 1877; Paris, 1905a;
Brotherus, 1925).

Meteoriump

M. ambiguum (Hornsch.) Mitt.

Meteoriopsis ambigua

M. ancistrodes (Mont.) Par.

Ancistrodes genuflexa

M. billardieri (Hamp.) Mitt.

Weynouthia cochlearifolia* var. *billardieri

M. decurrens Broth.

Meteoriopsis decurrens

M. excavatum Mitt.

Orthostichidium pentagonum

- M. filipendulum* Hook. f. et Wils. **Papillaria flexicaulis**
- M. flexicaule* Wils. **Papillaria flexicaulis**
- M. genuflexum* (C. Muell.) Mitt. **Ancistrodes genuflexa**
- M. illecebrinum* (C. Muell.) Broth.** (Brotherus, 1925).
 Syn. *Neckera illecebrina* C. Muell. (Mueller, 1879a).
Pilotrichella illecebrina (C. Muell.) Kindb. (Kindberg, 1888).
 Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938; Herzog, 1952a).
- M. implanatum* Mitt. **Meteoriopsis implanata**
- M. lorentzii* C. Muell. **Squamidium lorentzii**
- M. medium* (Aongstr.) Broth.** (Brotherus, 1906b).
 Syn. *Papillaria media* Aongstr. (Ångström, 1876a).
 Distr. Southern South America (Herter, 1933a).
- M. molle* Wils. **Weymouthia mollis**
- M. onustum* Spruc. ex Mitt. **Meteoriopsis onusta**
- M. piligerum* C. Muell. **Meteoriopsis piligera**
- M. stramineum* (C. Muell.) Mitt. **Meteoriopsis straminea**
- M. striatum* Lindb.** (Lindberg, 1864b [1865]).
 Distr. Southern South America (Lindberg, 1864b; Paris, 1905a).
- Microdus***
- M. euchlorus* (Mont.) Besch.** (*Bescherelle* in Paris, 1896).
 Syn. *Dicranum euchlorum* Mont. (Montagne, 1845a).
Aongstroemia euchlora (Mont.) C. Muell. (Mueller, 1848a).
Dicranella euchlora (Mont.) Lor. (Lorentz, 1866).
 Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Lorentz, 1866; Mitten, 1869; Jaeger, 1872; Paris, 1896, 1905a).
- M. hilarianus* (Mont.) Besch. **Dicranella hilariana**
- M. kunzeanus* (C. Muell.) Broth.** (Brotherus, 1924a).
 Syn. *Leptotrichum kunzeanum* C. Muell. (Mueller, 1847c).
Seligeria kunzeana (C. Muell.) C. Muell. (Mueller, 1848a).
Dicranella kunzeana (C. Muell.) Mitt. (Mitten, 1869).
Seligeria kunzei Kindb. (Kindberg, 1888) nom. illeg.
Weisia tenuis Kunz. ex Mont. (Kunze ex Montagne, 1850) hom. illeg.
Microdus tenuis Par. (Paris, 1896) nom. illeg.
 Distr. Southern South America (Mueller, 1847c, 1848a; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Paris, 1896, 1905; Brotherus, 1924a).
- M. tenuirostris* (Schwaegr.) Besch. **Dicranella hilariana**
- M. tenuis* Par. **Microdus kunzeanus**
- Micromitrium***
- M. tomentosum* (Hornschr.) Par. **Groutiella tomentosa**
- Microthamnium***
- M. flavidum* Aongstr. **Mittenothamnium simorhynchum**
- M. gracile* (Hook. f. et Wils.) Jaeg. **Camptochaete gracilis**
- M. humile* Besch. **Mittenothamnium humile**
- M. hylophilum* C. Muell. **Mittenothamnium hylophilum**
- M. longo-reptans* C. Muell. **Mittenothamnium longoreptans**

<i>M. plinthophilum</i> (C. Muell.) Par.	<i>Mittenothamnium plinthophilum</i>
<i>M. pseudo-elegans</i> C. Muell.	<i>Mittenothamnium pseudo-elegans</i>
<i>M. pumilum</i> (Hook. f. et Wils.) Jaeg.	<i>Thamnobryum pumilum</i>
<i>M. sellovii</i> (Hornschr.) Mitt.	<i>Mittenothamnium sellovii</i>
<i>M. tapes</i> C. Muell.	<i>Mittenothamnium tapes</i>
<i>M. thelistegum</i> (C. Muell.) Mitt.	<i>Mittenothamnium diminutivum</i>
<i>M. widgrenii</i> Aongstr.	<i>Mittenothamnium camptorryhnchium</i>

Mielichhoferia***M. acuminata*** C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).***M. auriseta*** C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).**var. *alpina*** C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a).***M. austro-georgica*** C. Muell. (Mueller, 1890a).*Distr.* South Georgia (Mueller, 1890a; Paris, 1905a; Cardot, 1908a; Steere, 1961b; Clarke, 1973a).***M. brevicaulis*** Hornsch. (Hornschuch, 1840).*Distr.* Southern South America (Hornschuch, 1840; Mueller, 1848a; Montagne, 1850; Mitten, 1869; Gibert, 1873; Jaeger, 1875; Krieger, 1904; Paris, 1905a; Brotherus, 1924a; Herter, 1933a).***M. chilensis*** (Mont.) Mitt. (Mitten, 1869).*Syn.* *Leptochlaena chilensis* Mont. (Montagne, 1845a).*Distr.* Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Mitten, 1869; Jaeger, 1875; Paris, 1905a; Brotherus, 1924a; Espinosa B., 1924).***M. coarctata*** C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).***M. cygnicolla*** Hamp.***Anomobryum cygnicollum******M. demissa*** C. Muell. (Mueller, 1858).*Distr.* Southern South America (Mueller, 1858; Jaeger, 1875; Paris, 1905a; Brotherus, 1924a).***M. emergens*** (C. Muell.) Broth. (Brotherus, 1904).*Syn.* *Bryum emergens* C. Muell. (Mueller, 1879a).*Webera emergens* (C. Muell.) Kindb. (Kindberg, 1889).*Distr.* Southern South America (Mueller, 1879a; Brotherus, 1904; Paris, 1905a; Kühnemann, 1938).***M. fulvonitens*** Card. et Broth. (Cardot and Brotherus, 1923).*Distr.* Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).***M. gayana*** Card.***Mielichhoferia pleurogyna******M. humipetens*** (C. Muell.) Par.***Haplodontium humipetens******M. immersidens*** C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

M. leptoclada C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

M. longiseta C. Muell. (Mueller, 1848a).

Syn. *Bryum nutans* (Hedw.) Turn. var. *minus* Hook. (Hooker in Kunth, 1822).
Webera nutans Hedw. var. *minor* (Hook.) Brid. (Bridel, 1827a).

Distr. Juan Fernandez (Mitten, 1885; Johow, 1896; Paris, 1905a; Brotherus, 1924b; Robinson, 1975).

M. lorentziana C. Muell. (Mueller, 1879a).

Syn. *Mielichhoferia lorentzii* C. Muell. ex Kindb. (Mueller ex Kindberg, 1889) *nom. illeg.*

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905a; Brotherus, 1924a; Hosseus, 1936, 1937; Kühnemann, 1938).

M. lorentzii C. Muell. ex Kindb.***Mielichhoferia lorentziana******M. manca*** (C. Muell.) Broth. (Brotherus, 1903).

Syn. *Bryum mancum* C. Muell. (Mueller, 1879a).
Webera manca (C. Muell.) Kindb. (Kindberg, 1889).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1903, 1924a; Paris, 1906; Kühnemann, 1938).

M. micropoma C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

var. *brunnescens* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a).

M. multiflora Bartr. (Bartram, 1946).

Distr. Southern South America (Rovainen, 1934; Bartram, 1946).

M. ochracea C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

M. pampae C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

M. patagonica C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a).

M. pernana (C. Muell.) Par.***Haplodontium sanguinolentum***var. *robustior* (C. Muell.) Par.***Haplodontium pernanum* var. *robustior******M. pleurogena*** Mont. ex C. Muell.***Mielichhoferia pleurogyna******M. pleurogyna*** Mont. (Montagne, 1845a).

Syn. *Mielichhoferia pleurogena* Mont. ex C. Muell. (Montagne ex Mueller, 1848c) *err.*
Mielichhoferia gayana Card. (Cardot in Thériot, 1921b) *nom. nud.*

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a, c; Lorentz, 1866; Mitten, 1869; Jaeger, 1875; Paris, 1905a; Thériot, 1921b, 1934b; Brotherus, 1924a; Looser, 1932; Bartram, 1952).

M. pohlioidea C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905a; Brotherus, 1924a; Hosseus, 1936, 1937, 1938c, d; Kühnemann, 1938).

M. pulchra Broth. (Brotherus in Felippone, 1909).

Distr. Southern South America (Felippone, 1909; Brotherus, 1924a; Herter, 1933a; Herzog, 1952a).

M. sanguinolenta (C. Muell.) Par.

Haplodontium sanguinolentum

M. serridens C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

M. skottsbergii Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

M. spegazzinii C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Paris, 1905a; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

M. subpohlioidea Broth. (Brotherus, 1918).

Distr. Southern South America (Brotherus, 1918, 1924a; Hosseus, 1936, 1937; Kühnemann, 1938).

M. ulei C. Muell. (Mueller, 1898c).

Distr. Southern South America (Herter, 1943).

M. uruguensis Besch. (Bescherelle in Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

Mittenothamnium

M. camptorrhynchium (Hamp.) Card. (Cardot, 1913b).

Syn. *Hypnum camptorrhynchium* Hamp. (Hampe, 1870).
Microthamnium widgrenii Aongstr. (Ångström, 1876a) *nom. illeg.*

Distr. Southern South America (Krieger, 1904).

M. diminutivum (Hamp.) Britt. (Britton, 1914).

Syn. *Hypnum diminutivum* Hamp. (Hampe, 1847).
Hypnum thelistegum C. Muell. (Mueller, 1851a).
Microthamnium thelistegum (C. Muell.) Mitt. (Mitten, 1869).

Distr. Southern South America (Hosseus, 1940; Herter, 1943; Herzog, 1952a).

M. humile (Besch.) Card. (Cardot, 1913b).

Syn. *Microthamnium humile* Besch. (Bescherelle, 1877).
Stereohypnum humile (Besch.) Herz. (Herzog, 1909).

Distr. Southern South America (Brotherus, 1918, 1925; Kühnemann, 1938).

M. hylophilum (C. Muell.) Card. (Cardot, 1913b).

Syn. *Microthamnium hylophilum* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1925; Kühnemann, 1938; Herzog, 1952a).

M. longoreptans (C. Muell.) Card. (Cardot, 1913b).

Syn. *Microthamnium longo-reptans* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1925; Kühnemann, 1938).

M. plinthophilum (C. Muell.) Card. (Cardot, 1913b).

Syn. *Hypnum plinthophilum* C. Muell. (Mueller, 1879a).

Microthamnium plinthophilum (C. Muell.) Par. (Paris, 1896).

Distr. Southern South America (Mueller, 1879a; Paris, 1896, 1905a; Brotherus, 1925; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939).

***M. pseudo-elegans* (C. Muell.) Card. (Cardot, 1913b).**

Syn. *Microthamnium pseudo-elegans* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1925; Kühnemann, 1938).

***M. sellovii* (Hornschr.) Card. (Cardot, 1913b).**

Syn. *Hookeria sellovii* Hornsch. (Hornschuch, 1840).

Microthamnium sellovii (Hornsch.) Mitt. (Mitten, 1869).

Distr. Southern South America (Herter, 1928, 1933a, b).

***M. simorhynchum* (Hamp.) Card. (Cardot, 1913b).**

Syn. *Hypnum simorhynchum* Hamp. (Hampe, 1870).

Microthamnium flavidum Aongstr. (Ångström, 1876a) *nom. illeg.*

Distr. Southern South America (Krieger, 1904).

***M. tapes* (C. Muell.) Card. (Cardot, 1913b).**

Syn. *Microthamnium tapes* C. Muell. (Mueller, 1897a).

Stereohypnum tapes (C. Muell.) Fleisch. (Fleischer, 1908a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

Mniadelphus* *nom. illeg.

M. apiculatus (Hook. f. et Wils.) Mitt.

Eriopus apiculatus

M. assimilis Broth.

Distichophyllum assimile

M. cavifolius Card.

Distichophyllum eremitae

M. dicksonii (Hook. et Grev.) Hamp.

Distichophyllum dicksonii

M. eremitae Jaeg.

Distichophyllum eremitae

M. flaccidus (Hook. f. et Wils.) Hamp.

Distichophyllum flaccidum

M. hahnianus C. Muell.

Eriopus apiculatus

M. hahnii C. Muell.

Eriopus apiculatus

M. krausei Lor.

Distichophyllum kraussei

M. kraussei Lor.

Distichophyllum kraussei

M. mollis C. Muell.

Eriopus bescherellei

M. nigricans C. Muell.

Distichophyllum kraussei

M. procumbens C. Muell.

Distichophyllum eremitae

M. rotundifolius (Hook. f. et Wils.) C. Muell.

Distichophyllum rotundifolium

Mniobryum

M. albicans (Wahlenb.) Limpr.

Pohlia wahlenbergii

M. alticaule (C. Muell.) Broth.

Pohlia wahlenbergii* var. *glacialis

var. crassinerve Broth. (Brotherus in Cardot and Brotherus, 1923).

Syn. *Webera alticaulis* (C. Muell.) Kindb. var. *crassinervis* Card. (Cardot in Cardot and Brotherus, 1923) *nom. nud.*

Distr. Southern South America (Cardot and Brotherus, 1923).
Falkland Islands (Cardot and Brotherus, 1923).

var. robustius (C. Muell.) Par. (Paris, 1905a).

Syn. *Bryum alticaule* C. Muell. var. *robustius* C. Muell. (Mueller, 1889).

Webera alticaulis (C. Muell.) Kindb. var. *robustior* (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1889; Paris, 1898, 1905a; Cardot, 1908a; Kühnemann, 1938).

M. amplirete (C. Muell.) Broth.

Pohlia wahlenbergii* var. *glacialis

M. aspillacae Thér. (Thériot, 1924).

Distr. Southern South America (Thériot, 1924; Brotherus, 1925).

M. austro-albicans (C. Muell.) Broth.

Pohlia wahlenbergii

M. philonoteum (C. Muell.) Broth. (Brotherus, 1903).

Syn. *Bryum philonoteum* C. Muell. (Mueller, 1885).

Webera philonotea (C. Muell.) Par. (Paris, 1906).

Webera philonotidea Par. (Paris, 1898) err.

Distr. Southern South America (Mueller, 1885; Paris, 1898, 1905a; Brotherus, 1903, 1924a; Cardot, 1908a; Kühnemann, 1938; Herzog, 1954).

M. pulchellum (Hedw.) Loesk. (Loeske, 1910).

Syn. *Bryum pulchellum* Hedw. (Hedwig, 1801).

Distr. Southern South America (Hornschuch, 1820b; Montagne, 1850).

M. wahlenbergii (Web. et Mohr) Jenn.

Pohlia wahlenbergii

var. *glaciale* (Brid.) Wijket Marg.

Pohlia wahlenbergii* var. *glacialis

Mnium

<i>M. affine</i> Bland. ex Funck var. <i>magellanicum</i> Card. et Broth.	<i>Plagiomnium rugicum</i>
var. <i>rugicum</i> (Laur.) B.S.G.	<i>Plagiomnium rugicum</i>
<i>M. albicans</i> Wahlenb.	<i>Pohlia wahlenbergii</i>
<i>M. aubertii</i> Schwaegr.	<i>Rhodobryum aubertii</i>
<i>M. beyrichianum</i> Hornsch.	<i>Rhodobryum beyrichianum</i>
<i>M. bimum</i> Schreb.	<i>Bryum pseudotriquetrum</i> ssp. <i>bimum</i>
var. <i>turbinatum</i> (Hedw.) Wahlenb.	<i>Bryum turbinatum</i>
<i>M. brevifolium</i> (Broth.) C. Muell.	<i>Rhizogonium brevifolium</i>
<i>M. caespiticium</i> (Hedw.) With.	<i>Bryum caespiticium</i>
<i>M. chloroloma</i> C. Muell.	<i>Mnium rostratum</i>
<i>M. cristatum</i> (Hedw.) P. Beauv.	<i>Eriopus cristatus</i>
<i>M. crudum</i> Hedw.	<i>Pohlia cruda</i>
<i>M. fontanum</i> Hedw.	<i>Philonotis fontana</i>
var. <i>magellanicum</i> Brid.	<i>Philonotis fontana</i> var. <i>magellanicica</i>
<i>M. lanatum</i> P. Beauv.	<i>Bryum argenteum</i> var. <i>lanatum</i>
<i>M. lepto-limbatum</i> C. Muell.	<i>Plagiomnium rugicum</i>
<i>M. lindigii</i> Hamp.	<i>Rhizogonium lindigii</i>
<i>M. longicolle</i> C. Muell. <i>in</i> Par.	<i>Rhizogonium longicolle</i>
<i>M. longicolle</i> C. Muell. (Mueller, 1901a [1900]) <i>hom. illeg.</i>	
<i>Distr.</i> Southern South America (Mueller, 1901a).	
<i>M. longirostre</i> Brid.	<i>Mnium rostratum</i>
<i>M. palustre</i> Hedw.	<i>Aulacomnium palustre</i>
<i>M. polycarpum</i> C. Muell.	<i>Rhizogonium mnioides</i>
<i>M. pseudo-triquetrum</i> Hedw.	<i>Bryum pseudotriquetrum</i>
<i>M. punctulatum</i> Brid.	<i>Tortula robusta</i>
<i>M. reticulatum</i> (Hook. f. et Wils.) C. Muell.	<i>Philonotis vagans</i>

- M. rigidum* Hornsch. *Bryum rigidum*
M. roseum Hedw. *Rhodobryum roseum*
- M. rostratum*** Schrad. (Schrader in Schreibius, 1802).
 Syn. *Hypnum rostratum* (Schrad.) Web. et Mohr (Weber and Mohr, 1803).
Bryum rostratum (Schrad.) Sm. (Smith, 1804b).
Mnium longirostre Brid. (Bridel, 1803) *nom. illeg.*
Bryum longirostre Brid. (Bridel, 1819).
Mnium chloroloma C. Muell. (Mueller, 1882).
 Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1848a, 1882; Montagne, 1850; Gibert, 1873; Dusén, 1903a; Paris, 1905a; Cardot, 1908a; Brotherus, 1924a; Herter, 1933a, 1939a; Kühnemann, 1938; Hosseus, 1940; Herzog, 1952a).
- M. rugicum* Laur. *Plagiomnium rugicum*
M. schleicheri (Cand.) Brid. *Bryum schleicheri*
M. stygium (Sw.) B.S.G. *Cinclidium stygium*
M. subbasilare (Hook.) C. Muell. *Goniobryum subbasilare*
M. subpunctatum Card. et Broth. *Cinclidium stygium*
M. truncorum Brid. *Bryum truncorum*
M. turbinatum Hedw. *Bryum turbinatum*
- M. undulatum*** Hedw. (Hedwig, 1801).
 Distr. Southern South America (Kühnemann, 1938).
- Molendoa***
- M. excelsa* (C. Muell.) Broth. *Molendoa sendtneriana*
M. fuegiana Bartr. (Bartram, 1946).
 Distr. Southern South America (Bartram, 1946).
M. sendtneriana (B.S.G.) Limpr. (Limpicht, 1886).
 Syn. *Anoectangium sendtnerianum* B.S.G. (Bruch and others, 1846).
Zygodon excelsus C. Muell. (Mueller, 1879a) *fide* Zander, 1977.
Anoectangium excelsum (C. Muell.) Kindb. (Kindberg, 1889) *fide* Zander, 1977.
Molendoa excelsa (C. Muell.) Broth. (Brotherus, 1902) *fide* Zander, 1977.
Anoectangium lechlerianum Mitt. (Mitten, 1869) *fide* Zander, 1977.
 Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Paris, 1905a; Kühnemann, 1938; Bartram, 1965; Zander, 1977).
- Mollia hom. illeg.*
M. brachydontia (Bruch) Lindb. *Trichostomum brachydontium*
M. humilis (Hedw.) Lindb. *Tortella humilis*
M. tortuosa (Hedw.) Schrank ex Lindb. *Tortella tortuosa*
- Monocranum***
- M. dicranoides*** C. Muell. (Mueller, 1901a [1900]).
 Distr. Southern South America (Mueller, 1901a).
- Muelleriella hom. illeg.*
M. crassifolia (Hook. f. et Wils.) Dus. (Dusén, 1905b).
 Syn. *Orthotrichum crassifolium* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Ulota crassifolia (Hook. f. et Wils.) Hook. f. (Hooker, 1867).
Muelleriella crassifolia (Hook. f. et Wils.) Dus. var. *lingulata* Dus. (Dusén, 1905b).
 Distr. Southern South America (Wilson and Hooker, 1847 (as vars 2 and 3); Mueller, 1849, 1885; Sullivant, 1859; Ångström, 1872; Jaeger, 1874; Bescherelle, 1889; Dusén, 1905b; Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Vitt, 1976).

Falkland Islands (Hooker, 1847 (as var. 1); Jaeger, 1874; Mueller, 1885; Bescherelle, 1889; Dusén, 1905b; Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Vitt, 1976).
South Georgia (Cardot, 1906a, 1908a; Taylor, 1914; Cardot and Brotherus, 1923; Steere, 1961b; Vitt, 1976).

var. *lingulata* Dus.

Muelleriella crassifolia

Muscoflorschuetzia

M. pilmaiwen (Crosby) Crosby (Crosby, 1978).

Syn. *Florschuetzia pilmaiwen* Crosby (Crosby, 1977b) nom. illeg.

Distr. Southern South America (Crosby, 1977b).

Myurella

M. julacea (Schwaegr.) B.S.G. (Bruch and others, 1853).

Syn. *Leskea julacea* Schwaegr. (Schwaegrichen, 1804).

Distr. South Georgia (Newton, 1977a).

Neckera

N. amblyoglossa C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

N. appressa C. Muell.

Papillaria nigrescens

N. argentinica Lor. ex C. Muell. (Lorentz ex Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1925; Williams, 1930; Kühnemann, 1938; Herzog, 1952a).

N. avellaneda C. Muell.

Orthostichopsis avellaneda

N. berteroana Schimp. ex Mitt. (Schimper ex Mitten, 1869).

Syn. *Neckera berteroana* Kindb. (Kindberg, 1888) nom. illeg.

Distr. Southern South America (Mitten, 1869; Jaeger, 1877; Paris, 1905a; Thériot, 1915; Brotherus, 1925).

N. berteroana Kindb.

Neckera berteroana

N. beyrichii Schwaegr.

Entodon beyrichii

N. billardieri Hamp.

Weymouthia cochlearifolia* var. *billardieri

N. characea C. Muell.

Meteoriopsis characea

var. *nigrescens* C. Muell.

Meteoriopsis characea* var. *nigrescens

var. *tenuis* C. Muell.

Meteoriopsis characea* var. *tenuis

N. chilensis Schimp. (Schimper, 1836).

Distr. Southern South America (Schimper, 1836; Montagne, 1850; Mueller, 1850; Mitten, 1869; Jaeger, 1877; Neger, 1899; Paris, 1905a; Cardot, 1908a; Campo, 1915; Thériot, 1915, 1917b; Costes, 1921; Herzog, 1923, 1938, 1940, 1954, 1957; Brotherus, 1924c, 1925; Reimers, 1926; Kühnemann, 1938; Espinosa B., 1941; Skottsberg, 1950; Seki, 1974).

N. coronata (Mont.) C. Muell.

Forsstroemia coronata

N. cumingii C. Muell.

Weymouthia mollis

N. curtipendula Hedw.

Antitrichia curtipendula

N. cyathipoma C. Muell.

Pilotrichella cyathipoma

N. dillenii Lindb.

Homalia glabella

N. disticha Hedw.

Neckeropsis disticha

N. eriodon C. Muell.

Eriodon conostomus

<i>N. fasciculata</i> (Hedw.) Arnott	<i>Porothamnium fasciculatum</i>
<i>N. glabella</i> (Hedw.) Web. et Mohr	<i>Homalia glabella</i>
<i>N. glauca</i> C. Muell.	<i>Lepyrodon hexastichus</i>
<i>N. humboldtii</i> (Hook.) C. Muell.	<i>Rhacocarpus purpurascens</i>
<i>N. illecebrina</i> C. Muell.	<i>Meteoriump illecebrinum</i>
<i>N. imberbis</i> (Sm.) C. Muell.	<i>Hedwigidium integrifolium</i>
<i>N. implexa</i> C. Muell.	<i>Lepyrodon hexastichus</i>
<i>N. jamesonii</i> Tayl. (Taylor, 1846).	
<i>Distr.</i> Southern South America (Mitten, 1869; Brotherus, 1925).	
<i>N. lagura</i> (Hook.) C. Muell.	<i>Lepyrodon lagurus</i>
<i>N. leucocaulon</i> C. Muell.	<i>Porothamnium leucocaulon</i>
<i>N. leucocolea</i> Mitt.	<i>Cryphidium leucocoleum</i>
<i>N. mollis</i> (Hedw.) C. Muell.	<i>Weymouthia mollis</i>
<i>N. patagonica</i> Brid.	<i>Homalia glabella</i>
<i>N. pennata</i> Hedw. (Hedwig, 1801).	
<i>Distr.</i> Southern South America (Montagne, 1850).	
<i>N. poeppigiana</i> C. Muell.	<i>Glyphothecium gracile</i>
<i>N. polytrichoides</i> (Hedw.) Web. et Mohr	<i>Lepidopilum polytrichoides</i>
<i>N. porteri</i> Thér.	<i>Neckera scabridens</i> var. <i>porteri</i>
<i>N. praelonga</i> Lor. (Lorentz, 1866).	
<i>Distr.</i> Southern South America (Lorentz, 1866; Paris, 1905a).	
<i>N. rotundata</i> Broth. (Brotherus, 1924b).	
<i>Distr.</i> Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Robinson, 1975).	
<i>N. scabridens</i> C. Muell. (Mueller, 1847c).	
<i>Distr.</i> Southern South America (Mueller, 1847c, 1850; Sullivant, 1859; Lorentz, 1866; Mitten, 1869; Jaeger, 1877; Paris, 1905a; Cardot, 1908a; Herzog, 1923, 1954, 1960; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938; Bartram, 1952; Seki, 1974).	
var. <i>porteri</i> Thér. (Thériot, 1915).	
<i>Syn.</i> <i>Neckera porteri</i> Thér. (Thériot, 1915) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Thériot, 1915, 1917b; Costes, 1921).	
<i>N. scabriseta</i> Schwaegr.	<i>Lepidopilum scabrisetum</i>
<i>N. schnyderi</i> C. Muell.	<i>Neckeropsis villaे-ricaе</i>
<i>N. smithii</i> (Hedw.) C. Muell.	<i>Leptodon smithii</i>
<i>N. suborthosticha</i> C. Muell.	<i>Lepyrodon tomentosus</i>
<i>N. tenella</i> Schwaegr.	<i>Cryphaea tenella</i>
<i>N. tomentosa</i> (Hook.) C. Muell.	<i>Lepyrodon tomentosus</i>
<i>N. trichoblepharis</i> Schimp. (Schimper in Paris, 1896) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Paris, 1896, 1905a).	
<i>N. trichophora</i> Schimp. (Schimper in Lorentz, 1866) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Lorentz, 1866; Jaeger, 1877; Paris, 1905a).	
<i>N. truncata</i> C. Muell.	<i>Entodon truncatus</i>
<i>N. turgidula</i> C. Muell.	<i>Squamidium turgidulum</i>
var. <i>tristis</i> C. Muell.	<i>Squamidium turgidulum</i> var. <i>triste</i>

<i>N. undulata</i> Hedw.	<i>Neckeropsis undulata</i>
<i>N. uruguensis</i> C. Muell.	<i>Neckeropsis villaे-ricaе</i>
<i>N. villaे-ricaе</i> Besch.	<i>Neckeropsis villaе-ricaе</i>
<i>N. viticulosoides</i> P. Beauv.	<i>Leptodontium viticulosoides</i> var. <i>viticulosoides</i>
<i>Neckeropsis</i>	
<i>N. disticha</i> (Hedw.) Kindb. (Kindberg, 1894).	
Syn. <i>Neckera disticha</i> Hedw. (Hedwig, 1801).	
Distr. Southern South America (Paris, 1905a).	
<i>N. undulata</i> (Hedw.) Reichdt. (Reichardt, 1870).	
Syn. <i>Neckera undulata</i> Hedw. (Hedwig, 1801).	
<i>Hypnum harrisonia</i> Poir. (Poiret in Bridel, 1827b) nom. nud.	
Distr. Southern South America (Bridel, 1801b, 1817, 1827b; Paris, 1905a; Hosseus, 1940; Herzog, 1952a).	
<i>N. villaе-ricaе</i> (Besch.) Broth. ex Par. (Brotherus ex Paris, 1909).	
Syn. <i>Neckera villaе-ricaе</i> Besch. (Bescherelle, 1877).	
<i>Neckera uruguensis</i> C. Muell. (Mueller, 1879a).	
<i>Neckera schnyderi</i> C. Muell. (Mueller, 1882).	
Distr. Southern South America (Mueller, 1879a, 1882; Paris, 1905a; Brotherus, 1925; Herter, 1933a, 1939b; Kühnemann, 1938; Hosseus, 1940; Herzog, 1952a).	
<i>Neobarbula</i>	
<i>N. magellanica</i> Dus.	<i>Tortula densifolia</i>
<i>Neuroloma</i> hom. illeg.	
<i>N. fuegianum</i> Card.	<i>Andreaea fuegiana</i>
<i>Notarisia</i> hom. illeg.	
<i>N. crispata</i> (Hedw.) Mont.	<i>Ptychomitrium crispatum</i>
<i>Octoblepharum</i>	
<i>O. albidum</i> Hedw. (Hedwig, 1801).	
Syn. <i>Bryum albidum</i> (Hedw.) P. Beauv. (Palisot de Beauvois, 1805a).	
Distr. Southern South America (Hornschorch, 1820b; Montagne, 1850; Mueller, 1879a; Kühnemann, 1938).	
<i>Octodiceras</i>	
<i>O. berterii</i> (Mont.) Jaeg.	<i>Fissidens fontanus</i>
<i>O. dillenii</i> (Pyl.) Brid.	<i>Fissidens fontanus</i>
<i>O. fissidentoides</i> Brid.	<i>Fissidens semicompletus</i>
<i>O. fontanum</i> (Pyl.) Lindb.	<i>Fissidens fontanus</i>
<i>O. hedwigii</i> Lindb.	<i>Fissidens semicompletus</i>
<i>O. julianum</i> (Cand.) Brid.	<i>Fissidens fontanus</i>
<i>Oedipodium</i>	
<i>O. griffithianum</i> (Dicks.) Schwaegr. (Schwaegrichen, 1823).	
Syn. <i>Bryum griffithianum</i> Dicks. (Dickson, 1801).	
<i>Gymnostomum griffithianum</i> (Dicks.) Sm. (Smith, 1804b).	
<i>Hymenostomum griffithianum</i> (Dicks.) Spreng. (Sprengel, 1827).	
Distr. Southern South America (Matteri, 1979).	
Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).	

Oligotrichum*O. antarcticum* (C. Muell.) Kindb.***Psilopilum trichodon****O. atrichopsis* C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

O. austro-aligerum G. L. Smith (Smith, 1969b).Syn. *O. magellanicum* Card. et Broth. (Cardot and Brotherus, 1923) *hom. illeg.*

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938; Herzog, 1954; Seki, 1974).

O. canaliculatum (Hook. et Arnott) Mitt. (Mitten, 1869).Syn. *Polytrichum canaliculatum* Hook. et Arnott (Hooker and Arnott, 1832).*Catharinea canaliculata* (Hook. et Arnott) C. Muell. (Mueller, 1848a).*Pogonatum liliputanum* Thér. (Thériot, 1917b).

Distr. Juan Fernandez (Bartram, 1959; Robinson, 1975).

Southern South America (Hooker and Arnott, 1832; Schwaegrichen, 1842; Mueller, 1848a; Montagne, 1850; Mitten, 1869; Jaeger, 1875; Krieger, 1904; Paris, 1905a; Thériot, 1915, 1917b, 1925a, 1935a; Costes, 1921; Herzog, 1923, 1938, 1939, 1954, 1960; Brotherus, 1924c, 1925; Reimers, 1926; Espinosa B., 1941; Mullen and Frye, 1947; Bartram, 1952; Seki, 1974).

var. *majus* (Lor.) Par. (Paris, 1896).Syn. *Catharinea canaliculata* (Hook. et Arnott) C. Muell. var. *major* Lor. (Lorentz, 1866).

Distr. Southern South America (Lorentz, 1866; Paris, 1896, 1905a).

var. *minus* Broth. (Brotherus, 1924b).Syn. *Oligotrichum subcanaliculatum* Dus. (Dusén in Mullen and Frye, 1947) *nom. nud.*

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Mullen and Frye, 1947; Bartram, 1959).

Southern South America (Herzog, 1938, 1939; Mullen and Frye, 1947).

var. *molinae* (Mont.) Thér. (Thériot, 1924).Syn. *Polytrichum molinae* Mont. (Montagne, 1845a).*Catharinea molinae* (Mont.) C. Muell. (Mueller, 1848a).*Oligotrichum molinae* (Mont.) Lor. (Lorentz, 1866).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Lorentz, 1866; Jaeger, 1875; Paris, 1905a; Thériot, 1924).

O. compressum (Hook. f. et Wils.) Kindb.***Atrichopsis compressa****O. crispatissimum* C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

O. dendroides (Hedw.) Kindb.***Dendroligotrichum dendroides****O. gymnostomulum* (C. Muell.) Kindb.***Psilopilum gymnostomulum****O. horridum* (Mitt.) Kindb.***Polytrichadelphus magellanicus****O. laevifolium* Thér. (Thériot, 1924).

Distr. Southern South America (Thériot, 1924; Brotherus, 1925; Herzog, 1938).

O. magellanicum (Hedw.) Kindb.***Polytrichadelphus magellanicus****O. magellanicum* Card. et Broth.***Oligotrichum austro-aligerum****O. molinae* (Mont.) Lor.***Oligotrichum canaliculatum* var. *molinae****O. obtusifolium* Thér. (Thériot, 1934b [1935]).

Distr. Southern South America (Thériot, 1934b).

O. riedelianum (Mont.) Mitt. (Mitten, 1869).Syn. *Polytrichum riedelianum* Mont. (Montagne, 1841).

Distr. Southern South America (Krieger, 1904).

O. rigidum (Lor.) Broth. (Brotherus, 1905).

Syn. *Atrichum rigidum* Lor. (Lorentz, 1864).

Catharinea rigida (Lor.) C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Lorentz, 1864, 1866; Jaeger, 1875; Mueller, 1901a; Paris, 1904a; Brotherus, 1905, 1925; Mullen and Frye, 1947).

O. squamosum (Hook. f. et Wils.) Kindb.

Dendroligotrichum squamosum

O. subcanaliculatum Dus.

Oligotrichum canaliculatum var. *minus*

O. tapes (C. Muell.) Kindb.

Psilopilum tapes

O. umbrosum (Mitt.) Kindb.

Polytrichadelphus umbrosus

Oncophorus

O. billardieri (Brid.) Brid.

Dicranoloma billardieri

O. cyathicarpus (Mont.) Mitt.

Amphidium tortuosum

O. dichotomus (P. Beauv.) Brid.

Dicranoloma billardieri

O. fuegianus Card. (Cardot, 1900).

Syn. *Cynodontium fuegianum* (Card.) Card. (Cardot, 1901).

Dichodontium brotheri Dix. (Dixon in Paris, 1904b) nom. nud.

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Robinson, 1975).

Southern South America (Cardot, 1900, 1901, 1905a, 1908a; Paris, 1904b, 1905a; Brotherus, 1924a; Kühnemann, 1938).

O. luteo-virens Bartr. (Bartram in Roivainen and Bartram, 1937).

Distr. Southern South America (Roivainen and Bartram, 1937).

O. starkei (Web. et Mohr) Brid.

Kiaeria starkei

O. striatus (Schrad.) Lindb.

Rhabdoweisia fugax

Oreoweisia

O. auridens (C. Muell.) Kindb.

Oreoweisia bogotensis

var. *robusta* (C. Muell.) Par. (Paris, 1896).

Syn. *Weisia auridens* C. Muell. var. *robusta* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1896, 1905a).

O. bogotensis (Hamp.) Mitt. (Mitten, 1869).

Syn. *Weisia bogotensis* Hamp. (Hampe, 1863).

Weisia auridens C. Muell. (Mueller, 1882).

Oreoweisia auridens (C. Muell.) Kindb. (Kindberg, 1889).

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1924a; Hosseus, 1938c, d; Kühnemann, 1938).

O. brevifolia Broth. (Brotherus, 1918).

Distr. Southern South America (Brotherus, 1918, 1924a; Kühnemann, 1938).

O. chilensis (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Weisia chilensis* Hamp. ex C. Muell. (Hampe ex Mueller, 1862).

Distr. Southern South America (Mueller, 1862; Jaeger, 1870; Paris, 1905a; Brotherus, 1924a).

O. patagonica Dus. (Dusén in Paris, 1905a) nom. nud.

Distr. Southern South America (Paris, 1905a).

Orthodontium

O. australe Hook. f. et Wils.

Orthodontium lineare

var. *brevisetum* Card. et Broth.

Orthodontium lineare

var. *sinuosifolium* Card. et Broth.

Orthodontium lineare

O. gracile Schwaegr. ex B.S.G. (Schwaegrichen *ex* Bruch and others, 1844).

Syn. *Orthodontium tenellum* Mitt. (Mitten, 1869).

Stableria tenella (Mitt.) Broth. (Brotherus, 1903).

Orthodontium laxirete Dus. (Dusén *in* Paris, 1905a) *nom. nud.*

Distr. Juan Fernandez (Bartram, 1959; Robinson, 1975).

Southern South America (Paris, 1905a; Thériot, 1923; Meijer, 1952; Herzog, 1954).

O. humile (Mont.) Mitt.

Pohlia humilis

O. laxirete Dus.

Orthodontium gracile

O. lineare Schwaegr. (Schwaegrichen, 1827a).

Syn. *Apalodium linare* (Schwaegr.) Mitt. (Mitten, 1885) *hom. illeg.*

Orthodontium australe Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Apalodium australe (Hook. f. et Wils.) Mitt. (Mitten, 1869).

Orthodontium australe Hook. f. et Wils. var. *brevisetum* Card. et Broth. (Cardot and Brotherus, 1923).

Orthodontium australe Hook. f. et Wils. var. *sinuosifolium* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a; Mitten, 1869; Jaeger, 1875; Paris, 1905a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Meijer, 1952).

Falkland Islands (Hooker J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a; Jaeger, 1875; Paris, 1905a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Meijer, 1952).

O. pellucens (Hook.) B.S.G. (Bruch and others, 1844).

Syn. *Bryum pellucens* Hook. (Hooker, 1836).

Apalodium pellucens (Hook.) Mitt. (Mitten, 1869).

Apalodium confine (Hamp.) Mitt. (Mitten, 1885).

Distr. Southern South America (Herzog, 1954).

O. tenellum Mitt.

Orthodontium gracile

O. tenue C. Muell.

Pohlia humilis

Orthopyxis nom. rejic.

O. palustris (Hedw.) P. Beauv.

Aulacomnium palustre

Orthorrhynchium

O. chilense Williams

Catagonium politum

Orthostichidium

O. excavatum (Mitt.) Broth.

Orthostichidium pentagonum

O. pentagonum (Hamp. et Lor.) C. Muell. (Mueller, 1897c).

Syn. *Pilotrichum pentagonum* Hamp. et Lor. (Hampe and Lorentz *in* Lorentz, 1868).

Meteoriump excavatum Mitt. (Mitten, 1869).

Orthostichidium excavatum (Mitt.) Broth. (Brotherus, 1906b).

Distr. Southern South America (Hosseus, 1940; Herzog, 1952a).

Orthostichopsis

O. avellanedaee (C. Muell.) Broth. (Brotherus, 1906b).

Syn. *Neckera avellanedaee* C. Muell. (Mueller, 1879a).

Pilotrichella avellanedaee (C. Muell.) Kindb. (Kindberg, 1891).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1906b, 1918, 1925; Kühnemann, 1938).

Orthotheca

O. berteroana (Brid.) Brid.

Syrrhopodon incompletus var. **berteroanus**

*Orthotheciella nom. nud.**O. austro-chilense* C. Muell.*Hygroamblystegium chilense**Orthothecium**O. filum* (C. Muell.) Kindb.*Hygroamblystegium filum**O. vittatum* Card. ex Syd.*Orthotrichum vittatum**Orthotrichum**O. affine* Schrad. ex Brid. var. *rupestre* Brid.*Orthotrichum rupestre**O. anaglyptodon* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

O. anderssonii C. Muell. ex Aongstr.*Orthotrichum leiothecium**O. angustissimum* Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a).

O. antarcticum Card. (Cardot, 1900).

Distr. Antarctic, peninsula region (Cardot, 1900, 1901; Paris, 1905a; Cardot, 1908a; Brotherus, 1925; Steere, 1961a).

O. apiculatum Hook.*Groutiella apiculata**O. argentinicum* Herz. (Herzog, 1957).

Distr. Southern South America (Herzog, 1957; Seki, 1974).

O. aristoblepharum Herz. (Herzog, 1954).

Distr. Southern South America (Herzog, 1954).

O. assimile C. Muell. (Mueller, 1849).

Distr. Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1874; Paris, 1905a; Thériot, 1921a, b, 1928; Brotherus, 1925; Herzog, 1954).

O. aurantiacum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

O. bellum C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

O. bicolor Thér. (Thériot, 1921a).

Distr. Southern South America (Thériot, 1921a, 1928, 1935a; Brotherus, 1925).

O. brevicolle Mitt.*Ulota rufula**O. carinatum* (Mitt.) Mitt.*Ulota carinata**O. chilense* Mitt.*Ulota rufula**O. compactum* Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938; Herzog, 1960).

O. conceptionense Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926).

O. crassifolium Hook. f. et Wils.*Muelleriella crassifolia**O. crenato-erosa* C. Muell. (Mueller, 1885).Syn. *Ulota crenato-erosa* (C. Muell.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Mueller, 1885; Paris, 1905a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

O. crispatum (Hedw.) Hook. et Grev.

Ptychomitrium crispatum

O. cupulatum Hoffm. ex Brid. (Hoffmann ex Bridel, 1801).

Distr. Southern South America (Montagne, 1850).

O. darwinii (Mitt.) Mitt.

Ulota fuegiana* ssp. *darwinii

O. diaphanum Schrad. ex Brid. var. *brevipilum* Warnst. (Warnstorf, 1904).

Distr. Southern South America (Warnstorf, 1904).

O. elegantulum Schimp. ex Mitt. (Schimper ex Mitten, 1869).

Distr. Southern South America (Lechler, 1857; Mitten, 1869; Jaeger, 1874; Mueller, 1885; Bescherelle, 1889; Cardot, 1905a (as *Ulota magellanica* fide Cardot, 1908a), 1908a; Paris, 1905a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Bartram, 1952; Bizot, 1967; Seki, 1974).

O. eremitense (Mitt.) Mitt.

Ulota glabella

O. erpodiaceum C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905a; Brotherus, 1925; Hosseus, 1935b and 1937 (both as *O. expodiaceum*); Kühnemann, 1938).

O. filiforme Hook. et Grev.

Macrocoma orthotrichoides

O. fimbriatum P. Beauv.

Macromitrium fimbriatum

O. fuegianum (Mitt.) Mitt.

Ulota fuegiana

O. fulvellum Mitt.

Ulota luteola

O. germanum Mont.

Ulota germana

O. glabellum (Mitt.) Mitt.

Ulota glabella

O. gymnomitrium (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Ulota gymnomitria* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Brotherus, 1902, 1925; Paris, 1905a; Cardot, 1908a; Kühnemann, 1938).

O. herteri Herz. (Herzog in Herter, 1937).

Distr. Southern South America (Herter, 1933a, 1937).

O. hymenomitrium Dus. (Dusén, 1903a) nom. nud.

Distr. Southern South America (Dusén, 1903a).

O. incanum C. Muell. (Mueller, 1885).

Syn. *Ulota incana* (C. Muell.) Kindb. (Kindberg, 1888).

Distr. South South America (Mueller, 1885; Paris, 1905a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

O. inclinatum C. Muell. (Mueller, 1885).

Syn. *Ulota inclinata* (C. Muell.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Mueller, 1885; Bescherelle, 1885a; Paris, 1905a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

O. jamesonii Arnott

Schlotheimia jamesonii

O. johnstonii Bartr. (Bartram, 1942).

Distr. Southern South America (Bartram, 1942, 1943).

O. jutlandicum Brid.

Ulota phyllantha

O. lebrunii Besch. (Bescherelle, 1885a).

Distr. Southern South America (Bescherelle, 1885a; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

O. leiothecium C. Muell. (Mueller, 1862).

Syn. *Ulota leiothecia* (C. Muell.) Jaeg. (Jaeger, 1874).

Orthotrichum anderssonii C. Muell. ex Aongstr. (Mueller ex Ångström, 1872).

Ulota anderssonii (Aongstr.) Jaeg. (Jaeger and Sauerbeck, 1879b).

Distr. Southern South America (Mueller, 1862, 1885; Ångström, 1872, 1876b; Jaeger, 1874; Bescherelle, 1885a; Paris, 1905a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

O. ligulatum Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a).

O. lobbianum (Mitt.) Mitt.

Ulota lobbiana

O. longipes Hook.

Macromitrium longipes

O. longirostre Hook.

Macromitrium longirostre

O. lorentzii C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1925; Herzog, 1938; Kühnemann, 1938).

O. luteolum Hook. f. et Wils.

Ulota luteola, *U. fuegiana* and *U. glabella*

O. macloskii Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

O. macrocalycinum (Mitt.) Mitt.

Ulota macrocalycina

O. magellanicum Mont.

Ulota magellanica

O. malacothecium C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

O. marginatum Aongstr.

Ulota macrocalycina

O. microstomum Hook. et Grev.

Macromitrium microstomum

O. nigrillum Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1905a; Seki, 1974).

O. nutans C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905a; Brotherus, 1925; Hosseus, 1935b, 1937; Kühnemann, 1938).

O. paleomitrium Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1905a; Herzog, 1957).

O. perpusillum Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1905a).

O. phyllanthum (Brid.) Steud.

Ulota phyllantha

O. podocarpi C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

O. polytrichoides (Hedw.) Brid.

Lepidopilum polytrichoides

O. pygmaeothecium C. Muell.

Ulota pygmaeothecia

O. quenoae C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1925; Hosseus, 1938a, d; Kühnemann, 1938).

O. rufulum Mitt.*Ulota rufula**O. rupestre* Schleich. ex Schwaegr. (Schleicher ex Schwaegrichen, 1816).Syn. *Dorcadion rupestre* (Schwaegr.) Lindb. (Lindberg, 1879).*Orthotrichum affine* Schrad. ex Brid. var. *rupestre* Brid. (Bridel, 1812) nom. nud.

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Costes, 1921; Kühnemann, 1938).

O. rupicola C. Muell. (Mueller, 1884 [1883]).

Distr. Antarctic, peninsula region (Cardot, 1900, 1901, 1908a; Paris, 1905a; Steere, 1961a).

O. schnyderi C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

O. smithii (Hedw.) Brid.*Leptodon smithii**O. subassimile* Dus. (Dusén, 1903a) nom. nud.

Distr. Southern South America (Dusén, 1903a).

O. subrupestre Thér. (Thériot, 1923b).Distr. Southern South America (Thériot, 1921a (as *O. rupestre*), 1923b; Brotherus, 1925).*O. torquatum* (Hedw.) Arnott*Schlotheimia torquata**O. truncato-dentatum* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Brotherus, 1925; Kühnemann, 1938).

O. ulotooides Herz. (Herzog, 1923).

Distr. Southern South America (Herzog, 1923).

O. urceolatum Hook.*Macromitrium urceolatum**O. verrucosum* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1925; Kühnemann, 1938).

O. vittatum Card. (Cardot, 1905a) hom. illeg.Syn. *Orthothecium vittatum* Card. ex Syd. (Cardot ex Sydow, 1910) nom. inval.

Distr. Southern South America (Cardot, 1905a, 1908a; Brotherus, 1925; Kühnemann, 1938).

Oxyrrhynchium

O. altisetum (C. Muell.) Broth. (Brotherus, 1909).Syn. *Rhynchosstegium altisetum* C. Muell. (Mueller, 1897a).*Hypnum altisetum* (C. Muell.) Broth. (Brotherus, 1900).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1909, 1925; Kühnemann, 1938).

O. clinocarpum (Tayl.) Broth. (Brotherus, 1909).Syn. *Hypnum clinocarpum* Tayl. (Taylor, 1848a).*Brachythecium clinocarpum* (Tayl.) Jaeg. (Jaeger, 1878).*Hypnum remotifolium* Grev. (Greville, 1826).*Eurhynchium remotifolium* (Grev.) Jaeg. (Jaeger, 1878).*Thuidium remotifolium* (Grev.) Ren. et Card. (Renauld and Cardot, 1893b).*Oxyrrhynchium remotifolium* (Grev.) Steere (Steere, 1948) hom. illeg.

Distr. Southern South America (Herzog, 1952a).

O. corralense (Lor.) Fleisch. ex Reim. (Fleischer ex Reimers, 1926).Syn. *Hypnum corralense* Lor. (Lorentz, 1866).*Eurhynchium corralense* (Lor.) Jaeg. (Jaeger, 1878).

Eurhynchium confusum Thér. (Thériot, 1918).
Rhynchosstegium subsquarrosum Herz. (Herzog, 1923).
Eurhynchium acanthophyllum (Mont.) Dus. ex Thér. (Dusén ex Thériot, 1934b). *nom. inval. fide* Thériot, 1934a.

Distr. Southern South America (Lorentz, 1866; Jaeger, 1878; Paris, 1904b; Thériot, 1918, 1934b; Campo, 1921; Herzog, 1923, 1938; Brotherus, 1924c, 1925; Reimers, 1926; Herter, 1933a, 1939b; Espinosa B., 1941; Skottsberg, 1950).

O. fuegianum (Card.) Herz. (Herzog, 1939).

Syn. *Eurhynchium fuegianum* Card. (Cardot, 1905a).
Rhynchosstegiella fuegiana (Card.) Broth. (Brotherus, 1909).
Eurhynchiella fuegiana (Card.) Fleisch. (Fleischer, 1923).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Fleischer, 1923; Brotherus, 1925; Herzog, 1938, 1939, 1954; Kühnemann, 1938; Seki, 1974).

O. remotifolium (Grev.) Steere

Oxrrhynchium clinocarpum

Pancovia hom. illeg.

P. denticulata (Hedw.) Kickx
P. rutabula (Hedw.) Kickx

Plagiothecium denticulatum
Brachythecium rutabulum

Papillaria

P. appressa (C. Muell.) Aongstr.
P. cardotii Broth.
P. filipendula (Hook. f. et Wils.) Jaeg.

Papillaria nigrescens
Floribundaria cardotii
Papillaria flexicaulis

P. flexicaulis (Wils.) Jaeg. (Jaeger, 1877).

Syn. *Meteoriump flexicaule* Wils. (Wilson, 1854).
Trachypus flexicaulis (Wils.) Mitt. (Mitten, 1860).
Meteoriump filipendulum Hook. f. et Wils. (Hooker and Wilson in Wilson, 1859).
Papillaria filipendula (Hook. f. et Wils.) Jaeg. (Jaeger, 1877).
Pilotrichella filipendula (Hook. f. et Wils.) Hamp. (Hampe, 1880b).

Distr. Juan Fernandez (Bartram, 1959; Robinson, 1975).
Southern South America (Paris, 1905a; Brotherus, 1925; Reimers, 1926; Herzog, 1938, 1940).

P. laeviuscula Broth.
P. media Aongstr.

Papillaria tijucae
Meteoriump medium

P. nigrescens (Hedw.) Jaeg. (Jaeger, 1877).

Syn. *Hypnum nigrescens* Hedw. (Hedwig, 1801).
Neckera appressa C. Muell. (Mueller, 1850).
Papillaria appressa (C. Muell.) Aongstr. (Ångström, 1876a).

Distr. Southern South America (Herzog, 1952a, b).

P. pilifolia C. Muell. (Mueller, 1901b).

Distr. Southern South America (Herzog, 1952a).

P. tijucae C. Muell. (Mueller, 1901b).

Syn. *Papillaria laeviuscula* Broth. (Brotherus in Ule, 1899) *nom. nud.*

Distr. Southern South America (Herzog, 1952a).

P. usneoides Broth.

Floribundaria usneoides

Pentastichella nom. nud.

P. aurea Dus.
P. bartramoides Dus.
P. jaffuelii Thér.
P. pentasticha (Mont.) C. Muell.
var. *strictifolia* Thér.

Stenomitrium aureum
Zygodon bartramoides
Zygodon porteri
Zygodon pentastichus
Zygodon pentastichus var. *strictifolius*

<i>P. robustula</i> Dus.	<i>Zygodon bartramoides</i>
<i>Perisiphorus</i>	
<i>P. ciliatus</i> (Hedw.) P. Beauv.	<i>Hedwigia ciliata</i>
<i>Peromnion</i>	
<i>P. magellanicum</i> Sull.	<i>Brachymenium indicum</i>
<i>Pharomitrium</i> nom. rejic.	
<i>P. subsessile</i> (Brid.) Schimp.	<i>Pterygoneurum subsessile</i>
<i>Phasconica</i>	
<i>P. lorentzii</i> C. Muell. (Mueller, 1882).	
Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1924a; Hilpert, 1933; Kühnemann, 1938; Hosseus, 1939).	
<i>Phascum</i>	
<i>P. brevipes</i> (Hook.) Schwaegr.	<i>Bruchia brevipes</i>
<i>P. calodictyon</i> C. Muell. (Mueller, 1888).	
Distr. Southern South America (Gibert, 1873 (as <i>P. turgidum</i> ?); Mueller, 1888; Paris, 1905a; Roth, 1911; Felippone, 1912; Brotherus, 1924a; Herter, 1928, 1933a; Kühnemann, 1938).	
<i>P. fruchartii</i> C. Muell.	<i>Astomum fruchartii</i>
<i>P. lamprocarpum</i> C. Muell.	<i>Tetrapterum lamprocarpum</i>
<i>P. lamprothecium</i> C. Muell.	<i>Tetrapterum lamprothecium</i>
<i>P. nervosum</i> Hook.	<i>Pleuridium nervosum</i>
var. <i>robinsonii</i> Mont.	<i>Pleuridium robinsonii</i>
<i>P. recurvirostre</i> C. Muell.	<i>Tetrapterum recurvirostre</i>
<i>P. robinsonii</i> Mont. ex C. Muell.	<i>Pleuridium robinsonii</i>
<i>Philibertiella</i>	
<i>P. ditrichoidea</i> Card. (Cardot, 1914b).	
Distr. Southern South America (Cardot, 1914b; Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1938, 1957).	
<i>P. nitens</i> Herz. (Herzog, 1960 [1961]).	
Distr. Southern South America (Herzog, 1960).	
<i>Philonotis</i>	
<i>P. acicularis</i> (C. Muell.) Kindb. (Kindberg, 1891).	
Syn. <i>Bartramia acicularis</i> C. Muell. (Mueller, 1890a).	
<i>Philonotis varians</i> Card. (Cardot, 1906a) <i>fide</i> Clarke, 1973a.	
<i>Philonotis elegantissima</i> Broth. ex Dism. (Brotherus ex Dismier, 1910) <i>hom. illeg.</i>	
<i>Meesia austro-georgica</i> C. Muell. (Mueller, 1890a) <i>fide</i> Clarke, 1973a.	
Distr. Southern South America (Dismier, 1910; Thériot, 1924; Matteri, 1968).	
South Georgia (Mueller, 1890a; Paris, 1905a; Cardot, 1906a, 1908a; Dismier, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Dixon, 1932; Kabiersch, 1937; Steere, 1961b; Matteri, 1968; Clarke, 1973a; Bell, 1977b).	
<i>P. acutissima</i> (C. Muell.) Kindb. (Kindberg, 1889).	
Syn. <i>Bartramia acutissima</i> C. Muell. (Mueller, 1882).	
Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1924a; Hosseus, 1938c, d, 1939; Kühnemann, 1938; Herzog, 1952a).	
<i>P. australis</i> Mitt.	<i>Conostomum pentastichum</i>

- P. balansae* Besch. ex Kindb. *Philonotis curvata*
P. balansaena Besch. *Philonotis curvata*
- P. bartramii*** Matteri (Matteri, 1973c).
 Syn. *Philonotis luteola* Bartr. (Bartram, 1946) hom. illeg.
 Distr. Southern South America (Bartram, 1946; Matteri, 1973c).
- P. brevifolia* Herz. *Philonotis krausei*
P. buenos-ayrensis (C. Muell.) Par. *Philonotis curvata*
- P. crenatula*** (C. Muell.) Kindb. (Kindberg, 1889).
 Syn. *Bartramia crenatula* C. Muell. (Mueller, 1882).
 Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1924a; Hosseus, 1935a, b, 1937; Kühnemann, 1938).
- P. curvata*** (Hamp.) Jaeg. (Jaeger, 1875).
 Syn. *Bartramia curvata* Hamp. (Hampe, 1862a).
Philonotis balansaena Besch. (Bescherelle, 1877).
Philonotis balansae Besch. ex Kindb. (Bescherelle ex Kindberg, 1888) nom. illeg.
Philonotula buenos-ayrensis C. Muell. (Mueller, 1897a).
Philonotis buenos-ayrensis (C. Muell.) Par. (Paris, 1900a).
Philonotula bonariensis C. Muell. (Mueller in Kuntze, 1898) err.
 Distr. Southern South America (Mueller, 1897a; Kuntze, 1898; Paris, 1900a, 1905a; Krieger, 1904; Felippone, 1909; Dismier, 1910; Brotherus, 1924a; Herter, 1933a).
- P. defecta* (C. Muell.) Kindb. *Philonotis glaucescens*
P. dimorpha Schimp. *Philonotis vagans*
P. elegantissima Broth. ex Dism. *Philonotis acicularis*
P. elegantula (Tayl.) Jaeg. *Philonotis rufiflora*
- P. elongatula*** (C. Muell.) Kindb. (Kindberg, 1889).
 Syn. *Bartramia elongatula* C. Muell. (Mueller, 1882).
Philonotis subelegantula Par. (Paris, 1897) nom. illeg.
Bartramia elegantula Par. (Paris, 1905a) nom. inval.
 Distr. Southern South America (Mueller, 1882; Paris, 1897, 1900a, 1905a; Brotherus, 1924a; Hosseus, 1938c, d, 1939; Kühnemann, 1938).
- P. esquelensis*** Matt. (Matteri, 1968).
 Distr. Southern South America (Matteri, 1968).
- P. exigua* (Sull.) Broth. *Philonotis scabrifolia*
- P. flexipes*** (C. Muell.) Par. (Paris, 1900a).
 Syn. *Philonotula flexipes* C. Muell. (Mueller, 1897a).
Bartramia flexipes (C. Muell.) C. Muell. (Mueller, 1901a [1900]).
 Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Felippone, 1912; Brotherus, 1924a; Herter, 1933a, 1943).
- P. fontana*** (Hedw.) Brid. (Bridel, 1827b).
 Syn. *Mnium fontanum* Hedw. (Hedwig, 1801).
Bryum fontanum (Hedw.) Gaertn., Meyer et Scherb. (Gärtner and others, 1802).
Bartramia fontana (Hedw.) Turn. (Turner, 1804a).
 Distr. Southern South America (Mueller, 1848a; Montagne, 1850).
- var. *magellanica*** (Brid.) Brid. (Bridel, 1827b).
 Syn. *Mnium fontanum* Hedw. var. *magellanicum* Brid. (Bridel, 1803).
Bartramia fontana (Hedw.) Turn. var. *magellanica* (Brid.) P. Beauv. (Palisot de Beauvois, 1805a).
 Distr. Southern South America (Bridel, 1803, 1827b).

P. fontanoides (Gill. et Grev.) Jaeg. (Jaeger, 1875).

Syn. *Bartramia fontanoides* Gill. et Grev. (Gillies and Greville *in Greville*, 1830).

Distr. Southern South America (Greville, 1830; Mitten, 1869; Jaeger, 1875; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

P. glabrata Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Robinson, 1975).

P. glaucescens (Hornsch.) Broth. (Brotherus, 1895).

Syn. *Bartramia glaucescens* Hornsch. (Hornschuch, 1840).

Philonotis muhlenbergii (Schwaegr.) Brid. var. *tenella* Brid. (Bridel, 1827b).

Bartramia tenella (Brid.) C. Muell. (Mueller, 1849) *nom. illeg.*

Philonotis tenella (Brid.) Jaeg. (Jaeger, 1875).

Philonotis sphaerocarpa (Hedw.) Brid. ssp. *tenella* (Brid.) Dism. (Dismier, 1910).

Bartramia defecta C. Muell. (Mueller, 1882).

Philonotis defecta (C. Muell.) Kindb. (Kindberg, 1889).

Bartramidula defecta (C. Muell.) Broth. (Brotherus, 1904).

Philonotis weberoides Dus. (Dusén *in Dismier*, 1910).

Distr. Southern South America (Mueller, 1882; Brotherus, 1904, 1924a; Paris, 1905a; Dismier, 1910; Hosseus, 1938b, c, d, 1939; Kühnemann, 1938; Bizot and Piovano, 1953; Piovano, 1954).

P. gourdonii Card. (Cardot, 1911b).

Distr. Antarctic, peninsula region (Cardot, 1911b, d, 1913a; Brotherus, 1924a; Steere, 1961a; Matteri, 1968; Robinson, 1972).

P. inflexa (C. Muell.) Broth.

Pohlia inflexa

P. kerguelensis Par.

Breutelia integrifolia

P. krauseana Broth.

Philonotis krausei

var. longinervis Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b).

P. krausei (C. Muell.) Broth. (Brotherus, 1904).

Syn. *Bartramia krausei* C. Muell. (Mueller, 1874).

Glyphocarpus krausei (C. Muell.) Jaeg. (Jaeger, 1874).

Bartramidula krausei (C. Muell.) Par. (Paris, 1875).

Philonotis krauseana Broth. (Brotherus, 1894).

Philonotis brevifolia Herz. (Herzog, 1924b) *nom. illeg.*

Distr. Juan Fernandez (Brotherus, 1924b) *fide* Seki, 1974.

Southern South America (Mueller, 1874; Bartram, 1894; Matteri, 1968; Robinson, 1975, 1904, 1924a; Thériot, 1924, 1934b; Jaeger, 1875; Paris, 1894, 1904a; Brotherus, 1940, 1954; Matteri, 1968; Seki, 1974).

P. litorea Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Juan Fernandez (Skottsberg, 1914). Falkland Islands (Cardot and Brotherus, 1923).

var. prolifera Card. (Cardot *in Skottsberg*, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914) *nom. nud.*

P. luteola Bartr.

philonotis bartramii

P. macropyxis Dus. (Dusén *in Paris*, 1905a) *nom. nud.*

Distr. Southern South America (Paris, 1905a).

P. muhlenbergii (Schwaegr.) Brid. var. *tenella* Brid.

Philonotis glaucescens

P. nigricans Herz. (Herzog, 1916b).

Distr. Southern South America (Herzog, 1916b).

P. nigroflava C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Brotherus, 1924a; Paris, 1900a, 1905a; Cardot, 1908a; Cardot, 1938; Matteri, 1968).

P. parallela Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Paris, 1905a; Cardot, 1908a; Dismier, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1957; Kühnemann, 1938; Matteri, 1968; Seki, 1974).

P. platensis Par. (Paris, 1900a).

Syn. *Philonotula secunda* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1924a; Kühnemann, 1938).

P. pomangium (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Bartramia pomangium* C. Muell. (Mueller, 1882).

Bartramia declivium C. Muell. (Mueller, 1882) nom. nud.

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

P. riparia Dus. (Dusén in Paris, 1905a) nom. nud.

Distr. Southern South America (Paris, 1905a; Dismier, 1910).

P. rufiflora (Hornschr.) Reichdt. (Reichardt, 1870).

Syn. *Bartramia rufiflora* Hornsch. (Hornschuch, 1840).

Bartramia elegantula Tayl. (Taylor, 1847).

Philonotis elegantula (Tayl.) Jaeg. (Jaeger, 1875).

Distr. Southern South America (Taylor, 1847; Sullivant, 1859; Jaeger, 1875).

P. scabrifolia (Hook. f. et Wils.) Braithw. (Braithwaite, 1893).

Syn. *Hypnum scabrifolium* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Bartramia scabrifolia (Hook. f. et Wils.) C. Muell. (Mueller, 1901a [1900]).

Bartramia pusilla Sull. (Sullivant, 1850) hom. illeg.

Bartramia exigua Sull. (Sullivant, 1859).

Glyphocarpus exiguus (Sull.) Jaeg. (Jaeger, 1875).

Bartramidula exigua (Sull.) Par. (Paris, 1894).

Philonotis exigua (Sull.) Broth. (Brotherus, 1904).

Bartramia willii C. Muell. (Mueller, 1890a).

Philonotis willii (C. Muell.) Broth. (Brotherus, 1904).

Bartramia ventanae C. Muell. (Mueller, 1897a).

Philonotis ventanae (C. Muell.) Par. (Paris, 1900a).

Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Sullivant, 1850, 1859; Mueller, 1851a, 1885, 1897a, 1901a; Jaeger, 1875; Paris, 1894, 1900a, 1904a, 1905a; Neger, 1899; Dusén, 1903a; Brotherus, 1904, 1924a; Cardot, 1905a, 1908a, 1913a; Dismier, 1910; Cardot and Brotherus, 1923; Herzog, 1923, 1939, 1954, 1957; Roivainen, 1934; Hosseus, 1938c, d, 1939; Kühnemann, 1938; Bartram, 1952; Matteri, 1968; Seki, 1974).

Falkland Islands (Cardot, 1905a, 1908a; Brotherus, 1924a; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Brotherus, 1904, 1924a; Paris, 1904a; Cardot, 1906a, 1908a; Dismier, 1910; Steere, 1961b; Matteri, 1968; Clarke, 1973a).

P. simplicissima (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Bartramia simplicissima* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Dismier, 1910; Kühnemann, 1938).

P. sphaerocarpa (Hedw.) Brid. ssp. *tenella* (Brid.) Dism.**Philonotis glaucescens****P. striata** (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Bartramia striata* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905a; Dismier, 1910; Brotherus, 1924a; Kühnemann, 1938).

P. strictiuscula (C. Muell.) Par. (Paris, 1900a).

Syn. *Philonotula strictiuscula* C. Muell. (Mueller, 1897a).

Bartramia strictiuscula (C. Muell.) C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1897a, 1901a; Paris, 1900a; Brotherus, 1924a; Herter, 1933a).

P. subelegantula Pär.

Philonotis elongatula

P. tenella (Brid.) Jaeg.

Philonotis glaucescens

P. trichophylla Herz. (Herzog, 1954) hom. illeg.

Distr. Southern South America (Herzog, 1954).

P. vagans (Hook. f. et Wils.) Mitt. (Mitten, 1860).

Syn. *Bryum vagans* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Meesia vagans (Hook. f. et Wils.) C. Muell. (Mueller, 1848a).

Bartramia vagans (Hook. f. et Wils.) Mitt. (Mitten, 1869).

Philonotis vagans (Hook. f. et Wils.) Mitt. var. *inundata* Card. (Cardot, 1906a) *fide* Clarke, 1973a.

Hypnum reticulatum Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844) hom. illeg.

Goniobryum reticulatum Lindb. (Lindberg, 1868b).

Rhizogonium reticulatum (Hook. f. et Wils.) Mitt. (Mitten, 1869) hom. illeg.

Mnium reticulatum (Hook. f. et Wils.) C. Muell. (Mueller, 1885).

Philonotis dimorpha Schimp. (Schimper in Mitten, 1860) nom. nud.

Bartramia dimorpha Schimp. (Schimper in Mueller, 1901a [1900]) nom. nud.

Distr. Juan Fernandez (Espinosa B., 1941; Matteri, 1968; Robinson, 1975).

Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Sullivant, 1859; Mitten, 1860, 1869; Ångström, 1872; Jaeger, 1875; Bescherelle, 1889; Neger, 1899; Dusén, 1903a, 1905d; Paris, 1904b, 1905a; Cardot, 1905a, 1908a, 1913a; Dismier, 1910; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen, 1934; Thériot, 1934b; Kühnemann, 1938; Herzog, 1939, 1954; Bartram, 1952; Matteri, 1968; Seki, 1974).

South Georgia (Cardot, 1906a, 1908a; Dismier, 1910; Cardot and Brotherus, 1923; Dixon, 1935; Steere, 1961b; Matteri, 1968; Clarke, 1973a).

var. *evanidinervis* Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959).

var. *inundata* Card.

Philonotis vagans

P. varians Card.

Philonotis acicularis

P. ventanae (C. Muell.) Par.

Philonotis scabrifolia

P. weberoides Dus.

Philonotis glaucescens

P. willii (C. Muell.) Broth.

Philonotis scabrifolia

Philonotula

P. bonariensis C. Muell.

Philonotis curvata

P. buenos-ayrensis C. Muell.

Philonotis curvata

P. flexipes C. Muell.

Philonotis flexipes

P. secunda C. Muell.

Philonotis platensis

P. strictiuscula C. Muell.

Philonotis strictiuscula

Photinophyllum hom. illeg.

P. subbasilare (Hook.) Mitt.

Goniobryum subbasilare

Phyllogonium

P. callichroum Mont.

Catagonium politum var. **phyllogonium**

Physcomitrium

P. acutifolium Broth. (Brotherus, 1900).

Distr. Southern South America (Herter, 1933a).

P. argentinicum Par. (Paris, 1897).

Syn. *Physcomitrium pusillum* C. Muell. (Mueller, 1882) *hom. illeg.*
Physcomitrium subpusillum C. Muell. (Mueller *in Kindberg*, 1889) *nom. nud.*

Distr. Southern South America (Mueller, 1882; Paris, 1897, 1905a; Brotherus, 1924a; Kühnemann, 1938).

P. badium Broth. (Brotherus, 1900).

Distr. Southern South America (Herter, 1933a; Herzog, 1954).

P. bonplandii (Hook.) De Not.**Funaria bonplandii****P. chlorodictyon** C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

P. cryptotheca C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

P. cupulare C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938; Herter, 1939b).

var. *latifolium* Broth. (Brotherus, 1918).

Distr. Southern South America (Brotherus, 1918; Kühnemann, 1938).

P. densirete Besch. (Bescherelle *in Mueller*, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

P. excurrens Broth. (Brotherus *in Krieger*, 1904) *nom. nud.*

Distr. Southern South America (Krieger, 1904).

P. felipponei Thér. (Thériot, 1939 [1941]).

Distr. Southern South America (Herter, 1939b; Thériot, 1939).

P. germanillae C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905a; Brotherus, 1924a; Hosseus, 1935a; Kühnemann, 1938; Herter, 1939b).

P. gibertii Mitt. (Mitten, 1885).

Distr. Southern South America (Mitten, 1885).

P. herteri Thér. (Thériot, 1939 [1941]).

Distr. Southern South America (Herter, 1939b; Thériot, 1939).

P. integrifolium C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

P. lorentzii C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Krieger, 1904; Kurtz, 1904; Paris, 1905a; Brotherus, 1918, 1924a; Herter, 1933a; Thériot, 1934b; Hosseus, 1935b, 1937, 1938a, c, d; Kühnemann, 1938).

P. macroglobum C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

P. mahui H. Robinson (Robinson, 1970b).

Distr. Southern South America (Robinson, 1970b).

P. orbignyanum (Mont.) Mont. (Montagne, 1839c).

Syn. *Gymnostomum orbignyanum* Mont. (Montagne, 1838).

Distr. Southern South America (Montagne, 1838, 1856; Paris, 1905a; Brotherus, 1924a; Kühnemann, 1938).

P. pusillum C. Muell.

Physcomitrium argentinicum

P. pyriforme (Hedw.) Hamp. (Hampe, 1837b).

Syn. *Gymnostomum pyriforme* Hedw. (Hedwig, 1801).

Distr. Southern South America (Montagne, 1850).

P. roseae Williams (Williams, 1915).

Distr. Southern South America (Williams, 1915; Brotherus, 1924a).

P. serrifolium C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905a; Brotherus, 1924a; Hosseus, 1935b, 1937, 1938a, c, d; Kühnemann, 1938).

P. spegazzinii C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

P. subpusillum C. Muell.

Physcomitrium argentinicum

P. thieleanum Hamp. (Hampe, 1844).

Syn. *Physcomitrium thielei* Hamp. ex Kindb. (Hampe ex Kindberg, 1888) *nom. illeg.*

Distr. Southern South America (Hampe, 1844; Mueller, 1848a; Mitten, 1869; Gibert, 1873; Jaeger, 1874; Herter, 1928, 1933a; Herzog, 1952a).

P. thielei Hamp. ex Kindb.

Physcomitrium thieleanum

P. umbonatum Mitt. (Mitten, 1869).

Distr. Southern South America (Felippone, 1909; Herter, 1933a).

Pilopogon

P. gracilis (Hook.) Brid. (Bridel, 1826).

Syn. *Didymodon gracilis* Hook. (Hooker, 1818).

Thysanomitrion gracile (Hook.) Arnott (Arnott, 1827).

Weisia gracilis (Hook.) Spreng. (Sprengel, 1827).

Dicranum strictum Schwaegr. (Schwaegrichen in Gaudichaud, 1826 [1828]) *hom.*

illeg.

Didymodon gracilis Hook. var. *robustum* Hook. (Hooker in Gaudichaud, 1826 [1828])

nom. nud.

Thysanomitrion gracile (Hook.) Arnott var. *robustum* Arnott (Arnott, 1827).

Distr. Falkland Islands (Dumont D'Urville, 1825; Gaudichaud, 1825, 1826; Bridel, 1827b).

P. leptodus (Mont.) Broth.

Campylopus clavatus

P. richardii (Brid.) Broth.

Campylopus richardii

P. schilleri Herz. et Thér. (Herzog and Thériot in Herzog and others, 1933).

Distr. Southern South America (Herzog and others, 1933).

Pilotrichella

P. avellanedae (C. Muell.) Kindb.

Orthostichopsis avellanedae

P. billardieri (Hamp.) Jaeg.

Weymouthia cochlearifolia* var. *billardieri

P. characea (C. Muell.) Kindb.

Meteoriopsis characea

var. *nigrescens* (C. Muell.) Par.

Meteoriopsis characea* var. *nigrescens

var. *tenuis* (C. Muell.) Par.

Meteoriopsis characea* var. *tenuis

P. cumingii (C. Muell.) Lor.

Weymouthia mollis

P. cyathipoma (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Neckera cyathipoma* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

<i>P. filipendula</i> (Hook. f. et Wils.) Hamp.	<i>Papillaria flexicaulis</i>
<i>P. illecebrina</i> (C. Muell.) Kindb.	<i>Meteoriopsis illecebrinum</i>
<i>P. kraussei</i> Lor.	<i>Weymouthia cochlearifolia</i> var. <i>billardieri</i>
<i>P. mollis</i> (Hedw.) Jaeg.	<i>Weymouthia mollis</i>
<i>P. turgidula</i> (C. Muell.) Par.	<i>Squamidium turgidulum</i>
var. <i>tristis</i> (C. Muell.) Par.	<i>Squamidium turgidulum</i> var. <i>triste</i>

Pilotrichum

<i>P. ambiguum</i> Hornsch.	<i>Meteoriopsis ambigua</i>
<i>P. ancistrodes</i> (Mont.) Lor.	<i>Ancistrodes genuflexa</i>
<i>P. ciliatum</i> (Hedw.) C. Muell.	<i>Hedwigia ciliata</i>
<i>P. consimile</i> (Mont.) C. Muell.	<i>Cryphaea tenella</i>
<i>P. flagelliferum</i> Brid.	<i>Pseudocryphaea flagellifera</i>
<i>P. genuflexum</i> C. Muell.	<i>Ancistrodes genuflexa</i>
<i>P. gorveanum</i> (Mont.) C. Muell.	<i>Dendrocryphaea gorveana</i>
<i>P. patagonicum</i> (Brid.) P. Beauv.	<i>Homalia glabella</i>
<i>P. pentagonum</i> Hamp. et Lor.	<i>Orthostichidium pentagonum</i>
<i>P. polytrichoides</i> (Hedw.) Brid.	<i>Lepidopilum polytrichoides</i>
<i>P. scabrisetum</i> (Schwaegr.) Brid.	<i>Lepidopilum scabrisetum</i>
<i>P. smithii</i> (Dicks.) P. Beauv.	<i>Leptodon smithii</i>
<i>P. stramineum</i> C. Muell.	<i>Meteoriopsis straminea</i>
<i>P. tenellum</i> (Schwaegr.) C. Muell.	<i>Cryphaea tenella</i>
<i>P. thieleeanum</i> (C. Muell.) Hamp.	<i>Pinnatella thieleana</i>

Pinnatella

P. callicostelloides (Thér.) Broth. (Brotherus, 1925).

Syn. *Porotrichum callicostelloides* Broth. ex Thér. (Brotherus ex Thériot, 1917b).

Distr. Southern South America (Paris, 1905b; Thériot, 1917b, 1926; Costes, 1921; Brotherus, 1925; Espinosa B., 1941; Herzog, 1954).

var. ***distichum*** Thér. (Thériot, 1924).

Distr. Southern South America (Thériot, 1924).

P. macrosticta Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Robinson, 1975).

P. pinnatelloides (C. Muell.) Broth. ex Par.

Pinnatella thieleana

P. thieleana (C. Muell.) Broth. ex Par. (Brotherus ex Paris, 1909).

Syn. *Hypnum thieleanum* C. Muell. (Mueller, 1851a).

Porotrichum thieleanum (C. Muell.) Mitt. (Mitten, 1869).

Pilotrichum thieleanum (C. Muell.) Hamp. (Hampe, 1879).

Porotrichum thielei Aongstr. (Ångström, 1876a) hom. illeg.

Thamnium thielei Kindb. (Kindberg, 1902) nom. illeg.

Porotrichum pinnatelloides C. Muell. (Mueller, 1879a).

Pinnatella pinnatelloides (C. Muell.) Broth. ex Par. (Brotherus ex Paris, 1909).

Distr. Southern South America (Hampe, 1879; Mueller, 1879a; Kindberg, 1902; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

Plagiobryum

P. lamprocarpum (C. Muell.) Par.

Bryum lamprocarpum

Plagiommium

P. rugicum (Laur.) Koponen (Koponen, 1968).

Syn. *Mnium rugicum* Laur. (Laurer, 1827).

Mnium affine Bland. ex Funck var. *rugicum* (Laur.) B.S.G. (Bruch and others, 1838).

Mnium affine Bland. ex Funck var. *magellanicum* Card. et Broth. (Cardot and Brotherus, 1923).

Mnium lepto-limbatum C. Muell. (Mueller, 1897a) *fide* Koponen, 1971.

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Cardot, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938; Koponen, 1971).

Plagiopus

P. ithyphyllus (Brid.) Mach.

Bartramia ithyphylla

P. pomiformis (Hedw.) Mach.

Bartramia pomiformis

P. strictus (Brid.) Mach.

Bartramia stricta

Plagiothecium

P. bellirete C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905b; Brotherus, 1925; Kühnemann, 1938).

P. cacti (C. Muell.) Kindb.

Isopterygium cacti

P. denticulatum (Hedw.) B.S.G. (Bruch and others, 1851).

Syn. *Hypnum denticulatum* Hedw. (Hedwig, 1801).

Stereodon denticulatus (Hedw.) Mitt. (Mitten, 1863).

Pancovia denticulata (Hedw.) Kickx (Kickx, 1867).

Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1851a, 1885; Sullivant, 1859; Cardot, 1908a, 1913a; Roivainen, 1934).

var. ***obtusifolium*** (Turn.) Moore (Moore, 1873).

Syn. *Hypnum denticulatum* Hedw. var. *obtusifolium* Turn. (Turner, 1804a).

Hypnum obtusifolium (Turn.) Brid. (Bridel, 1812).

Plagiothecium obtusifolium (Turn.) Amann (Amann, 1928) *nom. illeg.*

Hypnum donnianum Sm. (Smith, 1804b).

Plagiothecium donnianum (Sm.) Mitt. (Mitten, 1869).

Distr. Southern South America (Mitten, 1869; Jaeger, 1878; Paris, 1905b).

P. donnianum (Sm.) Mitt.

Plagiothecium denticulatum* var. *obtusifolium

P. elegans (Brid.) Sull. var. *nanum* (Jur.) Mol.

Isopterygium elegans* var. *nanum

P. fuegianum (Besch.) Dus.

Isopterygium fuegianum

P. georgico-antarcticum (C. Muell.) Kindb. (Kindberg, 1891).

Syn. *Hypnum georgico-antarcticum* C. Muell. (Mueller, 1890a).

Distr. South Georgia (Mueller, 1890a; Paris, 1905b; Cardot, 1908a; Brotherus, 1925; Steere, 1961b).

P. leptoplumosum Dus.

Isopterygium leptoplumosum

P. leucophyllum Hamp.

Isopterygium leucophyllum

P. lucidulum (Hook. f. et Wils.) Mitt. (Mitten, 1869).

Syn. *Hypnum lucidulum* Hook. f. et Wils. (Hooker and Wilson *in* Wilson and Hooker, 1847).

Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1851a, 1885; Mitten, 1869; Jaeger, 1878; Paris, 1905b; Kühnemann, 1938).

Falkland Islands (Paris, 1905b; Kühnemann, 1938).

P. lucidum (Hook. f. et Wils.) Par. (Paris, 1896).

Syn. *Hypnum lucidum* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Amblystegium lucidum (Hook. f. et Wils.) Jaeg. (Jaeger *in* Jaeger and Sauerbeck, 1879a).

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Mueller, 1851a; Jaeger and Sauerbeck, 1879a; Paris, 1896, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).
Falkland Islands (Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

<i>P. magellanicum</i> Schimp.	<i>Acrocladium auriculatum</i>
<i>P. magellanicum</i> Par.	<i>Isopterygium fuegianum</i>
<i>P. nanum</i> Jur.	<i>Isopterygium elegans</i> var. <i>nanum</i>
<i>P. nitidum</i> (Hook. f. et Wils.) Bartr.	<i>Catagonium nitidum</i>
<i>P. obtusifolium</i> (Turn.) Amann	<i>Plagiothecium denticulatum</i> var. <i>obtusifolium</i>

***P. orthocarpum* Mitt. (Mitten, 1869).**

Syn. *Isothecium orthocarpum* Brid. (Bridel, 1827b) *nom. nud.*
Hypnum orthocarpon Hornsch. ex C. Muell. (Hornschuch ex Mueller, 1851a) *hom. illeg.*

Distr. Southern South America (Hornschuch, 1820b; Bridel, 1827b; Mueller, 1851a; Mitten, 1869; Jaeger, 1878; Paris, 1905b).

***P. ovatifolium* Card. (Cardot, 1905a).**

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Seki, 1974).

***P. roseanum* B.S.G. var. *falklandicum* Card. et Broth. (Cardot and Brotherus, 1923).**

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

P. schimperi* Jur. et Mild. var. *nanum* (Jur.) Walth. et Mol. *Isopterygium elegans* var. *nanum

***P. simonovii* Savicz. et Smirn. (Savich-Ljubitskaya and Smirnova, 1964a).**

Distr. Antarctic, continental region (Savich-Ljubitskaya and Smirnova, 1964a; Greene, 1968a; Savich-Ljubitskaya, 1978).

P. subsimplex* (Hedw.) Besch. *Sematophyllum subsimplex

Platyhypnidium

***P. aquaticum* (Jaeg.) Fleisch. (Fleischer, 1923).**

Syn. *Rhynchostegium aquaticum* Jaeg. (Jaeger, 1878).

Distr. Southern South America (Hosseus, 1938c, 1939).

***P. riparioides* (Hedw.) Dix. (Dixon, 1933 [1934]).**

Syn. *Hypnum riparioides* Hedw. (Hedwig, 1801).

Hypnum rusciforme Brid. (Bridel, 1801) *nom. illeg.*

Eurhynchium riparioides (Hedw.) Richs. (Richards, 1937).

Eurhynchium rusciforme Mild. (Milde, 1869) *nom. illeg.*

Distr. Southern South America (Bridel, 1827b).

Platylomidium* nom. *nud.

P. crispatula* C. Muell. *Calyptopogon mnioides

Platyneurum

***P. laticostatum* (Card.) Broth. (Brotherus, 1924a).**

Syn. *Dicranum laticostatum* Card. (Cardot, 1900).

Dicranum cirrhifolium Schimp. ex Dus. (Schimper ex Dusén, 1903a).

Distr. Southern South America (Mueller, 1885; Neger, 1899; Cardot, 1900, 1901, 1905a, 1908a; Dusén, 1903a, 1905a; Paris, 1904b; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen and Bartram, 1937; Herzog, 1938, 1940, 1954, 1957; Kühnemann, 1938; Bartram, 1952; Bizot, 1967; Seki, 1974).

South Georgia (Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b; Bell, 1973b).

Pleuridiopsis nom. nud.**P. amblyocalyx** (C. Muell.) Par.**Astomiopsis amblyocalyx****P. subulata** (C. Muell.) Par.**Astomiopsis subulata****Pleuridium****P. costesii** Thér. (Thériot, 1921a).*Distr.* Southern South America (Costes, 1921; Thériot, 1921a; Brotherus, 1924a).**P. macrothecium** Dus. (Dusén, 1905a).*Distr.* Southern South America (Dusén, 1905a; Roth, 1911; Campo, 1921; Thériot, 1921a; Brotherus, 1924a).**P. nervosum** (Hook.) Mitt. (Mitten, 1856).*Syn.* *Phascum nervosum* Hook. (Hooker, 1819).*Astomum nervosum* (Hook.) C. Muell. (Mueller, 1847b).*Distr.* Juan Fernandez (Mueller, 1856; Johow, 1896; Roth, 1911).

Southern South America (Mueller, 1847b, 1848a, 1856; Montagne, 1850; Roth, 1911).

P. robinsonii (Mont.) Mitt. (Mitten, 1869).*Syn.* *Phascum nervosum* Hook. var. *robinsonii* Mont. (Montagne, 1835).*Phascum robinsonii* Mont. ex C. Muell. (Montagne ex Mueller, 1848a) err.*Astomum robinsonii* (Mont.) C. Muell. (Mueller, 1856).*Distr.* Juan Fernandez (Montagne, 1835; Mueller, 1856; Mitten, 1869; Gibert, 1873; Jaeger, 1873; Paris, 1905b; Brotherus, 1924b; Robinson, 1975).

Southern South America (Montagne, 1835; Mueller, 1856; Jaeger, 1869, 1873; Mitten, 1869; Gibert, 1873; Dusén, 1905a; Paris, 1905b; Felippone, 1909; Thériot, 1921a, b; Brotherus, 1924a; Herter, 1928, 1933a, 1939a; Herzog, 1938, 1954; Bartram, 1952).

P. subnervosum (C. Muell.) Jaeg. (Jaeger in Paris, 1896).*Syn.* *Astomum subnervosum* C. Muell. (Mueller, 1882).*Distr.* Southern South America (Jaeger and Sauerbeck, 1879b; Mueller, 1882; Paris, 1896, 1905b; Roth, 1911; Brotherus, 1924a; Herter, 1933a; Kühnemann, 1938).**Pleurochaete****P. ecuadorensis** Broth. (Brotherus, 1920).*Distr.* Southern South America (Hilpert, 1933).**Pleurorthotrichum****P. chilense** Broth. (Brotherus, 1905c).*Distr.* Southern South America (Brotherus, 1905c, 1924c, 1925; Espinosa B., 1941; Skottsberg, 1950; Herzog, 1954).**Pleurozium****P. schreberi** (Brid.) Mitt. (Mitten, 1869).*Syn.* *Hypnum schreberi* Brid. (Bridel, 1801b).*Hylocomium schreberi* (Brid.) De Not. (De Notaris, 1867).*Hypnopsis schreberi* (Brid.) Podp. (Podpěra, 1921).*Entodon schreberi* (Brid.) Moenk. (Mönkermeyer, 1927).*Cuspidaria morenoi* C. Muell. (Mueller, 1897a).*Hypnum morenoi* (C. Muell.) Par. (Paris, 1900a).*Acrocladium morenoi* (C. Muell.) Dus. (Dusén, 1903a).*Hypnum parietinum* Wahlenb. (Wahlenberg, 1812) hom. illeg.*Distr.* Southern South America (Mueller, 1897a; Paris, 1900a, 1904c; Dusén, 1903a; Cardot, 1908a; Brotherus, 1924a, 1925).**Pogonatum****P. alpinum** (Hedw.) Roehl.**Polytrichum alpinum**

<i>P. brevifolium</i> (R. Brown) Brid.	<i>Polytrichum alpinum</i>
<i>P. austro-georgicum</i> (C. Muell.) Par.	<i>Polytrichum alpinum</i>
<i>P. dendroides</i> (Hedw.) Brid.	<i>Dendroligotrichum dendroides</i>
<i>P. liliputanum</i> Thér.	<i>Oligotrichum canaliculatum</i>
<i>P. magellanicum</i> (Hedw.) P. Beauv.	<i>Polytrichadelphus magellanicus</i>
<i>P. oligodus</i> (C. Muell.) Mitt. (Mitten, 1869).	
<i>Syn.</i> <i>Polytrichum oligodus</i> Kunz. ex C. Muell. (Kunze ex Mueller, 1848a). <i>Cephalotrichum oligodus</i> B.S.G. (Bruch and others, 1844) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Bruch and others, 1844; Mueller, 1848a; Montagne, 1850; Mitten, 1869; Jaeger, 1875; Paris, 1905b; Thériot, 1915, 1917b, 1921b; Campo, 1921; Costes, 1921; Brotherus, 1925; Bartram, 1952).	
<i>P. plurisetum</i> (C. Muell.) Broth. (Brotherus, 1905a).	
<i>Syn.</i> <i>Polytrichum plurisetum</i> C. Muell. (Mueller, 1879a).	
<i>Distr.</i> Southern South America (Mueller, 1879a; Kurtz, 1904; Brotherus, 1905a, 1925; Hosseus, 1936, 1937; Kühnemann, 1938).	
<i>var. brevipes</i> (C. Muell.) O. Kühnemann (Kühnemann, 1938).	
<i>Syn.</i> <i>Polytrichum plurisetum</i> C. Muell. var. <i>brevipes</i> C. Muell. (Mueller, 1879a). <i>Pogonatum polycarpum</i> Par. var. <i>brevipes</i> (C. Muell.) Par. (Paris, 1896).	
<i>Distr.</i> Southern South America (Mueller, 1879a; Paris, 1896, 1905b; Kühnemann, 1938).	
<i>P. polycarpum</i> Par. var. <i>brevipes</i> (C. Muell.) Par.	<i>Pogonatum plurisetum</i> var. <i>brevipes</i>
<i>P. robustum</i> (Lindb.) Jaeg.	<i>Polytrichadelphus robustus</i>
<i>P. semi-angulatum</i> Brid.	<i>Polytrichadelphus semi-angulatus</i>
<i>P. spurio-cirratum</i> Broth. (Brotherus, 1910a).	
<i>Distr.</i> Southern South America (Thériot, 1926).	
<i>Pohlia</i>	
<i>P. albicans</i> Lindb.	<i>Pohlia wahlenbergii</i>
<i>P. calopyxis</i> (C. Muell.) Broth. (Brotherus, 1903).	
<i>Syn.</i> <i>Bryum calopyxis</i> C. Muell. (Mueller, 1882). <i>Webera calopyxis</i> (C. Muell.) Kindb. (Kindberg, 1889).	
<i>Distr.</i> Southern South America (Mueller, 1882; Brotherus, 1903, 1924a; Paris, 1906; Kühnemann, 1938).	
<i>P. clavata</i> Schimp.	<i>Bryum clavatum</i>
<i>P. commutata</i> (Schimp.) Lindb.	<i>Pohlia drummondii</i>
<i>P. cruda</i> (Hedw.) Lindb. (Lindberg, 1879).	
<i>Syn.</i> <i>Mnium crudum</i> Hedw. (Hedwig, 1801). <i>Hypnum crudum</i> (Hedw.) Web. et Mohr (Weber and Mohr, 1803). <i>Bryum crudum</i> (Hedw.) Turn. (Turner, 1804a). <i>Webera cruda</i> (Hedw.) Fuernr. (Fünnrohr, 1829). <i>Lamprophyllum crudum</i> (Hedw.) Lindb. (Lindberg, 1871). <i>Bryum synoico-crudum</i> C. Muell. (Mueller, 1884 [1883]). <i>Webera synoico-cruda</i> (C. Muell.) Kindb. (Kindberg, 1889). <i>Bryum viridatum</i> C. Muell. (Mueller, 1890a). <i>Webera viridata</i> (C. Muell.) Kindb. (Kindberg, 1891).	
<i>Distr.</i> Southern South America (Mitten, 1869; Mueller, 1884, 1885, 1889; Neger, 1899; Cardot, 1900, 1901, 1905a, 1908a, 1913a; Dusén, 1903a; Paris, 1906; Cardot and Brotherus, 1923; Herzog, 1923, 1954, 1957; Brotherus, 1924a; Roivainen, 1934; Kühnemann, 1938; Bartram, 1965; Seki, 1974). South Georgia (Mueller, 1890a; Cardot, 1906a, 1908a; Paris, 1906; Dixon, 1935; Steere, 1961b; Clarke, 1973a).	

var. *imbricata* (Card.) Bartr. (Bartram, 1957).

Syn. *Webera cruda* (Hedw.) Fuernr. var. *imbricata* Card. (Cardot, 1900).

Distr. South Georgia (Clarke, 1973a).

Antarctic, peninsula region (Cardot, 1900, 1901, 1906b, 1907c, 1908a, 1911b, d, 1913a; Paris, 1906; Bartram, 1957; Steere, 1961a; Greene and others, 1970; Robinson, 1972 (as *P. cruda*)).

P. cruegeri (Hamp.) Andrews

Bryum cruegeri

P. drummondii (C. Muell.) Andrews (Andrews, 1935).

Syn. *Bryum drummondii* C. Muell. (Mueller, 1862).

Webera drummondii (C. Muell.) Jaeg. (Jaeger, 1875).

Webera commutata Schimp. (Schimper, 1876).

Pohlia commutata (Schimp.) Lindb. (Lindberg, 1879).

Bryum commutatum (Schimp.) Boul. (Boulay, 1884) *hom. illeg.*

Distr. Southern South America (Cardot and Brotherus, 1923).

P. filiformis (Dicks.) Andrews

Anomobryum filiforme

P. gilliesii (Hook.) Mont.

Bryum gilliesii

P. humilis (Mont.) Broth. (Brotherus, 1903).

Syn. *Bryum humile* Mont. (Montagne, 1845a).

Orthodontium humile (Mont.) Mitt. (Mitten, 1869).

Webera humilis (Mont.) Broth. (Brotherus, 1924a).

Orthodontium tenue C. Muell. (Mueller, 1848a) *nom. illeg.*

Apalodium tenue Mitt. (Mitten, 1885) *nom. illeg.*

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1848a; Mitten, 1869, 1885; Jaeger, 1875; Brotherus, 1903, 1924a; Paris, 1905a).

P. inclinata Sm. ex Brid.

Bryum inclinatum

P. inflexa (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1965).

Syn. *Bryum inflexum* C. Muell. (Mueller, 1890a).

Webera inflexa (C. Muell.) Kindb. (Kindberg, 1891).

Philonotis inflexa (C. Muell.) Broth. (Brotherus, 1904).

Distr. South Georgia (Mueller, 1890a; Brotherus, 1904; Paris, 1905a; Cardot, 1906a, 1908a; Steere, 1961b; Clarke, 1973a, b).

P. lonchochaete (Dus.) Broth. (Brotherus, 1909).

Syn. *Webera lonchochaete* Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Brotherus, 1909, 1924a; Kühnemann, 1938).

P. nevadensis (C. Muell.) Broth. (Brotherus, 1903).

Syn. *Bryum nevadense* C. Muell. (Mueller, 1879a).

Webera nevadensis (C. Muell.) Kindb. (Kindberg, 1889).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1903, 1924a; Paris, 1906; Kühnemann, 1938).

P. nutans (Hedw.) Lindb. (Lindberg, 1879).

Syn. *Webera nutans* Hedw. (Hedwig, 1801).

Bryum nutans (Hedw.) Turn. (Turner, 1804a).

Lamprophyllum nutans (Hedw.) Lindb. (Lindberg, 1871).

Bryum sphagnadelphus C. Muell. (Mueller, 1885).

Webera sphagnadelphus (C. Muell.) Besch. (Bescherelle, 1889).

Bryum pulvinatum C. Muell. (Mueller, 1890a) *hom. illeg. fide* Clarke, 1973a.

Webera pulvinata Kindb. (Kindberg, 1891) *fide* Clarke, 1973a.

Pohlia pulvinata (Kindb.) Broth. (Brotherus, 1903) *fide* Clarke, 1973a.

Bryum caespitosum (Hopp. et Hornsch.) Brid. (Bridel, 1826).

Distr. Southern South America (Wilson and Hooker, 1847; Montagne, 1850; Sullivant, 1859; Mitten, 1869; Mueller, 1885; Bescherelle, 1889; Paris, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Thériot, 1924; Kühnemann, 1938; Herzog, 1954; Seki, 1974).
 Falkland Islands (Wilson and Hooker, 1847; Cardot and Brotherus, 1923).
 South Georgia (Mueller, 1890a; Brotherus, 1903, 1924a; Cardot, 1906a, 1908a; Paris, 1906; Dixon, 1934; Steere, 1961b; Clarke, 1973a).
 Antarctic, peninsula region (Cardot, 1901, 1907c, 1908a, 1911b, d, 1913a; Brotherus, 1924a; Bartram, 1957; Steere, 1961a; Robinson, 1972).
 Antarctic, continental region (Savich-Ljubitskaya and Smirnova, 1971; Savich-Ljubitskaya, 1978).

P. papilloosa (Jaeg.) Broth. (Brotherus, 1903).

Syn. *Bryum papillosum* C. Muell. (Mueller, 1848a) *hom. illeg.*
Brachymenium papillosum Jaeg. (Jaeger, 1875).

Distr. Southern South America (Bartram, 1965).

P. platyphylla Schwaegr.

Bryum platyphyllum

P. pluriseta Herz. (Herzog, 1909).

Distr. Southern South America (Robinson, 1964).

P. polycarpa (Mitt.) Broth. (Brotherus, 1903).

Syn. *Bryum polycarpum* Mitt. (Mitten, 1869).
Webera polycarpa (Mitt.) Jaeg. (Jaeger, 1875).
Webera costesii Card. et Thér. (Cardot and Thériot *in* Thériot, 1917b).

Distr. Southern South America (Thériot, 1917b, 1924; Costes, 1921; Brotherus, 1924a).

P. pulvinata (Kindb.) Broth.

Pohlia nutans

P. racovitzae (Card.) Broth. (Brotherus, 1903).

Syn. *Webera racovitzae* Card. (Cardot, 1900).

Distr. South Georgia (Cardot, 1900).

Antarctic, peninsula region (Cardot, 1901, 1906a, b, 1907c, 1908a, 1911b, c, d, 1912b, 1913a; Brotherus, 1903, 1924a; Wright, 1905 (*as Bryum sp.*); Paris, 1906; Dixon, 1935; Steere, 1961a).

var. *laxiretis* (Card.) Steere (Steere, 1961a).

Syn. *Webera racovitzae* Card. var. *laxiretis* Card. (Cardot, 1906b).

Distr. Antarctic, peninsula region (Cardot, 1906b, 1907c, 1908a; Horikawa and Ando, 1961 (*as Bryum inconnexum*); Steere, 1961a).

P. wahlenbergii (Web. et Mohr) Andrews (Andrews, 1935).

Syn. *Hypnum wahlenbergii* Web. et Mohr (Weber and Mohr, 1807).
Bryum wahlenbergii (Web. et Mohr) Schwaegr. (Schwaegrichen, 1816).
Webera wahlenbergii (Web. et Mohr) Fuernr. (Fürnrohr, 1829).
Mniobryum wahlenbergii (Web. et Mohr) Jenn. (Jennings, 1913).
Hypnum albicans Web. et Mohr (Weber and Mohr, 1803) *nom. nud.*
Mnium albicans Wahlenb. (Wahlenberg, 1812) *nom. illeg.*
Bryum albicans Roehl. (Röhling, 1813) *hom. illeg.*
Webera albicans Schimp. (Schimper, 1855 [1856]) *nom. illeg.*
Pohlia albicans Lindb. (Lindberg, 1879) *nom. illeg.*
Mniobryum albicans (Wahlenb.) Limpr. (Limprecht, 1892) *nom. illeg.*
Bryum austro-albicans C. Muell. (Mueller, 1884 [1883]).
Mniobryum austro-albicans (C. Muell.) Broth. (Brotherus, 1903).

Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1848a; Paris, 1905a; Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1938, 1939, 1954, 1957; Hosseus, 1938c, d, 1939; Kühnemann, 1938; Bartram, 1952; Seki, 1974).
 Falkland Islands (Wilson and Hooker, 1847; Mueller, 1848a; Cardot, 1905a, 1908a; Paris, 1905a; Cardot and Brotherus, 1923; Brotherus, 1924a; Rövainen, 1934; Kühnemann, 1938).
 South Georgia (Cardot and Brotherus, 1923; Steere, 1961b).

var. *glacialis* (Brid.) E. F. Warburg (Warburg, 1962).

Syn. *Mniobryum wahlenbergii* (Web. et Mohr) Jenn. var. *glaciale* (Brid.) Wijk et Marg. (van der Wijk and Margadant, 1959).

Bryum glaciale Brid. (Bridel, 1827a).

Bryum alticaule C. Muell. (Mueller, 1884 [1883]) *fide* Clarke, 1973a.

Webera alticaulis (C. Muell.) Kindb. (Kindberg, 1889) *fide* Clarke, 1973a.

Mniobryum alticaule (C. Muell.) Broth. (Brotherus, 1903) *fide* Clarke, 1973a.

Bryum amplirete C. Muell. (Mueller, 1890a) *fide* Clarke, 1973a.

Webera ampliretis (C. Muell.) Kindb. (Kindberg, 1891) *fide* Clarke, 1973a.

Mniobryum amplirete (C. Muell.) Broth. (Brotherus, 1903) *fide* Clarke, 1973a.

Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1884, 1885, 1889; Brotherus, 1903, 1924a; Cardot, 1905a, 1908a, 1913a; Paris, 1905a; Cardot and Brotherus, 1923; Herzog, 1923, 1954; Reimers, 1926; Kühnemann, 1938; Bartram, 1952).

Falkland Islands (Wilson and Hooker, 1847; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Brotherus, 1903, 1924a; Paris, 1905a; Cardot, 1906a, 1908a; Dixon, 1932 (as *Webera inflexa* *fide* Dixon, 1934), 1934; Steere, 1961b; Clarke, 1973a; Bell, 1977b).

Polytrichadelphus

P. croceus Mitt.

Polytrichadelphus semi-angulatus

P. dendroides (Hedw.) Mitt.

Dendroligotrichum dendroides

P. horridus Mitt.

Polytrichadelphus magellanicus

P. magellanicus (Hedw.) Mitt. (Mitten, 1860).

Syn. *Polytrichum magellanicum* L. ex Hedw. (Hedwig, 1801).

Pogonatum magellanicum (Hedw.) P. Beauv. (Palisot de Beauvois, 1805a).

Catharinea magellanica (Hedw.) Brid. (Bridel, 1819).

Oligotrichum magellanicum (Hedw.) Kindb. (Kindberg, 1888).

Gymnopilum magellanicum (Hedw.) Schimp. (Schimper *in* Paris, 1895) *nom. inval.*

Polytrichadelphus horridus Mitt. (Mitten, 1869).

Oligotrichum horridum (Mitt.) Kindb. (Kindberg, 1888).

Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Menzies, 1798; Hedwig, 1801; Bridel, 1803, 1806, 1827b; Schwaegrichen, 1816; Montagne, 1845b, 1850; Wilson and Hooker, 1847; Mueller, 1848a, 1885; Sullivant, 1859; Mitten, 1860, 1869; Duby, 1875a; Jaeger, 1875; Bescherelle, 1889; Neger, 1899; Dusén, 1903a; Cardot, 1905a, 1908a; Paris, 1905b; Rowlee, 1916; Cardot and Brotherus, 1923; Herzog, 1923, 1939, 1940, 1954; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938; Seki, 1974).

Falkland Islands (Dumont D'Urville, 1825; Bridel, 1827b; Wilson and Hooker, 1847; Jaeger, 1875; Mueller, 1885; Bescherelle, 1889; Cardot, 1905a, 1908a; Paris, 1905b; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

P. minimus Card.

Psilotilum minimum

P. robustus (Lindb.) Broth. (Brotherus, 1905a).

Syn. *Polytrichum robustum* Lindb. (Lindberg, 1868c).

Pogonatum robustum (Lindb.) Jaeg. (Jaeger, 1875) *hom. illeg.*

Distr. Southern South America (Lindberg, 1868c; Ångström, 1872; Jaeger, 1875; Brotherus, 1905a, 1925; Cardot, 1908a, 1913a; Cardot and Brotherus, 1923; Bartram, 1952).

P. semi-angulatus (Brid.) Mitt. (Mitten, 1869).

Syn. *Pogonatum semi-angulatum* Brid. (Bridel, 1827b).

Polytrichadelphus croceus Mitt. (Mitten, 1860).

Distr. Southern South America (Krieger, 1904).

P. squamosus (Hook. f. et Wils.) Mitt.

Dendroligotrichum squamosum

P. stricticaulis Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1905b).

P. umbrosus Mitt. (Mitten, 1869).

Syn. *Oligotrichum umbrosum* (Mitt.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Mitten, 1869; Jaeger, 1875; Paris, 1905b; Brotherus, 1925).

Polytrichastrum

P. alpinum (Hedw.) G. L. Smith

Polytrichum alpinum

P. longisetum (Brid.) G. L. Smith

Polytrichum longisetum

Polytrichum

P. affine Funck

Polytrichum alpestre

P. alpestre Hopp. (Hoppe, 1801).

Syn. *Polytrichum strictum* Menz. ex Brid. (Menzies ex Bridel, 1801a).

Polytrichum strictum Menz. ex Brid. var. *alpestre* (Hopp.) Rabenh. (Rabenhorst, 1848).

Polytrichum affine Funck (Funck, 1802).

Polytrichum juniperinum Hedw. var. *affine* (Funck) Brid. (Bridel, 1806).

Polytrichum juniperinum Hedw. var. *alpestre* (Hopp.) Roehl. (Röhling, 1813).

Polytrichum juniperinum Hedw. var. *strictum* (Brid.) Roehl. (Röhling, 1813) *nom. illeg.*

Polytrichum juniperinum Hedw. ssp. *strictum* (Brid.) Nyl. et Sael. (Nylander and Saelan, 1859).

Polytrichum macrорaphis C. Muell. (Mueller, 1890a) pro parte *fide* Greene, 1973.

Distr. Southern South America (Montagne, 1850; Mueller, 1885; Bescherelle, 1889; Dusén, 1903a, 1905d; Cardot, 1905a, 1908a; Paris, 1905b; Spegazzini, 1922; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Martin, 1949).

Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Paris, 1905b; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Steere, 1961b; Greene, 1973; Bell, 1977b).

Antarctic, peninsula region (Cardot, 1900, 1901, 1906a, 1908a, 1911b, d, 1913a; Paris, 1905b; Brotherus, 1925; Steere, 1961a; Greene and others, 1970; Robinson, 1972).

P. alpinum Hedw. (Hedwig, 1801).

Syn. *Pogonatum alpinum* (Hedw.) Roehl. (Röhling, 1814).

Polytrichastrum alpinum (Hedw.) G. L. Smith (Smith, 1971).

Polytrichum austro-georgicum C. Muell. (Mueller, 1890a) *fide* Greene, 1973.

Pogonatum austro-georgicum (C. Muell.) Par. (Paris, 1898).

Polytrichum nano-cephalum C. Muell. (Mueller, 1890a).

Polytrichum plurirameum C. Muell. (Mueller, 1890a) *fide* Greene, 1973.

Polytrichum timmioides C. Muell. (Mueller, 1908a) *fide* Greene, 1973.

Pogonatum alpinum (Hedw.) Roehl. var. *brevifolium* (R. Brown) Brid. (Bridel, 1927b).

Polytrichum macrорaphis C. Muell. (Mueller, 1890a) pro parte *fide* Greene, 1973.

Distr. Southern South America (Cardot, 1905a, 1908a; Thériot, 1935a; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Paris, 1898, 1905b; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Dixon, 1932; Steere, 1961b; Greene, 1973).

Antarctic, peninsula region (Cardot, 1900*, 1901*, 1906a*, b, 1907c, 1908a*, 1911b*, d*, 1913a*; Paris, 1905b*; Dixon, 1920, 1935; Brotherus, 1925*; Bartram, 1957*; Steere, 1961a*; Greene, 1968a; Greene and others, 1970; Robinson, 1972).

var. *integritifolium* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

P. antarcticum Card.

Polytrichum piliferum

P. antillarum Rich.

Polytrichum juniperinum

ssp. *prionotum* (C. Muell.) Walth.

Polytrichum juniperinum var. *tumescens*

ssp. *secundulum* (C. Muell.) Walth.

Polytrichum juniperinum

ssp. *substrictum* Walth.

Polytrichum juniperinum

P. australe Hook. f. et Wils.

Psilotum australe

*and as *Pogonatum alpinum* (Hedw.) Roehl. var. *brevifolium* (R. Brown) Brid. *fide* Greene and others, 1970.

<i>P. austro-georgicum</i> C. Muell.	<i>Polytrichum alpinum</i>
<i>P. breve</i> C. Muell.	<i>Polytrichum juniperinum</i> var. <i>tumescens</i>
<i>P. canaliculatum</i> Hook. et Arnott	<i>Oligotrichum canaliculatum</i>
<i>P. compressum</i> Hook. f. et Wils.	<i>Atrichopsis compressa</i>
<i>P. dendroides</i> Brid. ex Hedw.	<i>Dendroligotrichum dendroides</i>
<i>P. elongatum</i> P. Beauv. (Palisot de Beauvois, 1805a).	
<i>Distr.</i> Southern South America (Palisot de Beauvois, 1805a; Bridel, 1806, 1827b; Schwaegrichen, 1816; Mueller, 1848a; Mitten, 1869; Jaeger, 1875; Paris, 1905b; Cardot, 1908a).	
<i>P. gracile</i> Dicks.	<i>Polytrichum longisetum</i>
<i>P. juniperifolium</i> Schimp. ex Par.	<i>Polytrichum juniperinum</i>
<i>P. juniperiforme</i> Schimp. ex Mitt.	<i>Polytrichum juniperinum</i>
<i>P. juniperinum</i> Hedw. (Hedwig, 1801).	
<i>Syn.</i> <i>Polytrichum juniperinum</i> Hedw. var. <i>alpinum</i> Schimp. (Schimper, 1860). <i>Polytrichum juniperinum</i> Hedw. var. <i>integrum</i> C. Muell. (Mueller, 1848a). <i>Polytrichum antillarum</i> Rich. (Richard in Bridel, 1827b) <i>nom. illeg.</i> . <i>Polytrichum antillarum</i> Rich. ssp. <i>substrictum</i> Walth. (Walther, 1934). <i>Polytrichum antillarum</i> Rich. ssp. <i>secundulum</i> (C. Muell.) Walth. (Walther, 1934). <i>Polytrichum secundulum</i> C. Muell. (Mueller, 1897b). <i>Polytrichum prionotrichum</i> C. Muell. (Mueller, 1897a). <i>Polytrichum substrictum</i> Hamp. (Hampe in Paris, 1897) <i>nom. nud.</i> . <i>Polytrichum juniperiforme</i> Schimp. ex Mitt. (Schimper ex Mitten, 1869). <i>Polytrichum juniperifolium</i> Schimp. ex Par. (Schimper ex Paris, 1905b) <i>nom. inval.</i>	
<i>Distr.</i> Southern South America (Mueller, 1848a, 1897a; Montagne, 1850; Lorentz, 1866; Mitten, 1869; Jaeger, 1875; Paris, 1897, 1905b; Dusén, 1903a, 1905d; Kurtz, 1904; Cardot, 1905a, 1908a; Thériot, 1921b, 1935a; Cardot and Brotherus, 1923; Espinosa B., 1924; Brotherus, 1925; Reimers, 1926; Roivainen, 1934; Walther, 1934; Hosseus, 1936, 1937, 1938c, d; Kühnemann, 1938; Herter, 1939b; Bartram, 1952; Bizot, 1967; Seki, 1974). Falkland Islands (Kühnemann, 1938). South Georgia (Cardot, 1906a, 1908a; Steere, 1961b; Greene, 1973). Antarctic, peninsula region (Greene, 1968a; Greene and others, 1970).	
ssp. <i>strictum</i> (Brid.) Nyl. et Sael.	<i>Polytrichum alpestre</i>
var. <i>affine</i> (Funck) Brid.	<i>Polytrichum alpestre</i>
var. <i>alpestre</i> (Hopp.) Roehl.	<i>Polytrichum alpestre</i>
var. <i>alpinum</i> Schimp.	<i>Polytrichum juniperinum</i>
var. <i>australe</i> Walth. (Walther, 1934) <i>hom. illeg.</i>	
<i>Syn.</i> <i>Polytrichum spegazzinii</i> C. Muell. (Mueller, 1885).	
<i>Distr.</i> Southern South America (Mueller, 1885; Neger, 1899; Paris, 1905b; Cardot, 1908a; Brotherus, 1925; Walther, 1934, 1940; Kühnemann, 1938; Herzog, 1939, 1957).	
var. <i>integrum</i> C. Muell.	<i>Polytrichum juniperinum</i>
var. <i>strictum</i> (Brid.) Roehl.	<i>Polytrichum alpestre</i>
var. <i>tumescens</i> (C. Muell.) Herz. (Herzog, 1916a).	
<i>Syn.</i> <i>Polytrichum tumescens</i> C. Muell. (Mueller, 1897a). <i>Polytrichum prionotum</i> C. Muell. (Mueller, 1897a). <i>Polytrichum breve</i> C. Muell. (Mueller, 1897a). <i>Polytrichum antillarum</i> Rich. ssp. <i>prionotum</i> (C. Muell.) Walth. (Walther, 1934).	
<i>Distr.</i> Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Walther, 1934).	
<i>P. ionchopelma</i> C. Muell. (Mueller, 1901a [1900]) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Mueller, 1901a).	

P. longisetum Sw. ex Brid. (Swartz ex Bridel, 1801).

Syn. *Polytrichastrum longisetum* (Brid.) G. L. Smith (Smith, 1971).
Polytrichum gracile Dicks. (Dickson in Menzies in Schrebier, 1802).

Distr. Southern South America (Herzog, 1939).

P. macrорaphis C. Muell. **Polytrichum alpestre** pro parte and **Polytrichum alpinum** pro parte

P. magellanicum L. ex Hedw.

Polytrichadelphus magellanicus

P. molinae Mont.

Oligotrichum canaliculatum var. **molinae**

P. nano-cephalum C. Muell.

Polytrichum alpinum

P. noduliferum Dus.

Polytrichum piliferum

P. oligodus Kunz. ex C. Muell.

Polygonatum oligodus

P. patagonicum C. Muell.

Polytrichum piliferum

P. piliferum Hedw. (Hedwig, 1801).

Syn. *Polytrichum pilifolium* C. Muell. (Mueller, 1897a) hom. illeg.

Polytrichum patagonicum C. Muell. (Mueller, 1897a).

Polytrichum antarcticum Card. (Cardot, 1900) fide Greene and others, 1970.

Polytrichum subpiliferum Card. (Cardot, 1900).

Polytrichum trachynotum C. Muell. (Mueller, 1885).

Polytrichum noduliferum Dus. (Dusén in Paris, 1905b) nom. nud.

Distr. Juan Fernandez (Montagne, 1835; Paris, 1905b; Brotherus, 1925).

Southern South America (Mitten, 1869; Ångström, 1872; Mueller, 1885, 1897a; Cardot, 1900, 1901, 1905a, 1908a, 1913a; Dusén, 1903a; Kurtz, 1904; Paris, 1905b; Spegazzini, 1922; Brotherus, 1925; Reimers, 1926; Hosseus, 1936, 1937; Herzog, 1938, 1939, 1957; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

Falkland Islands (Cardot, 1905a, 1908a; Paris, 1905b; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

South Georgia (Cardot, 1906a, 1908a; Brotherus, 1925; Steere, 1961b; Greene, 1973).

Antarctic, peninsula region (Cardot, 1900, 1901, 1906a, b, 1907c, 1908a, 1911b, c, d, 1912b, 1913a; Wright, 1905; Paris, 1905b; Dixon, 1920, 1935; Brotherus, 1925; Bartram, 1957; Steere, 1961a; Greene and others, 1970; Robinson, 1972).

P. pilifolium C. Muell.

Polytrichum piliferum

P. plurirameum C. Muell.

Polytrichum alpinum

P. plurisetum C. Muell.

Polygonatum plurisetum

var. *brevipes* C. Muell.

Polygonatum plurisetum var. *brevipes*

P. prionotrichum C. Muell.

Polytrichum juniperinum

P. prionotum C. Muell.

Polytrichum juniperinum var. *tumescens*

P. pygmaeocaulon C. Muell. (Mueller, 1901a [1900]) nom. nud.

Distr. Southern South America (Mueller, 1901a).

P. riedelianum Mont.

Oligotrichum riedelianum

P. robustum Lindb.

Polytrichadelphus robustus

P. secundulum C. Muell.

Polytrichum juniperinum

P. spegazzinii C. Muell.

Polytrichum juniperinum var. *australe*

P. squamosum Hook. f. et Wils.

Dendroligotrichum squulosum

P. stolonigerum C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Mueller, 1901a).

P. strictum Menz. ex Brid.

Polytrichum alpestre

var. *alpestre* (Hopp.) Rabenh.

Polytrichum alpestre

P. subpiliferum Card.

Polytrichum piliferum

P. substrictum Hamp.

Polytrichum juniperinum

P. timmiaceum C. Muell. (Mueller, 1901a [1900]) nom. nud.

Distr. Southern South America (Mueller, 1901a).

- P. timmioides* C. Muell. *Polytrichum alpinum*
P. trachynotum C. Muell. *Polytrichum piliferum*
P. trichodon Hook. f. et Wils. *Psilopilum trichodon*
P. tumescens C. Muell. *Polytrichum juniperinum* var. *tumescens*
var. *pygmaeocaulon* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b).

- P. umbrorus* Mitt. (Mitten in Brotherus, 1925).

Distr. Southern South America (Brotherus, 1925).

Porothamnium

- P. arbusculans* (C. Muell.) Fleisch. (Fleischer in Cardot and Brotherus, 1923).

- Syn. *Hypnum arbusculans* C. Muell. (Mueller, 1874).
Thamnium arbusculans (C. Muell.) Jaeg. (Jaeger, 1877).
Thamnium arbusculans (C. Muell.) Jaeg. var. *complanatum* C. Muell. (Mueller in Kindberg, 1902) nom. nud.
Porothamnium arbusculans (C. Muell.) Fleisch. var. *complanatum* Card. (Cardot in Cardot and Brotherus, 1923).
Hypnum naumanii C. Muell. (Mueller, 1884 [1883]).
Hypnodendron naumanii (C. Muell.) Kindb. (Kindberg, 1891).
Thamnium naumanii (C. Muell.) Kindb. (Kindberg, 1902).
Thamnium decumbens Besch. (Bescherelle, 1885a).
Porothamnium decumbens (Besch.) Fleisch. (Fleischer, 1908b).
Porotrichum corralense Broth. (Brotherus in Kindberg, 1902) nom. nud.
Thamnium corralense Broth. (Brotherus in Reimers, 1926) nom. nud.

Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b; Robinson, 1975).
Southern South America (Mueller, 1874, 1884, 1885, 1889; Jaeger, 1877; Bescherelle, 1885a, 1889; Neger, 1899; Kindberg, 1902; Paris, 1904b, 1905b; Cardot, 1908a; Fleischer, 1908b; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1960; Brotherus, 1925; Reimers, 1926; Gunckel, 1938; Kühnemann, 1938; Bizot, 1967; Seki, 1974).

var. *complanatum* Card.

Porothamnium arbusculans

- P. decumbens* (Besch.) Fleisch.

Porothamnium arbusculans

- P. explanatum* (Mitt.) Fleisch. (Fleischer in Brotherus, 1925).

- Syn. *Porotrichum explanatum* Mitt. (Mitten, 1869).
Thamnium explanatum (Mitt.) Kindb. (Kindberg, 1902).

Distr. Southern South America (Kindberg, 1902).

- P. fasciculatum* (Hedw.) Fleisch. (Fleischer, 1908b).

- Syn. *Hypnum fasciculatum* Hedw. (Hedwig, 1801).
Neckera fasciculata (Hedw.) Arnott (Arnott, 1827).
Porotrichum fasciculatum (Hedw.) Mitt. (Mitten, 1869).
Dendro-Hypnum fasciculatum (Hedw.) Hamp. (Hampe, 1870) nom. inval.
Thamnium fasciculatum (Hedw.) C. Muell. (Mueller, 1898b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959).
Southern South America (Montagne, 1850; Fleischer, 1908b).

- P. leucocaulon* (C. Muell.) Fleisch. (Fleischer in Brotherus, 1925).

- Syn. *Neckera leucocaulon* C. Muell. (Mueller, 1847c).
Porotrichum leucocaulon (C. Muell.) Mitt. (Mitten, 1869).
Thamnium leucocaulon (C. Muell.) Kindb. (Kindberg, 1902).

Distr. Southern South America (Mueller, 1847c; Montagne, 1850; Mitten, 1869; Jaeger, 1877; Kindberg, 1902; Paris, 1905b; Cardot, 1908a; Herzog, 1923, 1938, 1940, 1957; Brotherus, 1925; Roivainen, 1934; Gunckel, 1938; Kühnemann, 1938; Bizot, 1967).

- P. lorentzii* (C. Muell.) Fleisch.

Porothamnium neckeraeforme

P. matogrossense (Broth.) Fleisch. (Fleischer in Brotherus, 1925).

Syn. *Thamnium matogrossense* Broth. (Brotherus, 1900).

Distr. Southern South America (Brotherus, 1925; Kühnemann, 1938).

P. neckeraeforme (Hamp.) Fleisch. (Fleischer in Brotherus, 1925).

Syn. *Hypnum neckeraeforme* Hamp. (Hampe, 1866).

Porotrichum neckeraeforme (Hamp.) Mitt. (Mitten, 1869).

Thamnium neckeraeforme (Hamp.) Kindb. (Kindberg, 1902).

Porotrichum lorentzii C. Muell. (Mueller, 1879a).

Porothamnium lorentzii (C. Muell.) Fleisch. (Fleischer, 1908b).

Distr. Southern South America (Mueller, 1879a; Kindberg, 1902; Paris, 1905b; Fleischer, 1908b; Brotherus, 1925; Kühnemann, 1938).

P. panduraefolium (C. Muell.) Fleisch. (Fleischer, 1908b).

Syn. *Hypnum panduraefolium* C. Muell. (Mueller, 1855).

Porotrichum panduraefolium (C. Muell.) Mitt. (Mitten, 1869).

Thamnium panduraefolium (C. Muell.) Broth. (Brotherus, 1906b).

Porotrichum panduraeforme Kindb. (Kindberg, 1891) err.

Thamnium angustifolium Schimp. (Schimper in Lechler, 1857) nom. nud. fide Mitten, 1869.

Distr. Southern South America (Mueller, 1855; Lechler, 1857; Mitten, 1869; Jaeger, 1877; Paris, 1905b; Brotherus, 1906b, 1925; Fleischer, 1908b; Campo, 1915, 1921; Thériot, 1918; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1954, 1960; Reimers, 1926; Gunckel, 1938; Seki, 1974).

P. porrectulum (C. Muell.) Fleisch. (Fleischer in Brotherus, 1925).

Syn. *Porotrichum porrectulum* C. Muell. (Mueller, 1879a).

Thamnium porrectulum (C. Muell.) Broth. (Brotherus, 1906b).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1906b, 1925; Kühnemann, 1938).

P. valdiviae (C. Muell.) Fleisch. (Fleischer in Brotherus, 1925).

Syn. *Hypnum valdiviae* C. Muell. (Mueller, 1855).

Porotrichum valdiviae (C. Muell.) Mitt. (Mitten, 1869).

Thamnium valdiviae (C. Muell.) Kindb. (Kindberg, 1902).

Thamnium valdivicum C. Muell. (Mueller in Kindberg, 1902) nom. nud.

Hypnum neckera Schimp. (Schimper in Mueller, 1855) nom. nud.

Thamnium neckera Schimp. (Schimper in Jaeger, 1877) nom. nud.

Distr. Juan Fernandez (Robinson, 1975).

Southern South America (Mueller, 1855; Lorentz, 1866; Mitten, 1869; Jaeger, 1877; Kindberg, 1902; Paris, 1905b; Cardot, 1908a; Thériot, 1918; Cardot and Brotherus, 1923; Brotherus, 1925; Reimers, 1926; Roivainen, 1934; Gunckel, 1938; Herzog, 1938, 1954, 1957, 1960; Kühnemann, 1938; Bartram, 1952; Bizot, 1967; Seki, 1974).

var. pallidum (Lor.) Wijk et Marg. (van der Wijk and Margadant, 1961).

Syn. *Hypnum valdiviae* C. Muell. var. *pallidum* Lor. (Lorentz, 1866).

Porotrichum valdiviae (C. Muell.) Mitt. var. *pallidum* (Lor.) Jaeg. (Jaeger, 1877).

Distr. Southern South America (Lorentz, 1866; Paris, 1900a, 1905b).

Porotrichodendron

P. porteri (Thér.) Broth. (Brotherus, 1925).

Syn. *Porotrichum porteri* Thér. (Thériot, 1917b).

Distr. Southern South America (Thériot, 1917b; Brotherus, 1925).

P. robustum Broth. (Brotherus in Herzog, 1916a).

Distr. Southern South America (Bartram, 1965).

Porotrichum

P. callicostelloides Broth. ex Thér.

Pinnatella callicostelloides

P. chilense Thér. (Thériot, 1924).

Distr. Southern South America (Thériot, 1918 (as *Porothamnium valdiviae forma minor* Thér.), 1924; Costes, 1921; Brotherus, 1925).

<i>P. cochlearifolium</i> (Schwaegr.) Mitt.	Weymouthia cochlearifolia
<i>P. confertum</i> Mitt.	Thamnobryum confertum
<i>P. corralense</i> Broth.	Porothamnium arbusculans
<i>P. crassinervium</i> Mitt. ex Broth.	Thamnobryum rigidum
<i>P. explanatum</i> Mitt.	Porothamnium explanatum
<i>P. fasciculatum</i> (Hedw.) Mitt.	Porothamnium fasciculatum
<i>P. gracile</i> (Hook. f. et Wils.) Mitt.	Camptochaete gracilis
<i>P. latinerve</i> Mitt.	Thamnobryum rigidum
<i>P. leucocaulon</i> (C. Muell.) Mitt.	Porothamnium leucocaulon
<i>P. lorentzii</i> C. Muell.	Porothamnium neckeraeforme
<i>P. neckeraeforme</i> (Hamp.) Mitt.	Porothamnium neckeraeforme
<i>P. neckeroides</i> (Hook.) Williams	Thamnium neckeroides
<i>P. panduraeforme</i> Kindb.	Porothamnium panduraefolium
<i>P. panduraefolium</i> (C. Muell.) Mitt.	Porothamnium panduraefolium
<i>P. pinnatelloides</i> C. Muell.	Pinnatella thieleana
<i>P. porrectulum</i> C. Muell.	Porothamnium porrectulum
<i>P. porteri</i> Thér.	Porotrichodendron porteri
<i>P. rigidum</i> Mitt.	Thamnobryum rigidum
<i>P. thieleanum</i> (C. Muell.) Mitt.	Pinnatella thieleana
<i>P. thieli</i> Aongstr.	Pinnatella thieleana
<i>P. valdiviae</i> (C. Muell.) Mitt.	Porothamnium valdiviae
var. <i>pallidum</i> (Lor.) Jaeg.	Porothamnium valdiviae var. <i>pallidum</i>

Pottia**P. altipes** Broth. (Brotherus, 1918).

Distr. Southern South America (Brotherus, 1918, 1924a; Kühnemann, 1938).

<i>P. amblyophylla</i> (Hook.) C. Muell.	Didymodon amblyophyllus
<i>P. anderssonii</i> Sauerb.	Pottia antarctica

P. antarctica Schimp. (Schimper in Jaeger, 1873) nom. nud.

Distr. Southern South America (Jaeger, 1873).

P. antarctica (Aongstr.) C. Muell. (Mueller, 1885).

Syn. *Gymnostomum antarcticum* Aongstr. (Ångström, 1872).
Gymnostomum magellanicum Aongstr. ex Par. (Ångström ex Paris, 1896) nom. inval.
Pottia magellanica Schimp. ex Par. (Schimper ex Paris, 1898) nom. illeg.
Pottia anderssonii Sauerb. (Sauerbeck in Jaeger and Sauerbeck, 1879b) nom. illeg.

Distr. Southern South America (Lechler, 1857; Ångström, 1872; Jaeger and Sauerbeck, 1879b; Mueller, 1885; Paris, 1898, 1905b; Dusén, 1906; Cardot, 1908a; Warnstorff, 1916; Kühnemann, 1938).

P. austro-georgica Card. (Cardot, 1906a).

Syn. *P. austro-georgica* Card. var. *microphylla* Card. et Broth. (Cardot and Brotherus, 1923) fide Matteri, 1977a.

Distr. Southern South America (Warnstorff, 1916).

South Georgia (Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b; Greene, 1968a; Matteri, 1977a).
Antarctic, peninsula region (Matteri, 1977b).

- var. *microphylla* Card. et Broth. *Pottia austro-georgica*
- P. brunnea* (C. Muell.) Par. *Didymodon brunneus*
- P. cavifolia* Ehrh. ex Fuernr. *Pterygoneurum ovatum*
- P. charcotii* Card. *Pottia heimii*
- P. chubutensis* Card. et Broth. (Cardot and Brotherus, 1923).**
- Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).
- P. flavipes* Mont. (Montagne, 1845a).**
- Syn. *Pottia heimii* (Hedw.) Hamp. var. *flavipes* (Mont.) Warnst. (Warnstorf, 1916).
Pottia flaviseta Mont. (Montagne, 1888) err.
Tortula flaviseta Mitt. (Mitten, 1869).
- Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Mitten, 1869; Reichardt, 1870; Paris, 1905b; Warnstorf, 1916; Brotherus, 1924a; Bartram, 1942, 1943).
- P. flaviseta* Mont. *Pottia flavipes*
- P. glauco-viridis* (C. Muell.) Par. *Didymodon glauco-viridis*
- P. gracillima* (C. Muell.) Par. *Trichostomum gracillimum*
- P. guessfeldtii* Schleph. *Pottia heimii* var. *guessfeldtii*
- P. gymna* (C. Muell.) Par. *Didymodon gymnus*
- P. heimii* (Hedw.) Hamp. (Hampe, 1837a).**
- Syn. *Gymmnostomum heimii* Hedw. (Hedwig, 1801).
Bryum heimii (Hedw.) Dicks. ex With. (Dickson ex Withering, 1801).
Desmatodon heimii (Hedw.) Mitt. (Mitten, 1864).
Tortula heimii (Hedw.) Mitt. (Mitten, 1869).
Didymodon heimii (Hedw.) Kindb. (Kindberg, 1897).
Pottia charcotii Card. (Cardot, 1911d) fide Matteri, 1977b.
Pottia heimii (Hedw.) Hamp. var. *charcotii* (Card.) Savicz. et Smirn. (Savich-Ljubitskaya and Smirnova, 1963a) fide Matteri, 1977b.
- Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1849; Sullivant, 1859; Mitten, 1869; Jaeger, 1873; Dusén, 1907; Cardot, 1908a; Warnstorf, 1916; Cardot and Brotherus, 1923; Herzog, 1923; Brotherus, 1924a; Kühnemann, 1938).
Falkland Islands (Wilson and Hooker, 1847; Cardot, 1905a, 1908a).
Antarctic, peninsula region (Cardot, 1911b, d, 1913a; Brotherus, 1924a; Bartram, 1957; Steere, 1961a; Savich-Ljubitskaya and Smirnova, 1963a, 1965b; Robinson, 1972; Matteri, 1977b).
- var. *brevinervis* Savicz. et Smirn. (Savich-Ljubitskaya and Smirnova, 1965b).**
- Distr. Antarctic, continental region (Savich-Ljubitskaya and Smirnova, 1965b; Greene, 1968a; Kuc, 1969; Savich-Ljubitskaya, 1978). *Pottia heimii*
- var. *charcotii* (Card.) Savicz. et Smirn.**
- var. *eurystoma* Card. et Broth. (Cardot and Brotherus, 1923).**
- Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938). *Pottia heimii*
- var. *flavipes* (Mont.) Warnst.**
- var. *guessfeldtii* (Schleph.) Warnst. (Warnstorf, 1916).**
- Syn. *Pottia guessfeldtii* Schleph. (Schlephacke, 1884).
Stegonia guessfeldtii (Schleph.) Hag. (Hagen in Hagen and Printz, 1929).
- Distr. Southern South America (Schlephacke, 1884; Paris, 1905b; Warnstorf, 1916; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938). *Pottia flavipes*
- var. *magellanica* Warnst. (Warnstorf, 1916).**
- Syn. *Pottia magellanica* Schimp. (Schimper in Mueller, 1885) nom. nud.
Pottia megapoda C. Muell. (Mueller, 1897a).
Pottia systyliopsis C. Muell. (Mueller, 1897a).

Distr. Southern South America (Jaeger, 1873; Mueller, 1897a; Paris, 1905b; Cardot, 1908a; Warnstorf, 1916; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

var. *maxima* Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923).

var. *spegazzinii* (C. Muell.) Warnst. (Warnstorf, 1916).

Syn. *Pottia spegazzinii* C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Dusén, 1903a, 1906; Paris, 1905b; Cardot, 1908a; Warnstorf, 1916; Brotherus, 1924a; Kühnemann, 1938).

var. *thaxteri* Card. et Thér. (Cardot and Thériot in Thériot, 1929c [1930]).

Distr. Southern South America (Thériot, 1929c).

P. imperfecta (C. Muell.) Par.

Barbula imperfecta

P. kunzeana C. Muell.

Hymenostylium kunzeanum

P. lamprocarpa (C. Muell.) Broth.

Tetrapherum lamprocarpum

P. lamprothecium (C. Muell.) Broth.

Tetrapherum lamprothecium

P. longirostris C. Muell.

Hymenostylium kunzeanum

P. lorentziana C. Muell.

Didymodon lorentzianus

P. lorentzii C. Muell.

Stegonia lorentzii

P. macrocarpa Schimp. (Schimper, 1836).

Syn. *Pottia macropoda* Schimp. ex Broth. (Schimper ex Brotherus, 1902) err.

Tortula macrocarpa (Schimp.) Mitt. (Mitten, 1869).

Hyophila macrocarpa (Schimp.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Schimper, 1836; Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Brotherus, 1902; Paris, 1905b; Warnstorf, 1916).

P. macropoda Schimp. ex Broth.

Pottia macrocarpa

P. magellanica Schimp.

Pottia heimii* var. *magellanica

P. magellanica Schimp. ex Par.

Pottia antarctica

P. megapoda C. Muell.

Pottia heimii* var. *magellanica

P. microthecia (C. Muell.) Par.

Didymodon microthecius

P. mucronata Warnst. (Warnstorf, 1916) *nom. nud.*

Distr. Southern South America (Warnstorf, 1916).

P. nuda (C. Muell.) Par.

Didymodon brunneus

P. orbigniana C. Muell.

Didymodon orbignyanus

P. ovata (Hedw.) Fuernr. ex Lindb.

Pterygoneurum ovatum

P. pellucida Card. (Cardot, 1911d) *nom. nud.*

Distr. Southern South America (Cardot, 1911d).

P. physcomitrioides C. Muell.

Pottia truncata

P. poeppigiana (C. Muell.) C. Muell.

Trichostomopsis australasiae

P. recurvirostra Ehrh. ex Hedw.

Hymenostylium recurvirostre

P. spathulato-linearis (C. Muell.) Par.

Didymodon spathulato-linearis

P. spegazzinii C. Muell.

Pottia heimii* var. *spegazzinii

P. subsessilis (Brid.) B.S.G.

Pterygoneurum subsessile

P. systyliopsis C. Muell.

Pottia heimii* var. *magellanica

P. truncata (Hedw.) B.S.G. (Bruch and others, 1843).

Syn. *Gymnostomum truncatum* Hedw. (Hedwig, 1801).

Bryum truncatum (Hedw.) Brotero (Brotero, 1804).

Tortula truncata (Hedw.) Mitt. (Mitten in Godman, 1870).

Bryum truncatulum L. ex With. (Withering, 1801) *nom. illeg.*
Pottia truncatula (With.) Bus. (Buse, 1858) *hom. illeg.*
Tortula truncatula (With.) Lindb. (Lindberg, 1879) *nom. illeg.*
Pottia physcomitrioides C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Krieger, 1904; Paris, 1905b; Felippone, 1909;
Warnstorff, 1916; Brotherus, 1924a; Herter, 1933a; Kühnemann, 1938).

P. truncatula (With.) Bus.

Pottia truncata

Pseudocrossidium

P. chilense Williams (Williams, 1915).

Distr. Southern South America (Williams, 1915; Brotherus, 1924a; Hilpert, 1933).

P. leucocalyx (Mont.) Thér. (Thériot, 1923a [1921]).

Syn. *Tortula leucocalyx* Mont. (Montagne, 1838).
Barbula leucocalyx (Mont.) C. Muell. (Mueller, 1849).
Barbula pachyneura Dus. (Dusén, 1906).
Pseudocrossidium pachyneuron (Dus.) Thér. (Thériot, 1921a).

Distr. Southern South America (Montagne, 1838, 1839c, 1845a, 1850, 1856; Mueller, 1849;
Mitten, 1869; Jaeger, 1873; Paris, 1904a; Dusén, 1906; Thériot, 1917b, 1921a, 1922,
1923a; Costes, 1921; Brotherus, 1924a, c; Espinosa B., 1941; Skottsberg, 1950).

P. pachyneuron (Dus.) Thér.

Pseudocrossidium leucocalyx

Pseudocryphaea

P. flagellifera (Brid.) Britt. (Britton in Brotherus, 1925).

Syn. *Pilotrichum flagellifera* Brid. (Bridel, 1827b).

Distr. Southern South America (Hosseus, 1940; Herzog, 1952a).

Pseudodistichium

P. atlanticum* Dix. (Dixon in Dixon and Fleischer, 1937).

Distr. Southern South America (Rovainen and Bartram, 1937; Robinson, 1964).

P. austro-georgicum Card. (Cardot, 1905b).

Distr. Southern South America (Rovainen and Bartram, 1937; Kühnemann, 1938).
Falkland Islands (Brotherus, 1924a; Kühnemann, 1938).
South Georgia (Cardot, 1905b, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus,
1924a; Steere, 1961b).

var. *longifolium* Broth. (Brotherus in Cardot and Brotherus, 1923).

Syn. *Pseudodistichium falklandicum* Card. (Cardot in Cardot and Brotherus, 1923) *nom.*
nud.

Distr. Southern South America (Rovainen and Bartram, 1937; Kühnemann, 1938).
Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

P. falklandicum Card.

Pseudodistichium austro-georgicum* var. *longifolium

P. fuegianum Roiv. (Rovainen in Rovainen and Bartram, 1937).

Distr. Southern South America (Rovainen and Bartram, 1937; Kühnemann, 1938).
Antarctic, peninsular region (Robinson, 1972).

P. taitaoense* Froehl. (Froehlich, 1953).

Distr. Southern South America (Froehlich, 1953; Seki, 1974).

Pseudoleskea

<i>P. antarctica</i> Card.	<i>Hygroamblystegium antarcticum</i>
<i>P. calochroa</i> Card.	<i>Hygroamblystegium calochroum</i>
<i>P. catenulatula</i> C. Muell.	<i>Leskeadelphus catenulatus</i>

*According to Robinson (1972), both of these species are synonyms of *Ditrichum conicum* (Mont.) Mitt.

<i>P. filum</i> (C. Muell.) Par.	<i>Hygroamblystegium filum</i>
<i>P. fuegiana</i> (Besch.) Broth.	<i>Hygroamblystegium fuegianum</i>
var. <i>gracilis</i> Card. et Broth.	<i>Hygroamblystegium fuegianum</i> var. <i>gracilis</i>
var. <i>skottsbergii</i> Card.	<i>Hygroamblystegium fuegianum</i> var. <i>skottsbergii</i>
<i>P. laplatae</i> C. Muell.	<i>Haplocladium microphyllum</i>
<i>P. lurida</i> Card.	<i>Hygroamblystegium luridum</i>
<i>P. microphylla</i> (Hedw.) Sauerb.	<i>Haplocladium microphyllum</i>
<i>P. platyphylla</i> Card.	<i>Hygroamblystegium platyphyllum</i>
<i>P. pseudogracilis</i> (C. Muell.) Par.	<i>Haplocladium microphyllum</i>
<i>P. siambonica</i> C. Muell.	<i>Rauiella siambonica</i>
<i>P. sordidoviridis</i> Card. et Broth.	<i>Hygroamblystegium sordidoviride</i>
<i>P. strictula</i> Card.	<i>Hygroamblystegium strictulum</i>
<i>P. uruguensis</i> C. Muell.	<i>Haplocladium microphyllum</i>

Pseudoscleropodium***P. pseudopurum*** (C. Muell.) Broth. (Brotherus, 1925).

Syn. *Cuspidaria pseudo-pura* C. Muell. (Mueller, 1897a).
Hypnum pseudo-purum (C. Muell.) Par. (Paris, 1900a).
Scleropodium pseudo-purum (C. Muell.) Broth. (Brotherus, 1908).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1905a; Brotherus, 1908, 1925; Kühnemann, 1938).

Psilotum***P. angulatum*** Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Seki, 1974).

P. antarcticum (C. Muell.) Par.***Psilotum trichodon******var. densifolium*** Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).
Falkland Islands (Cardot and Brotherus, 1923).

P. australe (Hook. f. et Wils.) Mitt. (Mitten, 1860).

Syn. *Polytrichum australe* Hook. f. et Wils. (Hooker and Wilson in Wilson, 1854).

Distr. Southern South America (Wilson, 1854).

P. compressum (Hook. f. et Wils.) Mitt.***Atrichopsis compressa******P. cuspidatum*** Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

P. gymnostomulum (C. Muell.) Par. (Paris, 1898).

Syn. *Catharinea gymnostomula* C. Muell. (Mueller, 1879a).
Oligotrichum gymnostomulum (C. Muell.) Kindb. (Kindberg, 1889).
Catharinea gymnostoma C. Muell. ex Par. (Mueller ex Paris, 1894) err.

Distr. Southern South America (Mueller, 1879a, 1885; Paris, 1898, 1905b; Brotherus, 1925; Hosseus, 1936, 1937; Kühnemann, 1938).
Falkland Islands (Mueller, 1885).

P. magellanicum Dus.***Psilotum trichodon******P. minimum*** (Card.) G. L. Smith (Smith, 1969a).

Syn. *Polytrichadelphus minimus* Card. (Cardot, 1905a).

Distr. Juan Fernandez (Brotherus, 1924b (as *Psilotum antarcticum* fide Smith, 1969a); Smith, 1969a).

Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923 (as *Psilopilum compressum* fide Smith, 1969a); Brotherus, 1925; Smith, 1969a).

P. subtapes Dus.

Psilopilum tapes var. *apiculatum*

P. tapes (C. Muell.) Par. (Paris, 1898).

Syn. *Catharinea tapes* C. Muell. (Mueller, 1890a).

Oligotrichum tapes (C. Muell.) Kindb. (Kindberg, 1891).

Distr. Southern South America (Cardot, 1913a; Brotherus, 1925).

South Georgia (Mueller, 1890a; Paris, 1898, 1905b; Cardot, 1908a; Brotherus, 1925; Steere, 1961b; Greene, 1973).

var. *apiculatum* Card. (Cardot, 1908a).

Syn. *Psilopilum subtapes* Dus. (Dusén in Cardot, 1908a) *nom. nud.*

Distr. Southern South America (Cardot, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938).

P. trichodon (Hook. f. et Wils.) Mitt. (Mitten, 1869).

Syn. *Polytrichum trichodon* Hook. f. et Wils. (Hooker and Wilson in Wilson, 1847).

Catharinea antarctica C. Muell. (Mueller, 1884 [1883]).

Oligotrichum antarcticum (C. Muell.) Kindb. (Kindberg, 1889).

Psilopilum antarcticum (C. Muell.) Par. (Paris, 1898) *fide* Smith, 1971.

Psilopilum trichodon (Hook. f. et Wils.) Mitt. var. *antarcticum* Mitt. ex Par. (Mitten ex Paris, 1898) *err.*

Psilopilum magellanicum Dus. (Dusén, 1905b).

Distr. Southern South America (Cardot, 1905a, 1908a; Dusén, 1905b; Brotherus, 1925; Kühnemann, 1938).

Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Paris, 1898, 1905b; Cardot, 1906a, 1908a; Dixon, 1920; Brotherus, 1925; Steere, 1961b; Greene, 1973).

Antarctic, peninsula region (Greene, 1968a; Greene and others, 1970).

var. *antarcticum* Mitt.

Psilopilum trichodon

Pterigynandrum

P. austro-alpinum C. Muell.

Leptopterigynandrum austro-alpinum

P. bicolor Lindb. ex Aongstr.

Erythrodontium squarrosum

P. julaceum Hedw.

Leucodon julaceus

P. longirostre Brid.

Diplostichum longirostre

P. smithii (Hedw.) Lam. et Cand.

Leptodon smithii

Pterobryon

P. densus Hornsch. (Hornschuch, 1840).

Syn. *Pterobryon densus* Hornsch. var. *condensatum* Lor. (Lorentz in Arzeni, 1954) *nom. nud.*

Pterobryon lorentzii C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

var. *condensatum* Lor.

Pterobryon densus

P. lorentzii C. Muell.

Pterobryon densus

P. stolonaceum C. Muell.

Pterobryopsis stolonacea

Pterobryopsis

P. stolonacea (C. Muell.) Broth. (Brotherus, 1906).

Syn. *Pterobryon stolonaceum* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1906, 1925; Herzog, 1952a).

var. *dilatata* Herz. (Herzog, 1952a) *nom. nud.*

Distr. Southern South America (Herzog, 1952a).

Pterogoniadelphus

P. montevidensis (C. Muell.) Fleisch.

Felipponea montevidensis

Pterogoniopsis

P. cylindrica C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

Pterogonium

P. julaceum (Hedw.) Schwaegr.

Leucodon julaceus

Pterygoneurum

P. cavifolium Jur.

Pterygoneurum ovatum

P. ovatum (Hedw.) Dix. (Dixon, 1933 [1934]).

Syn. *Gymnostomum ovatum* Hedw. (Hedwig, 1801).

Barbula ovata (Hedw.) Schimp. (Schimper in Lindberg, 1864a) *nom. inval.*

Pottia ovata (Hedw.) Fuernr. ex Lindb. (Fürnrohr ex Lindberg, 1864a) *nom. inval.*

Pottia cavifolia Ehrh. ex Fuernr. (Fürnrohr, 1829) *hom. illeg.*

Barbula cavifolia Schimp. (Schimper, 1860).

Desmatodon cavifolius Mitt. (Mitten, 1864) *nom. illeg.*

Pterygoneurum cavifolium Jur. (Juratzka, 1882) *nom. illeg.*

Pterygophyllum cavifolium Amann (Amann, 1916) *err.*

Tortula pusilla Mitt. (Mitten, 1869).

Distr. Southern South America (Dusén, 1906; Warnstorff, 1916).

P. roseae Williams

Aloina roseae

P. subsessile (Brid.) Jur. (Juratzka, 1882).

Syn. *Gymnostomum subsessile* Brid. (Bridel, 1806).

Schistidium subsessile (Brid.) Brid. (Bridel, 1819).

Anictangium subsessile (Brid.) Grev. et Arnott (Greville and Arnott, 1824).

Anoectangium subsessile (Brid.) Spreng. (Sprengel, 1827).

Pottia subsessilis (Brid.) B.S.G. (Bruch and others, 1843).

Pharomitrium subsessile (Brid.) Schimp. (Schimper, 1860).

Tortula subsessilis (Brid.) Mitt. (Mitten, 1869).

Anoectangium acaule (Web. et Mohr) Roehl. (Röhling, 1809).

Distr. Southern South America (Mitten, 1869; Jaeger, 1873; Paris, 1905b).

Pterygophyllum hom. illeg.

P. anomalum (Schwaegr.) Mitt.

Achrophyllum anomalum

var. *pallidum* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Welch, 1969; Seki, 1974).

P. apiculatum (Hook. f. et Wils.) Jaeg.

Eriopus apiculatus

P. arcuatum (Hedw.) Brid.

Hypnodendron arcuatum

P. cavifolium Amann

Pterygoneurum ovatum

P. chonoticum Mitt. (Mitten, 1885).

Distr. Southern South America (Mitten, 1885; Paris, 1905b; Cardot, 1908a; Brotherus, 1925; Welch, 1969).

P. crassirete Matteri (Matteri, 1972).

Syn. *Hookeria obscura* Mont. (Montagne, 1845a).

Pterygophyllum obscurum (Mont.) Mitt. (Mitten, 1869) *hom. illeg.*

Distr. Juan Fernandez (Brotherus, 1924b).

Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1851a; Lorentz, 1866; Mitten, 1869; Jaeger, 1877; Neger, 1899; Paris, 1905b; Cardot, 1908a; Herzog, 1923, 1939, 1960; Brotherus, 1925; Reimers, 1926; Welch, 1969; Matteri, 1972; Seki, 1974).

Falkland Islands (Cardot, 1905a, 1908a; Kühnemann, 1938).

P. dentatum (Hook. f. et Wils.) Dix.

Achrophyllum dentatum

P. denticulatum Mitt.

Achrophyllum dentatum

P. fragile Mitt. (Mitten, 1885).

Distr. Southern South America (Mitten, 1885; Paris, 1905b; Brotherus, 1925; Welch, 1969).

P. haesselianum Matteri (Matteri, 1972).

Distr. Southern South America (Matteri, 1972, 1975a; Seki, 1974).

P. lamellatum Dus. (Dusén, 1903a) *nom. nud.*

Syn. Hepaticina lamellata Fleisch. (Fleischer, 1922) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Fleischer, 1922).

P. magellanicum Besch. (Bescherelle, 1889).

Distr. Southern South America (Bescherelle, 1889; Hariot, 1891; Paris, 1905b; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938; Welch, 1969; Matteri, 1972, 1975a; Seki, 1974). Falkland Islands (Matteri, 1972).

var. oligodontium Matteri (Matteri, 1972).

Distr. Southern South America (Matteri, 1972, 1975a; Seki, 1974).

P. nigellum (Hook. f. et Wils.) Mitt.

Achrophyllum dentatum

P. obscurum (Mont.) Mitt.

Pterygophyllum crassirete

P. peruncinatum Fleisch. (Fleischer, 1922) *nom. nud.*

Syn. Hepaticina peruncinata Dus. (Dusén in Fleischer, 1922) *nom. nud.*

Distr. Southern South America (Fleischer, 1922).

P. rigidum (Schwaegr.) Brid.

Lepidopilum polytrichoides

P. sublimbatum (C. Muell.) Mitt.

Cyclodictyon sublimbatum

P. tenuinerve Broth.

Achrophyllum tenuinerve

Ptychomitrium

P. aligrimmioides Broth. (Brotherus, 1918).

Distr. Southern South America (Brotherus, 1918, 1925; Kühnemann, 1938).

P. balansae Besch. (Bescherelle, 1877).

Syn. *Glyphomitrium balansae* (Besch.) Broth. (Brotherus, 1902).

Brachysteleum brevifolium C. Muell. (Mueller, 1879a).

Ptychomitrium brevifolium (C. Muell.) Kindb. (Kindberg, 1889).

Glyphomitrium brevifolium (C. Muell.) Broth. (Brotherus, 1902).

Brachysteleum uruguense C. Muell. (Mueller, 1882).

Ptychomitrium uruguense (C. Muell.) Kindb. (Kindberg, 1889).

Glyphomitrium uruguense (C. Muell.) Broth. (Brotherus, 1902).

Distr. Southern South America (Mueller, 1879a, 1882; Brotherus, 1902, 1918, 1925; Kurtz, 1904; Paris, 1905b; Williams, 1930; Herter, 1933a, 1943; Hosseus, 1935b, 1936, 1939; Kühnemann, 1938; Herzog, 1952a; Bizot and Piovano, 1953; Piovano, 1954).

P. brevifolium (C. Muell.) Kindb.

Ptychomitrium balansae

P. chimbazense (Mitt.) Jaeg. var. *obtusifolium* (Thér.) Broth. (Brotherus, 1925).

Syn. *Brachysteleum chimbazense* (Mitt.) Thér. var. *obtusifolium* Thér. (Thériot, 1923b).

Distr. Southern South America (Thériot, 1923b; Brotherus, 1925).

P. crispatum (Hedw.) Jaeg. (Jaeger, 1874).

Syn. *Encalypta crispata* Hedw. (Hedwig, 1801).

Orthotrichum crispatum (Hedw.) Hook. et Grev. (Hooker and Greville, 1824a).

Ulota crispata (Hedw.) Sw. (Swartz in Hooker and Greville, 1824a) *nom. inval.*

Brachypodium crispatum (Hedw.) Brid. (Bridel, 1826).

Brachysteleum crispatum (Hedw.) Hornsch. (Hornschuch, 1840).

Notarisia crispata (Hedw.) Mont. (Montagne, 1840b).

Distr. Juan Fernandez (Montagne, 1850; Johow, 1896).

Southern South America (Hornschuch, 1840; Montagne, 1850).

P. cummingii Dub.**Ptychomitrium fernandesianum****P. deltiorii** (Thér.) Broth. (Brotherus, 1925).

Syn. *Brachysteleum deltiorii* Thér. (Thériot, 1921a).

Distr. Southern South America (Thériot, 1921a; Brotherus, 1925).

P. emersum (C. Muell.) Kindb. (Kindberg, 1889).

Syn. *Brachysteleum emersum* C. Muell. (Mueller, 1882).

Glyphomitrium emersum (C. Muell.) Broth. (Brotherus, 1902).

Distr. Southern South America (Mueller, 1882; Brotherus, 1902, 1918, 1925; Paris, 1905b; Kühnemann, 1938).

P. felipponei Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933a).

P. fernandesianum (Mitt.) Jaeg. (Jaeger, 1874).

Syn. *Glyphomitrium fernandesianum* Mitt. (Mitten, 1860).

Ptychomitrium cummingii Dub. (Duby, 1880a).

Brachysteleum cummingii (Dub.) Hamp. (Hampe, 1880a).

Glyphomitrium cummingii (Dub.) Broth. (Brotherus, 1902).

Distr. Juan Fernandez (Mitten, 1860, 1885; Jaeger, 1874; Johow, 1896; Paris, 1905b; Skottsberg, 1914; Brotherus, 1924b, 1925; Espinosa B., 1941; Robinson, 1975). Southern South America (Duby, 1880a, b; Brotherus, 1902, 1925; Paris, 1905b; Thériot, 1935a).

var. *majus* Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b).

P. hieronymi Besch. (Bescherelle, 1891).

Syn. *Glyphomitrium hieronymi* (Besch.) Broth. (Brotherus, 1902).

Distr. Southern South America (Bescherelle, 1891; Brotherus, 1902, 1925; Paris, 1905b; Kühnemann, 1938).

P. ligulatum (Mitt.) Jaeg. (Jaeger, 1874).

Syn. *Glyphomitrium ligulatum* Mitt. (Mitten, 1869).

Brachysteleum ligulatum (Mitt.) C. Muell. (Mueller, 1885).

Distr. Southern South America (Mitten, 1869; Jaeger, 1874; Mueller, 1885; Paris, 1905b; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

P. sellowianum (C. Muell.) Jaeg. (Jaeger, 1874).

Syn. *Brachysteleum sellowianum* C. Muell. (Mueller, 1849).

Glyphomitrium sellowianum (C. Muell.) Mitt. (Mitten, 1869).

Brachysteleum sellowii Aongstr. (Ångström, 1876a) *hom. illeg.*

Ptychomitrium sellowii C. Muell. ex Kindb. (Mueller ex Kindberg, 1888) *hom. illeg.*

Distr. Southern South America (Mueller, 1849; Mitten, 1869; Gibert, 1873; Jaeger, 1874; Krieger, 1904; Paris, 1905b; Brotherus, 1925; Herter, 1933a).

var. *uruguense* Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929).

- P. sellowii* C. Muell. ex Kindb. *Ptychomitrium sellowianum*
P. uruguense (C. Muell.) Kindb. *Ptychomitrium balansae*
- P. vernicosum* (C. Muell.) Kindb.** (Kindberg, 1889).
 Syn. *Brachysteleum vernicosum* C. Muell. (Mueller, 1882).
Glyphomitrium vernicosum (C. Muell.) Broth. (Brotherus, 1902).
 Distr. Southern South America (Mueller, 1882; Brotherus, 1902, 1925; Paris, 1905b; Kühnemann, 1938).
- Ptychomnion***
- P. aciculare* (Brid.) Mitt.** (Mitten, 1869).
 Syn. *Hypnum aciculare* Brid. (Bridel, 1801b).
Stereodon acicularis (Brid.) Mitt. (Mitten, 1860).
Hypnum cucullifolium P. Beauv. (Palisot de Beauvois, 1805a).
 Distr. Juan Fernandez (Montagne, 1835; Mitten, 1869, 1885; Johow, 1896; Paris, 1905b).
 Southern South America (Palisot de Beauvois, 1805a; Bridel, 1817, 1819, 1827b; Wilson and Hooker, 1847; Montagne, 1850; Mueller, 1851a; Sullivant, 1859; Lorentz, 1866; Mitten, 1869; Ångström, 1872; Jaeger and Sauerbeck, 1879a; Eaton, 1892; Paris, 1905b; Cardot, 1908a; Kühnemann, 1938; Bartram, 1952; Kühnemann and Gonçalves, 1975).
- P. cygnisetum* (C. Muell.) Kindb.** (Kindberg, 1888).
 Syn. *Hypnum cygnisetum* C. Muell. (Mueller, 1885).
Ectropothecium cygnisetum C. Muell. (Mueller in Mitten, 1883) nom. nud.
Eurhynchium cygnisetum C. Muell. (Mueller in Jaeger, 1878) nom. nud.
Hylocomium splendidissimum De Not. (De Notaris in Bescherelle, 1889) nom. nud.
 Distr. Juan Fernandez (Robinson, 1975).
 Southern South America (Mueller, 1885; Bescherelle, 1889; Hariot, 1891; Neger, 1899; Paris, 1905b; Cardot, 1908a; Campo, 1915; Thériot, 1915; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1940, 1954, 1957, 1960; Brotherus, 1924c, 1925; Reimers, 1926; Kühnemann, 1938; Skottsberg, 1950; Robinson, 1970b; Seki, 1974; Kühnemann and Gonçalves, 1975).
 var. *chilense* Broth. (Brotherus in Espinosa B., 1941) nom. nud.
 Distr. Southern South America (Espinosa B., 1941).
- P. densifolium* (Brid.) Jaeg.** (Jaeger and Sauerbeck, 1879a).
 Syn. *Hypnum densifolium* Brid. (Bridel, 1812).
Stereodon densifolius (Brid.) Mitt. (Mitten, 1860).
 Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Roivainen, 1934; Kühnemann, 1938; Robinson, 1970b; Seki, 1974; Kühnemann and Gonçalves, 1975).
 var. *gracile* Card. et Broth. (Cardot and Brotherus, 1923).
 Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann and Gonçalves, 1975).
- P. falcatulum* Broth.** (Brotherus, 1924b).
 Distr. Juan Fernandez (Brotherus, 1924b, 1925; Robinson, 1975; Kühnemann and Gonçalves, 1975).
 var. *gracilescens* Broth. (Brotherus, 1924b).
 Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Kühnemann and Gonçalves, 1975).
- P. fruticetorum* C. Muell.** (Mueller, 1898c).
 Distr. Southern South America (Bizot, 1967).
- P. horridum* Card. et Broth.** (Cardot and Brotherus, 1923).
 Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann and Gonçalves, 1975).

P. ptychocarpum (Schwaegr.) Mitt. (Mitten, 1869).

Syn. *Hypnum ptychocarpum* Schwaegr. (Schwaegrichen, 1844 [1845]).
Stereodon ptychocarpum (Schwaegr.) Mitt. (Mitten, 1860).
Leskeia gayana Mont. (Montagne, 1845a).
Hypnum gayanum (Mont.) Lor. (Lorentz, 1866).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).
 Southern South America (Mueller, 1844b, 1851a; Schwaegrichen, 1844; Montagne, 1845a, 1850, 1856; Lorentz, 1866; Mitten, 1869; Jaeger and Sauerbeck, 1879a; Bescherelle, 1889; Paris, 1905b; Cardot, 1908a; Thériot, 1918, 1935a; Campo, 1921; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1954, 1957, 1960; Brotherus, 1925; Reimers, 1926; Roivainen, 1934; Kühnemann, 1938; Bartram, 1952; Bizot, 1967; Robinson, 1970b; Seki, 1974; Kühnemann and Gonçalves, 1975).

P. subaciculare Besch. (Bescherelle, 1885a).

Distr. Juan Fernandez (Paris, 1905b; Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1970b; Kühnemann and Gonçalves, 1975).
 Southern South America (Bescherelle, 1885a, 1889; Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Robinson, 1970b; Seki, 1974).

Pulvinella**P. albicans** Broth. et Herz. (Brotherus and Herzog *in Herzog*, 1928).

Distr. Southern South America (Herzog, 1928).

Racomitrium

R. aciculare (Hedw.) Brid. var. *aquaticum* (Schrad.) Hueb.

Racomitrium aquaticum

R. andreaeoides Herz. (Herzog, 1954).

Distr. Southern South America (Herzog, 1954; Seki, 1974).

R. aquaticum (Schrad.) Brid. (Bridel, 1819).

Syn. *Trichostomum aquaticum* Brid. ex Schrad. (Bridel ex Schrader, 1803a).
Grimmia aquatica (Schrad.) C. Muell. (Mueller, 1849).
Trichostomum aciculare (Hedw.) P. Beauv. var. *aquaticum* (Schrad.) Web. et Mohr ex Brid. (Weber and Mohr ex Bridel, 1826) *nom. inval.*
Racomitrium aciculare (Hedw.) Brid. var. *aquaticum* (Schrad.) Hueb. (Hübener, 1833).
Trichostomum protensum Braun (Braun *in Schultz*, 1828).
Racomitrium protensum (Braun) Hueb. (Hübener, 1833).
Grimmia protensum (Braun) Mitt. (Mitten, 1879).

Distr. Southern South America (Paris, 1905b; Cardot, 1908a).

R. austro-canescens Dus. (Dusén, 1907).

Distr. Southern South America (Dusén, 1907; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

R. austro-georgicum Par. (Paris, 1895).

Syn. *Grimmia austro-patens* C. Muell. (Mueller, 1890a).
Racomitrium substenocladum Card. (Cardot, 1911d) *fide* Bell, 1973c.
Racomitrium skottsbergii Card. et Broth. (Cardot and Brotherus, 1923).
Grimmia austro-georgica C. Muell. ex Par. (Mueller ex Paris, 1905b) *nom. inval.*

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Roivainen, 1955a).
 South Georgia (Mueller, 1890a; Paris, 1895, 1905b; Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Steere, 1961b; Bell, 1974).
 Antarctic, peninsula region (Cardot, 1911b, d, 1913a; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961a; Bell, 1973c).

var. kranckii Roiv. (Roivainen, 1955a).

Distr. Southern South America (Roivainen, 1955a).

R. bartramii (Rov.) H. Robinson (Robinson, 1974).

Syn. *Bucklandia bartramii* Rov. (Rovainen, 1955b).
Bucklandiella bartramii Rov. (Rovainen, 1972).

Distr. Southern South America (Rovainen, 1955b, 1972; Robinson, 1974).

R. convolutum Mont.

Racomitrium crispulum

R. crispipilum (Tayl.) Jaeg. (Jaeger, 1874).

Syn. *Trichostomum crispipilum* Tayl. (Taylor, 1846).
Distr. Juan Fernandez (Robinson, 1975).

R. crispulum (Hook. f. et Wils.) Hook. f. et Wils. (Hooker and Wilson *in* Wilson, 1854).

Syn. *Dryptodon crispulus* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Grimmia crispula (Hook. f. et Wils.) C. Muell. (Mueller, 1849) *hom. illeg.*
Racomitrium convolutum Mont. (Montagne, 1845a).
Grimmia convoluta (Mont.) C. Muell. (Mueller, 1851a).
Grimmia nigrita C. Muell. (Mueller, 1849) *hom. illeg. fide* Bell, 1974.
Racomitrium nigritum Jaeg. (Jaeger, 1874) *fide* Bell, 1974.

Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b; Bartram, 1959; Robinson, 1975).
Southern South America (Montagne, 1845a, 1856; Hooker, 1847; Mueller, 1849, 1851a, 1885; Mitten, 1869; Jaeger, 1874; Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1938, 1954; Kühnemann, 1938; Bartram, 1952; Rovainen, 1955a; Seki, 1974).
South Georgia (Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b; Bell, 1974).
Antarctic, peninsula region (Robinson, 1972; Bell, 1973c).

var. rupestre (Hook. f. et Wils.) Dix. (Dixon, 1926).

Syn. *Dryptodon rupestris* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Grimmia rupestris (Hook. f. et Wils.) C. Muell. (Mueller, 1849).
Racomitrium rupestre (Hook. f. et Wils.) Hook. f. et Wils. (Hooker and Wilson *in* Wilson, 1854).

Distr. Southern South America (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1849, 1885; Sullivant, 1859; Mitten, 1869; Jaeger, 1874; Bescherelle, 1889; Eaton, 1892; Cardot, 1905a, 1908a; Paris, 1905b; Dusén, 1907; Cardot and Brotherus, 1923; Brotherus, 1924a; Thériot, 1925a; Rovainen, 1934, 1955a; Bartram, 1952; Seki, 1974).
South Georgia (Cardot, 1906a; Bell, 1974).

R. didymum (Mont.) Lor. (Lorentz, 1866).

Syn. *Grimmia didyma* Mont. (Montagne, 1845a).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Lorentz, 1866; Mitten, 1869; Jaeger, 1874; Paris, 1905b; Brotherus, 1924a; Reimers, 1926; Thériot, 1935a; Herzog, 1938, 1939, 1954, 1960; Seki, 1974).

R. flavescentia Card.

Racomitrium symphyodontum

R. flavo-pallidum Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1905b).

R. geminatum Rov. (Rovainen, 1955a).

Distr. Southern South America (Rovainen, 1955a).

R. geronticum C. Muell.

Racomitrium lanuginosum

R. glaciale Kindb.

Racomitrium lanuginosum

R. grimmiooides Herz. (Herzog, 1957).

Distr. Southern South America (Herzog, 1957).

R. heterostichoides Card. (Cardot, 1905a).

Syn. *Racomitrium horridum* Dus. (Dusén *in* Rovainen, 1955a) *nom. nud.*

Distr. Southern South America (Cardot, 1905a, 1908a; Dusén, 1907; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Roivainen, 1955a).
 Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).
 South Georgia (Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b; Bell, 1974).

var. acutifolium Dus.

Racomitrium subulifolium

***R. heterostichum* (Hedw.) Brid. (Bridel, 1819).**

Syn. *Trichostomum heterostichum* Hedw. (Hedwig, 1801).
Bryum heterostichum (Hedw.) Dicks. (Dickson, 1801).
Grimmia heterosticha (Hedw.) C. Muell. (Mueller, 1849).

Distr. Southern South America (Herzog, 1938).
 Falkland Islands (Wilson and Hooker, 1847; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

R. horridum Dus.

Racomitrium heterostichoides

R. hyalino-cuspidatum (C. Muell.) Kindb.

Grimmia hyalino-cuspidata

R. hypnoides Lindb.

Racomitrium lanuginosum

var. senile Lindb.

Racomitrium lanuginosum

***R. integrripilum* Dus. (Dusén, 1907).**

Distr. Southern South America (Dusén, 1907; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1954).

***R. laevigatum* Jaeg. (Jaeger, 1874).**

Syn. *Grimmia laevigata* Mitt. (Mitten, 1869) *hom. illeg.*

Distr. Juan Fernandez (Robinson, 1975).
 Southern South America (Mitten, 1869; Jaeger, 1874; Dusén, 1903a, 1907; Paris, 1905b; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

***R. lamprocarpum* (C. Muell.) Jaeg. (Jaeger, 1874).**

Syn. *Grimmia lamprocarpa* C. Muell. (Mueller, 1849).

Distr. Falkland Islands (Wilson and Hooker, 1847 (as *R. protensum* var. *fide* Mueller, 1849); Mueller, 1849; Jaeger, 1874; Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

***R. lanuginosum* (Hedw.) Brid. (Bridel, 1819).**

Syn. *Trichostomum lanuginosum* Hedw. (Hedwig, 1801).

Bryum lanuginosum (Hedw.) Brotero (Brotero, 1804).

Grimmia lanuginosa (Hedw.) C. Muell. (Mueller, 1849).

Bryum hypnoides L. ex With. (Linnaeus ex Withering, 1801) *nom. illeg.*

Trichostomum hypnoides Willd. ex P. Beauv. (Willdenow ex Palisot de Beauvois, 1822) *nom. illeg.*

Racomitrium hypnoides Lindb. (Lindberg, 1866 [1867]) *nom. illeg.*

Racomitrium hypnoides Lindb. *var. senile* Lindb. (Lindberg, 1866 [1867]) *nom. nud.*

Racomitrium senile Schimp. (Schimper in Jaeger, 1874) *nom. nud.*

Racomitrium puccioanum De Not. (De Notaris, 1859).

Grimmia glacialis C. Muell. (Mueller, 1890a) *hom. illeg.*

Racomitrium glaciale Kindb. (Kindberg, 1891).

Racomitrium geronticum C. Muell. (Mueller, 1869).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Bridel, 1819; Montagne, 1845b, 1850; Wilson and Hooker, 1847; Sullivant, 1859; De Notaris, 1859; Lorentz, 1866; Mitten, 1869; Mueller, 1869, 1885; Jaeger, 1874; Bescherelle, 1889; Hariot, 1891; Neger, 1899; Cardot, 1900, 1901, 1905a, 1908a; Dusén, 1903a, 1907; Krieger, 1904; Paris, 1905b; Middleton, 1909; Thériot, 1921a; Cardot and Brotherus, 1923; Brotherus, 1924a; Reimers, 1926; Herzog, 1938, 1940, 1954, 1957, 1960; Kühnemann, 1938; Bartram, 1952; Roivainen, 1955a; Gunckel, 1971; Seki, 1974).

Falkland Islands (Dumont d'Urville, 1825; Gaudichaud, 1825, 1826; Wilson and Hooker, 1847; Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938).

- South Georgia (Mueller, 1890a; Paris, 1905b; Cardot, 1906a, 1908a; Dixon, 1920, 1932; Brotherus, 1924a; Steere, 1961b; Bell, 1974, 1977b).
Antarctic, peninsula region (Dixon, 1935; Greene, 1968a; ? error see Bell, 1973c).
- R. limbatum* Bartr. *Racomitrium subnigritum*
- R. looseri* Thér.** (Thériot, 1934b [1935]).
Distr. Southern South America (Thériot, 1934b).
- R. loriforme* Dus. *Racomitrium willii*
- R. nigritum* Jaeg. *Racomitrium crispulum*
- R. pachydictyon* Card.** (Cardot, 1908a).
Syn. *Racomitrium symphyodontum* (C. Muell.) Par. var. *muticum* Card. (Cardot, 1905a).
Distr. Southern South America (Cardot, 1905a (and as *R. flavescentia* pro parte and *R. rupestre* pro parte fide Cardot, 1908a), 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1938; Kühnemann, 1938; Roivainen, 1955a; Seki, 1974).
South Georgia (Cardot, 1908a; Brotherus, 1924a; Steere, 1961b; Bell, 1974).
- R. pachyneuron* Herz.** (Herzog, 1957).
Distr. Southern South America (Herzog, 1939, 1954, 1957, 1960; Seki, 1974).
- R. patagonicum* Par. (Paris, 1905b) *nom. nud.*
Syn. *Racomitrium subnigritum* Dus. (Dusén in Paris, 1905b) *nom. nud.*
Distr. Southern South America (Paris, 1905b).
- R. plicatum* Herz.** (Herzog in Donat, 1936a).
Distr. Southern South America (Donat, 1936a; Seki, 1974).
- R. protensum* (Braun) Hueb. *Racomitrium aquaticum*
- R. ptychophyllum* (Mitt.) Mitt.** (Mitten in Hooker, 1867).
Syn. *Grimmia ptychophylla* Mitt. (Mitten in Lindsay, 1866).
Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938; Bartram, 1952).
Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).
South Georgia (Cardot, 1906a, 1908a; Steere, 1961b; Bell, 1974).
- R. puccioanum* De Not. *Racomitrium lanuginosum*
- R. rupestre* (Hook. f. et Wils.) Hook. f. et Wils. *Racomitrium crispulum* var. *rupestre*
- R. scabrifolium* Bartr. *Racomitrium willii*
- R. senile* Schimp. *Racomitrium lanuginosum*
- R. skottsbergii* Card. et Broth. *Racomitrium austro-georgicum*
- R. stenocladooides* Roiv.** (Roivainen, 1955a).
Distr. Southern South America (Roivainen, 1955a).
- R. stenocladum* Dus.** (Dusén, 1907).
Distr. Southern South America (Dusén, 1907; Cardot, 1908a; Brotherus, 1925; Herzog, 1938; Kühnemann, 1938; Seki, 1974).
- var. *obtusum*** Dus. (Dusén, 1907).
Distr. Southern South America (Dusén, 1907; Cardot, 1908a; Kühnemann, 1938).
- R. striatipilum* Card.** (Cardot, 1905a).
Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941).
Southern South America (Cardot, 1905a, 1908a; Dusén, 1907; Cardot and Brotherus, 1923; Brotherus, 1924a; Thériot, 1935a; Herzog, 1938; Kühnemann, 1938; Roivainen, 1955a; Seki, 1974).
Falkland Islands (Cardot and Brotherus, 1923).
South Georgia (Cardot, 1906a, 1908a; Dixon, 1934; Steere, 1961b; Bell, 1974).

R. subcrispipilum (C. Muell.) Jaeg. (Jaeger, 1874).

Syn. *Grimmia subcrispila* C. Muell. (Mueller, 1862).

Distr. Southern South America (Mueller, 1862; Paris, 1905b; Brotherus, 1924a).

R. sublamprocarpum (C. Muell.) Par. (Paris, 1898).

Syn. *Grimmia sublamprocarpa* C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Paris, 1898, 1905b).

R. subnigritum (C. Muell.) Par. (Paris, 1898).

Syn. *Grimmia subnigrita* C. Muell. (Mueller, 1885).

Racomitrium limbatum Bartr. (Bartram, 1946).

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959).

Southern South America (Mueller, 1885; Cardot, 1905a, 1908a; Paris, 1905b; Dusén, 1907; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Bartram, 1946, 1952; Herzog, 1954; Rovainen, 1955a; Seki, 1974).

Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

R. subnigritum Dus.

Racomitrium patagonicum

R. subrupestre Rov. (Rovainen, 1955a).

Distr. Southern South America (Rovainen, 1955a).

R. substenocladum Card.

Racomitrium austro-georgicum

R. substriatipilum Rov. (Rovainen, 1955a).

Distr. Southern South America (Rovainen, 1955a).

R. subulifolium Card. (Cardot, 1908a).

Syn. *Racomitrium heterostichoides* Card. var. *acutifolium* Dus. (Dusén, 1907).

Distr. Southern South America (Dusén, 1907; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Rovainen, 1955a; Herzog, 1957).

R. symphyodon Jaeg. (Jaeger, 1874).

Syn. *Grimmia symphyodon* Mitt. (Mitten, 1882) *nom. illeg.*

Distr. Southern South America (Jaeger, 1874).

R. symphyodontum (C. Muell.) Par. (Paris, 1897).

Syn. *Grimmia symphydonta* C. Muell. (Mueller, 1849).

Racomitrium flavescentia Card. (Cardot, 1900).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941).

Southern South America (Wilson and Hooker, 1847 (as *R. fasciculare* vars 2 and 3 *fide* Mueller 1849); Mueller, 1849, 1855; Cardot, 1900, 1905a *pro parte*, 1908a, 1910; Paris, 1905b; Dusén, 1907; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Herzog, 1954, 1957; Rovainen, 1955a; Seki, 1974).

Falkland Islands (Cardot, 1905a, 1908a; Paris, 1905b; Brotherus, 1924a; Kühnemann, 1938).

var. *dusenii* Card. (Cardot, 1908a) *nom. nud.*

Distr. Southern South America (Dusén, 1907 (as *R. symphyodontum* var. *muticum* *fide* Cardot, 1908a); Cardot, 1908a).

var. *muticum* Card.

Racomitrium pachydictyon

R. willii (C. Muell.) Kindb. (Kindberg, 1891).

Syn. *Grimmia willii* C. Muell. (Mueller, 1890a).

Racomitrium loriforme Dus. (Dusén, 1907).

Racomitrium scabrifolium Bartr. (Bartram, 1946).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941).

Southern South America (Paris, 1905b; Dusén, 1907; Cardot, 1908a; Cardot and

Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Bartram, 1946; Roivainen, 1955a; Seki, 1974).
South Georgia (Mueller, 1890a; Paris, 1905b; Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b; Bell, 1974).

Racopilum

- R. anomalum* Schwaegr. **Achrophyllum anomalum**
R. arcuatum (Hedw.) Jaeg. **Hypnodendron arcuatum**

R. chilense Dus. ex Herz. (Dusén ex Herzog, 1938).

Distr. Southern South America (Paris, 1905b; Brotherus, 1925; Herzog, 1938).

R. fernandezianum Card. (Cardot in Thériot, 1921b).

Distr. Juan Fernandez (Skottsberg, 1914; Thériot, 1921b, 1927b; Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).
Southern South America (Herzog, 1954).

R. leptocarpum Wils. (Wilson, 1854).

Distr. Juan Fernandez (Wilson, 1854).

- R. ornithopodioides* (Brid.) Lindb. **Racopilum tomentosum**

R. tomentosum (Hedw.) Brid. (Bridel, 1827b).

Syn. *Hypnum tomentosum* Hedw. (Hedwig, 1801).
Hypopterygium tomentosum (Hedw.) C. Muell. (Mueller, 1850).
Hypnum ornithopodioides Brid. (Bridel, 1801b).
Leskeia ornithopodioides (Brid.) Brid. (Bridel, 1812).
Racopilum ornithopodioides (Brid.) Lindb. (Lindberg, 1868d).

Distr. Juan Fernandez (Montagne, 1835; Mitten, 1885; Johow, 1896; Paris, 1905b; Skottsberg, 1914; Robinson, 1975).
Southern South America (Bridel, 1801b, 1812, 1817, 1827b; Palisot de Beauvois, 1805a; Montagne, 1850; Mueller, 1879a; Krieger, 1904; Paris, 1905b; Brotherus, 1918; Kühnemann, 1938; Hosseus, 1940; Herzog, 1952a).

Rauia hom. illeg.

- R. firmula* (C. Muell.) Broth. **Rauiella firmula**
R. microcalycina Broth. ex Par. **Rauiella niveo-calycina**
R. niveo-calycina (C. Muell.) Broth. **Rauiella niveo-calycina**
R. siambonica (C. Muell.) Broth. **Rauiella siambonica**

Rauiella

R. firmula (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1962).

Syn. *Thuidium firmulum* C. Muell. (Mueller, 1897a).
Rauia firmula (C. Muell.) Broth. (Brotherus, 1907).

Distr. Southern South America (Mueller, 1897a; Paris, 1906; Brotherus, 1907, 1925; Hosseus, 1938a, d, 1939; Kühnemann, 1938).

R. niveo-calycina (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1962).

Syn. *Thuidium niveo-calycinum* C. Muell. (Mueller, 1897a).
Rauia niveo-calycina (C. Muell.) Broth. (Brotherus, 1907).
Rauia microcalycina Broth. ex Par. (Brotherus ex Paris, 1909) err.

Distr. Southern South America (Mueller, 1897a; Paris, 1906; Brotherus, 1907, 1925; Kühnemann, 1938; Herzog, 1952a).

R. siambonica (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1962).

Syn. *Pseudoleskea siambonica* C. Muell. (Mueller, 1897a).
Rauia siambonica (C. Muell.) Broth. (Brotherus, 1907).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1907, 1925; Kühnemann, 1938).

Renauldia

R. chilensis Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926).

Rhabdowesia

R. crispata (With.) Lindb. (Lindberg, 1871).

Syn. *Bryum crispatum* Dicks. ex With. (Dickson ex Withering, 1801).

Distr. Juan Fernandez (Robinson, 1975).

R. cyathicarpa (Mont.) Mitt.

Amphidium tortuosum

R. fugax (Hedw.) B.S.G. (Bruch and others, 1846).

Syn. *Weisia fugax* Hedw. (Hedwig, 1801).

Grimmia striata Schrad. (Schrader, 1803b) nom. illeg.

Oncophorus striatus (Schrad.) Lindb. (Lindberg, 1879).

Distr. Juan Fernandez (Mitten, 1885; Johow, 1896; Paris, 1905b; Lawton, 1961).

R. sphaerothecia (C. Muell.) Kindb.

Amphidium tortuosum

Rhachithecium

R. demissum (C. Muell.) Broth.

Rhachithecium perpusillum

R. perpusillum (Thwait. et Mitt.) Broth. (Brotherus, 1909).

Syn. *Zygodon perpusillus* Thwait. et Mitt. (Thwaites and Mitten in Mitten, 1873).

Hypnodon perpusillus (Thwait. et Mitt.) C. Muell. (Mueller, 1897a).

Hypnodon demissus C. Muell. (Mueller, 1897a).

Decodon demissus C. Muell. (Mueller, 1897a) nom. nud.

Rhachithecium demissum (C. Muell.) Broth. (Brotherus, 1909).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1909, 1925; Kühnemann, 1938).

Rhacocarpus

R. australis (Hook. f. et Wils.) Par.

Rhacocarpus purpurascens

R. crasso-limbatus (C. Muell.) Par.

Rhacocarpus purpurascens

R. humboldtii (Hook.) Lindb.

Rhacocarpus purpurascens

var. *argentinicus* (C. Muell.) Par.

Rhacocarpus purpurascens* var. *argentinicus

R. orbiculatus (Mitt.) Par. (Paris, 1898).

Syn. *Hedwigia orbiculata* Mitt. (Mitten, 1869).

Harrisonia orbiculata (Mitt.) Jaeg. (Jaeger, 1876),

Distr. Southern South America (Mitten, 1869; Jaeger, 1876; Mueller, 1897d; Paris, 1898, 1905b; Brotherus, 1925).

R. patagonicus Broth. (Brotherus in Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

R. purpurascens (Brid.) Par. (Paris, 1900a).

Syn. *Hypnum purpurascens* Brid. (Bridel, 1812).

Harrisonia purpurascens (Brid.) C. Muell. (Mueller, 1897d).

Anictangium humboldtii Hook. (Hooker, 1816).

Hedwigia humboldtii (Hook.) Hook. (Hooker, 1819).

Anoectangium humboldtii (Hook.) Brid. (Bridel, 1827b).

Harrisonia humboldtii (Hook.) Spreng. (Sprengel, 1827).

Neckera humboldtii (Hook.) C. Muell. (Mueller, 1850).

Rhacocarpus humboldtii (Hook.) Lindb. (Lindberg, 1862 [1863]).

Braunia humboldtii (Hook.) Hook. f. (Hooker, 1867).

Anoectangium humboldtii (Hook.) Brid. var. *australis* Hook. f. et Wils. (Hooker and Wilson in Wilson and Hooker, 1845).

Hedwigia humboldtii (Hook.) Hook. var. *australis* (Hook. f. et Wils.) Wils. (Wilson, 1854).

Harrisonia australis (Hook. f. et Wils.) Hamp. (Hampe, 1860).

Rhacocarpus australis (Hook. f. et Wils.) Par. (Paris, 1898).

Harrisonia crasso-limbata C. Muell. (Mueller, 1897d).

Rhacocarpus crasso-limbatus (C. Muell.) Par. (Paris, 1900a).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Robinson, 1975).
 Southern South America (Wilson and Hooker, 1847; Mitten, 1869; Mueller, 1879a, 1885, 1897d; Paris, 1900a, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938; Seki, 1974).
 Falkland Islands (Paris, 1900a, 1905b; Cardot, 1908a; Brotherus, 1925).

var. *argentinicus* (Brid.) Wijk et Marg. (van der Wijk and Margadant, 1961).

Syn. *Harrisonia humboldtii* (Hook.) Spreng. var. *argentinica* C. Muell. (Mueller, 1879a).
Rhacocarpus humboldtii (Hook.) Lindb. var. *argentinicus* (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1879a; Paris, 1898, 1905b; Kühnemann, 1938).

Rhamphidium

R. laetum (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Trichostomum laetum* Kunz. ex C. Muell. (Kunze ex Mueller, 1849).
Tortula laeta (C. Muell.) Mitt. (Mitten, 1869).

Distr. Juan Fernandez (Montagne, 1850).
 Southern South America (Mueller, 1849; Mitten, 1869; Jaeger, 1873; Brotherus, 1902, 1924a; Paris, 1905b).

Raphidorrhynchium

R. aberrans (Broth.) Broth.

Sematophyllum aberrans

R. amoenum (Hedw.) Fleisch. (Fleischer, 1923).

Syn. *Hypnum amoenum* Hedw. (Hedwig, 1801).
Isothecium amoenum (Hedw.) Brid. (Bridel, 1827b).
Sematophyllum amoenum (Hedw.) Mitt. (Mitten, 1869).
Raphidostegium amoenum (Hedw.) Jaeg. (Jaeger, 1878).
Raphidostegium amaenum (Hedw.) Par. (Paris, 1898) err.
Hypnum leptorhynchum Brid. (Bridel, 1812) hom. illeg.

Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1851a; Mitten, 1869; Jaeger, 1878; Paris, 1898, 1905b; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).
 Falkland Islands (Kühnemann, 1938).

R. berberidis (Dus.) Broth. (Brotherus, 1925).

Syn. *Ectropothecium berberidis* Dus. (Dusén, 1905b).
Raphidostegium berberidis (Dus.) Broth. (Brotherus, 1908).

Distr. Southern South America (Dusén, 1903a, 1905b; Brotherus, 1908, 1925; Cardot, 1008a; Seki, 1974).

R. brachypus (Hamp.) Broth. (Brotherus, 1925).

Syn. *Hypnum brachypus* Hamp. (Hampe, 1879).

Distr. Southern South America (Herzog, 1952a).

R. callidum (Mont.) Broth. (Brotherus, 1925).

Syn. *Hypnum callidum* Mont. (Montagne, 1845a).
Sematophyllum callidum (Mont.) Mitt. (Mitten, 1869).
Raphidostegium callidum (Mont.) Jaeg. (Jaeger, 1878).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1851a; Lorentz, 1866; Mitten, 1869; Jaeger, 1878; Paris, 1905b; Brotherus, 1908, 1924c; Cardot, 1908a; Campo, 1915; Thériot, 1917b, 1918; Costes, 1921; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1954, 1957, 1960; Reimers, 1926; Kühnemann, 1938; Espinosa B., 1941; Skottsberg, 1950; Bartram, 1952; Seki, 1974).

R. dendroligotrichum (Dus.) Broth. (Brotherus, 1925).

Syn. *Rhaphidostegium dendroligotrichum* Dus. (Dusén, 1905b).
Rhaphidostegium polytrichadelphus Dus. (Dusén, 1903a) nom. nud.

Distr. Southern South America (Dusén, 1903a, 1905b; Paris, 1905b; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938; Herzog, 1954, 1960; Seki, 1974).

R. leptophyllum (Mitt.) Broth. (Brotherus, 1925).

Syn. *Sematophyllum leptophyllum* Mitt. (Mitten, 1869).
Rhaphidostegium leptophyllum (Mitt.) Jaeg. (Jaeger, 1878).

Distr. Southern South America (Mitten, 1869; Jaeger, 1878; Paris, 1905b; Brotherus, 1925).

R. masafuerae (Broth.) Broth.

Sematophyllum masafuerae

R. minutum (Broth.) Broth. (Brotherus, 1925).

Syn. *Sematophyllum minutum* Broth. (Brotherus, 1900).

Distr. Southern South America (Herzog, 1952a).

R. noduliferum (Mitt.) Broth. (Brotherus, 1925).

Syn. *Sematophyllum noduliferum* Mitt. (Mitten, 1869).
Rhaphidostegium noduliferum (Mitt.) Jaeg. (Jaeger, 1878).

Distr. Southern South America (Mitten, 1869; Jaeger, 1878; Paris, 1905b; Cardot, 1908a; Brotherus, 1925).

R. pallido-nitidum (C. Muell.) Broth. (Brotherus, 1925).

Syn. *Cupressina pallido-nitida* C. Muell. (Mueller, 1897a).
Ectropothecium pallido-nitidum (C. Muell.) Par. (Paris, 1900a).
Rhaphidostegium pallido-nitidum (C. Muell.) Broth. (Brotherus, 1908).

Distr. Southern South America (Mueller, 1897a; Paris, 1900a, 1904b; Brotherus, 1908, 1925; Kühnemann, 1938).

R. scorpiurus (Mont.) Broth. (Brotherus, 1925).

Syn. *Hypnum scorpiurus* Mont. (Montagne, 1845a).
Sematophyllum scorpiurus (Mont.) Mitt. (Mitten, 1869).
Rhaphidostegium scorpiurus (Mont.) Jaeg. (Jaeger, 1878).
Sematophyllum aureo-nitidum Bartr. (Bartram, 1946) fide Seki, 1974.

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1851a; Lorentz, 1866; Mitten, 1869; Jaeger, 1878; Paris, 1905b; Herzog, 1923, 1938, 1954, 1960; Brotherus, 1925; Bartram, 1946; Seki, 1974).

R. subsimplex (Hedw.) Broth.

Sematophyllum subsimplex

Rhaphidostegium hom. illeg.

- | | |
|---|--------------------------------------|
| <i>R. aberrans</i> Broth. | Sematophyllum aberrans |
| <i>R. amaenum</i> (Hedw.) Par. | Rhaphidorrhynchium amoenum |
| <i>R. amoenum</i> (Hedw.) Jaeg. | Rhaphidorrhynchium amoenum |
| <i>R. ampullatum</i> Par. | Sematophyllum ampullulatum |
| <i>R. ampullulatum</i> (C. Muell.) Broth. | Sematophyllum ampullulatum |
| <i>R. aureo-viride</i> (C. Muell.) Par. | Sematophyllum aureo-viride |
| <i>R. berberidis</i> (Dus.) Broth. | Rhaphidorrhynchium berberidis |
| <i>R. brachycladulum</i> Broth. | Sematophyllum brachycladulum |
| <i>R. bracteatum</i> (C. Muell.) Kindb. | Sematophyllum bracteatum |
| <i>R. caespitosoides</i> Broth. | Sematophyllum aberrans |
| <i>R. caespitosum</i> (Hedw.) Besch. | Sematophyllum caespitosum |
| <i>R. callidum</i> (Mont.) Jaeg. | Rhaphidorrhynchium callidum |
| <i>R. catilliforme</i> (C. Muell.) Par. | Sematophyllum bracteatum |

R. cochleatulum (C. Muell.) Par. (Paris, 1898).

Syn. *Aptychus cochleatus* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b).

R. condensatum (C. Muell.) Par. (Paris, 1898).

Syn. *Aptychus condensatus* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b).

R. dendroligotrichum Dus.

Rhaphidorrhynchium dendroligotrichum

R. diaphanodictyon (C. Muell.) Par. (Paris, 1898).

Syn. *Aptychus diaphanodictus* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b).

R. grandicellulosum (C. Muell.) Par.

Sematophyllum grandicellulosum

R. laxo-alare (C. Muell.) Par.

Entodon laxo-alaris

R. leptophyllum (Mitt.) Jaeg.

Rhaphidorrhynchium leptophyllum

R. leucocytus (C. Muell.) Jaeg.

Sematophyllum leucocytus

R. loxense (Hook.) Jaeg.

Sematophyllum caespitosum

R. macrocytus (C. Muell.) Par.

Sematophyllum macrocytus

R. masafuerae Broth.

Sematophyllum masafuerae

R. micrangingium (C. Muell.) Par.

Sematophyllum micrangingium

R. microcarpoides (C. Muell.) Kindb.

Sematophyllum microcarpoides

R. micropyxis (C. Muell.) Kindb.

Trichosteleum sp.

R. nanocephalum (C. Muell.) Par.

Sematophyllum nanocephalum

var. *subglauculum* (C. Muell.) Par.

Sematophyllum nanocephalum var. *subglauculum*

R. nanum Kindb. (Kindberg, 1891) *nom. nud.*

Syn. *Hypnum nanum* C. Muell. (Mueller in Balansa and Savès, 1887) *nom. nud.*
Sematophyllum nanum Hert. (Herter, 1933a) *nom. nud.*

Distr. Southern South America (Felippone, 1912; Herter, 1933a).

R. noduliferum (Mitt.) Jaeg.

Rhaphidorrhynchium noduliferum

R. pallens Dus.

Hypnum pallens

R. pallido-nitidum (C. Muell.) Broth.

Rhaphidorrhynchium pallido-nitidum

R. paraguariense C. Muell. ex Broth. (Mueller ex Brotherus, 1900).

Distr. Southern South America (Brotherus, 1918; Kühnemann, 1938).

R. patagonicum Broth. (Brotherus in Cardot and Brotherus, 1923).

Distr. Southern South America (Dusén, 1903a; Paris, 1905b; Cardot and Brotherus, 1923;
Kühnemann, 1938; Seki, 1974).

R. polytrichadelphus Dus.

Rhaphidorrhynchium dendroligotrichum

R. prostratum Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

R. purpureo-axis Dus. (Dusén in Paris, 1905b) *nom. nud.*

Distr. Southern South America (Paris, 1905b).

R. scorpiurus (Mont.) Jaeg.

Rhaphidorrhynchium scorpiurus

R. secundifolium (C. Muell.) Jaeg.

Drepanocladus secundifolius

R. serifolium (C. Muell.) Par.

Sematophyllum serifolium

R. stenopyxidium (C. Muell.) Kindb.

Sematophyllum stenopyxidium

R. strictipes Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a).

R. subsimplex (Hedw.) Besch.

Sematophyllum subsimplex

R. temperatum (C. Muell.) Par.

Sematophyllum temperatum

R. tenerifolium (C. Muell.) Par.

Sematophyllum tenerifolium

R. tristifolium Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a).

Rhaphidostichum

R. gunckelii Thér. (Thériot, 1930).

Distr. Southern South America (Thériot, 1930).

Rhexophyllum

R. lacinatum Herz. (Herzog, 1916a).

Distr. Southern South America (Bartram, 1965).

Rhizogonium

R. brevifolium Broth. (Brotherus, 1891).

Syn. *Mnium brevifolium* (Broth.) C. Muell. (Mueller, 1901a [1900]).

Distr. Southern South America (Thériot, 1935a).

R. lindigii (Hamp.) Mitt. (Mitten, 1869).

Syn. *Mnium lindigii* Hamp. (Hampe, 1865).

Distr. Southern South America (Herzog, 1960).

R. longicolle C. Muell. (Mueller in Kindberg, 1891) *nom. nud.*

Syn. *Mnium longicolle* C. Muell. (Mueller in Paris, 1898) *nom. nud.*

Distr. Southern South America (Paris, 1898).

R. mnioides (Hook.) Wils. (Wilson, 1854).

Syn. *Hypnum mnioides* Hook. (Hooker, 1820).

Mnium polycarpum C. Muell. (Mueller, 1848a) *nom. illeg.*

Rhizogonium polycarpum Schimp. (Schimper in Mueller, 1889) *nom. illeg.*

Aulacomnium chilense C. Muell. (Mueller, 1843b).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Robinson, 1975).

Southern South America (Hooker, 1820; Bridel, 1827b; Schwaegrichen, 1829; Mueller, 1843b, 1848a, 1885, 1889; Montagne, 1845b, 1850; Wilson and Hooker, 1847; Lorentz, 1866; Mitten, 1869; Ångström, 1872; Bescherelle, 1889; Neger, 1899; Dusén, 1903a; Cardot, 1905a, 1908a; Paris, 1905b; Thériot, 1915, 1918, 1934b; Campo, 1921; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1940, 1954, 1957, 1960; Brotherus, 1924a; Reimers, 1926; Roivainen, 1934; Kühnemann, 1938; Bartram, 1952; Bizot, 1967; Seki, 1974).

R. novae-hollandiae (Brid.) Brid. var. *patagonicum* Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Robinson, 1975).

Southern South America (Cardot and Brotherus, 1923).

R. polycarpum Schimp.

Rhizogonium mnioides

R. reticulatum (Hook. f. et Wils.) Mitt.

Philonotis vagans

R. spiniforme (Hedw.) Bruch (Bruch in Krauss, 1846).

Syn. *Hypnum spiniforme* Hedw. (Hedwig, 1801).

Distr. Southern South America (Herter, 1943).

R. spininervium (Hook.) Schimp.

Hypnodendron arcuatum

R. subbasilare (Hook.) Schimp.

Goniobryum subbasilare

Rhodobryum*R. andicola* (Hook.) Par.*Bryum andicola**R. aubertii* (Schwaegr.) Thér. (Thériot, 1922).

Syn. *Mnium aubertii* Schwaegr. (Schwaegrichen, 1816).
Bryum aubertii (Schwaegr.) Brid. (Bridel, 1819).

Distr. Southern South America (Montagne, 1839c; Kühnemann, 1938 (as *M. augerii*)).

R. beyrichianum (Hornschr.) C. Muell. (Mueller in Hampe, 1874 [1875]).

Syn. *Mnium beyrichianum* Hornsch. (Hornschuch, 1840).
Bryum beyrichianum (Hornschr.) C. Muell. (Mueller, 1848a).
Bryum beyrichii Aongstr. (Ångström, 1876a) *nom. illeg.*
Bryum beyrichiae Hornsch. ex Kindb. (Hornschuch ex Kindberg, 1888) *nom. illeg.*

Distr. Southern South America (Mueller, 1848a; Montagne, 1850; Mitten, 1869; Paris, 1905b; Brotherus, 1924a; Reimers, 1926; Hosseus, 1940; Herter, 1943; Herzog, 1952a).

R. chilense Thér. (Thériot, 1924).

Distr. Southern South America (Thériot, 1924; Brotherus, 1925).

R. gracilescens C. Muell.*Bryum densifolium**R. hieronymi* (C. Muell.) Par. (Paris, 1898).

Syn. *Bryum hieronymi* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1898, 1905b; Brotherus, 1924a; Kühnemann, 1938).

R. lechleri (C. Muell.) Par.*Bryum andicola**R. lorentzianum* (C. Muell.) Par. (Paris, 1898).

Syn. *Bryum lorentzianum* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1898, 1905b; Brotherus, 1924a; Hosseus, 1938b, c, d; Kühnemann, 1938).

R. platense (C. Muell.) Par. (Paris, 1898).

Syn. *Bryum platense* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1924a; Kühnemann, 1938).

R. rigidum (Hornschr.) Par.*Bryum rigidum**R. roseolum* (C. Muell.) Par.*Bryum roseolum**R. roseum* (Hedw.) Limpr. (Limprecht, 1892).

Syn. *Mnium roseum* Hedw. (Hedwig, 1801).

Distr. Southern South America (Montagne, 1839c; Kühnemann, 1938).

R. subintegrum Broth. (Brotherus in Krieger, 1904) *nom. nud.*

Distr. Southern South America (Krieger, 1904).

Rhyncho-hypnum*R. leucostegum* (Brid.) Hamp.*Stereophyllum leucostegum**R. sellowii* (Hornschr.) Hamp.*Rhynchosstegium sellowii***Rhynchosstegiella***R. acanthophylla* (Mont.) Broth.*Eurhynchiella acanthophylla*

var. robusta Thér. (Thériot, 1917b).

Distr. Southern South America (Thériot, 1917b).

R. chilensis Thér. (Thériot, 1934b [1935]).

Distr. Southern South America (Thériot, 1934b).

R. fuegiana (Card.) Broth.

Oxyrrhynchium fuegianum

R. leptoneura Card. (Cardot in Skottsberg, 1914) *nom nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

Rhynchosstegium

R. acanthophyllum (Mont.) Jaeg.

Eurhynchiella acanthophylla

R. altisetum C. Muell.

Oxyrrhynchium altisetum

R. aquaticum Jaeg.

Platyhypnidium aquaticum

R. berteroanum (Mont.) Jaeg.

Catagoniopsis berteroana

R. berteroii Kindb.

Catagoniopsis berteroana

R. beskeanum (C. Muell.) Jaeg. (Jaeger, 1878).

Syn. *Hypnum beskeanum* C. Muell. (Mueller, 1857).

Distr. Southern South America (Herter, 1933a).

R. brevicuspis C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Hosseus, 1938a, b, d; Kühnemann, 1938).

R. byssicladum Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a).

R. cacticola (C. Muell.) Par. (Paris, 1898).

Syn. *Hypnum cacticola* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1898, 1905b; Brotherus, 1925; Kühnemann, 1938).

R. campylocladulum C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Hosseus, 1938a, b, d; Kühnemann, 1938).

R. capillirameum Dus. (Dusén in Paris, 1905b) *nom. nud.*

Distr. Southern South America (Paris, 1905b).

R. complanum (Mitt.) Jaeg. (Jaeger, 1878).

Syn. *Hypnum complanum* Mitt. (Mitten, 1869).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Mitten, 1869; Jaeger, 1878; Paris, 1905b; Brotherus, 1925; Herzog, 1954, 1960).

R. concavifolium Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

R. confertum (Dicks.) B.S.G. (Bruch and others, 1852).

Syn. *Hypnum confertum* Dicks. (Dickson, 1801).

Distr. Southern South America (Montagne, 1850).

R. drepanocladoides (C. Muell.) Kindb. (Kindberg, 1891).

Syn. *Hypnum drepanocladoides* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

R. fissidens (C. Muell.) Kindb. (Kindberg, 1891).

Syn. *Hypnum fissidens* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

R. glauco-virescens (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum glauco-virescens* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905b; Hosseus, 1935b, 1937; Kühnemann, 1938).

R. globipyxis (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum globipyxis* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

R. homaliocaulon (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum homaliocaulon* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905b; Brotherus, 1925; Hosseus, 1935a, 1937; Kühnemann, 1938).

R. leptopteridium C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

R. leucodictyon C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

R. mollissimum Dus.

Brachythecium paradoxum

R. pampae (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum pampae* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905b; Brotherus, 1925; Hosseus, 1935a, b, 1937; Kühnemann, 1938).

R. pendulum (Brid.) Jaeg.

Helicodontium pendulum

R. pinnicaule (C. Muell.) Kindb. (Kindberg, 1888).

Syn. *Hypnum pinnicaule* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905b; Brotherus, 1925; Hosseus, 1935b; Kühnemann, 1938).

R. plagioteciella C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

R. planirameum Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

R. sellowii (Hornschr.) Jaeg. (Jaeger, 1878).

Syn. *Hypnum sellowii* Hornsch. (Hornschuch, 1840).

Hypnum sellowianum C. Muell. (Mueller, 1847a [1846]) *nom. illeg.*

Rhyncho-hypnum sellowii (Hornschr.) Hamp. (Hampe, 1870) *comb. inval.*

Distr. Southern South America (Hornschuch, 1840; Mueller, 1851a; Mitten, 1869; Gibert, 1873; Jaeger, 1878; Paris, 1905b; Felippone, 1912; Brotherus, 1925; Herter, 1933a; Hosseus, 1938a, c, d; Kühnemann, 1938).

R. semi-tortulum Kindb. (Kindberg, 1891).

Syn. *Hypnum semi-tortulum* C. Muell. (Mueller, 1882) *hom. illeg.*

Distr. Southern South America (Mueller, 1882; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

R. serrulatum (Hedw.) Jaeg. (Jaeger, 1878).

Syn. *Hypnum serrulatum* Hedw. (Hedwig, 1801).

Distr. Juan Fernandez (Montagne, 1835, 1839c; Johow, 1896).

Southern South America (Hornschuch, 1820b; Bridel, 1827b; Montagne, 1850; Kühnemann, 1938).

R. striatellum Schimp. ex Par.

Rigodium toxarion

R. subsquarrosum Herz.

Oxyrrhynchium corralense

R. taphrophilum C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).

R. tenuifolium (Hedw.) Reichdt. (Reichardt, 1870).

Syn. *Hypnum tenuifolium* Hedw. (Hedwig, 1801).

Isothecium tenuifolium (Hedw.) Brid. (Bridel, 1827b).

Amblystegiella tenuifolia (Hedw.) Fleisch. (Fleischer, 1920).

Helicodontium tenuifolium (Hedw.) C. Muell. (Mueller in Fleischer, 1920) *nom. inval.*

Distr. Juan Fernandez (Mueller, 1885; Johow, 1896; Brotherus, 1924b; Herter, 1928, 1933b; Robinson, 1975).

Southern South America (Mitten, 1869; Gibert, 1873; Jaeger, 1878; Paris, 1905b; Felippone, 1912; Brotherus, 1925; Herter, 1933a, b; Kühnemann, 1938; Herzog, 1954).

var. *ledifolium* Jaeg. (Jaeger, 1878) *nom. nud.*

Distr. Southern South America (Jaeger, 1878).

var. *minus* Jaeg. (Jaeger, 1878) *nom. nud.*

Distr. Southern South America (Jaeger, 1878).

R. trieblingii C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925).

Rigodium

R. acuminatum Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914).

R. arborescens (C. Muell.) Broth. (Brotherus, 1909).

Syn. *Hypnum arborescens* C. Muell. (Mueller, 1858).

Rigodium lechleri Schimp. (Schimper in Lorentz, 1866) *nom. illeg.*

Heterocladium lechleri Schimp. (Schimper in Lorentz, 1866) *nom. nud. pro parte.*

Heterocladium prolixum Schimp. (Schimper in Lorentz, 1866) *nom. nud. pro parte.*

Distr. Juan Fernandez (Espinosa B., 1941).

Southern South America (Lechler, 1857; Mueller, 1858; Lorentz, 1866; Jaeger, 1878; Neger, 1899; Paris, 1905b; Brotherus, 1909, 1925; Campo, 1915, 1921; Thériot, 1917b, 1918; Costes, 1921; Herzog, 1923, 1954, 1960; Roivainen, 1934; Seki, 1974).

R. argentinicum (C. Muell.) Kindb. (Kindberg, 1891).

Syn. *Hypnum argentinicum* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938; Seki, 1974).

R. brachypodium (C. Muell.) Par. (Paris, 1898).

Syn. *Hypnum brachypodium* C. Muell. (Mueller, 1851a).

Hypnum brachypelma C. Muell. (Mueller, 1882) *nom. inval.*

Distr. Southern South America (Mueller, 1851a, 1882; Paris, 1898, 1905b; Brotherus, 1925; Bartram, 1952).

R. carnosulum Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1905b; Seki, 1974).

R. elegantulum Card.

Rigodium tamarix

var. fernandezianum Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Southern South America (Skottsberg, 1914).

R. hylocomioides Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Juan Fernandez (Brotherus, 1924b; Robinson, 1975).

Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Thériot, 1926; Kühnemann, 1938; Herzog, 1939, 1940, 1960).

var. gracilius Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923).

R. implexum Kunz. ex Schwaegr. (Kunze ex Schwaegrichen, 1844 [1845]).

Syn. *Hypnum implexum* (Schwaegr.) C. Muell. (Mueller, 1844b [1845]) *hom. illeg.*

Heterocladium implexum (Schwaegr.) Lor. (Lorentz, 1866).

Hypnum neei Mohr (Mohr in Mueller, 1851a) *nom. nud.*

Distr. Southern South America (Mueller, 1844b, 1851a; Schwaegrichen, 1844; Sullivant, 1859; Lorentz, 1866; Paris, 1905b; Campo, 1915 (as *R. inflexum* Kurz); Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1940, 1954, 1960; Brotherus, 1925; Roivainen, 1934; Kühnemann, 1938; Bartram, 1952; Bizot, 1967; Seki, 1974).

R. lechleri Schimp.

Rigodium arborescens

R. looseri Thér.

Rigodium robustum

R. nano-fasciculare C. Muell. ex Kindb.

Rigodium toxarion

R. nano-fasciculatum C. Muell. ex Thér.

Rigodium toxarion

R. pendulum Herz. et Thér. (Herzog and Thériot in Herzog, 1939).

Distr. Southern South America (Herzog, 1939, 1954; Seki, 1974).

R. pseudo-thuidium Dus. (Dusén, 1905b).

Syn. *Rigodium ptychomnioides* Broth. (Brotherus, 1909) *nom. nud.*

Distr. Southern South America (Dusén, 1903a, 1905b; Paris, 1905b; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938; Herzog, 1954, 1960; Seki, 1974).

R. ptychomnioides Broth.

Rigodium pseudo-thuidium

R. robustum Broth. (Brotherus, 1924b).

Syn. *Rigodium looseri* Thér. (Thériot, 1927b) *fide* Robinson, 1975.

Distr. Juan Fernandez (Brotherus, 1924b (and as *R. arborescens* *fide* Robinson, 1975); Thériot, 1927b; Bartram, 1959 (and as *R. arborescens* *fide* Robinson, 1975); Robinson, 1975).

R. tamarix C. Muell. (Mueller, 1897a).

Syn. *Rigodium elegantulum* Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Juan Fernandez (Skottsberg, 1914; Thériot, 1921b).

Southern South America (Mueller, 1897a; Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Roivainen, 1934; Thériot, 1934b; Kühnemann, 1938; Seki, 1974).

R. toxarion (Schwaegr.) Jaeg. (Jaeger, 1878).

Syn. *Hypnum toxarion* Schwaegr. (Schwaegrichen, 1816).

Hypnum taxarion Schwaegr. ex Mont. (Schwaegrichen ex Montagne, 1835) *err.*

Eurhynchium striatellum Schimp. (Schimper in Mitten, 1869) *nom. nud.*

Rhynchostegium striatellum Schimp. ex Par. (Schimper ex Paris, 1898) *nom. nud.*

Heterocladium lechleri Schimp. (Schimper in Lorentz, 1866) *nom. nud. pro parte.*

Heterocladium prolixum Schimp. (Schimper in Lorentz, 1866) *nom. nud. pro parte.*

Rigodium nano-fasciculare C. Muell. ex Kindb. (Mueller ex Kindberg, 1891) *err.*

Rigodium nano-fasciculatum C. Muell. ex Thér. (Mueller ex Thériot, 1929d [1930]) *fide* Robinson, 1975.

Distr. Juan Fernandez (Montagne, 1835; Mitten, 1885; Johow, 1896; Brotherus, 1924b, (and as *R. arborescens* *fide* Robinson, 1975); Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Montagne, 1850; Mitten, 1869; Jaeger, 1878; Bescherelle, 1889; Neger, 1899; Cardot, 1908a; Campo, 1915, 1921; Thériot, 1918, 1929d, 1934b; Brotherus, 1924c; Roivainen, 1934; Kühnemann, 1938; Herzog, 1939, 1954, 1960; Espinosa B., 1941; Skottsberg, 1950; Seki, 1974).

R. varium (Hedw.) Kindb.

Amblystegium varium

Rudia

R. clarazii (Dub.) Jaeg.

Helicodontium clarazii

Sanionia

S. orthotheciooides (Lindb.) Loesk.

Drepanocladus uncinatus var. *subjulaceus*

S. symmetrica (Ren. et Card.) Wheld.

Drepanocladus uncinatus var. *symmetricus*

S. uncinata (Hedw.) Loesk.

Drepanocladus uncinatus

Sarconeurum

S. antarcticum Bryhn

Sarconeurum glaciale

S. glaciale (C. Muell.) Card. et Bryhn (Cardot and Bryhn *in* Cardot, 1907a).

Syn. *Didymodon glacialis* Hook. f. et Wils. (Hooker and Wilson *in* Wilson and Hooker, 1847) *hom. illeg.*

Leptotrichum glaciale C. Muell. (Mueller, 1851a).

Ditrichum glaciale (C. Muell.) O. Kuntze (Kuntze, 1891).

Sarconeurum antarcticum Bryhn (Bryhn, 1902).

Tortula saxicola Dus. (Dusén, 1905b) *hom. illeg. fide* Greene, 1975.

Tortula lithophila Dus. (Dusén, 1906) *fide* Greene, 1975.

Tortula pygmaea Dus. (Dusén, 1907) *fide* Greene, 1975.

Distr. Southern South America (Dusén, 1905b, 1906, 1907; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Bizot, 1967; Seki, 1974; Greene, 1975).

Antarctic, peninsula region (Wilson and Hooker, 1847; Mueller, 1851a; Jaeger, 1873; Paris, 1904b; Cardot, 1908a; Steere, 1961a; Greene, 1967; Greene and others, 1970).

Antarctic, continental region (Bryhn, 1902; Cardot, 1907a, 1908a, 1910; Dixon and Watts, 1918; Hilpert, 1933; Bartram, 1938, 1957; Steere, 1961a; Savich-Ljubitskaya and Smirnova, 1962; Greene, 1967; Greene and others, 1970; Robinson, 1972; Savich-Ljubitskaya, 1978).

*S. tortelloides** S. W. Greene (Greene *in* Greene and others, 1970).

Distr. Antarctic, peninsula region (Greene and others, 1970).

Sauloma

S. tenella (Hook. f. et Wils.) Mitt. (Mitten, 1860).

Syn. *Hookeria tenella* Hook. f. et Wils. (Hooker and Wilson *in* Wilson, 1854).

Distr. Southern South America (Bartram, 1946; Welch, 1969; Matteri, 1972, 1975a).

Schimperobryum

S. splendidissimum (Mont.) Marg. (Margadant, 1959).

Syn. *Hookeria splendidissima* Mont. (Montagne, 1835).

Cyathophorum splendidissimum (Mont.) Hamp. et Lor. (Hampe and Lorentz *in* Lorentz, 1866).

Lepidopilum splendidissimum (Mont.) Mitt. (Mitten, 1869).

Lamprophyllum splendidissimum (Mont.) Broth. (Brotherus, 1907).

Hookeria speciosissima Mont. ex Schwaegr. (Montagne ex Schwaegrichen, 1842) *err.*

*Transferred to *Tortella* by Robinson (1972).

Distr. Juan Fernandez (Montagne, 1835, 1856; Schwaegrichen, 1842; Mueller, 1851a; Mitten, 1869, 1885; Jaeger, 1876; Johow, 1896; Skottsberg, 1914; Brotherus, 1924b, 1925; Thériot, 1927b; Espinosa B., 1941; Bartram, 1959; Matteri, 1972; Robinson, 1975).
 Southern South America (Montagne, 1850; Mueller, 1851a; Lorentz, 1866; Mitten, 1869; Jaeger, 1876; Neger, 1899; Paris, 1904a; Brotherus, 1907, 1925; Cardot, 1908a; Campo, 1915; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1954; Thériot, 1924; Reimers, 1926; Kühnemann, 1938; Welch, 1969).

var. *perdentatum* Matteri (Matteri, 1972).

Distr. Juan Fernandez (Matteri, 1972).
 Southern South America (Matteri, 1972; Seki, 1974).

Schistidium

S. alpicola (Hedw.) Limpr. (Limpicht, 1889).

Syn. *Grimmia alpicola* Sw. ex Hedw. (Hedwig, 1801).
Weisia alpicola (Hedw.) Poir. (Poiret, 1808) *nom. inval.*
Grimmia apocarpa Hedw. ssp. *alpicola* (Hedw.) C. Hartm. (Hartman in Lindberg, 1879).
Schistidium apocarpum (Hedw.) B.S.G. ssp. *alpicola* (Hedw.) Meyl. (Meylan, 1905).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938; Herzog, 1957).

var. *rivulare* (Brid.) Limpr.

Schistidium apocarpum

S. amblyophyllum (C. Muell.) Herz.

Grimmia amblyophylla

S. antarctici (Card.) Savicz. et Smirn. (Savich-Ljubitskaya and Smirnova, 1965a).

Syn. *Grimmia antarctici* Card. (Cardot, 1906a) *fide* Savich-Ljubitskaya and Smirnova, 1965a.
Grimmia antarctici Card. var. *percompacta* Bartr. (Bartram, 1938) *fide* Savich-Ljubitskaya and Smirnova, 1965a.
Grimmia antarctici Card. var. *pilifera* Bartr. (Bartram, 1938) *fide* Savich-Ljubitskaya and Smirnova, 1965a.

Distr. Antarctic, peninsula region (Wilson and Hooker, 1847 (as *Tortula gracilis* see Cardot, 1908a, p. 241); Wright, 1905 (as *G. amblyophylla* *fide* Cardot, 1908a); Cardot, 1906a, 1907c, 1908a, 1911b, c, d, 1912b, 1913a; Dixon, 1920; Brotherus, 1924a; Steere, 1961a; Greene, 1968a (as *Tortula gracilis* see Cardot, 1908a, p. 241); Robinson, 1972).
 Antarctic, continental region (Bartram, 1938, 1957; Steere, 1961a; Savich-Ljubitskaya and Smirnova, 1965a; Robinson, 1972; Savich-Ljubitskaya, 1978).

S. apocarpum (Hedw.) B.S.G. (Bruch and others, 1845).

Syn. *Grimmia apocarpa* Hedw. (Hedwig, 1801).
Bryum apocarpum (Hedw.) L. ex With. (Withering, 1801).
Weisia apocarpa (Hedw.) Poir. (Poiret, 1808).
Grimmia rivularis Brid. (Bridel, 1801a).
Schistidium rivulare (Brid.) Podp. (Podpéra, 1911).
Grimmia apocarpa Hedw. var. *rivularis* (Brid.) Roehl. (Röhling, 1812).
Schistidium apocarpum (Hedw.) B.S.G. var. *rivulare* (Brid.) B.S.G. (Bruch and others, 1845).
Grimmia alpicola Sw. ex Hedw. var. *rivularis* (Brid.) Wahlenb. (Wahlenberg, 1812).
Schistidium alpicola (Hedw.) Limpr. var. *rivulare* (Brid.) Limpr. (Limpicht, 1889).

Distr. Southern South America (Wilson and Hooker, 1847; Mueller, 1849; 1885; Montagne, 1850; Mitten, 1869; Paris, 1904b; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen, 1934; Kühnemann, 1938).
 Falkland Islands (Wilson and Hooker, 1847; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a).
 Antarctic, peninsula region (Wright, 1905; Cardot, 1908a, 1911c, 1912b; Brotherus, 1924a; Bartram, 1957; Steere, 1961a; Robinson, 1972).

ssp. *alpicola* (Hedw.) Meyl.

Schistidium alpicola

ssp. *confertum* (Funck) Loesk. (Loeske, 1913).

Syn. *Grimmia conferta* Funck (Funck, 1820).

- Schistidium confertum* (Funck) B.S.G. (Bruch and others, 1845).
Grimmia apocarpa Hedw. var. *conferta* (Funck) Spreng. (Sprengel, 1827).
Grimmia apocarpa Hedw. ssp. *conferta* (Funck) Boul. (Boulay, 1884).
- Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).
Falkland Islands (Cardot and Brotherus, 1923).
- var. *rivulare* (Brid.) B.S.G. *Schistidium apocarpum*
S. ciliatum (Hedw.) Brid. var. *imberbe* (Sm.) Hueb. *Hedwigidium integrifolium*
S. confertum (Funck) B.S.G. *Schistidium apocarpum* ssp. *confertum*
S. imberbe (Sm.) Nees et Hornsch. *Hedwigidium integrifolium*
- S. maritimum*** (Turn.) B.S.G. (Bruch and others 1845).
Syn. *Grimmia maritima* Turn. (Turner, 1804a).
Grimmia alpicola Sw. ex Hedw. var. *maritima* (Turn.) Wahlenb. (Wahlenberg, 1812).
Distr. Southern South America (Wilson and Hooker, 1847).
- S. plagiopodium* (Hedw.) Loesk. *Grimmia plagiopodia*
S. praemorsum (C. Muell.) Herz. *Grimmia praemorsa*
S. rivulare (Brid.) Podp. *Schistidium apocarpum*
S. subsessile (Brid.) Brid. *Pterygoneurum subsessile*
- Schistophyllum hom. illeg.***
S. acacioides (Schrad.) Lindb. *Fissidens osmundoides*
S. adianthoides (Hedw.) Lindb. *Fissidens adianthoides*
S. aspleniooides (Hedw.) Brid. *Fissidens aspleniooides*
S. bryoides (Hedw.) Lindb. *Fissidens bryoides*
S. julianum (Cand.) Lindb. *Fissidens fontanus*
S. osmundoides (Hedw.) Lindb. *Fissidens osmundoides*
var. *microcarpum* (B.S.G.) Kindb. *Fissidens osmundoides*
S. semicompletum (Hedw.) Lindb. *Fissidens semicompletus*
S. taxifolium (Hedw.) Lindb. *Fissidens taxifolius*
- Schlotheimia***
S. aequifolia Dus. (Dusén in Naveau, 1928) nom. nud.
Distr. Southern South America (Naveau, 1928).
- S. argentinica*** Lor. et C. Muell. (Lorentz and Mueller in Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938).
- S. asperrima*** Broth. (Brotherus, 1918).
Distr. Southern South America (Brotherus, 1918, 1925; Kühnemann, 1938).
S. gracillima Besch. *Macromitrium gracillimum*
- S. jamesonii*** (Arnott) Brid. (Bridel, 1826).
Syn. *Orthotrichum jamesonii* Arnott (Arnott, 1824b).
Macromitrium jamesonii (Arnott) C. Muell. (Mueller, 1845).
Distr. Southern South America (Hornschuch, 1840; Herter, 1933a).
- S. linearifolia*** (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1960b).
Syn. *Macromitrium linearifolium* C. Muell. (Mueller, 1845).
Schlotheimia longifolia Hornsch. (Hornschuch, 1840) hom. illeg.
Distr. Southern South America (Hornschuch, 1840; Mueller, 1849; Gibert, 1873; Paris, 1905b; Herter, 1933a).

- S. longifolia* Hornsch.
S. nitida Schwaegr.
S. ottonis Schwaegr.
- S. striata*** Herz. (Herzog, 1952a).
Distr. Southern South America (Herzog, 1952a).
- S. torquata*** (Hedw.) Brid. (Bridel, 1812).
Syn. *Hypnum torquatum* Hedw. (Hedwig, 1801).
Orthotrichum torquatum (Hedw.) Arnott (Arnott, 1827).
Macromitrium torquatum (Hedw.) C. Muell. (Mueller, 1845).
Schlotheimia nitida Schwaegr. (Schwaegrichen, 1826).
Schlotheimia ottonis Schwaegr. (Schwaegrichen, 1826).
Macromitrium ottonis (Schwaegr.) C. Muell. (Mueller, 1845).
Distr. Southern South America (Schwaegrichen, 1826; Bridel, 1827a; Hornschuch, 1840; Mueller, 1849; Gibert, 1873; Jaeger, 1874; Herter, 1933a).
- S. trichomitria*** Schwaegr. (Schwaegrichen, 1826).
Syn. *Macromitrium trichomitrium* (Schwaegr.) C. Muell. (Mueller, 1845).
Distr. Southern South America (Schwaegrichen, 1826, 1827a; Bridel, 1827a; Hornschuch, 1840; Mueller, 1849; Gibert, 1873; Jaeger, 1874; Paris, 1905b; Felippone, 1912; Herter, 1933a).
- Sciaromium***
- S. confluens*** (C. Muell.) Par. (Paris, 1898).
Syn. *Hypnum confluens* C. Muell. (Mueller, 1885).
Brachythecium confluens (C. Muell.) Kindb. (Kindberg, 1888).
Limbella confluens (C. Muell.) C. Muell. (Mueller, 1889) *comb. inval.*
Distr. Southern South America (Mueller, 1885, 1889; Paris, 1898, 1905b; Dusén, 1903a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938; Seki, 1974).
- S. conspissatulum*** Par. (Paris, 1898).
Syn. *Limbella conspissatula* C. Muell. (Mueller, 1897a) *comb. inval.*
Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925; Kühnemann, 1938).
- S. conspissatum*** (Hook. f. et Wils.) Mitt. (Mitten, 1869).
Syn. *Hypnum conspissatum* Hook. f. et Wils. (Hooker, J.D. and Wilson, 1844).
Limbella conspissata (Hook. f. et Wils.) C. Muell. (Mueller, 1889) *comb. inval.*
Distr. Southern South America (Mueller, 1851a, 1889; Sullivant, 1859; Lorentz, 1866; Jaeger and Sauerbeck, 1879a; Paris, 1905b; Cardot, 1908a; Thériot, 1915; Brotherus, 1925; Kühnemann, 1938; Herzog, 1939, 1954, 1960; Bartram, 1952; Seki, 1974). Falkland Islands (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mitten, 1869; Jaeger and Sauerbeck, 1879a; Paris, 1905b; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938). South Georgia (Cardot, 1906a, 1908a; Brotherus, 1925; Steere, 1961b).
*var. *maritimum** (Card.) Herz.
- Sciaromium obscurifolium***
- S. crassinervatum*** Mitt. (Mitten, 1869).
Distr. Southern South America (Herzog, 1939).
- S. depastum*** Dus. (Dusén, 1903a).
Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).
- S. drepanophyllopsis*** Par. (Paris, 1898).
Syn. *Limbella drepanophyllopsis* C. Muell. (Mueller, 1897a) *comb. inval.*

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925; Kühnemann, 1938).

S. flavidulum Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b; Cardot, 1908a; Brotherus, 1925).

S. gracile Dus. (Dusén, 1903a).

Distr. Southern South America (Dusén, 1903a; Cardot, 1908a; Brotherus, 1925; Kühnemann, 1938).

S. krauseanum Par. (Paris, 1898).

Syn. *Limbella krauseana* C. Muell. (Mueller, 1897a) *comb. inval.*

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

S. maritimum Card.

Sciaromium obscurifolium

S. nigritum Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b; Cardot, 1908a; Brotherus, 1925).

S. obscurifolium Mitt. (Mitten, 1869).

Syn. *Sciaromium maritimum* Card. (Cardot, 1905a).

Sciaromium conspissatum (Hook. f. et Wils.) Mitt. var. *maritimum* (Card.) Herz. (Herzog, 1954).

Distr. Southern South America (Mitten, 1869; Jaeger and Sauerbeck, 1879a; Paris, 1905b; Cardot and Brotherus, 1923; Thériot, 1924, 1934b; Brotherus, 1925; Kühnemann, 1938; Herzog, 1954; Seki, 1974).

Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

S. pachyloma (Mont.) Par. (Paris, 1898).

Syn. *Gymnostomum pachyloma* Mont. (Montagne, 1838).

Grimmia pachyloma (Mont.) Mitt. (Mitten, 1869).

Hypnum pachyloma (Mont.) C. Muell. (Mueller, 1889).

Limbella pachyloma (Mont.) C. Muell. (Mueller, 1897a) *comb. inval.*

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Montagne, 1838, 1850; Mueller, 1851a, 1889, 1897a; Mitten, 1869; Jaeger, 1874; Paris, 1898, 1905b; Cardot and Brotherus, 1923; Espinosa B., 1924; Brotherus, 1925; Reimers, 1926; Kühnemann, 1938; Bartram, 1952).

var. *brevifolium* Thér. (Thériot, 1924).

Distr. Southern South America (Costes, 1921 (as fo. *brevifolium*); Thériot, 1924).

var. *gracilescens* Herz. (Herzog, 1954).

Distr. Southern South America (Herzog, 1954).

S. pachylomatum Par. (Paris, 1898).

Syn. *Limbella pachylomata* C. Muell. (Mueller, 1897a) *comb. inval.*

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Kurtz, 1904; Brotherus, 1925; Kühnemann, 1938).

S. platylomatum Par. (Paris, 1898).

Syn. *Limbella platylomata* C. Muell. (Mueller, 1897a) *comb. inval.*

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925; Hosseus, 1936, 1937).

Sclerodontium nom. rejic.

S. pellucidum Schwaegr.

Dimerodontium pellucidum

Scleropodium

S. pseudo-purum (C. Muell.) Broth.

Pseudoscleropodium pseudopurum

Scouleria

S. patagonica (Mitt.) Jaeg. (Jaeger, 1876).

Syn. *Grimmia patagonica* Mitt. (Mitten, 1869).

Distr. Southern South America (Mitten, 1869; Jaeger, 1876; Paris, 1905b; Dusén, 1907; Cardot, 1908a; Brotherus, 1924a; Thériot, 1935a (as *fo. squarrosa*); Kühnemann, 1938; Bartram, 1952; Herzog, 1954, 1960; Seki, 1974; Matteri, 1979).

Seligeria

S. kunzeana (C. Muell.) C. Muell.

Microdus kunzeanus

S. kunzei Kindb.

Microdus kunzeanus

Sematophyllum

S. aberrans (Broth.) Bartr. (Bartram, 1959) *fide* Robinson, 1975.

Syn. *Rhaphidostegium aberrans* Broth. (Brotherus, 1924b).

Rhaphidorrhynchium aberrans (Broth.) Broth. (Brotherus, 1925).

Rhaphidostegium caespitosoides Broth. (Brotherus, 1924b) *fide* Robinson, 1975.

Sematophyllum caespitosoides (Broth.) Broth. (Brotherus, 1925).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

S. amoenum (Hedw.) Mitt.

Rhaphidorrhynchium amoenum

S. ampullulatum (C. Muell.) Broth. (Brotherus, 1925).

Syn. *Aptychus ampullulatus* C. Muell. (Mueller, 1897a).

Rhaphidostegium ampullulatum (C. Muell.) Broth. (Brotherus, 1908).

Aptychus ampullulatus C. Muell. ex Par. (Mueller ex Paris, 1898) *err.*

Rhaphidostegium ampullulatum Par. (Paris, 1898) *hom. illeg.*

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1908, 1925; Kühnemann, 1938).

S. aquicola (C. Muell.) Kindb.

***Bryum* sp.**

S. aureo-nitidum Bartr.

Rhaphidorrhynchium scorpiurus

S. aureo-viride (C. Muell.) Broth. (Brotherus, 1925).

Syn. *Aptychus aureo-viridis* C. Muell. (Mueller, 1897a).

Rhaphidostegium aureo-viride (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925; Kühnemann, 1938).

S. brachycladulum (Broth.) Broth. (Brotherus, 1925).

Syn. *Rhaphidostegium brachycladulum* Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b (and as *S. caespitosum* *fide* Robinson, 1975), 1925; Bartram, 1959; Robinson, 1975).

S. bracteatum (C. Muell.) Broth. (Brotherus, 1925).

Syn. *Hypnum bracteatum* C. Muell. (Mueller, 1879a).

Rhaphidostegium bracteatum (C. Muell.) Kindb. (Kindberg, 1888).

Aptychus catilliformis C. Muell. (Mueller, 1897a).

Rhaphidostegium catilliforme (C. Muell.) Par. (Paris, 1898).

Sematophyllum catilliforme (C. Muell.) Broth. (Brotherus, 1925).

Distr. Southern South America (Mueller, 1879a, 1897a; Paris, 1898, 1900a, 1905b; Brotherus, 1925; Felippone, 1929; Herter, 1933a, 1943; Kühnemann, 1938; Hosseus, 1939).

S. caespitosoides (Broth.) Broth.

Sematophyllum aberrans

S. caespitosum (Hedw.) Mitt. (Mitten, 1869).

- Syn.* *Leskea caespitosa* Hedw. (Hedwig, 1801).
Rhaphidostegium caespitosum (Hedw.) Besch. (Bescherelle, 1876).
Hypnum loxense Hook. (Hooker *in* Kunth, 1822).
Rhaphidostegium loxense (Hook.) Jaeg. (Jaeger, 1874).

Distr. Juan Fernandez (Montagne, 1835).
 Southern South America (Krieger, 1904; Hosseus, 1938c, d).

S. callidum (Mont.) Mitt.**Rhaphidorrhynchium callidum****S. catilliforme** (C. Muell.) Broth.**Sematophyllum bracteatum****S. cirrhifolium** (Arnott) Mitt. (Mitten, 1885).

- Syn.* *Hookeria cirrhifolium* Arnott (Arnott, 1827).
Hypnum cirrifolium (Arnott) Schwaegr. (Schwaegrichen *in* Gaudichaud, 1826 [1828])
hom. illeg.

Distr. Juan Fernandez (Montagne, 1835).
 Falkland Islands (Gaudichaud, 1826).

S. crassiusculum (Brid.) Broth. (Brotherus, 1925).

- Syn.* *Hypnum crassiusculum* Brid. (Bridel, 1812).

Distr. Juan Fernandez (Johow, 1896).
 Southern South America (Montagne, 1850).

S. grandicellulosum (C. Muell.) Broth. (Brotherus, 1925).

- Syn.* *Aptychus grandicellulosus* C. Muell. (Mueller, 1897a).

Rhaphidostegium grandicellulosum (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925;
 Kühnemann, 1938).

S. kunkelii H. Robinson (Robinson, 1974).

Distr. Juan Fernandez (Robinson, 1974, 1975).

S. leptophyllum Mitt.**Rhaphidorrhynchium leptophyllum****S. leucocytus** (C. Muell.) Sains. (Sainsbury, 1955).

- Syn.* *Hypnum leucocytus* C. Muell. (Mueller, 1851a).

Rhaphidostegium leucocytus (C. Muell.) Jaeg. (Jaeger, 1878).

Distr. Southern South America (Cardot, 1908a).

S. macrocytus (C. Muell.) Broth. (Brotherus, 1925).

- Syn.* *Aptychus macrocytus* C. Muell. (Mueller, 1897a).

Rhaphidostegium macrocytus (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925;
 Kühnemann, 1938).

S. masafuerae (Broth.) Bartr. (Bartram, 1959) *fide* Robinson, 1975.

- Syn.* *Rhaphidostegium masafuerae* Broth. (Brotherus, 1924b) *fide* Robinson, 1975.
Rhaphidorrhynchium masafuerae (Broth.) Broth. (Brotherus, 1925).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Bartram, 1959; Robinson, 1975).

S. micrangium (C. Muell.) Broth. (Brotherus, 1925).

- Syn.* *Aptychus micrangius* C. Muell. (Mueller, 1897a).

Rhaphidostegium micrangium (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Kühnemann,
 1938).

S. microcarpoides (C. Muell.) Broth. (Brotherus, 1925).

- Syn.* *Hypnum microcarpoides* C. Muell. (Mueller, 1879a).

Rhaphidostegium microcarpoides (C. Muell.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1925; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939).

S. minutum Broth.

Rhaphidorrhynchium minutum

***S. nanocephalum* (C. Muell.) Broth.** (Brotherus, 1925).

Syn. *Aptychus nano-cephalus* C. Muell. (Mueller, 1897a).

Rhaphidostegium nano-cephalum (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925; Kühnemann, 1938).

***var. subglauculum* (C. Muell.) Wijk et Marg.** (van der Wijk and Margadant, 1959).

Syn. *Aptychus nano-cephalus* C. Muell. var. *subglauculus* C. Muell. (Mueller, 1897a).

Rhaphidostegium nano-cephalum (C. Muell.) Par. var. *subglauculum* (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925; Kühnemann, 1938).

S. noduliferum Mitt.

Rhaphidorrhynchium noduliferum

***S. perichaetiale* Thér.** (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933a).

S. scorpiurus (Mont.) Mitt.

Rhaphidorrhynchium scorpiurus

S. secundifolium (C. Muell.) Mitt.

Drepanocladus secundifolius

***S. serifolium* (C. Muell.) Broth.** (Brotherus, 1925).

Syn. *Aptychus serifolius* C. Muell. (Mueller, 1897a).

Rhaphidostegium serifolium (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925; Kühnemann, 1938).

***S. stenopyxidium* (C. Muell.) Broth.** (Brotherus, 1925).

Syn. *Hypnum stenopyxidium* C. Muell. (Mueller, 1879a).

Rhaphidostegium stenopyxidium (C. Muell.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1925; Herter, 1933a; Kühnemann, 1938; Hosseus, 1939).

***S. subsimplex* (Hedw.) Mitt.** (Mitten, 1869).

Syn. *Hypnum subsimplex* Hedw. (Hedwig, 1801).

Isothecium subsimplex (Hedw.) Brid. (Bridel, 1827b).

Plagiothecium subsimplex (Hedw.) Besch. (Bescherelle, 1871–72 [1872]).

Rhaphidorrhynchium subsimplex (Hedw.) Broth. (Brotherus, 1925).

Rhaphidostegium subsimplex (Hedw.) Besch. (Bescherelle, 1876).

Hypnum simplex Brid. (Bridel, 1812) *nom. illeg.*

Distr. Southern South America (Bescherelle, 1889; Paris, 1905b; Cardot, 1908a).

***S. temperatum* (C. Muell.) Broth.** (Brotherus, 1925).

Syn. *Aptychus temperatus* C. Muell. (Mueller, 1897a).

Rhaphidostegium temperatum (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925; Kühnemann, 1938).

***S. tenerifolium* (C. Muell.) Broth.** (Brotherus, 1925).

Syn. *Aptychus tenerifolius* C. Muell. (Mueller, 1897a).

Rhaphidostegium tenerifolium (C. Muell.) Par. (Paris, 1898).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b; Brotherus, 1925; Kühnemann, 1938).

***S. tucumanense* Bartr.** (Bartram, 1965).

Distr. Southern South America (Bartram, 1965).

<i>Sigmatella hom. illeg.</i>	
<i>S. microthamnioides</i> C. Muell.	<i>Taxithelium microthamnioides</i>
<i>Simophyllum nom. illeg.</i>	
<i>S. viridulum</i> Lindb.	<i>Weisia controversa</i>
<i>Skitophyllum nom. illeg.</i>	
<i>S. acacioides</i> (Schrad.) Pyl.	<i>Fissidens osmundoides</i>
<i>S. adianthoides</i> (Hedw.) Pyl.	<i>Fissidens adianthoides</i>
<i>S. asplenoides</i> (Hedw.) Pyl.	<i>Fissidens asplenoides</i>
<i>S. bryoides</i> (Hedw.) Pyl.	<i>Fissidens bryoides</i>
<i>S. dillenii</i> Pyl.	<i>Fissidens fontanus</i>
<i>S. fontanum</i> Pyl.	<i>Fissidens fontanus</i>
<i>S. julianum</i> (Cand.) Brid.	<i>Fissidens fontanus</i>
<i>S. osmundoides</i> (Hedw.) Pyl.	<i>Fissidens osmundoides</i>
<i>S. semi-completum</i> (Hedw.) Pyl.	<i>Fissidens semicompletus</i>
<i>S. taxifolium</i> (Hedw.) Pyl.	<i>Fissidens taxifolius</i>
Skottsbergia	
<i>S. paradoxa</i> Card. (Cardot, 1905b).	
<i>Distr.</i> Southern South America (Rovainen and Bartram, 1937; Kühnemann, 1938). South Georgia (Cardot, 1905a, 1906a, 1907b, 1908a; Brotherus, 1924a; Steere, 1961b).	
<i>Sphaerangium nom. illeg.</i>	
<i>S. lorentzii</i> (C. Muell.) Par.	<i>Astomum lorentzii</i>
<i>S. vesiculosum</i> (C. Muell.) Kindb.	<i>Acaulon vesiculosum</i>
Sphaerocephalus	
<i>S. palustris</i> (Hedw.) Lindb.	<i>Aulacomnium palustre</i>
Sphagnum	
<i>S. acutifolium</i> Schrad. (Schrader, 1794).	
<i>Distr.</i> Southern South America (Paris, 1905b (as forma <i>diblastum</i> (C. Muell.) Par.)). Falkland Islands (Gaudichaud, 1825).	
var. <i>tenellum</i> Schwaegr. (Schwaegrichen in Gaudichaud, 1826 [1828]) <i>nom. nud.</i>	
<i>Distr.</i> Falkland Islands (Gaudichaud, 1826).	
<i>S. amblyphyllum</i> (Russ.) Warnst. (Warnstorff, 1911).	
<i>Distr.</i> Southern South America (Thériot, 1934b).	
<i>S. andinum</i> Hamp. (Hampe, 1866).	
<i>Distr.</i> Southern South America (Lorentz, 1866; Jaeger, 1870).	
<i>S. bicolor</i> Besch. (Bescherelle, 1885a).	
<i>Distr.</i> Southern South America (Mueller, 1885; Bescherelle, 1885a, 1889; Hariot, 1891).	
<i>S. capillaceum</i> (Weiss) Schrank (Schrank, 1789).	
<i>Distr.</i> Southern South America (Andrews, 1947).	
<i>S. chilense</i> Lor. (Lorentz, 1866).	
<i>Distr.</i> Southern South America (Lorentz, 1866).	
<i>S. cuspidatum</i> Ehrh. ex Hoffm. (Ehrhardt ex Hoffmann, 1795 [1796]).	
<i>Distr.</i> Southern South America (Wilson and Hooker, 1847; Sullivant, 1859; Cardot, 1908a; Warnstorff, 1911; Cardot and Brotherus, 1923; Thériot, 1934b; Andrews, 1949b). Falkland Islands (Wilson and Hooker, 1847; Kühnemann, 1938).	

var. falcatulum (Besch.) Par. (Paris, 1897).

Distr. Southern South America (Paris, 1897, 1905b).

var. falcatum Russ. (Russow, 1867).

Distr. Southern South America (Dusén, 1903b; Paris, 1905b; Cardot, 1908a).

var. microporum (Warnst.) Par. (Paris, 1897).

Distr. Falkland Islands (Paris, 1897, 1905b).

var. miquelonense (Ren. et Card.) Ren. et Card. (Renauld and Cardot *in* Cardot, 1887).

Distr. Southern South America (Dusén, 1903b; Cardot, 1908a).

var. serratum (Aust.) Aust. (Austin, 1880).

Distr. Southern South America (Dusén, 1903b; Cardot 1908a).

var. submersum Schimp. (Schimper, 1858 [1857]).

Distr. Southern South America (Dusén, 1903b; Paris, 1905b).

S. cymbifolium (Ehrh.) Hedw. (Hedwig, 1782).

Distr. Southern South America (Sullivant, 1859; Mitten, 1869; Jaeger, 1870; Paris, 1905b; Spegazzini, 1922).

Falkland Islands (Bridel, 1827a; Wilson and Hooker, 1847).

var. condensatum Hook. f. (Hooker *in* Wilson and Hooker, 1847) *hom. illeg.*

Distr. Southern South America (Wilson and Hooker, 1847; Ångström, 1872; Paris, 1905b).

var. magellanicum (Brid.) P. Beauv. (Palisot de Beauvois, 1805a).

Distr. Southern South America (Palisot de Beauvois, 1805a; Bridel, 1826).

var. paradisi Besch. (Bescherelle *in* Warnstorff, 1890a) *nom. nud.*

Distr. Southern South America (Warnstorff, 1890a).

S. diblastum C. Muell. (Mueller, 1887a).

Distr. Southern South America (Mueller, 1887a; Warnstorff, 1911; Brotherus, 1924a; Kühnemann, 1938).

S. dissimile Roiv. (Rovainen *in* Rovainen and Bartram, 1937).

Distr. Southern South America (Rovainen and Bartram, 1937).

S. dusenioides Roiv. (Rovainen *in* Rovainen and Bartram, 1937).

Distr. Southern South America (Rovainen and Bartram, 1937).

S. falcatulum Besch. (Bescherelle, 1885a).

Distr. Southern South America (Bescherelle, 1885a, 1889; Mueller, 1885; Hariot, 1891; Warnstorff, 1894, 1895, 1911; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Rovainen and Bartram, 1937; Kühnemann, 1938; Andrews, 1949b; Seki, 1974).

var. microporum Warnst. (Warnstorff, 1895).

Distr. Falkland Islands (Warnstorff, 1895, 1911; Cardot, 1908a; Kühnemann, 1938).

S. fimbriatum Wils. (Wilson *in* Wilson and Hooker, 1845).

Distr. Southern South America (Wilson and Hooker, 1847; Sullivant, 1859; Mitten, 1869; Jaeger, 1870; Mueller, 1885; Cardot, 1900, 1901, 1905a, 1908a; Paris, 1905b; Warnstorff, 1911; Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1939; Andrews, 1947; Seki, 1974).

Falkland Islands (Wilson and Hooker, 1847; Jaeger, 1870; Mueller, 1885; Cardot, 1905a, 1908a; Paris, 1905b; Andrews, 1947).
South Georgia (Greene, 1964, 1968a).

var. molluscoides Card. (Cardot *in* Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

var. robustum Braithw. ex Warnst. (Braithwaite ex Warnstorff, 1884a) *nom. illeg.*

Distr. Southern South America (Dusén, 1903a, b (and as forma *brachydasyyclada* and forma *laxifolia*), 1905d (as forma *brachydasyyclada*); Paris, 1905b; Cardot, 1908a; Warnstorff, 1911; Cardot and Brotherus, 1923; Kühnemann, 1938; Herzog, 1939).

var. *validus* Card. (Cardot, 1884).

Distr. Southern South America (Warnstorff, 1911; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923).

S. hahnianum C. Muell. (Mueller in Warnstorff, 1890a) *nom. nud.*

Distr. Southern South America (Warnstorff, 1890a).

S. imbricatum Hornsch. ex Russ. (Hornschuch ex Russow, 1867).

Distr. Southern South America (Warnstorff, 1911).

S. lechleri Warnst. (Warnstorff, 1907).

Distr. Southern South America (Warnstorff, 1907, 1911 (but see Andrews, 1947, p. 183)).

S. magellanicum Brid. (Bridel, 1798).

Distr. Southern South America (Bridel, 1801b, 1803, 1806; Rovainen and Bartram, 1937; Herzog, 1938, 1939; Kühnemann, 1938; Seki, 1974).

var. *congestum* (Schimp.) Roell (Röll, 1915).

Distr. Falkland Islands (Kühnemann, 1938).

var. *versicolor* (Warnst.) Wijk et Marg. (van der Wijk and Margadant, 1965).

Distr. Southern South America (Kühnemann, 1938).

S. medium Limpr. (Limpricht, 1881).

Distr. Southern South America (Warnstorff, 1894, 1911 (as forma *brachy-orthocladum* Warnst.); Paris, 1905b; Cardot, 1908a; Brotherus, 1924a; Herter, 1933a, 1943; Martin, 1949).

var. *congestum* (Schimp.) Warnst. (Warnstorff, 1883).

Distr. Southern South America (Röll, 1886 (as forma *bicolor* Besch.); Cardot, 1905a, 1913a; Cardot and Brotherus, 1923).

Falkland Islands (Cardot and Brotherus, 1923).

var. *fusco-rubellum* Warnst. (Warnstorff in Dusén, 1903b) *nom. nud.*

Distr. Southern South America (Dusén, 1903a, b (as forma *brachydasyyclada*); Kühnemann, 1938).

var. *laeve* (Russ.) Warnst. (Warnstorff, 1890b).

Distr. Southern South America (Warnstorff, 1890b (as forma *albescens* Warnst.); Paris, 1905b).

var. *pallido-carneum* Warnst. (Warnstorff in Dusén, 1903b) *nom. nud.*

Distr. Southern South America (Dusén, 1903b (as forma *brachydasyyclada*).

var. *purpurascens* (Russ.) Warnst. (Warnstorff, 1884b).

Distr. Southern South America (Dusén, 1903b, 1905d (and as forma *brachylomatoclada*); Cardot and Brotherus, 1923; Kühnemann, 1938).
Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

var. *roseum* ((Roell) Warnst. (Warnstorff, 1888).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

var. *violascens* Warnst. (Warnstorff in Horrell, 1900).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

var. *virescens* Warnst. (Warnstorff, 1888).

Distr. Southern South America (Cardot and Brotherus, 1923).

S. nano-porosum Warnst. (Warnstorff, 1898).

Distr. Falkland Islands (Warnstorff, 1898, 1911; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

- var. *skottsbergii*** Card. (Cardot in Cardot and Brotherus, 1923).
Distr. Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).
- S. patagoniense*** Warnst. (Warnstorf, 1911).
Distr. Southern South America (Warnstorf, 1911; Brotherus, 1924a; Herzog, 1939 (as *S. patagonicum* Warnst.)).
- var. *submersum*** Warnst. (Warnstorf, 1911).
Distr. Southern South America (Warnstorf, 1911).
- S. personatum*** Roiv. (Rovainen in Rovainen and Bartram, 1937).
Distr. Southern South America (Rovainen and Bartram, 1937).
- S. plumulosum*** Roell (Röll, 1886).
Distr. Southern South America (Warnstorf, 1911; Brotherus, 1924a; Kühnemann, 1938; Andrews, 1947).
- S. pulchricoma*** C. Muell. (Mueller, 1848a).
Distr. Southern South America (Mitten, 1869; Jaeger, 1870; Herzog, 1938).
- S. recurvum*** P. Beauv. (Palisot de Beauvois, 1805a).
Distr. Southern South America (Paris, 1905b; Cardot, 1908a; Andrews, 1949b).
- S. rigescens*** Warnst. (Warnstorf, 1898).
Distr. Southern South America (Warnstorf, 1898, 1911; Dusén, 1903a, b; Paris, 1905b; Cardot, 1908a; Brotherus, 1924a; Andrews, 1941).
- S. schwabeanum*** Paul (Paul in Herzog, 1939).
Distr. Southern South America (Herzog, 1939).
- S. squarrosum*** Crom. (Crome in Hoppe, 1801).
Distr. Southern South America (Hariot, 1891).
- S. subfalcatum*** Roiv. (Rovainen in Rovainen and Bartram, 1937).
Distr. Southern South America (Rovainen and Bartram, 1937; Kühnemann, 1938).
- S. subnitens*** Russ. et Warnst. var. *purpurascens* Schlieph. ex C. Jens. (Schliephacke ex Jensen, 1890).
Distr. Southern South America (Dusén, 1903b; Paris, 1905b; Cardot, 1908a).
- S. subrigidum*** Hamp. et Lor. (Hampe and Lorentz in Lorentz, 1868).
Distr. Southern South America (Paris, 1905b; Warnstorf, 1911; Brotherus, 1924a (but see Andrews, 1947)).
- S. subserratum*** Roiv. (Rovainen in Rovainen and Bartram, 1937).
Distr. Southern South America (Rovainen and Bartram, 1937).
- S. tenerum*** Sull. et Lesq. (Sullivan and Lesquereux, 1856).
Distr. Southern South America (Andrews, 1947).
- S. teres*** (Schimp.) Aongstr. var. *laxum* Braithw. ex Warnst. (Braithwaite ex Warnstorf, 1881a).
Distr. Southern South America (Paris, 1905b).
- S. torreyanum*** Sull. (Sullivan, 1849).
Distr. Southern South America (Warnstorf, 1911; Brotherus, 1924a).
- S. trinitense*** C. Muell. (Mueller, 1848a).
Distr. Southern South America (Cardot and Brotherus, 1923).
Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).
- S. undulatum*** Warnst. (Warnstorf, 1894).
Distr. Southern South America (Warnstorf, 1894, 1895, 1911; Dusén, 1903b; Paris, 1905b; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

Splachnobryum***S. lorentzii*** C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1905b; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

S. valdiviae C. Muell. (Mueller, 1897c).

Distr. Southern South America (Mueller, 1897c; Paris, 1905b; Brotherus, 1924a).

Splachnum***S. magellanicum*** Brid.***Tayloria magellanica******S. mnioides*** Hedw.***Tetrapodon mnioides******S. octoblepharum*** Hook.***Tayloria octoblepharum******Sporledera******S. hampeana*** (C. Muell.) Mitt.***Bruchia hampeana******S. schwaegrichenii*** Hamp.***Bruchia hampeana******Squamidium******S. lorentzii*** (C. Muell.) Broth. (Brotherus, 1906b).Syn. ***Meteorium lorentzii*** C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905a; Brotherus, 1906b, 1925; Kühnemann, 1938).

S. turgidulum (C. Muell.) Broth. (Brotherus, 1906b).Syn. ***Neckera turgidula*** C. Muell. (Mueller, 1879a).***Pilotrichella turgidula*** (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1879a; Paris, 1897, 1905b; Brotherus, 1906b, 1925; Kühnemann, 1938).

var. *triste* (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1961).Syn. ***Neckera turgidula*** C. Muell. var. ***tristis*** C. Muell. (Mueller, 1879a).***Pilotrichella turgidula*** (C. Muell.) Par. var. ***tristis*** (C. Muell.) Par. (Paris, 1897).

Distr. Southern South America (Mueller, 1879a; Paris, 1897, 1905b).

Stableria***S. tenella*** (Mitt.) Broth.***Orthodontium gracile******Stegonia******S. guessfeldtii*** (Schliep.) Hag.***Pottia heimii* var. ***guessfeldtii********S. latifolia*** (Schwaegr.) Vent. ex Broth. (Brotherus, 1923).Syn. ***Weisia latifolia*** Schwaegr. (Schwaegrichen, 1804).

Distr. Antarctic, peninsula region (Smith, 1972, p. 31).

S. lorentzii (C. Muell.) Hag. (Hagen in Hagen and Printz, 1929).Syn. ***Pottia lorentzii*** C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Warnstorf, 1916; Brotherus, 1924a; Kühnemann, 1938).

Stenocarpidium***S. leucodon*** C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Hosseus, 1935b, 1938c, d; Kühnemann, 1938).

Stenomitrium***S. aureum*** Par. (Paris, 1905b) nom. nud.

- Syn.* *Pentastichella aurea* Dus. (Dusén, 1903a) *nom. nud.*
Distr. Southern South America (Dusén, 1903a; Paris, 1905b).
- S. pentastichum* (Mont.) Broth. *Zygodon pentastichus*
S. robustulum Par. *Zygodon bartramoides*
- Stereodon** *nom. illeg.*
- S. acicularis* (Brid.) Mitt. *Ptychomnion aciculare*
S. auriculatus (Mont.) Mitt. *Acrocladium auriculatum*
S. campoanus Thér. *Hypnum campoanum*
S. cochlearifolius (Schwaegr.) Mitt. *Weymouthia cochlearifolia*
S. cypressiformis (Hedw.) Brid. ex Mitt. *Hypnum cypressiforme*
S. densifolius (Brid.) Mitt. *Ptychomnion densifolium*
S. denticulatus (Hedw.) Mitt. *Plagiothecium denticulatum*
S. gracilis (Hook. f. et Wils.) Mitt. *Camptochaete gracilis*
S. gracilis Dus. *Hypnum gracile*
S. lagurus (Hook.) Mitt. *Lepyrodon lagurus*
S. lechleri (C. Muell.) Mitt. *Hypnum chrysogaster*
S. mollis (Hedw.) Mitt. *Weymouthia mollis*
S. nitidus (Hook. f. et Wils.) Mitt. *Catagonium nitidum*
S. pallens Broth. *Hypnum skottsbergii*
S. pseudo-pallens Dus. (Dusén in Naveau, 1928) *nom. nud.*
Distr. Southern South America (Naveau, 1928).
- S. ptychocarpum* (Schwaegr.) Mitt. *Ptychomnion ptychocarpum*
S. revolutus Mitt. *Hypnum revolutum*
S. spegazzinii (C. Muell.) Broth. *Hypnum spegazzinii*
S. varius (Hedw.) Mitt. *Amblystegium varium*
- Stereohypnum**
- S. humile* (Besch.) Herz. *Mittenothamnium humile*
S. tapes (C. Muell.) Fleisch. *Mittenothamnium tapes*
- Stereophyllum**
- S. aptychopsis* C. Muell. (Mueller, 1897a).
Distr. Southern South America (Mueller, 1897a; Paris, 1905b; Brotherus, 1925; Kühnemann, 1938; Herzog, 1952a).
- S. argentinicum* Broth. (Brotherus, 1918).
Distr. Southern South America (Brotherus, 1918, 1925; Kühnemann, 1938).
- S. brevipes* (C. Muell.) Mitt. (Mitten, 1869).
Syn. *Hypnum brevipes* C. Muell. (Mueller, 1851a).
Euglossophyllum brevipes (C. Muell.) Hamp. (Hampe, 1870) *comb. inval.*
Glossophyllum brevipes (C. Muell.) Hamp. (Hampe, 1879).
Distr. Southern South America (Brotherus, 1918; Kühnemann, 1938).
- S. fuegianum* Besch. *Brachythecium fuegianum*
- S. homaliooides* Besch. (Bescherelle, 1891).
Distr. Southern South America (Herzog, 1952a).
var. *bellirete* Herz. (Herzog, 1952a).
Distr. Southern South America (Herzog, 1952a).

<i>S. krausei</i> (C. Muell.) Broth.	<i>Juratzkaea krausei</i>
<i>S. lepidopilooides</i> (C. Muell.) Wijk et Marg. (van der Wijk and Margadant, 1959).	
Syn. <i>Hypnum lepidopilooides</i> C. Muell. (Mueller, 1879a). <i>Brachythecium lepidopilooides</i> (C. Muell.) Kindb. (Kindberg, 1888).	
Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904a; Hosseus, 1935a, 1937).	
<i>S. leucostegum</i> (Brid.) Mitt. (Mitten, 1869).	
Syn. <i>Leskea leucostega</i> Brid. (Bridel, 1827b). <i>Hypnum leucostegum</i> (Brid.) C. Muell. (Mueller, 1851a). <i>Rhyncho-hypnum leucostegum</i> (Brid.) Hamp. (Hampe, 1870) <i>comb. inval.</i>	
Distr. Southern South America (Brotherus, 1918, 1925; Kühnemann, 1938; Herzog, 1952a).	
<i>S. seminerve</i> (Schwaegr.) Mitt.	<i>Juratzkaea seminervis</i>
<i>Streblotrichum</i>	
<i>S. gracile</i> Par.	<i>Tortella humilis</i>
<i>S. humile</i> (Hedw.) P. Beauv.	<i>Tortella humilis</i>
<i>S. piliferum</i> (Hook.) Hilp.	<i>Tortula pilifera</i>
<i>S. unguiculatum</i> (Hedw.) Loesk.	<i>Barbula unguiculata</i>
<i>Streptocalypta</i>	
<i>S. fernandeziana</i> Card. (Cardot in Skottsberg, 1914) <i>nom. nud.</i>	
Distr. Southern South America (Skottsberg, 1914).	
<i>S. lorentziana</i> C. Muell. (Mueller, 1879a).	
Syn. <i>Streptocalypta lorentzii</i> C. Muell. ex Kindb. (Mueller ex Kindberg, 1889) <i>nom. illeg.</i>	
Distr. Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1924a; Hilpert, 1933; Kühnemann, 1938).	
<i>S. lorentzii</i> C. Muell. ex Kindb.	<i>Streptocalypta lorentziana</i>
<i>Streptopogon</i>	
<i>S. crispatus</i> (C. Muell.) Par.	<i>Calyptopogon mnioides</i>
<i>S. heterophyllus</i> Herz. (Herzog, 1916a).	
Distr. Southern South America (Herzog, 1952a).	
<i>S. mnioides</i> (Schwaegr.) Mitt.	<i>Calyptopogon mnioides</i>
var. <i>prostratus</i> (Mont.) Jaeg.	<i>Tortula prostrata</i>
<i>Symblepharis</i>	
<i>S. pumila</i> Mitt.	<i>Holodontium pumilum</i>
<i>Syntrichia</i>	
<i>S. anderssonii</i> (Aongstr.) Herz.	<i>Tortula anderssonii</i>
<i>S. apiculato-pilosa</i> Dus.	<i>Tortula apiculato-pilosa</i>
<i>S. arenae</i> (Besch.) Herz.	<i>Tortula arenae</i>
<i>S. brachychaete</i> Dus.	<i>Tortula brachychaete</i>
<i>S. brevicaulis</i> Dus.	<i>Tortula brevicaulis</i>
<i>S. costesii</i> (Thér.) Herz.	<i>Tortula costesii</i>
<i>S. crispatula</i> C. Muell.	<i>Calyptopogon mnioides</i>
<i>S. ericetorum</i> (With.) Brid.	<i>Barbula unguiculata</i> var. <i>patagonica</i>

<i>S. flavidopilosa</i> Dus.	<i>Tortula flavidopilosa</i>
<i>S. fuegiana</i> Mitt.	<i>Tortula fuegiana</i>
<i>S. hyperborea</i> Brid.	<i>Tortula mucronifolia</i> var. <i>arctica</i>
<i>S. intermedia</i> Brid.	<i>Tortula intermedia</i>
<i>S. laevipila</i> Brid.	<i>Tortula laevipila</i>
<i>S. lepto-syntrichia</i> C. Muell.	<i>Tortula leptosyntrichia</i>
<i>S. levipila</i> Brid. ex Moenk.	<i>Tortula levipila</i>
<i>S. lingulaefolia</i> (Card. et Broth.) Herz.	<i>Tortula lingulaefolia</i>
<i>S. marginato-serrata</i> Dus.	<i>Tortula marginato-serrata</i>
<i>S. micro-runcinata</i> Dus.	<i>Leptodontium longicaule</i> var. <i>microruncinatum</i>
<i>S. mniadelphus</i> (C. Muell.) Herz. var. <i>cochlearifolia</i> Herz.	<i>Tortula mniadelphus</i> <i>Tortula cochlearifolia</i>
<i>S. muelleri</i> Bruch	<i>Tortula princeps</i>
<i>S. muralis</i> (Hedw.) Raab	<i>Tortula muralis</i>
<i>S. papillosa</i> (Wils.) Jur.	<i>Tortula papillosa</i>
<i>S. perichaetialis</i> Herz. (Herzog, 1957) nom. nud. <i>Distr.</i> Southern South America (Herzog, 1957).	
<i>S. perrubiginosa</i> (Dus.) Par.	<i>Tortula anderssonii</i>
<i>S. princeps</i> (De Not.) Mitt.	<i>Tortula princeps</i>
<i>S. prostrata</i> (Mont.) Herz.	<i>Tortula prostrata</i>
<i>S. pseudo-robusta</i> (Dus.) Herz.	<i>Tortula pseudo-robusta</i>
<i>S. robusta</i> Dus.	<i>Tortula pseudo-robusta</i>
<i>S. robusta</i> (Hook. et Grev.) Herz.	<i>Tortula robusta</i>
<i>S. rubripungeus</i> Broth. (Brotherus in Neger, 1899) nom. nud. <i>Distr.</i> Southern South America (Neger, 1899).	
<i>S. runcinata</i> C. Muell.	<i>Tortula robusta</i> var. <i>runcinata</i>
<i>S. ruralis</i> (Hedw.) Web. et Mohr var. <i>laevipila</i> (Brid.) Spreng.	<i>Tortula ruralis</i> <i>Tortula laevipila</i>
<i>S. scabrella</i> (Dus.) Herz.	<i>Tortula scabrella</i>
<i>S. scabrinervis</i> (C. Muell.) Herz.	<i>Tortula scabrinervis</i>
<i>S. serripungens</i> (Lor. et C. Muell.) Herz.	<i>Tortula serripungens</i>
<i>S. squarripila</i> (Thér.) Herz.	<i>Tortula squarripila</i>

Syrrhopodon***S. argentinicus*** Lor. ex C. Muell. (Lorentz ex Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a; Paris, 1905b; Brotherus, 1924a; Kühnemann, 1938).

<i>S. berteroanus</i> (Brid.) C. Muell.	<i>Syrrhopodon incompletus</i> var. <i>berteroanus</i>
<i>S. berteroi</i> Kindb.	<i>Syrrhopodon incompletus</i> var. <i>berteroanus</i>
<i>S. decolorans</i> C. Muell.	<i>Syrrhopodon incompletus</i>
<i>S. fasciculatus</i> Hook. et Grev.	<i>Thyridium fasciculatum</i>

S. incompletus Schwaegr. (Schwaegrichen, 1824).*Syn.* *Syrrhopodon decolorans* C. Muell. (Mueller, 1897c).*Distr.* Southern South America (Hosseus, 1940).***var. berteroanus*** (Brid.) Reese (Reese, 1977).

- Syn.* *Dicranum berteroanum* Brid. (Bridel, 1826).
Orthotheca berteroana (Brid.) Brid. (Bridel, 1827a).
Syrrhopodon berteroanus (Brid.) C. Muell. (Mueller, 1849).
Syrrhopodon berteroii Kindb. (Kindberg, 1888) *nom. illeg.*
Syrrhopodon malouinensis C. Muell. (Mueller, 1849) *nom. illeg.*
- Distr.* Falkland Islands (Bridel, 1827a; Mueller, 1849; Jaeger, 1873; Paris, 1905b; Cardot, 1908a; Kühnemann, 1938).
- S. malouinensis* C. Muell. *Syrrhopodon incompletus* var. *berteroanus*
S. reinwardtii Hornsch. *Zygodon reinwardtii*
- S. rigescens* Schwaegr. (Schwaegrichen, 1827a).
Distr. Falkland Islands (Schwaegrichen, 1827a; Mueller, 1849; Jaeger, 1873; Paris, 1905b; Cardot, 1908a; Kühnemann, 1938).
- S. tortuosus* Hornsch. *Amphidium tortuosum*
- Systegium horn.* *illeg.*
- S. fruchartii* (C. Muell.) Kindb. *Astomum fruchartii*
S. recurvirostre (C. Muell.) Kindb. *Tetraphterum recurvirostre*
- Tamariscella**
- T. pseudo-aequatorialis* C. Muell. *Thuidium peruvianum*
var. *casuarina* C. Muell. *Thuidium pseudo-aequatoriale* var. *casuarinum*
- Taxicaulis**
- T. adflatum* C. Muell. *Isopterygium adflatum*
T. byssobolax C. Muell. *Isopterygium byssobolax*
T. pyrrhopus C. Muell. *Isopterygium pyrrhopus*
T. saprophilus C. Muell. *Isopterygium saprophilum*
- Taxithelium**
- T. microthamnioides* (C. Muell.) Par. (Paris, 1898).
Syn. *Sigmatella microthamnioides* C. Muell. (Mueller, 1897a).
Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1905b).
- Tayloria**
- T. dubyi* Broth. (Brotherus, 1903).
Syn. *Hymenocleiston magellanicum* Dub. (Duby, 1875b).
Dissodon dubyanus Besch. (Bescherelle in Brotherus, 1924a) *nom. nud.*
Distr. Southern South America (Duby, 1875a, b; Jaeger and Sauerbeck, 1879b; Mueller, 1885; Bescherelle, 1885a, 1889; Brotherus, 1903, 1924a; Cardot, 1905a, 1908a, b; Paris, 1905b; Cardot and Brotherus, 1923; Kühnemann, 1938; Bartram, 1952; Seki, 1974).
var. *edenensis* (Besch.) Par. (Paris, 1905b).
Syn. *Hymenocleiston magellanicum* Dub. var. *edenense* Besch. (Bescherelle, 1885a).
Distr. Southern South America (Bescherelle, 1885a, 1889; Paris, 1905b; Cardot, 1908a).
- T. magellanica* (Brid.) Mitt. (Mitten, 1869).
Syn. *Splachnum magellanicum* Brid. (Bridel, 1802).
Eremodon magellanicus (Brid.) Brid. (Bridel, 1826).
Dissodon magellanicus (Brid.) Hamp. (Hampe in Mueller, 1851a).
Distr. Southern South America (Bridel, 1803, 1806, 1826; Schwaegrichen, 1811; Mueller, 1848a, 1851b, 1885; Montagne, 1850; Mitten, 1869; Ångström, 1872; Jaeger, 1874; Bescherelle, 1889; Cardot, 1900, 1901, 1908a; Paris, 1904b; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).
var. *gigantea* Bartr. (Bartram, 1946).
Distr. Southern South America (Bartram, 1946).

T. mirabilis (Card.) Broth. (Brotherus, 1903).

Syn. *Dissodon mirabilis* Card. (Cardot, 1900).

Distr. Southern South America (Cardot, 1900, 1901, 1908a; Brotherus, 1903, 1924a; Paris, 1904b; Cardot and Brotherus, 1923; Kühnemann, 1938; Bartram, 1952).

T. octoblepharum (Hook.) Mitt. (Mitten, 1883).

Syn. *Splachnum octoblepharum* Hook. (Hooker, 1819).

Dissodon plagiopus (Mont.) C. Muell. (Mueller, 1851a).

Distr. Southern South America (Ångström, 1872).

T. patagonica Dus. (Dusén *in* Paris, 1905b) *nom. nud.*

Distr. Southern South America (Paris, 1905b).

T. spathulata (Hook. f. et Wils.) C. Muell. (Mueller, 1848a).

Syn. *Eremodon spathulatus* Hook. f. et Wils. (Hooker and Wilson *in* Wilson, 1846).

Distr. Southern South America (Wilson, 1846; Paris, 1905b).

Tetraplodon

T. fuegianus Besch. (Bescherelle, 1885a).

Distr. Southern South America (Bescherelle, 1885a, 1889; Paris, 1905b; Cardot, 1908a; Brotherus, 1924a; Seki, 1974).

T. mnioides (Hedw.) B.S.G. (Bruch and others, 1844).

Syn. *Splachnum mnioides* Hedw. (Hedwig, 1801).

Distr. Southern South America (Robinson, 1964).

T. stenophysatus Herz. (Herzog, 1939).

Distr. Southern South America (Herzog, 1939).

Tetrapterum

T. lamprocarpum (C. Muell.) Broth. (Brotherus, 1924a).

Syn. *Phascum lamprocarpum* C. Muell. (Mueller, 1879a).

Pottia lamprocarpa (C. Muell.) Broth. (Brotherus, 1902).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Roth, 1911; Kühnemann, 1938).

T. lamprothecium (C. Muell.) Broth. (Brotherus, 1924a).

Syn. *Phascum lamprothecium* C. Muell. (Mueller, 1879a).

Pottia lamprothecium (C. Muell.) Broth. (Brotherus, 1902).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Roth, 1911; Kühnemann, 1938).

T. recurvirostre (C. Muell.) Broth. (Brotherus, 1924a).

Syn. *Phascum recurvirostre* C. Muell. (Mueller, 1888).

Astomium recurvirostre (C. Muell.) Roth (Roth, 1911).

Systegium recurvirostre (C. Muell.) Kindb. (Kindberg, 1889).

Astomium chilense Williams (Williams, 1915).

Distr. Southern South America (Williams, 1915; Thériot, 1921a).

Thamnium

T. angustifolium Schimp.

Porothamnium pandraefolium

T. arbusculans (C. Muell.) Jaeg.

Porothamnium arbusculans

var. *complanatum* C. Muell.

Porothamnium arbusculans

T. assimile Broth.

Thamnobryum rigidum

T. caroli Broth.

Thamnobryum carolii

<i>T. confertum</i> (Mitt.) Par.	<i>Thamnobryum confertum</i>
<i>T. corralense</i> Broth.	<i>Porothamnium arbusculans</i>
<i>T. crassinervium</i> Broth.	<i>Thamnobryum rigidum</i>
<i>T. decumbens</i> Besch.	<i>Porothamnium arbusculans</i>
<i>T. explanatum</i> (Mitt.) Kindb.	<i>Porothamnium explanatum</i>
<i>T. fasciculatum</i> (Hedw.) C. Muell.	<i>Porothamnium fasciculatum</i>
<i>T. ingae</i> Broth.	<i>Thamnobryum ingae</i>
<i>T. latinerve</i> (Mitt.) Broth.	<i>Thamnium rigidum</i>
<i>T. leucocaulon</i> (C. Muell.) Kindb.	<i>Porothamnium leucocaulon</i>
<i>T. mattogrossense</i> Broth.	<i>Porothamnium mattogrossense</i>
<i>T. naumannii</i> (C. Muell.) Kindb.	<i>Porothamnium arbusculans</i>
<i>T. neckera</i> Schimp.	<i>Porothamnium valdiviae</i>
<i>T. neckeraeforme</i> (Hamp.) Kindb.	<i>Porothamnium neckeraeforme</i>
<i>T. neckeroides</i> (Hook.) Jaeg. (Jaeger, 1877).	
Syn. <i>Hypnum neckeroides</i> Hook. (Hooker, 1818).	
<i>Isothecium neckeroides</i> (Hook.) Brid. (Bridel, 1827b).	
<i>Porotrichum neckeroides</i> (Hook.) Williams (Williams, 1901).	
Distr. Juan Fernandez (Johow, 1896).	
Southern South America (Montagne, 1850; Paris, 1905b; Brotherus, 1925).	
<i>T. panduraefolium</i> (C. Muell.) Broth.	<i>Porothamnium panduraefolium</i>
<i>T. platycladum</i> Card. (Cardot in Skottsberg, 1914) nom. nud.	
Distr. Juan Fernandez (Skottsberg, 1914).	
<i>T. porrectulum</i> (C. Muell.) Broth.	<i>Porothamnium porrectulum</i>
<i>T. proboscideum</i> Broth.	<i>Thamnobryum proboscideum</i>
<i>T. pumilum</i> (Hook. f. et Wils.) Kindb.	<i>Thamnobryum pumilum</i>
<i>T. rigidum</i> (Mitt.) Broth.	<i>Thamnobryum rigidum</i>
<i>T. thielei</i> Kindb.	<i>Pinnatella thieleana</i>
<i>T. valdiviae</i> (C. Muell.) Kindb.	<i>Porothamnium valdiviae</i>
<i>T. valdivicum</i> C. Muell.	<i>Porothamnium valdiviae</i>
<i>Thamnobryum</i>	
<i>T. carolii</i> (Broth.) H. Robinson (Robinson, 1974).	
Syn. <i>Thamnium caroli</i> Broth. (Brotherus, 1924b).	
Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Bartram, 1959; Robinson, 1974, 1975).	
<i>T. confertum</i> (Mitt.) H. Robinson (Robinson, 1974).	
Syn. <i>Porotrichum confertum</i> Mitt. (Mitten, 1885).	
<i>Thamnium confertum</i> (Mitt.) Par. (Paris, 1898).	
Distr. Juan Fernandez (Mitten, 1885; Johow, 1896; Paris, 1898, 1905b; Brotherus, 1924b; Robinson, 1974, 1975).	
<i>T. ingae</i> (Broth.) H. Robinson (Robinson, 1974).	
Syn. <i>Thamnium ingae</i> Broth. (Brotherus, 1924b).	
Distr. Juan Fernandez (Brotherus, 1924b, 1925; Thériot, 1927b; Espinosa B., 1941; Bartram, 1959; Robinson, 1974, 1975).	
<i>T. pumilum</i> (Hook. f. et Wils.) Nieuwl. (Nieuwland, 1917).	
Syn. <i>Isothecium pumilum</i> Hook. f. et Wils. (Hooker and Wilson in Wilson, 1859).	

Microthamnium pumilum (Hook. f. et Wils.) Jaeg. (Jaeger, 1878).
Thamnium pumilum (Hook. f. et Wils.) Kindb. (Kindberg, 1891).

Distr. Juan Fernandez (Kindberg, 1902).

T. proboscideum (Broth.) H. Robinson (Robinson, 1974).

Syn. *Thamnium proboscideum* Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Bartram, 1959; Robinson, 1974, 1975).

T. rigidum (Mitt.) H. Robinson (Robinson, 1974).

Syn. *Porotrichum rigidum* Mitt. (Mitten, 1869).

Thamnium rigidum (Mitt.) Broth. (Brotherus, 1906).

Porotrichum latinerve Mitt. (Mitten, 1885).

Thamnium latinerve (Mitt.) Broth. (Brotherus, 1924b).

Thamnium crassinervium Broth. (Brotherus, 1906).

Porotrichum crassinervum Mitt. ex Broth. (Mitten ex Brotherus, 1924b) *nom. inval.*

Thamnium assimile Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Mitten, 1869, 1885; Jaeger, 1877; Johow, 1896; Brotherus, 1906, 1924b, 1925; Skottsberg, 1914; Thériot, 1921b; Espinosa B., 1941; Bartram, 1959; Robinson, 1974, 1975).

Thuidiopsis

T. chilensis (Mitt.) Broth. (Brotherus, 1925).

Syn. *Thuidium chilense* Mitt. (Mitten, 1869).

Distr. Southern South America (Mitten, 1869; Jaeger, 1878; Paris, 1906; Herzog, 1923, 1954; Brotherus, 1925).

T. corralensis (Broth.) Herz.

Thuidiopsis furfurosa

T. dusenii (Broth.) Broth. (Brotherus, 1925).

Syn. *Thuidium dusenii* Broth. (Brotherus in Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1925; Seki, 1974).

T. filaria (Mitt.) Broth.

Thuidiopsis furfurosa

T. furfurosa (Hook. f. et Wils.) Fleisch. (Fleischer, 1923).

Syn. *Hypnum furfurosum* Hook. f. et Wils. (Hooker and Wilson in Wilson, 1854).

Thuidium furfurosum (Hook. f. et Wils.) Reichdt. (Reichardt, 1870).

Thuidium filarium Mitt. (Mitten, 1869).

Cyto-Hypnum filarium (Mitt.) Hamp. (Hampe, 1878).

Hypnum filarium (Mitt.) Hamp. (Hampe, 1879).

Thuidium filare Dus. (Dusén in Paris, 1906) *nom. nud.*

Thuidiopsis filaria (Mitt.) Broth. (Brotherus, 1925).

Hypnum unguiculatum Hook. f. et Wils. (Hooker and Wilson in Wilson, 1859).

Thuidium unguiculatum (Hook. f. et Wils.) Fleisch. (Fleischer, 1923).

Thuidiopsis unguiculata (Hook. f. et Wils.) Fleisch. (Fleischer, 1923).

Thuidium corralense Broth. (Brotherus in Cardot and Brotherus, 1923).

Thuidiopsis corralensis (Broth.) Herz. (Herzog, 1939).

Thuidium valdiviae Broth. (Brotherus, 1924b) *fide* Robinson, 1975.

Thuidiopsis valdiviae (Broth.) Broth. (Brotherus, 1925).

Distr. Juan Fernandez (Skottsberg, 1914 (as *Thuidium valdiviense*); Brotherus, 1924b, 1925; Bartram, 1959; Robinson, 1975).

Southern South America (Dusén, 1903a; Paris, 1906; Thériot, 1918; Cardot and Brotherus, 1923; Brotherus, 1925; Reimers, 1926; Dixon, 1929b; Herzog, 1938, 1939, 1954, 1960; Hosseus, 1940; Bartram, 1952; Bizot, 1967; Seki, 1974).

T. masafuerae (Broth.) Broth. (Brotherus, 1925).

Syn. *Thuidium masafuerae* Broth. (Brotherus, 1924b).

Distr. Juan Fernandez (Brotherus, 1924b, 1925; Espinosa B., 1941; Robinson, 1975).

T. unguiculata (Hook. f. et Wils.) Fleisch.

Thuidiopsis furfurosa

T. valdiviae (Broth.) Broth.

Thuidiopsis furfurosa

Thuidium

<i>T. amblyostomum</i> (C. Muell.) Kindb.	<i>Haplocladium microphyllum</i>
var. <i>attenuatifolium</i> (C. Muell.) Par.	<i>Haplocladium microphyllum</i>
var. <i>paludicola</i> (C. Muell.) Par.	<i>Haplocladium microphyllum</i>
<i>T. austro-serpens</i> (C. Muell.) Kindb.	<i>Haplocladium microphyllum</i>
<i>T. brachypyxis</i> C. Muell. (Mueller, 1897a).	
Syn. <i>Thuidium lorentzii</i> Par. (Paris, 1898) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Mueller, 1897a; Paris, 1898, 1906; Brotherus, 1925; Kühnemann, 1938).	
<i>T. chacoanum</i> C. Muell. (Mueller, 1897a).	
<i>Distr.</i> Southern South America (Mueller, 1897a; Paris, 1906; Brotherus, 1925; Kühnemann, 1938).	
<i>T. chilense</i> Mitt.	<i>Thuidiopsis chilensis</i>
<i>T. condensatifolium</i> Dus. (Dusén in Paris, 1906) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Paris, 1906).	
<i>T. corralense</i> Broth.	<i>Thuidiopsis furfurosa</i>
<i>T. cylindrella</i> C. Muell. (Mueller, 1897a).	
<i>Distr.</i> Southern South America (Mueller, 1897a; Paris, 1906; Kühnemann, 1938).	
<i>T. dusenii</i> Broth.	<i>Thuidiopsis dusenii</i>
<i>T. exasperatum</i> Mitt. (Mitten, 1869).	
<i>Distr.</i> Southern South America (Herzog, 1952a).	
<i>T. filare</i> Dus.	<i>Thuidiopsis furfurosa</i>
<i>T. filarium</i> Mitt.	<i>Thuidiopsis furfurosa</i>
<i>T. firmulum</i> C. Muell.	<i>Rauiella firmula</i>
<i>T. furfurosum</i> (Hook. f. et Wils.) Reichdt.	<i>Thuidiopsis furfurosa</i>
<i>T. inconspicuum</i> Herz. (Herzog, 1923).	
<i>Distr.</i> Southern South America (Herzog, 1923).	
<i>T. laterale</i> Par.	<i>Haplocladium microphyllum</i>
<i>T. laterculi</i> (C. Muell.) Kindb.	<i>Haplocladium microphyllum</i>
<i>T. ligulifolium</i> Herz. (Herzog, 1916b).	
<i>Distr.</i> Southern South America (Herzog, 1916b).	
<i>T. lorentzii</i> Par.	<i>Thuidium brachypyxis</i>
<i>T. masafuerae</i> Broth.	<i>Thuidiopsis masafuerae</i>
<i>T. microphyllum</i> (Hedw.) Jaeg.	<i>Haplocladium microphyllum</i>
<i>T. niveo-calycinum</i> C. Muell.	<i>Rauiella niveo-calycina</i>
<i>T. occultirete</i> C. Muell. (Mueller, 1897a).	
<i>Distr.</i> Southern South America (Mueller, 1897a; Paris, 1906; Kühnemann, 1938).	
<i>T. persistens</i> (C. Muell.) Kindb.	<i>Haplocladium persistens</i>
<i>T. peruvianum</i> Mitt. (Mitten, 1869).	
Syn. <i>Tamariscella pseudo-aequatorialis</i> C. Muell. (Mueller, 1897a).	
<i>Thuidium pseudo-aequatoriale</i> (C. Muell.) Par. (Paris, 1898).	
<i>Distr.</i> Southern South America (Mueller, 1897a; Paris, 1898, 1906; Kühnemann, 1938).	
<i>T. pseudo-aequatoriale</i> (C. Muell.) Par.	<i>Thuidium peruvianum</i>

var. *casuarinum* (C. Muell.) Par. (Paris, 1898).

Syn. *Tamariscella pseudo-aequatorialis* C. Muell. var. *casuarina* C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1898, 1906; Brotherus, 1925).

T. pseudo-gracile (C. Muell.) Par.

Haplocladium microphyllum

T. pseudo-recognitum (Hamp.) Kindb.

Thuidium urceolatum

T. pulvinatulum C. Muell. (Mueller, 1897a).

Distr. Southern South America (Mueller, 1897a; Paris, 1906; Brotherus, 1925; Herter, 1933a; Hosseus, 1939).

T. raphidostegum (C. Muell.) Mitt. (Mitten, 1869).

Syn. *Hypnum raphidostegum* C. Muell. (Mueller, 1851a).

Hypnum raphidostegum C. Muell. ex Par. (Mueller ex Paris, 1895) nom. inval.

Thuidium raphidostegium Mitt. ex Par. (Mitten ex Paris, 1898) nom. inval.

Distr. Southern South America (Mueller, 1851a; Mitten, 1869; Jaeger, 1878; Paris, 1898, 1906; Brotherus, 1925; Herzog, 1938).

T. recognitum (Hedw.) Lindb. var. ***delicatulum*** (Hedw.) Warnst. (Warnstorff, 1881b).

Syn. *Hypnum delicatulum* Hedw. (Hedwig, 1801).

Hypnum tamariscinum Hedw. var. *delicatulum* (Hedw.) Brid. (Bridel, 1812).

Distr. Southern South America (Montagne, 1850).

T. remotifolium (Grev.) Ren. et Card.

Oxyrrhynchium clinocarpum

T. raphidostegium Mitt. ex Par.

Thuidium raphidostegum

T. semi-lunare C. Muell.

Haplocladium microphyllum

T. tamariscinum (Hedw.) B.S.G. (Bruch and others, 1852).

Syn. *Hypnum tamariscinum* Hedw. (Hedwig, 1801).

Distr. Southern South America (Montagne, 1850).

T. torskii Kiaer ex C. Muell.

Haplocladium microphyllum

T. unguiculatum (Hook. f. et Wils.) Fleisch.

Thuidiopsis furfurosa

T. urceolatum Lor. (Lorentz, 1864).

Syn. *Thuidium pseudo-recognitum* (Hamp.) Kindb. (Kindberg, 1891).

Distr. Southern South America (Herzog, 1952a).

T. valdiviae Broth.

Thuidiopsis furfurosa

Thyridium

T. fasciculatum (Hook. et Grev.) Mitt. (Mitten, 1868).

Syn. *Syrrhopodon fasciculatus* Hook. et Grev. (Hooker and Greville, 1825b).

Calymperes fasciculatum (Hook. et Grev.) Mitt. (Mitten, 1859) hom. illeg.

Distr. Southern South America (Mitten, 1859, 1869; Jaeger, 1873; Paris, 1905b).

Thysanomitrion

T. gracile (Hook.) Arnott

Pilopogon gracilis

var. *robustum* Arnott

Pilopogon gracilis

T. griseum Hornsch.

Campylopus griseus

T. introflexum (Hedw.) Arnott

Campylopus introflexus

T. laetevirens Broth.

Campylopus laetevirens

T. leptodus (Mont.) Broth.

Campylopus clavatus

T. richardii (Brid.) Schwaegr.

Campylopus richardii

T. scabrisetum Hamp.

Campylopus arctocarpus

T. stenocarpum Hamp.

Campylopus stenocarpus

Timmiella***T. argentinica*** Broth. (Brotherus, 1918).*Distr.* Southern South America (Brotherus, 1918, 1924a; Hosseus, 1936, 1937, 1938a, b, c, d, 1939; Thériot, 1936; Kühnemann, 1938).***T. umbrosa*** (C. Muell.) Broth. (Brotherus, 1902).*Syn.* *Trichostomum umbrosum* C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Kurtz, 1904; Paris, 1906; Williams, 1930; Hosseus, 1935a, b, 1937; Kühnemann, 1938).***Tortella******T. acaulon*** (C. Muell.) Broth. (Brotherus, 1902).*Syn.* *Trichostomum acaulon* C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Kurtz, 1904; Paris, 1906; Hosseus, 1935b, 1937; Kühnemann, 1938).***T. brachydontia*** (Bruch) C. Jens.***Trichostomum brachydontium******T. caespitosa*** (Schwaegr.) Limpr.***Tortella humilis******T. calycina*** (Schwaegr.) Dix. (Dixon, 1923).*Syn.* *Barbula calycina* Schwaegr. (Schwaegrichen, 1823).*Tortula calycina* (Schwaegr.) Hook. et Grev. (Hooker and Greville, 1824b).*Distr.* Southern South America (Mitten, 1869; Jaeger, 1873; Paris, 1904a; Brotherus, 1925).***T. hosseusii*** Herz. (Herzog, 1938).*Distr.* Southern South America (Herzog, 1938).***T. humilis*** (Hedw.) Jenn. (Jennings, 1913).*Syn.* *Barbula humilis* Hedw. (Hedwig, 1801).*Tortula humilis* (Hedw.) Turn. (Turner, 1804a).*Streblotrichum humile* (Hedw.) P. Beauv. (Palisot de Beauvois, 1805a).*Mollia humilis* (Hedw.) Lindb. (Lindberg, 1883).*Trichostomum humile* (Hedw.) Mach. (Machado, 1919).*Barbula caespitosa* Schwaegr. (Schwaegrichen, 1811).*Tortula caespitosa* (Schwaegr.) Hook. et Grev. (Hooker and Greville, 1824b).*Tortella caespitosa* (Schwaegr.) Limpr. (Limpricht, 1888).*Streblotrichum gracile* Par. (Paris, 1905b) err.*Distr.* Southern South America (Brotherus, 1918; Herter, 1933a, 1939b; Hosseus, 1938a, b, d, 1940; Kühnemann, 1938; Herzog, 1952a; Bizot and Piovano, 1953; Piovano, 1954).***T. perrufula*** (C. Muell.) Broth. (Brotherus, 1902).*Syn.* *Barbula perrufula* C. Muell. (Mueller, 1897a).*Tortula perrufula* (C. Muell.) Williams (Williams, 1930).*Distr.* Southern South America (Mueller, 1897a; Paris, 1900a, 1906; Brotherus, 1902, 1924a; Herter, 1933a; Kühnemann, 1938).***T. pseudo-caespitosa*** (C. Muell.) Broth. (Brotherus, 1902).*Syn.* *Barbula pseudo-caespitosa* C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Kurtz, 1904; Paris, 1906; Hosseus, 1935b, 1937; Kühnemann, 1938).***var. brachybasis*** (C. Muell.) Par. (Paris, 1904a).*Syn.* *Barbula pseudo-caespitosa* C. Muell. var. *brachybasis* C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a).***var. pungens*** (C. Muell.) Par. (Paris, 1904a).*Syn.* *Barbula pseudo-caespitosa* C. Muell. var. *pungens* C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a).

T. tortuosa (Hedw.) Limpr. (Limpicht, 1888).

Syn. *Tortula tortuosa* Hedw. (Hedwig, 1801).
Bryum tortuosum (Hedw.) L. ex With. (Linnaeus ex Withering, 1801).
Barbula tortuosa (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Mollia tortuosa (Hedw.) Schrank ex Lindb. (Schrank ex Lindberg, 1879).
Trichostomum tortuosum (Hedw.) Dix. (Dixon, 1896).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

T. viridula C. Jens.**Weisia controversa****Tortella** sp.

Syn. *Trichostomum tortella* C. Muell. (Mueller, 1879a).
Distr. Southern South America (Mueller, 1879a).

Tortula**T. aculeonervis** (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula aculeonervis* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Kurtz, 1904; Paris, 1906; Hosseus, 1935b, 1937; Kühnemann, 1938).

T. aequatoriale Spruc.**Trichostomum aequatoriale****T. aestiva** (Hedw.) P. Beauv.**Tortula muralis** ssp. **aestiva****T. amblyophylla** (Hook.) Mitt.**Didymodon amblyophyllus****T. amblyophylla** (Mont.) Thér.**Tortula platyphylla****T. amphidiifolia** (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula amphidiifolia* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Kurtz, 1904; Paris, 1906; Hosseus, 1935b, 1937; Kühnemann, 1938).

T. anacamptophylla (C. Muell.) Broth.**Tortula leptosyntrichia****T. anderssonii** Aongstr. (Ångström, 1872).

Syn. *Barbula anderssonii* (Aongstr.) Jaeg. (Jaeger, 1873).

Syntrichia anderssonii (Aongstr.) Herz. (Herzog, 1954).

Barbula conotricha C. Muell. (Mueller, 1885).

Tortula conotricha (C. Muell.) Par. (Paris, 1906).

Barbula perrubiginosa Dus. (Dusén, 1903a).

Tortula perrubiginosa (Dus.) Par. (Paris, 1906).

Syntrichia perrubiginosa (Dus.) Par. (Paris, 1906) *nom. nud.*

Distr. Southern South America (Ångström, 1872; Jaeger, 1873; Mueller, 1885; Cardot, 1900, 1901, 1905a, 1908a; Dusén, 1903a, 1906; Paris, 1906; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen, 1934; Kühnemann, 1938; Bartram, 1952; Herzog, 1954, 1957; Seki, 1974).

var. *fagicola* (C. Muell.) Card. (Cardot, 1908a).

Syn. *Barbula conotricha* C. Muell. var. *fagicola* C. Muell. (Mueller, 1885).

Tortula conotricha (C. Muell.) Par. var. *fagicola* (C. Muell.) Par. (Paris, 1906).

Distr. Southern South America (Mueller, 1885; Paris, 1906; Cardot, 1908a).

T. andicola Mont. (Montagne, 1838).

Distr. Southern South America (Bartram, 1965).

T. annulus (C. Muell.) Broth.**Tortula muralis****T. antarctica** (Hamp.) Wils.**Tortula princeps****T. apiculato-pilosa** Par. (Paris, 1906) *nom. nud.*

Syn. *Syntrichia apiculato-pilosa* Dus. (Dusén in Paris, 1906) *nom. nud.*

Distr. Southern South America (Paris, 1906).

T. appressa Mitt. (Mitten, 1869).

Syn. *Barbula appressa* (Mitt.) Jaeg. (Jaeger, 1873).

Distr. Southern South America (Mitten, 1869; Jaeger, 1873; Paris, 1906).

T. arenae (Besch.) Broth. (Brotherus, 1902).

Syn. *Barbula arenae* Besch. (Bescherelle, 1885a).

Syntrichia arenae (Besch.) Herz. (Herzog, 1957).

Distr. Southern South America (Bescherelle, 1885a, 1889; Brotherus, 1902, 1924a; Paris, 1906; Cardot, 1908a; Herzog, 1923, 1939, 1957; Roivainen, 1934).

T. arvensis Mont.

Tortula ruralis

var. *australis* Mont.

Tortula flagellaris

T. atrata Thér. (Thériot, 1921a).

Distr. Southern South America (Thériot, 1921a; Brotherus, 1924a).

T. atro-virens (Sm.) Lindb.

Desmatodon convolutus

var. *brevifolia* Thér. (Thériot, 1921a).

Distr. Southern South America (Thériot, 1921a).

T. australasiae Hook. et Grev.

Trichostomopsis australasiae

T. berteroana (C. Muell.) Broth. ex Par.

Tortula muralis

T. berthoana Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926).

T. brachychaete Dus. (Dusén, 1905b).

Syn. *Barbula brachychaete* Dus. (Dusén, 1903a) nom. nud.

Syntrichia brachychaete Dus. (Dusén in Paris, 1906) nom. nud.

Distr. Southern South America (Dusén, 1903a, 1905b, 1906; Paris, 1906; Brotherus, 1924a; Kühnemann, 1938).

T. brachyclada Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Dusén, 1906; Brotherus, 1924a; Kühnemann, 1938).

Falkland Islands (Brotherus, 1924a; Kühnemann, 1938).

T. brachypelma Dus. (Dusén, 1905b).

Syn. *Tortula pusilla* Aongstr. (Ångström, 1872) hom. illeg.

Barbula pusilla Jaeg. (Jaeger, 1873).

Distr. Southern South America (Ångström, 1872; Jaeger, 1873; Mueller, 1885; Dusén, 1905b; Paris, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Bartram, 1952).

T. brevicaulis Par. (Paris, 1906) nom. nud.

Syn. *Syntrichia brevicaulis* Dus. (Dusén in Paris, 1906) nom. nud.

Distr. Southern South America (Paris, 1906).

T. breviseta Mont. (Montagne, 1845a).

Syn. *Barbula breviseta* (Mont.) C. Muell. (Mueller, 1849).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Mitten, 1869; Jaeger, 1873; Paris, 1904a; Thériot, 1923a, b; Brotherus, 1924a; Herzog, 1938).

T. bullata Herz. (Herzog, 1923) hom. illeg.

Distr. Southern South America (Herzog, 1923).

T. caespitosa (Schwaegr.) Hook. et Grev.

Tortella humilis

T. calycina (Schwaegr.) Hook. et Grev.

Tortella calycina

T. calyculosa Mitt.

Barbula calyculosa

T. campestris Dus. (Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Cardot, 1908a; Thériot, 1921a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

T. carinata Gill. ex Grev. (Gillies ex Greville, 1830).

Distr. Southern South America (Greville, 1830).

T. characodonta (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula characodonta* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Brotherus, 1902, 1918, 1924a; Paris, 1906; Kühnemann, 1938).

T. chilensis Mitt.**Trichostomum aequatoriale****T. chrysopila** (C. Muell.) Par. (Paris, 1906).

Syn. *Barbula chrysopila* C. Muell. (Mueller, 1885).

Distr. Southern South America (Mueller, 1885; Paris, 1906; Cardot, 1908a).

T. chubutensis Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b, 1907; Brotherus, 1924a; Kühnemann, 1938).

T. chubutensis Broth. (Brotherus, 1931) *hom. illeg.*

Distr. Southern South America (Brotherus, 1931).

T. conferta Bartr. (Bartram, 1957).

Distr. Antarctic, peninsula region (Bartram, 1957; Steere, 1961a; Greene, 1967; Robinson, 1972 (as *T. grossiretis*)).

Antarctic, continental region (Greene, 1967).

T. conotricha (C. Muell.) Par.**Tortula anderssonii**

var. *fagicola* (C. Muell.) Par.

Tortula anderssonii var. *fagicola***T. contorta** (C. Muell.) Mont. (Montagne, 1850).

Syn. *Barbula contorta* Hamp. ex C. Muell. (Hampe ex Mueller, 1849).

Distr. Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Paris, 1906; Brotherus, 1924a).

T. costesii Thér. (Thériot, 1921a).

Syn. *Syntrichia costesii* (Thér.) Herz. (Herzog, 1954).

Tortula curta Card. et Broth. (Cardot and Brotherus, 1923) *hom. illeg.*

Distr. Southern South America (Thériot, 1921a, b; Cardot and Brotherus, 1923; Brotherus, 1924a; Herzog, 1938, 1954; Kühnemann, 1938).

T. curta Card. et Broth..**Tortula costesii****T. densifolia** (Hook. f. et Wils.) Hook. f. et Wils. (Hooker and Wilson in Wilson and Hooker, 1847).

Syn. *Barbula densifolia* Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).

Neobarbula magellanica Dus. (Dusén, 1905b).

Distr. Southern South America (Mueller, 1849; Dusén, 1905b, 1906; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Herzog, 1923; Kühnemann, 1938).

Falkland Islands (Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Jaeger, 1873; Paris, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

T. epilosa Broth. ex Dus. (Brotherus ex Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Brotherus, 1924a; Thériot, 1934b).

var. *pilifera* Thér. (Thériot, 1926).

Distr. Southern South America (Thériot, 1926).

T. excelsa Card. (Cardot, 1906a).

Distr. Antarctic, peninsula region (Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961a; Robinson, 1972).

T. felipponei Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Am. t in *Index muscorum*, 5, 82, but not found in Felippone (1929).

T. ferruginea Bartr. (Bartram, 1965).

Distr. Southern South America (Bartram, 1965).

T. filaris (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula filaris* C. Muell. (Mueller, 1890a).

Distr. South Georgia (Mueller, 1890a; Brotherus, 1902, 1924a; Paris, 1906; Cardot, 1906a, 1908a; Steere, 1961b).

T. flagellaris (Schimp.) Mont. (Montagne, 1850).

Syn. *Baarbula flagellaris* Schimp. (Schimper, 1836).

Barbula perflaccida Broth. (Brotherus in Paris, 1906) *nom. nud.*

Tortula perflaccida Broth. ex Dus. (Brotherus ex Dusén, 1906).

Tortula ruralis (Hedw.) Gaertn., Meyer et Scherb. var. *australis* Mont. (Montagne, 1835).

Tortula arvensis Mont. var. *australis* Mont. (Montagne, 1850) *err.*

Distr. Juan Fernandez (Montagne, 1835; Johow, 1896; Brotherus, 1924b; Robinson, 1975). Southern South America (Schimper, 1836; Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Dusén, 1903a, 1906; Paris, 1904a, 1906; Cardot, 1908a; Campo, 1915; Thériot, 1917a, b, 1921a; Costes, 1921; Herzog, 1923; Brotherus, 1924a, c; Kühnemann, 1938; Espinosa B., 1941).

var. *densiretis* Thér. (Thériot, 1917a).

Distr. Southern South America (Thériot, 1917a).

T. flavidо-пilosa Par. (Paris, 1906) *nom. nud.*

Syn. *Barbula flavidо-пilosa* Dus. (Dusén, 1903a) *nom. nud.*

Syntrichia flavidо-пilosa Dus. (Dusén in Paris, 1906) *nom. nud.*

Distr. Southern South America (Dusén, 1903a; Paris, 1906; Seki, 1974).

T. flavipes Broth. (Brotherus, 1924c) *hom. illeg.*

Distr. Southern South America (Brotherus, 1924c, 1925; Espinosa B., 1941; Skottsberg, 1950).

T. flaviseta Mitt.**Pottia flavipes****T. fontana** (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula fontana* C. Muell. (Mueller, 1890a).

Distr. South Georgia (Mueller, 1890a; Brotherus, 1902, 1924a; Paris, 1906; Cardot, 1908a).

T. fragilis Tayl. (Taylor, 1847).

Distr. Southern South America (Hosseus, 1936, 1937, 1938a, d; Kühnemann, 1938).

T. fuegiana (Mitt.) Mitt. (Mitten, 1869).

Syn. *Syntrichia fuegiana* Mitt. (Mitten, 1860).

Barbula fuegiana (Mitt.) Jaeg. (Jaeger, 1873).

Barbula magellanica C. Muell. (Mueller, 1862).

Tortula magellanica (C. Muell.) Aongstr. (Ångström, 1876b) *hom. illeg.*

Barbula patagonica C. Muell. (Mueller, 1885) *hom. illeg.*

Barbula australis Par. (Paris, 1894) *nom. nud.*

Distr. Southern South America (Mitten, 1860, 1869; Mueller, 1862, 1885, 1889; Jaeger, 1873; Ångström, 1876b; Paris, 1894, 1906; Dusén, 1903a, 1906; Cardot, 1908a; Herzog, 1923, 1957; Brotherus, 1924a; Kühnemann, 1938).

Falkland Islands (Mitten, 1860, 1869; Jaeger, 1873; Paris, 1906; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

- Antarctic, continental region (Gepp, 1902; Cardot, 1908a; Steere, 1961a).
- T. fusca** (C. Muell.) Mont. **Barbula fusca**
- T. fuscinervia** Mitt. **Barbula fuscinervia**
- T. fusco-viridis** Card. (Cardot, 1906a).
Distr. South Georgia (Cardot, 1906a, 1908a; Steere, 1961b).
- T. geniculata** Mont. **Barbula geniculata**
- T. glacialis** (C. Muell.) Mont. (Montagne, 1850).
Syn. *Barbula glacialis* Kunz. ex C. Muell (Kunze ex Mueller, 1849).
Distr. Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Paris, 1906; Costes, 1921; Brotherus, 1924a).
- T. graminicolor** (C. Muell.) Mont. **Barbula graminicolor**
- T. gracilis** (*sensu* Wilson and Hooker, 1847). see **schistidium antarctici**
- T. gromschii** Thér. (Thériot, 1934b [1935]).
Distr. Southern South America (Thériot, 1934b).
- T. grossiretis** Card. (Cardot, 1906a).
Distr. Southern South America (Cardot, 1900 and 1901 (as *Barbula ruralis* forma *fide* Cardot, 1908a), 1906a, 1908a; Thériot, 1918; Cardot and Brotherus, 1923; Brotherus, 1924a; Bartram, 1952); South Georgia (Cardot, 1906a, 1908a; Brotherus, 1924a; Steere, 1961b). Antarctic, peninsula region (Dixon, 1920; Steere, 1961a).
- var. *atrata*** Card. (Cardot, 1906a).
Distr. South Gerogia (Cardot, 1906a, 1908a; Steere, 1961b).
- T. heimii** (Hedw.) Mitt. **Pottia heimii**
- T. heteronema** Card. **Tortula heteroneura**
- T. heteroneura** Card. (Cardot, 1911d).
Syn. *Tortula heteronema* Card. (Cardot, 1911b) err.
Distr. Antarctic, peninsula region (Cardot, 1911b, d, 1913a; Brotherus, 1924a; Steere, 1961a).
- T. humilis** (Hedw.) Turn. **Tortella humilis**
- T. hyperborea** (Brid.) Mont. **Tortula mucronifolia** var. *arctica*
- T. intermedia** (Brid.) De Not. (De Notaris, 1838a).
Syn. *Syntrichia intermedia* Brid. (Bridel, 1826).
Tortula montana (Nees) Lindb. (Lindberg, 1879) *hom. illeg.*
Distr. Southern South America (Dusén, 1905d; Kühnemann, 1938).
- T. inundata** Mitt. **Didymodon inundatus**
- T. jaffuelii** Thér. (Thériot, 1923b).
Distr. Southern South America (Thériot, 1923b; Brotherus, 1925).
- T. kunzeana** (C. Muell.) Mont. (Montagne, 1850).
Syn. *Barbula kunzeana* C. Muell. (Mueller, 1843a).
Barbula kunzei C. Muell. ex Kindb. (Mueller ex Kindberg, 1888) *nom. illeg.*
Distr. Scuthern South America (Mueller, 1843a, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Paris, 1906; Williams, 1915; Costes, 1921; Thériot, 1921a; Brotherus, 1924a).
- T. laeta** (C. Muell.) Mitt. **Rhamphidium laetum**
- T. laevigata** Mitt. **Barbula laevigata**
- T. laevinervis** Broth. ex Dus. (Brotherus ex Dusén, 1907).

Distr. Southern South America (Paris, 1906; Dusén, 1907; Brotherus, 1924a).

T. laevipila (Brid.) Schwaegr. (Schwaegrichen, 1823).

Syn. *Syntrichia laevipila* Brid. (Bridel, 1819).

Barbula laevipila (Brid.) Garov. (Garovaglio, 1840).

Tortula ruralis (Hedw.) Gaertn., Meyer et Scherb. var. *laevipila* (Brid.) Hook. et Grev. (Hooker and Greville, 1824b).

Syntrichia ruralis (Hedw.) Web. et Mohr var. *laevipila* (Brid.) Spreng. (Sprengel, 1827).

Syntrichia levipila Brid. ex Moenk. (Bridel ex Mönkemeyer, 1927) err.

Distr. Falkland Islands (Wilson and Hooker, 1847 (as vars 1 and 2); Kühnemann, 1938).

Antarctic, peninsula region (Wilson and Hooker, 1847 (as vars 3 and 4, see also Cardot, 1908a, p. 241); Greene, 1968a (as vars 3 and 4)).

T. lamprocalyx (C. Muell.) Mont.

Barbula lamprocalyx

T. lechleri (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula lechleri* C. Muell. (Mueller, 1859).

Distr. Southern South America (Mueller, 1859, 1885; Jaeger, 1873; Brotherus 1902, 1924a; Paris, 1906; Cardot, 1908a; Herzog, 1923).

T. lepto-syntrichia (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula lepto-syntrichia* C. Muell. (Mueller, 1890a).

Syntrichia lepto-syntrichia C. Muell. (Mueller, 1890a) nom. nud.

Barbula anacamptophylla C. Muell. (Mueller, 1890a).

Tortula anacamptophylla (C. Muell.) Broth. (Brotherus, 1902).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Brotherus, 1902, 1924a; Cardot, 1906a, 1908a; Paris, 1906; Dixon, 1920, 1934; Steere, 1961b).

T. leucocalyx Mont.

Pseudocrossidium leucocalyx

T. lingulaefolia Card. et Broth. (Cardot and Brotherus, 1923).

Syn. *Syntrichia lingulaefolia* (Card. et Broth.) Herz. (Herzog, 1954) fide Seki, 1974.

Distr. Southern South America (Herzog, 1954; Seki, 1974).

South Georgia (Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b).

T. lithophila Dus.

Sarconeurum glaciale

T. litorea Card. (Cardot in Cardot and Brotherus, 1923).

Distr. Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

T. longipila Dus. (Dusén in Paris, 1906) nom. nud.

Distr. Southern South America (Paris, 1906).

T. lorentzii (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula lorentzii* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Paris, 1906; Hosseus, 1936, 1937, 1938a, b, d; Kühnemann, 1938).

T. macrocarpa (Schimp.) Mitt.

Pottia macrocarpa

T. magellanica Mont. (Montagne, 1850).

Syn. *Barbula magellanica* (Mont.) C. Muell. (Mueller, 1885) hom. illeg.

Distr. Southern South America (Montagne, 1850, 1856; Mueller, 1885; Paris, 1900a, 1906; Cardot, 1908a).

T. magellanica (C. Muell.) Aongstr.

Tortula fuegiana

T. marginato-serrata Par. (Paris, 1906) nom. nud.

Syn. *Barbula marginato-serrata* Dus. (Dusén, 1903a) nom. nud.

Syntrichia marginato-serrata Dus. (Dusén in Paris, 1906) nom. nud.

Distr. Southern South America (Dusén, 1903a; Paris, 1906).

T. mendozensis Mitt.

Barbula mendozensis

T. micro-runcinata Par.

Leptodontium lingicaule* var. *microruncinatum

***T. minutirostula* (C. Muell.) Broth.** (Brotherus, 1902).

Syn. *Barbula minutirostula* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Kurtz, 1904; Paris, 1906; Hosseus, 1935b, 1937; Kühnemann, 1938).

***T. mniadelphus* (C. Muell.) Broth.** (Brotherus, 1901).

Syn. *Barbula mniadelphus* C. Muell. (Mueller, 1897b).

Tortula uniadelphus C. Muell. (Mueller in Herzog, 1952c) err.

Syntrichia mniadelphus (C. Muell.) Herz. (Herzog, 1954).

Distr. Southern South America (Thériot, 1923b).

var. *cochlearifolia* (Herz.) Seki (Seki, 1974).

Syn. *Tortula unidelphus* C. Muell. var. *cochlearifolia* Herz. (Herzog, 1952c) nom. nud.

Syntrichia mniadelphus (C. Muell.) Herz. var. *cochlearifolia* Herz. (Herzog, 1954).

Distr. Southern South America (Herzog, 1952c, 1954; Seki, 1974).

T. mniooides (Schwaegr.) Mont.

Calyptopogon mniooides

T. monoica Card.

Tortula tenella

T. montana (Nees) Lindb.

Tortula intermedia

T. mucronifolia* Schwaegr. var. *arctica Hook. et Grev. (Hooker and Greville, 1824b).

Syn. *Syntrichia hyperborea* Brid. (Bridel, 1826).

Tortula hyperborea (Brid.) Mont. (Montagne, 1845b).

Barbula hyperborea Mont. ex Par. (Montagne ex Paris, 1894) err.

Distr. Southern South America (Montagne, 1839c, 1845b; Wilson and Hooker, 1847; Mueller, 1885; Kühnemann, 1938).

T. muelleri Hook. f. et Wils.

Tortula princeps

***T. muralis* Hedw.** (Hedwig, 1801).

Syn. *Bryum murale* (Hedw.) L. ex With. (Linnaeus ex Withering, 1801).

Barbula muralis (Hedw.) Crom. (Crome, 1805).

Syntrichia muralis (Hedw.) Raab (Raab, 1819).

Barbula muralis (Hedw.) Crom. var. *rupestris* Schultz (Schultz, 1823).

Tortula muralis Hedw. var. *rupestris* Chev. (Chevallier, 1827).

Barbula muralis (Hedw.) Crom. var. *australis* Hamp. (Hampe in Mueller, 1862) nom. nud.

Barbula annulus C. Muell. (Mueller, 1882).

Tortula annulus (C. Muell.) Broth. (Brotherus ex Paris, 1906).

Barbula berteroana C. Muell. (Mueller, 1862).

Tortula berteroana (C. Muell.) Broth. ex Par. (Brotherus ex Paris, 1906) hom. illeg.

Barbula berteroii C. Muell. ex Kindb. (Mueller ex Kindberg, 1888) hom. illeg.

Barbula muricola C. Muell. (Mueller, 1857).

Tortula muricola (C. Muell.) Mitt. (Mitten, 1869).

Distr. Southern South America (Montagne, 1850; Mueller, 1862, 1879a, 1882; Mitten, 1869;

Jaeger, 1873; Brotherus, 1900, 1924a, c; Krieger, 1904; Dusén, 1906; Paris, 1906;

Williams, 1915; Costes, 1921; Thériot, 1921a, b; Herter, 1933a, 1939a, b; Roivainen,

1934; Herzog, 1938, 1939; Hosseus, 1938c, d, 1940; Espinosa B., 1941; Piovano,

1954).

var. *aestiva* Hedw. (Hedwig, 1801).

Syn. *Tortula aestiva* (Hedw.) P. Beauv. (Palisot de Beauvois, 1805a).

Barbula muralis (Hedw.) Crom. var. *aestiva* (Hedw.) Roehl. (Röhling, 1813).

Barbula aestiva (Hedw.) Schultz (Schultz, 1823).

Barbula muralis (Hedw.) Crom. ssp. *aestiva* (Hedw.) Kindb. (Kindberg, 1897).

Distr. Southern South America (Thériot, 1921a).

var. *longipila* Dus. (Dusén, 1907).

Distr. Southern South America (Dusén, 1907; Kühnemann, 1938).

var. *rupestris* Chev.

Tortula muralis

T. muricola (C. Muell.) Mitt.

Tortula muralis

T. obscuriretis Thér. (Thériot, 1918).

Distr. Southern South America (Thériot, 1918; Campo, 1921; Brotherus, 1925).

T. occultilimbata Card.

Tortula platyphylla

T. papillosa Wils. (Wilson in Spruce, 1845).

Syn. ***Syntrichia papillosa*** (Wils.) Jur. (Juratzka, 1882).

Distr. Southern South America (Dusén, 1905b, 1906; Cardot, 1908a; Thériot, 1921a; Kühnemann, 1938; Herzog, 1954).

Falkland Islands (Paris, 1906; Cardot, 1908a; Brotherus, 1924a; Kühnemann, 1938).

var. *chilensis* Thér. (Thériot, 1921a).

Syn. ***Tortula subpapillosa*** Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Thériot, 1921a, 1930; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

T. patagonica Mitt.

Didymodon patagonicus

T. perangusta Dus. (Dusén, 1905d) *nom. nud.*

Distr. Southern South America (Dusén, 1905d; Kühnemann, 1938).

T. perarmata Broth. (Brotherus, 1924c).

Distr. Southern South America (Brotherus, 1924c, 1925; Espinosa B., 1941; Skottsberg, 1950).

T. percarnosa (C. Muell.) Broth. (Brotherus, 1902).

Syn. ***Barbula percarnosa*** C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Paris, 1906; Kühnemann, 1938).

T. perflaccida Broth. ex Dus.

Tortula flagellaris

T. perpusilla (C. Muell.) Broth. (Brotherus, 1902).

Syn. ***Barbula perpusilla*** C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Kurtz, 1904; Paris, 1906; Hosseus, 1935a; Kühnemann, 1938).

T. perrubiginosa (Dus.) Par.

Tortula anderssonii

T. perrufula (C. Muell.) Williams

Tortella perrufula

T. pilifera Hook. (Hooker, 1818).

Syn. ***Barbula pilifera*** (Hook.) Brid. (Bridel, 1826) *hom. illeg.*

Streblotrichum piliferum (Hook.) Hilp. (Hilpert, 1933).

Barbula crinita Schultz (Schultz, 1823) *nom. illeg.*

Barbula depressa Sull. (Sullivant, 1859).

Tortula subglacialis Thér. (Thériot, 1917b).

Distr. Juan Fernandez (Thériot, 1917b; Brotherus, 1924b).

Southern South America (Mueller, 1849; Montagne, 1850; Sullivant, 1859; Paris, 1904a; Thériot, 1917b, 1918, 1921a, b, 1923b, 1925b, 1928, 1934b; Campo, 1921; Costes, 1921; Cardot and Brotherus, 1923; Brotherus, 1924a; Reimers, 1926; Herzog, 1938, 1954, 1957, 1960; Bartram, 1952; Seki, 1974).

var. *denticulata* (Dus.) Wijk et Marg. (van der Wijk and Margadant, 1959).

Syn. ***Barbula flagellaris*** Schimp. var. ***denticulata*** Dus. (Dusén, 1906).

Barbula depressa Sull. var. ***denticulata*** (Dus.) Thér. (Thériot, 1917a).

Distr. Southern South America (Dusén, 1906; Cardot, 1908a; Thériot, 1917a; Campo, 1921; Kühnemann, 1938).

var. *gracilis* (Dus.) Wijk et Marg. (van der Wijk and Margadant, 1959).

Syn. *Barbula flagellaris* Schimp. var. *gracilis* Dus. (Dusén, 1906).
Barbula depressa Sull. var. *gracilis* (Dus.) Thér. (Thériot, 1917a).

Distr. Southern South America (Dusén, 1906; Thériot, 1917a; Kühnemann, 1938).

var. *oliviensis* (Card.) Wijk et Marg. (van der Wijk and Margadant, 1959).

Syn. *Barbula oliviensis* Card. (Cardot, 1905a).
Barbula depressa Sull. var. *oliviensis* (Card.) Thér. (Thériot, 1917a).

Distr. Southern South America (Cardot, 1905a, 1908a; Thériot, 1917a; Cardot and Brotherus, 1923; Kühnemann, 1938).

T. *planicosta* Herz. (Herzog, 1952b).

Distr. Southern South America (Herzog, 1952b).

T. *platyphylla* Mitt. (Mitten, 1869).

Syn. *Desmatodon amblyophyllum* Mont. (Montagne, 1845a).
Trichostomum amblyophyllum (Mont.) C. Muell. (Mueller, 1849).
Tortula amblyophylla (Mont.) Thér. (Thériot, 1923a) *hom. illeg.*
Tortula occultilimbata Card. (Cardot in Thériot, 1921b) *nom. nud.*

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Mitten, 1869; Jaeger, 1873; Paris, 1906; Thériot, 1921b, 1922, 1923a, 1934b; Brotherus, 1924a; Looser, 1932; Herzog, 1939, 1954).

T. *podocarpi* (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula podocarpi* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Paris, 1906; Kühnemann, 1938).

T. *poeppigiana* (C. Muell.) Mont.

Trichostomopsis australasiae

T. *polycarpa* Dus. (Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938; Seki, 1974).

T. *princeps* De Not. (De Notaris, 1838a).

Syn. *Barbula princeps* (De Not.) C. Muell. (Mueller, 1849).
Syntrichia princeps (De Not.) Mitt. (Mitten, 1859).
Barbula muelleri B.S.G. (Bruch and others, 1842) *nom. illeg.*
Syntrichia muelleri Bruch (Bruch in Bruch and others, 1842) *nom. nud.*
Tortula muelleri Hook. f. et Wils. (Hooker and Wilson in Wilson and Hooker, 1847)
nom. illeg.
Barbula antarctica Hamp. (Hampe in Mueller, 1849).
Tortula antarctica (Hamp.) Wils. (Wilson, 1859).

Distr. Southern South America (Mitten, 1869; Jaeger, 1873; Paris, 1906; Brotherus, 1925; Seki, 1974).

Falkland Islands (Wilson and Hooker, 1847; Mueller, 1849; Jaeger, 1873; Paris, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).

T. *prostrata* Mont. (Montagne, 1845a).

Syn. *Barbula prostrata* (Mont.) Jaeg. (Jaeger, 1873).
Syntrichia prostrata (Mont.) Herz. (Herzog, 1960 [1961]).
Barbula mnioides Schwaege. var. *prostrata* (Mont.) C. Muell. (Mueller, 1849).
Streptopogon mnioides (Schwaege.) Mitt. var. *prostrata* (Mont.) Jaeg. (Jaeger, 1873).
Calyptopogon mnioides (Schwaege.) Broth. var. *prostratus* (Mont.) Par. (Paris, 1904a).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Mitten, 1869; Jaeger, 1873; Dusén, 1906; Paris, 1906; Thériot, 1915, 1928; Brotherus, 1924a; Herzog, 1938, 1960; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

T. *pseudo-latifolia* Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Dusén, 1906; Brotherus, 1925; Kühnemann, 1938; Seki, 1974).

T. pseudo-robusta Dus. (Dusén, 1906).

Syn. *Syntrichia pseudo-robusta* (Dus.) Herz. (Herzog, 1954).
Syntrichia robusta Dus. (Dusén *in Cardot*, 1908a) *nom. nud.*
Barbula ventanae C. Muell. (Mueller *in Thériot*, 1932 [1933]) *nom. nud.*

Distr. Southern South America (Dusén, 1906; Cardot, 1908a; Thériot, 1921a, 1932; Cardot and Brotherus, 1923; Herzog, 1938, 1939, 1954; Kühnemann, 1938; Seki, 1974).

T. pulvinatula Dus. (Dusén, 1905b).

Distr. Southern South America (Dusén, 1905b, 1906; Brotherus, 1924a; Kühnemann, 1938).

T. punctulata (Brid.) Mitt.**Tortula robusta****T. purpureo-velutina** Herz. (Herzog, 1954).

Distr. Southern South America (Herzog, 1954).

T. pusilla Mitt.**Pterygoneurum ovatum****T. pusilla** Aongstr.**Tortula brachypelma****T. pygmaea** Dus.**Sarconeurum glaciale****T. replicata** (Tayl.) Wils.**Barbula replicata****T. rigidula** (Hedw.) Lindb.**Didymodon rigidulus****T. rivularis** Dus. (Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

T. robusta Hook. et Grev. (Hooker and Greville, 1824b).

Syn. *Barbula robusta* (Hook. et Grev.) Brid. (Bridel, 1827a).
Syntrichia robusta (Hook. et Grev.) Herz. (Herzog, 1954).
Barbula speciosa Hook. f. et Wils. (Hooker, J. D. and Wilson, 1844).
Mnium punctulatum Brid. (Bridel, 1827b).
Tortula punctulata (Brid.) Mitt. (Mitten, 1869).
Barbula punctulata (Brid.) Jaeg. (Jaeger, 1873).
Tortula serrulata Hook. et Grev. (Hooker and Greville, 1824b) *fide* Greene, 1964.
Barbula serrulata (Hook. et Grev.) Brid. (Bridel, 1827a) *fide* Greene, 1964.
Desmatodon serrulatus (Hook. et Grev.) Mitt. (Mitten, 1859) *fide* Greene, 1964.
Tortula rubra Mitt. (Mitten *in Hooker*, 1867) *fide* Greene, 1964.
Barbula rubra (Mitt.) Jaeg. (Jaeger, 1873) *fide* Greene, 1964.
Barbula dioica (R. Brown ter.) Par. (Paris, 1900a).
Desmatodon ruber (Mitt.) Hamp. (Hampe, 1880c).

Distr. Southern South America (Hooker and Greville, 1824b; Bridel, 1827a; Hooker, J. D. and Wilson, 1844; Wilson and Hooker, 1847; Mueller, 1849, 1885; Mitten, 1869; Jaeger, 1873; Bescherelle, 1889; Paris, 1900a, 1906; Cardot, 1905a, 1908a; Dusén, 1906; Cardot and Brotherus, 1923; Brotherus, 1924a; Roivainen, 1934; Kühnemann, 1938; Herzog, 1939, 1940, 1954; Bartram, 1952; Seki, 1974).

Falkland Islands (Bridel, 1827a; Wilson and Hooker, 1847 (*as T. robusta* var. β); Mueller, 1849, 1885; Hooker, 1867; Mitten, 1869; Jaeger, 1873; Bescherelle, 1889; Paris, 1900a, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

South Georgia (Cardot, 1906a, 1908a; Dixon, 1920, 1932, 1934; Cardot and Brotherus, 1923; Brotherus, 1924a; Steere, 1961b; Greene, 1964).

var. laxa Bartr. (Bartram, 1946).

Distr. Southern South America (Roivainen, 1934; Bartram, 1946).

var. runcinata (C. Muell.) Broth. (Brotherus *in Cardot and Brotherus*, 1923).

Syn. *Barbula runcinata* C. Muell. (Mueller, 1890a).
Syntrichia runcinata C. Muell. (Mueller, 1890a) *nom. nud.*
Tortula runcinata (C. Muell.) Broth. (Brotherus, 1902).

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

Falkland Islands (Cardot and Brotherus, 1923; Kühnemann, 1938).

South Georgia (Mueller, 1890a; Brotherus, 1902, 1924a; Paris, 1906).

T. robustula Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Dusén, 1906; Brotherus, 1924a; Kühnemann, 1938).
 Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).

T. rubra Mitt. **Tortula robusta**

T. runcinata (C. Muell.) Broth. **Tortula robusta** var. **runcinata**

T. ruralis (Hedw.) Gaertn., Meyer et Scherb. (Gärtner and others, 1802).

Syn. *Barbula ruralis* Hedw. (Hedwig, 1801).
Bryum rurale (Hedw.) L. ex With. (Linnaeus ex Withering, 1801).
Syntrichia ruralis (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Tortula arvensis Mont. (Montagne, 1850) *err.*

Distr. Southern South America (Jaeger, 1873; Dusén, 1905a, 1906; Paris, 1906; Brotherus, 1924a; Roivainen, 1934; Kühnemann, 1938; Bartram, 1965).

var. *australis* Mont. **Tortula flagellaris**

var. *laevipila* (Brid.) Hook. et Grev. **Tortula laevipila**

T. santiagensis Broth. (Brotherus, 1924c).

Distr. Southern South America (Brotherus, 1924c; Espinosa B., 1941).

T. saxicola Card. (Cardot, 1905a).

Distr. Southern South America (Cardot, 1905a, 1908a; Kühnemann, 1938; Bizot, 1967; Seki, 1974).

T. saxicola Dus. **Sarconeurum glaciale**

T. scabrella Dus. (Dusén, 1907).

Syn. *Barbula scabrella* Dus. (Dusén in Paris, 1906) *nom. nud.*
Syntrichia scabrella (Dus.) Herz. (Herzog, 1957).

Distr. Southern South America (Paris, 1906; Dusén, 1907; Thériot, 1921b, 1923b; Brotherus, 1924a; Herzog, 1957).

T. scabrinervis (C. Muell.) Mont. (Montagne, 1850).

Syn. *Barbula scabrinervis* C. Muell. (Mueller, 1849).
Syntrichia scabrinervis (C. Muell.) Herz. (Herzog, 1954).

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Robinson, 1975).
 Southern South America (Mueller, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Paris, 1906; Costes, 1921; Herzog, 1954).

T. schnyderi (C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula schnyderi* C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Brotherus, 1902, 1924a; Dusén, 1906; Paris, 1906; Cardot, 1908a; Kühnemann, 1938).

T. serrata Dix. (Dixon, 1923).

Distr. South Gerogia (Greene, 1964, 1968a).

T. serripungens (Lor. et C. Muell.) Broth. (Brotherus, 1902).

Syn. *Barbula serripungens* Lor. et C. Muell. (Lorentz and Mueller in Mueller, 1879a).
Syntrichia serripungens (Lor. et C. Muell.) Herz. (Herzog, 1954).

Distr. Southern South America (Mueller, 1879a; Brotherus, 1902, 1924a; Kurtz, 1904; Paris, 1906; Hosseus, 1935b, 1937, 1938b, c, d; Kühnemann, 1938; Herzog, 1952a, 1954).

var. *exesa* (C. Muell.) Herz. (Herzog, 1916a).

Syn. *Barbula serripungens* Lor. et C. Muell. var. *exesa* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kühnemann, 1938).

- T. serrulata* Hook. et Grev. **Tortula robusta**
- T. sinuata* Bartr. (Bartram, 1965).
Distr. Southern South America (Bartram, 1965).
- T. socialis* Dus. (Dusén, 1907).
Distr. Southern South America (Dusén, 1907; Brotherus, 1924a).
- T. squarripila* Thér. (Thériot, 1917b).
Syn. *Syntrichia squarripila* (Thér.) Herz. (Herzog, 1954).
Distr. Southern South America (Thériot, 1917b; Costes, 1921; Brotherus, 1924a; Bartram, 1952; Herzog, 1954, 1960).
- T. stenophylla* Card. et Broth. (Cardot and Brotherus, 1923) *hom. illeg.*
Distr. Southern South America (Cardot and Brotherus, 1923; Brotherus, 1924a; Thériot, 1935a; Kühnemann, 1938).
- T. subenervis* Dus. (Dusén in Paris, 1906) *nom. nud.*
Distr. Southern South America (Paris, 1906).
- T. subglacialis* Thér. **Tortula pilifera**
- T. subpapillosa* Card. et Broth. **Tortula papillosa** var. **chilensis**
- T. subsessilis* (Brid.) Mitt. **Pterygoneurum subsessile**
- T. tenella* Broth. (Brotherus, 1898).
Syn. *Barbula tenella* (Broth.) Par. (Paris, 1900a).
Tortula monoica Card. (Cardot, 1905a).
Distr. Falkland Islands (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Kühnemann, 1938).
South Georgia (Cardot, 1906a, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Dixon, 1934; Steere, 1961b).
Antarctic, peninsula region (Cardot, 1906a, 1908a; Brotherus, 1925; Steere, 1961a).
- T. tortuosa* Hedw. **Tortella tortuosa**
- T. truncata* (Hedw.) Mitt. **Pottia truncata**
- T. truncatula* (With.) Lindb. **Pottia truncata**
- T. umbrosa* Dus. (Dusén, 1907).
Distr. Southern South America (Dusén, 1907; Brotherus, 1924a).
- T. unguiculata* (Hedw.) A. Roth. ex P. Beauv. **Barbula unguiculata**
- T. uniadelphus* C. Muell. **Tortula mniadelphus**
var. *cochlearifolia* Herz. **Tortula mniadelphus** var. **cochlearifolia**
- T. vahliana* (Schultz) Mont. (Montagne, 1850).
Syn. *Barbula vahliana* Schultz (Schultz, 1823).
Distr. Southern South America (Bruch and others, 1842; Montagne, 1850; Reichardt, 1870).
- T. vinealis* (Brid.) Spruc. var. *flaccida* (B.S.G.) Spruc. **Barbula vinealis** var. **flaccida**
- Trachyloma**
T. subbasilare (Hook.) Mitt. **Goniobryum subbasilare**
- Trachypus**
T. flexicaulis (Wils.) Mitt. **Papillaria flexicaulis**
- Trematodon**
T. affinis B.S.G. **Trematodon reflexus**
T. boryanus (Schwaegr.) Spreng. **Dicranoloma billardieri**

T. crispatissimus Hamp.

Trematodon longicollis var. *crispatissimus*

T. crispifolius Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933b).

T. felipponei Thér. (Thériot in Felippone, 1929 [1930]).

Distr. Southern South America (Felippone, 1929; Herter, 1933b).

T. latifolius C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

T. longicollis Michx. var. *crispatissimus* Hornsch. (Hornschuch, 1840) *hom. illeg.*

Syn. *Trematodon crispatissimus* Hamp. (Hampe, 1879) *hom. illeg.*

Distr. Southern South America (Hornschuch, 1840; Gibert, 1873; Hampe, 1879; Paris, 1906; Roth, 1911; Herter, 1933a).

T. reflexus C. Muell. (Mueller, 1848a).

Syn. *Trematodon affinis* B.S.G. (Bruch and others, 1846) *nom. nud.*

Distr. Southern South America (Krieger, 1904; Bartram, 1965).

T. uruguensis Broth. (Brotherus in Felippone, 1909).

Distr. Southern South America (Felippone, 1909; Brotherus, 1924a; Herter, 1933a).

T. vaginatus C. Muell. (Mueller, 1857).

Distr. Southern South America (Krieger, 1904).

Trichosteleum

T. micropyxidium (C. Muell.) Broth. (Brotherus, 1908).

Syn. *Aptychus micropyxis* C. Muell. (Mueller, 1896b).

Distr. Southern South America (Hosseus, 1935b; Kühnemann, 1938).

Trichosteleum sp.

Syn. *Hypnum micropyxis* C. Muell. (Mueller, 1879a).

Rapidostegium micropyxis (C. Muell.) Kindb. (Kindberg, 1888).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1905b; Hosseus, 1937).

Trichostomopsis

*T. australasiae** (Hook. et Grev.) H. Robinson (Robinson, 1970a).

Syn. *Tortula australasiae* Hook. et Grev. (Hooker and Greville, 1824b).

Barbula australasiae (Hook. et Grev.) Brid. (Bridel, 1827a).

Trichostomum australasiae (Hook. et Grev.) Jaeg. (Jaeger, 1873).

Asteriscium australasiae (Hook. et Grev.) Hilp. (Hilpert, 1933).

Barbula poeppigiana C. Muell. (Mueller, 1843a).

Pottia poeppigiana (C. Muell.) C. Muell. (Mueller, 1849).

Tortula poeppigiana (C. Muell.) Mont. (Montagne, 1850).

Weisia poeppigiana (C. Muell.) Mitt. (Mitten, 1869).

Hyophila poeppigiana (C. Muell.) Jaeg. (Jaeger, 1873).

Asteriscium poeppigianum (C. Muell.) Hilp. (Hilpert, 1933).

Hyophila poeppigii C. Muell. ex Kindb. (Mueller ex Kindberg, 1888) *nom. illeg.*

Didymodon subtophaceus Williams (Williams, 1903a) *fide* Robinson, 1970a.

Desmatodon subtophaceus (Williams) Williams (Williams, 1915) *fide* Robinson, 1970a.

Asteriscium subtophaceum (Williams) Hilp. (Hilpert, 1933).

Distr. Southern South America (Mueller, 1843a, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1873; Paris, 1904a, b; Dusén, 1906; Williams, 1915; Warnstorff, 1916; Thériot, 1917b, 1921b, 1928; Costes, 1921; Brotherus, 1924a, c; Herzog, 1938, 1954, 1957; Espinosa B., 1941; Skottsberg, 1950; Robinson, 1970a).

* = *Didymodon australasii* (Hook. et Grev.) Zander *fide* Zander, 1978.

T. curvipes (C. Muell.) H. Robinson (Robinson, 1970a).

Syn. *Barbula curvipes* C. Muell. (Mueller, 1879a).

Asteriscium curvipes (C. Muell.) Hilp. (Hilpert, 1933).

Distr. Southern South America (Mueller, 1879a; Paris, 1904a; Brotherus, 1924a; Kühnemann, 1938; Robinson, 1970a).

T. umbrosat† (C. Muell.) H. Robinson (Robinson, 1970a).

Syn. *Barbula umbrosa* C. Muell. (Mueller, 1879a).

Asteriscium umbrorum (C. Muell.) Herz. (Herzog, 1952a).

Asteriscium flavidum Herz. (Herzog, 1952a).

Barbula flaviseta (Herz.) Wijk et Marg. (van der Wijk and Margadant, 1958).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1904a; Brotherus, 1924a; Hosseus, 1935a, b, 1937; Kühnemann, 1938; Herzog, 1952a; Robinson, 1970a).

Trichostomum

T. acaulon C. Muell.

Tortella acaulon

T. aciculare (Hedw.) P. Beauv. var. *aquaticum* (Schrad.) Web. et Mohr ex Brid.

Racomitrium aquaticum

T. aequatoriale Spruc. ex Dix. (Spruce ex Dixon, 1924a).

Syn. *Tortula aequatoriale* Spruc. (Spruce in Dixon, 1924a) nom. nud.

Tortula chilensis Mitt. (Mitten, 1869 non *Trichostomum chilense* Mont.).

Distr. Southern South America (Mitten, 1869).

T. affine (C. Muell.) Mont.

Ditrichum difficile

T. amblyophyllum (Mont.) C. Muell.

Tortula platyphylla

T. aquaticum Brid. ex Schrad.

Racomitrium aquaticum

T. australasiae (Hook. et Grev.) Jaeg.

Trichostomopsis australasiae

T. australe Mitt.

Ditrichum strictum

T. brachydontium Bruch (Bruch in F. A. Mueller, 1829).

Syn. *Didymodon brachydontius* (Bruch) Wils. (Wilson in Hooker, 1833).

Mollia brachydonta (Bruch) Lindb. (Lindberg, 1879).

Tortella brachydonta (Bruch) C. Jens. (Jensen, 1923).

Distr. Juan Fernandez (Brotherus, 1924b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

T. brunneum C. Muell.

Didymodon brunneus

T. calcareum (Nees et Hornsch.) Lindb.

Gymnostomum aeruginosum

T. campylocarpum C. Muell.

Bryoerthrophyllum campylocarpum

T. capillaceum (Hedw.) Sm.

Distichium capillaceum

T. chilense Mont.

Cheilotrichia chilensis

T. compactulum C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Kurtz, 1904; Paris, 1906; Brotherus, 1924a; Hosseus, 1935b, 1937, 1938c, d; Kühnemann, 1938).

T. convolutum Brid.

Desmatodon convolutus

T. crispipilum Tayl.

Racomitrium crispipilum

T. cylindricarpum (C. Muell.) Wils.

Ditrichum cylindricarpum

T. difficile Dub.

Ditrichum difficile

T. dusenii Par. (Paris, 1906) nom. nud.

Syn. *Trichostomum glauco-viride* Dus. (Dusén in Paris, 1906) nom. nud.

Distr. Southern South America (Paris, 1906).

† = *Didymodon umbrosus* (C. Muell.) Zander fide Zander, 1978.

T. elliotii Broth. ex Dus. (Brotherus ex Dusén, 1906).

Distr. Southern South America (Dusén, 1906; Brotherus, 1924a; Herzog, 1954).

T. elongatum Hook. f. et Wils.

Ditrichum cylindricarpum

T. fernandezianum Card. (Cardot in Skottsberg, 1914) *nom. nud.*

Distr. Southern South America (Skottsberg, 1914).

T. glauco-viride C. Muell.

Didymodon glauco-viridis

T. glauco-viride Dus.

Trichostomum dusenii

T. gracillimum C. Muell. (Mueller, 1879a).

Syn. *Anacalypta gracillima* (C. Muell.) C. Muell. (Mueller, 1882) *nom. inval.*

Pottia gracillima (C. Muell.) Par. (Paris, 1900a).

Distr. Southern South America (Mueller, 1879a; Paris, 1900a, 1906; Kurtz, 1904; Warnstorff, 1916; Brotherus, 1924a; Hosseus, 1935b, 1937; Kühnemann, 1938).

T. gymnum C. Muell.

Didymodon gymnos

T. heterostichum Hedw.

Racomitrium heterostichum

T. humile (Hedw.) Mach.

Tortella humilis

T. hypnoides Willd. ex P. Beauv.

Racomitrium lanuginosum

T. imperfectum C. Muell.

Barbula imperfecta

T. julaceum Dus. (Dusén in Paris, 1906) *nom. nud.*

Distr. Southern South America (Paris, 1906).

T. laetum Kunz. ex C. Muell.

Rhamphidium laetum

T. lambii Bartr. (Bartram, 1965).

Distr. Southern South America (Bartram, 1965).

T. lanuginosum Hedw.

Racomitrium lanuginosum

T. laxifolium Hook. f. et Wils.

Ditrichum difficile

T. leptodum (Mont.) Mitt.

Campylopus clavatus

T. ligulum C. Muell. (Mueller, 1901a [1900]) *nom. nud.*

Distr. Southern South America (Mueller, 1901a).

T. longifolium Brid.

Leucoloma longifolium

T. microthecium C. Muell.

Didymodon microthecius

T. nudum C. Muell.

Didymodon brunneus

T. orthotrichoides C. Muell.

Leptodontium orthotrichoides

T. plicatulum C. Muell. (Mueller, 1882).

Distr. Southern South America (Mueller, 1882; Paris, 1906; Kühnemann, 1938).

T. protensum Braun

Racomitrium aquaticum

T. pungens (Mitt.) Par.

Leptodontium pungens

T. rigidulum (Hedw.) Turn.

Didymodon rigidulus

T. schimperi Mont.

Didymodon schimperi

T. spathulato-lineare C. Muell.

Didymodon spathulato-linearis

T. strictum Web. et Mohr

Ditrichum rufescens

T. tortella C. Muell.

Tortella sp.

T. tortuosum (Hedw.) Dix.

Tortella tortuosa

T. tucumanense Bartr. (Bartram, 1965).

Distr. Southern South America (Bartram, 1965).

T. umbrosum C. Muell.

Timmiella umbrosa

Triquetrella***T. filicaulis*** Dus. (Dusén, 1906).*Distr.* Southern South America (Dusén, 1906; Brotherus, 1924a, c; Espinosa B., 1941).***T. patagonica*** C. Muell. (Mueller, 1897e).*Distr.* Southern South America (Mueller, 1897e; Paris, 1900a, 1906; Thériot, 1917b; Costes, 1921; Brotherus, 1924a; Herzog, 1938; Kühnemann, 1938).***Tristichiopsis******T. mirabilis*** C. Muell.***Tristichium mirabile******Tristichium******T. lorentzii*** C. Muell. (Mueller, 1879a).*Distr.* Southern South America (Mueller, 1879a; Paris, 1906; Roth, 1911; Brotherus, 1924a; Kühnemann, 1938).***T. mirabile*** (C. Muell.) Herz. (Herzog, 1914).*Syn.* ***Tristichiopsis mirabilis*** C. Muell. (Mueller, 1882).*Distr.* Southern South America (Mueller, 1882; Paris, 1906; Brotherus, 1924a; Kühnemann, 1938).***Ulea hom. illeg.******U. nitida*** Thér. (Thériot in Felippone, 1929 [1930]).*Distr.* Southern South America (Felippone, 1929; Herter, 1933a).***Ulota******U. anderssoni*** (Aongstr.) Jaeg.***Orthotrichum leiothecium******U. angustissima*** C. Muell. (Mueller, 1897a).*Distr.* Southern South America (Mueller, 1897a; Paris, 1906; Brotherus, 1925).***U. aurantiaca*** Dus. in Malta (Dusén in Malta, 1927).*Distr.* Southern South America (Malta, 1927).***U. brevicollis*** (Mitt.) Jaeg.***Ulota rufula******U. carinata*** Mitt. (Mitten, 1860).*Syn.* ***Orthotrichum carinatum*** (Mitt.) Mitt. (Mitten, 1869).*Distr.* Southern South America (Mitten, 1860, 1869; Paris, 1906; Brotherus, 1925; Malta, 1927; Herzog, 1938, 1939).***U. chilensis*** (Mitt.) Jaeg.***Ulota rufula******U. crassifolia*** (Hook. f. et Wils.) Hook. f.***Muelleriella crassifolia******U. crenato-erosa*** (C. Muell.) Kindb.***Orthotrichum crenato-erosa******U. crispata*** (Hedw.) Sw.***Ptychomitrium crispatum******U. darwinii*** Mitt.***Ulota fuegiana* ssp. *darwinii******U. eremitensis*** Mitt.***Ulota glabella******U. fernandeziana*** Malta (Malta, 1927).*Distr.* Juan Fernandez (Malta, 1927; Bartram, 1959; Robinson, 1975).***U. fuegiana*** Mitt. (Mitten, 1860).*Syn.* ***Orthotrichum fuegianum*** (Mitt.) Mitt. (Mitten, 1869).*Orthotrichum luteolum* Hook. f. et Wils. (Hooker and Wilson in Wilson and Hooker, 1847) pro parte.

Distr. Southern South America (Wilson and Hooker, 1847; Mitten, 1860, 1869; Ångström, 1872; Jaeger, 1874; Bescherelle, 1885a, 1889; Mueller, 1885; Cardot, 1900, 1901, 1905a, 1908a; Paris, 1906; Cardot and Brotherus, 1923; Brotherus, 1925; Malta, 1927; Kühnemann, 1938; Bartram, 1952; Herzog, 1957; Seki, 1974).

ssp. *darwinii* (Mitt.) Malta (Malta, 1927).

Syn. *Ulota darwinii* Mitt. (Mitten, 1860).

Orthotrichum darwinii (Mitt.) Mitt. (Mitten, 1869).

Distr. Southern South America (Mitten, 1860, 1869; Jaeger, 1874; Bescherelle, 1885a; Mueller, 1885; Paris, 1906; Cardot, 1908a; Brotherus, 1925; Malta, 1927; Kühnemann, 1938).

var. *crispata* Malta (Malta, 1927).

Distr. Southern South America (Malta, 1927; Bartram, 1952 (as var. *crispa* Malta); Herzog, 1954).

U. fulvella Mitt.

Ulota luteola

U. germana (Mont.) Mitt. (Mitten, 1860).

Syn. *Orthotrichum germanum* Mont. (Montagne, 1845a).

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Mitten, 1869; Jaeger, 1874; Paris, 1906; Cardot, 1908a; Herzog, 1923, 1954; Brotherus, 1925; Reimers, 1926; Malta, 1927; Thériot, 1935a; Kühnemann, 1938; Seki, 1974).

var. *brevisetata* Malta (Malta, 1927).

Distr. Southern South America (Malta, 1927; Herzog, 1938; Seki, 1974).

U. glabella Mitt. (Mitten, 1860).

Syn. *Orthotrichum glabellum* (Mitt.) Mitt. (Mitten, 1869).

Orthotrichum luteolum Hook. f. et Wils. var. β (Hooker and Wilson in Wilson and Hooker, 1847) pro parte.

Ulota eremitensis Mitt. (Mitten, 1860) *fide* Malta, 1927.

Orthotrichum eremitense Mitt. (Mitten, 1869) *fide* Malta, 1927.

Ulota hermitei Besch. (Bescherelle, 1889) *nom. illeg.*

Distr. Southern South America (Wilson and Hooker, 1847; Mitten, 1860, 1869; Jaeger, 1874; Bescherelle, 1885a, 1889; Mueller, 1885; Paris, 1906; Cardot, 1908a; Brotherus, 1925; Malta, 1927; Kühnemann, 1938; Seki, 1974).

U. gymnomitria C. Muell.

Orthotrichum gymnomitrium

U. hamata Dus.

Ulota magellanica

U. hermitei Besch.

Ulota glabella

U. immarginata Card.

Ulota pygmaeothecia

U. incana (C. Muell.) Kindb.

Orthotrichum incanum

U. inclinata (C. Muell.) Kindb.

Orthotrichum inclinatum

U. lativentrosa C. Muell. ex Malta (Mueller ex Malta, 1927).

Distr. Southern South America (Neger, 1899 (as *Ulota latavenosa*); Malta, 1927; Herzog, 1938).

U. leiothecia (C. Muell.) Jaeg.

Orthotrichum leiothecium

U. lobbiania Mitt. (Mitten, 1860).

Syn. *Orthotrichum lobbianum* (Mitt.) Mitt. (Mitten, 1869).

Ulota lobpii Mitt. ex Kindb. (Mitten ex Kindberg, 1888) *nom. illeg.*

Distr. Southern South America (Mitten, 1860, 1869; Jaeger, 1874; Paris, 1906; Cardot, 1908a; Brotherus, 1925; Malta, 1927).

U. lobpii Mitt. ex Kindb.

Ulota lobbiania

U. luteola (Hook. f. et Wils.) Wijk et Marg. (van der Wijk and Margadant, 1961).

Syn. *Orthotrichum luteolum* Hook. f. et Wils. (Hooker and Wilson in Wilson and Hooker, 1847) pro parte.

Ulota fulvella Mitt. (Mitten, 1860) *nom. illeg.*
Orthotrichum fulvellum Mitt. (Mitten, 1869) *nom. illeg.*
Ulota savatieri Besch. (Bescherelle, 1885a).

Distr. Southern South America (Wilson and Hooker, 1847; Sullivant, 1859; Mitten, 1860, 1869; Jaeger, 1874; Bescherelle, 1885a, 1889; Mueller, 1885; Eaton, 1892; Cardot, 1900, 1901, 1908a; Dusén, 1903a, 1905d; Paris, 1906; Cardot and Brotherus, 1923; Brotherus, 1925; Malta, 1927; Kühnemann, 1938; Herzog, 1957).

***U. macrocalycina* Mitt. (Mitten, 1860).**

Syn. *Orthotrichum macrocalycinum* (Mitt.) Mitt. (Mitten, 1869).
Orthotrichum marginatum Aongstr. (Ångström, 1872) *fide* Malta, 1927.
Ulota marginata (Aongstr.) Jaeg. (Jaeger in Jaeger and Sauerbeck, 1879b) *fide* Malta, 1927.
Ulota nothofagi Card. (Cardot, 1905a) *fide* Malta, 1927.

Distr. Southern South America (Mitten, 1860, 1869; Ångström, 1872, 1876b (as *U. fuegiana* Mitt.); Jaeger, 1874; Bescherelle, 1885a; Mueller, 1885; Cardot, 1905a, 1908a; Paris, 1906; Brotherus, 1925; Malta, 1927; Kühnemann, 1938; Seki, 1974).

***U. macrodontia* Dus. ex Malta (Dusén ex Malta, 1927).**

Distr. Southern South America (Dusén, 1903a; Malta, 1927; Herzog, 1954; Seki, 1974).

***U. magellanica* (Mont.) Jaeg. (Jaeger, 1874).**

Syn. *Orthotrichum magellanicum* Mont. (Montagne, 1843).
Ulota hamata Dus. (Dusén, 1903a).

Distr. Southern South America (Montagne, 1843, 1845b, 1850, 1856; Mueller, 1849, 1885; Sullivant, 1859; Mitten, 1869; Ångström, 1872; Jaeger, 1874; Bescherelle, 1885a; Dusén, 1903a, 1905d; Cardot, 1905a, 1908a; Paris, 1906; Brotherus, 1925; Malta, 1927; Herzog, 1938; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

U. marginata (Aongstr.) Jaeg.

Ulota macrocalycina

U. nothofagi Card.

Ulota macrocalycina

U. persubulata Dus.

Ulota rufula

***U. phyllantha* Brid. (Bridel, 1819).**

Syn. *Orthotrichum phyllanthum* (Brid.) Steud. (Steudel, 1824).
Weisia phyllantha (Brid.) Lindb. (Lindberg, 1879).
Orthotrichum jutlandicum Brid. (Bridel, 1826) *nom. illeg.*

Distr. Southern South America (Wilson and Hooker, 1847 (as *Orthotrichum crispum* *fide* Malta, 1927); Sullivant, 1859; Mitten, 1869; Bescherelle, 1885a; Mueller, 1885; Paris, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Malta, 1927; Bartram, 1952).

***U. pusilla* Malta (Malta, 1927).**

Distr. Southern South America (Malta, 1927).

***U. pycnophylla* Dus. ex Malta (Dusén ex Malta, 1927).**

Distr. Southern South America (Malta, 1927).

***U. pygmaeothecia* (C. Muell.) Kindb. (Kindberg, 1888).**

Syn. *Orthotrichum pygmaeothecium* C. Muell. (Mueller, 1885).
Ulota immarginata Card. (Cardot, 1905a).

Distr. Southern South America (Bescherelle, 1885a; Mueller, 1885; Paris, 1905a; Cardot, 1905a, 1908a; Cardot and Brotherus, 1923; Malta, 1927; Kühnemann, 1938; Bartram, 1952; Seki, 1974).

***U. rufula* (Mitt.) Jaeg. (Jaeger, 1874).**

Syn. *Orthotrichum rufulum* Mitt. (Mitten, 1869).
Orthotrichum brevicolle Mitt. (Mitten, 1869) *fide* Malta, 1927.
Ulota brevicollis (Mitt.) Jaeg. (Jaeger, 1874) *fide* Malta, 1927.
Orthotrichum chilense Mitt. (Mitten, 1869) *fide* Malta, 1927.

Ulota chilensis (Mitt.) Jaeg. (Jaeger, 1874) *fide* Malta, 1927.
Ulota persubulata Dus. (Dusén, 1903a) *nom. nud.* *fide* Malta, 1927.

Distr. Juan Fernandez (Brotherus, 1924b; Bartram, 1959; Robinson, 1975).
 Southern South America (Mitten, 1869; Jaeger, 1874; Dusén, 1903a; Paris, 1906;
 Herzog, 1923, 1938, 1954, 1957; Brotherus, 1925; Malta, 1927; Roivainen, 1934;
 Thériot, 1934b, 1935a; Seki, 1974).

var. *fagiola* Dus. ex Malta (Dusén ex Malta, 1927).

Distr. Southern South America (Malta, 1927).

var. *patagonica* Malta (Malta, 1927).

Distr. Southern South America (Malta, 1927; Seki, 1974).

U. savatieri Besch.

Ulota luteola

U. ventricosa (C. Muell.) Malta (Malta, 1927).

Syn. *Zygodon ventricosus* C. Muell. (Mueller, 1844b [1845]).

Distr. Southern South America (Mueller, 1844b, 1849; Montagne, 1850; Mitten, 1869; Jaeger,
 1874; Paris, 1906; Brotherus, 1925; Malta, 1927).

Valdiviella

V. pauperculata Thér. et Herz. (Thériot and Herzog *in Herzog*, 1928).

Distr. Southern South America (Herzog, 1928).

Verrucidens

V. intermedium Dix. (Dixon, 1920).

Syn. *Hymenoloma intermedium* (Dix.) Broth. (Brotherus, 1924a).

Distr. South Georgia (Dixon, 1920; Brotherus, 1924a; Steere, 1961b).

V. turpis (Card.) Card. (Cardot, 1908a).

Syn. *Blindia turpis* Card. (Cardot, 1905a).

Hymenoloma turpe (Card.) Card. et Broth. (Cardot and Brotherus, 1923).

Distr. Southern South America (Cardot, 1905a, 1908a; Cardot and Brotherus, 1923;
 Brotherus, 1924a; Donat, 1936a; Kühnemann, 1938; Herzog, 1957; Seki, 1974).

Vesicularia

V. argentinica Broth. (Brotherus, 1918).

Distr. Southern South America (Brotherus, 1918, 1925; Kühnemann, 1938; Herzog, 1952a).

V. vesicularis (Schwaegr.) Broth. (Brotherus, 1908).

Syn. *Hypnum vesiculosum* Schwaegr. (Schwaegrichen, 1827a).

Leskeia rutilans Brid. (Bridel, 1827b).

Ectropothecium rutilans (Brid.) Mitt. (Mitten, 1869).

Distr. Southern South America (Krieger, 1904; Herzog, 1952a).

Vesiculariopsis

V. spirifolium (Dus.) Broth. (Brotherus, 1908).

Syn. *Ectropothecium spirifolium* Dus. (Dusén, 1905a).

Vesiculariopsis spiripes Broth. ex Par. (Brotherus ex Paris, 1909) *err.*

Achrohypnella subenervis Herz. (Herzog, 1954) *fide* Seki, 1974.

Distr. Southern South America (Dusén, 1903a, 1905b; Paris, 1904b; Brotherus, 1908, 1925;
 Cardot, 1908a; Kühnemann, 1938; Herzog, 1954; Matteri, 1972; Seki, 1974).

V. spiripes Broth. ex Par.

Vesiculariopsis spirifolium

Webera hom. illeg.

W. albicans Schimp.

Pohlia wahlenbergii

W. alticaulis (C. Muell.) Kindb.

Pohlia wahlenbergii* var. *glacialis

<i>var. crassinervis</i> Card.	<i>Mniobryum alticaule</i> var. <i>crassinerve</i>
<i>var. robustior</i> (C. Muell.) Par.	<i>Mniobryum alticaule</i> var. <i>robustius</i>
<i>W. ampliretis</i> (C. Muell.) Kindb.	<i>Pohlia wahlenbergii</i> var. <i>glacialis</i>
<i>W. antarctica</i> (Hook. f. et Wils.) Jaeg.	<i>Bryum antarcticum</i>
<i>W. calopyxis</i> (C. Muell.) Kindb.	<i>Pohlia calopyxis</i>
<i>W. chilensis</i> Par. (Paris, 1906) <i>nom. nud.</i>	
Syn. <i>Webera paucifolia</i> Dus. (Dusén in Paris, 1906) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Paris, 1906).	
<i>W. commutata</i> Schimp.	<i>Pohlia drummondii</i>
<i>W. costesii</i> Card. et Thér.	<i>Pohlia polycarpa</i>
<i>W. cruda</i> (Hedw.) Fuernr.	<i>Pohlia cruda</i>
<i>var. imbricata</i> Card.	<i>Pohlia cruda</i> var. <i>imbricata</i>
<i>W. drummondii</i> (C. Muell.) Jaeg.	<i>Pohlia drummondii</i>
<i>W. emergens</i> (C. Muell.) Kindb.	<i>Mielichhoferia emergens</i>
<i>W. firmo-acuminata</i> Dus. (Dusén in Paris, 1906) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Paris, 1906).	
<i>W. gerlachei</i> Card.	<i>Bryum gerlachei</i>
<i>W. hariotiana</i> Besch.	<i>Goniobryum subbasilare</i>
<i>W. humilis</i> (Mont.) Broth.	<i>Pohlia humilis</i>
<i>W. inflexa</i> (C. Muell.) Kindb.	<i>Pohlia inflexa</i>
<i>W. leptoclada</i> Card. et Broth. (Cardot and Brotherus, 1923).	
<i>Distr.</i> Falkland Islands (Cardot and Brotherus, 1923; Brotherus, 1924a; Kühnemann, 1938).	
<i>W. lonchochaete</i> Dus.	<i>Pohlia lonchochaete</i>
<i>W. looseri</i> Thér. (Thériot, 1934b [1935]).	
<i>Distr.</i> Southern South America (Looser, 1932; Thériot, 1934b).	
<i>W. magnifica</i> Herz. (Herzog, 1954).	
<i>Distr.</i> Southern South America (Herzog, 1954).	
<i>W. manca</i> (C. Muell.) Kindb.	<i>Mielichhoferia manca</i>
<i>W. meyeniana</i> Hamp.	<i>Brachymenium meyenianum</i>
<i>W. nevadensis</i> (C. Muell.) Kindb.	<i>Pohlia nevadensis</i>
<i>W. nutans</i> Hedw.	<i>Pohlia nutans</i>
<i>var. minor</i> (Hook.) Brid.	<i>Mielichhoferia longiseta</i>
<i>W. paucifolia</i> Dus.	<i>Webera chilensis</i>
<i>W. philonotea</i> (C. Muell.) Par.	<i>Mniobryum philonoteum</i>
<i>W. philonotidea</i> Par.	<i>Mniobryum philonoteum</i>
<i>W. polycarpa</i> (Mitt.) Jaeg.	<i>Pohlia polycarpa</i>
<i>W. pulvinata</i> Kindb.	<i>Pohlia nutans</i>
<i>W. pyriformis</i> Hedw.	<i>Leptobryum pyriforme</i>
<i>W. racovitzae</i> Card.	<i>Pohlia racovitzae</i>
<i>var. laxiretis</i> Card.	<i>Pohlia racovitzae</i> var. <i>laxiretis</i>
<i>W. sphagnadelphus</i> (C. Muell.) Besch.	<i>Pohlia nutans</i>
<i>W. synoico-cruda</i> (C. Muell.) Kindb.	<i>Pohlia cruda</i>
<i>W. timmiaecaulon</i> Dus. (Dusén, 1903a) <i>nom. nud.</i>	
<i>Distr.</i> Southern South America (Dusén, 1903a).	

<i>W. viridata</i> (C. Muell.) Kindb.	<i>Pohlia cruda</i>
<i>W. wahlenbergii</i> (Web. et Mohr) Fuernr.	<i>Pohlia wahlenbergii</i>
<i>W. wilsonii</i> (Mitt.) Jaeg.	<i>Leptobryum wilsonii</i>

Weisa

<i>W. acuta</i> Hedw.	<i>Blindia acuta</i>
<i>W. alpicola</i> (Hedw.) Poir.	<i>Schistidium alpicola</i>
<i>W. amblyodon</i> Brid.	<i>Weisia controversa</i> var. <i>amblyodon</i>
<i>W. antarctica</i> (C. Muell.) Wils.	<i>Dicranoweisia antarctica</i>
<i>W. apocarpa</i> (Hedw.) Poir.	<i>Schistidium apocarpum</i>
<i>W. argentinica</i> C. Muell.	<i>Hymenostomum argentinicum</i>
<i>W. auridens</i> C. Muell. var. <i>robusta</i> C. Muell.	<i>Oreoweisia bogotensis</i>
<i>W. balansaeana</i> (Besch.) C. Muell.	<i>Oreoweisia auridens</i> var. <i>robusta</i>
<i>W. bogotensis</i> Hamp.	<i>Hymenostomum balansaeanum</i>
<i>W. calcarea</i> (Nees et Hornsch.) C. Muell.	<i>Oreoweisia bogotensis</i>
<i>W. chilensis</i> Hamp. ex C. Muell.	<i>Gymnostomum aeruginosum</i>
<i>W. cuncta</i> Hook. f. et Wils.	<i>Oreoweisia chilensis</i>
<i>W. controversa</i> Hedw. (Hedwig, 1801).	<i>Blindia cuncta</i>

Syn. *Grimmia controversa* (Hedw.) Web. et Mohr (Weber and Mohr, 1803).
Bryum controversum (Hedw.) P. Beauv. (Palisot de Beauvois, 1805a).
Weisia viridula Hedw. ex Brid. (Hedwig ex Bridel, 1819) nom. illeg.
Simophyllum viridulum Lindb. (Lindberg, 1871) nom. illeg.
Tortella viridula C. Jens. (Jensen, 1923) nom. illeg.
Weisia flavipes Hook. f. et Wils. (Hooker and Wilson in Wilson, 1854).

Distr. Juan Fernandez (Bartram, 1959; Robinson, 1975).
Southern South America (Herter, 1933a; Piovano, 1954).

var. *amblyodon* (Brid.) Sendtn. (Sendtner, 1848).

Syn. *Weisia amblyodon* Brid. (Bridel, 1827a).
Weisia viridula Hedw. ex Brid. var. *amblyodon* (Brid.) B.S.G. (Bruch and others, 1846).

Distr. Southern South America (Herzog, 1954).

<i>W. crispula</i> sensu Wils. et Hook. (Wilson and Hooker, 1847).	see <i>Dicranoweisia antarctica</i>
<i>W. cryptodon</i> Mont.	<i>Camptodontium cryptodon</i>
<i>W. curvirostris</i> C. Muell.	<i>Hymenostylium recurvirostre</i>
<i>W. donniana</i> (Sm.) Poir.	<i>Grimmia donniana</i>

***W. felipponei* Thér. (Thériot in Felippone, 1929 [1930]).**

Distr. Southern South America (Felippone, 1929; Herter, 1933a).

<i>W. flavipes</i> Hook. f. et Wils.	<i>Weisia controversa</i>
<i>W. fugax</i> Hedw.	<i>Rhabdoweisia fugax</i>
<i>W. gracilis</i> (Hook.) Spreng.	<i>Pilopogon gracilis</i>
<i>W. heteromalla</i> Hedw.	<i>Ditrichum heteromallum</i>
<i>W. kunzeana</i> C. Muell.	<i>Hymenostomum kunzeanum</i>
<i>W. latifolia</i> Schwaegr.	<i>Stegonia latifolia</i>
<i>W. longirostris</i> Mitt.	<i>Hymenostylium kunzeanum</i>

***W. patagonica* Card. et Broth. (Cardot and Brotherus, 1923).**

Distr. Southern South America (Cardot and Brotherus, 1923; Kühnemann, 1938).

<i>W. pentasticha</i> (Brid.) Hedw.	<i>Conostomum pentastichum</i>
<i>W. perichaetialis</i> Mont.	<i>Eucamptodon perichaetialis</i>
<i>W. phyllantha</i> (Brid.) Lindb.	<i>Ulota phyllantha</i>
<i>W. plagiopodia</i> (Hedw.) Poir.	<i>Grimmia plagiopodia</i>
<i>W. poeppigiana</i> (C. Muell.) Mitt.	<i>Trichostomopsis australasiae</i>
<i>W. recurvirostris</i> Hedw.	<i>Bryoerythrophyllum recurvirostre</i>
<i>W. recurvirostris</i> (Hedw.) Dix.	<i>Hymenostylium recurvirostre</i>
<i>W. richardii</i> (Brid.) Schwaegr.	<i>Campylopus richardii</i>
<i>W. semi-involuta</i> C. Muell.	<i>Hymenostomum semi-involutum</i>
<i>W. sphaerothecia</i> C. Muell.	<i>Amphidium tortuosum</i>
<i>W. tenuis</i> Kunz. ex Mont.	<i>Microdus kunzeanus</i>
<i>W. verticillata</i> Brid.	<i>Eucladium verticillatum</i>
<i>W. viridula</i> Hedw. ex Brid.	<i>Weisia controversa</i>
var. <i>amblyodon</i> (Brid.) B.S.G.	<i>Weisia controversa</i> var. <i>amblyodon</i>

Weymouthia

<i>W. billardieri</i> (Hamp.) Broth.	<i>Weymouthia cochlearifolia</i> var. <i>billardieri</i>
---	---

***W. cochlearifolia* (Schwaegr.) Dix. (Dixon, 1927).**

Syn. *Hypnum cochlearifolium* Schwaegr. (Schwaegrichen, 1816).
Isothecium cochlearifolium (Schwaegr.) Mitt. (Mitten, 1856).
Stereodon cochlearifolius (Schwaegr.) Mitt. (Mitten, 1860).
Coelidium cochlearifolium (Schwaegr.) Reichdt. (Reichardt, 1870).
Porotrichum cochlearifolium (Schwaegr.) Mitt. (Mitten, 1883).
Lembophyllum cochlearifolium (Schwaegr.) Lindb. (Lindberg in Paris, 1896).

Distr. Southern South America (Montagne, 1850; Eaton, 1892; Paris, 1896, 1905a; Thériot, 1935a).

var. *billardieri* (Hamp.) Dix. (Dixon, 1927).

Syn. *Neckera billardieri* Hamp. (Hampe, 1860).
Pilotrichella billardieri (Hamp.) Jaeg. (Jaeger, 1877).
Meteoriump billardieri (Hamp.) Mitt. (Mitten, 1882).
Weymouthia billardieri (Hamp.) Broth. (Brotherus, 1906b).
Pilotrichella kraussei Lor. (Lorentz, 1864).

Distr. Southern South America (Lorentz, 1864, 1866; Jaeger, 1877; Paris, 1905b; Brotherus, 1906b, 1925; Cardot, 1908a; Campo, 1915; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1940, 1954, 1960; Reimers, 1926; Bartram, 1952; Seki, 1974).

***W. mollis* (Hedw.) Broth. (Brotherus, 1906b).**

Syn. *Leskea mollis* Hedw. (Hedwig, 1801).
Neckera mollis (Hedw.) C. Muell. (Mueller, 1850).
Meteoriump molle Wils. (Wilson, 1854).
Isothecium molle (Hedw.) Mitt. (Mitten, 1856).
Stereodon mollis (Hedw.) Mitt. (Mitten, 1860) hom. illeg.
Pilotrichella mollis (Hedw.) Jaeg. (Jaeger, 1877).
Neckera cumingii C. Muell. (Mueller, 1850).
Pilotrichella cumingii (C. Muell.) Lor. (Lorentz in Jaeger, 1877).

Distr. Juan Fernandez (Montagne, 1835; Johow, 1896; Paris, 1905b; Skottsberg, 1914; Brotherus, 1924b; Thériot, 1927b; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Montagne, 1850; Mueller, 1850, 1855; Lorentz, 1866; Mitten, 1869; Jaeger, 1877; Bescherelle, 1889; Neger, 1899; Paris, 1905b; Brotherus, 1906b, 1924c, 1925; Cardot, 1908a; Rowlee, 1916; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1954, 1960; Reimers, 1926; Rovainen, 1934; Espinosa B., 1941; Skottsberg, 1950; Bartram, 1952; Bizot, 1967; Seki, 1974).

***W. orbiculata* Thér.**

Camptochaete orbiculata

Willia

W. austro-leucophaea (Besch.) Broth. (Brotherus, 1902).

- Syn. *Grimmia austro-leucophaea* Besch. (Bescherelle, 1885a).
Guembelia austro-leucophaea C. Muell. (Mueller in Paris, 1895) *nom. nud.*
Willia grimmoides C. Muell. (Mueller, 1890a) *fide* Bell, 1974.
Willia senilis C. Muell. (Mueller, 1901a [1900]).
Grimmia senilis Besch. (Bescherelle in Mueller, 1901a [1900]) *nom. nud.*
Distr. Southern South America (Bescherelle, 1885a, 1889; Mueller, 1901a; Brotherus, 1902, 1924a; Paris, 1906; Cardot, 1908a).
South Georgia (Mueller, 1890a; Cardot, 1906a, 1908a; Paris, 1906; Brotherus, 1924a; Steere, 1961b; Bell, 1974).

W. grimmoides C. Muell.

Willia austro-leucophaea

W. senilis C. Muell.

Willia austro-leucophaea

Wollyan

W. pottiaeae (Dus.) Herz.

Leptobryum pottiaeae

W. wilsonii (Mitt.) Herz.

Leptobryum wilsonii

Zieria hom. illeg.

Z. lamprocarpa (C. Muell.) Kindb.

Bryum lamprocarpum

Zygodon

Z. bartramoides Malta (Malta, 1924).

- Syn. *Pentastichella bartramoides* Dus. (Dusén in Malta, 1924) *nom. nud.*
Zygodon cylindropyxis C. Muell. (Mueller in Malta, 1926) *nom. nud.*
Stenomitrium robustulum Par. (Paris, 1905b) *nom. nud.*
Pentastichella robustula Dus. (Dusén in Paris, 1905b) *nom. nud.*
Zygodon robustulus Broth. (Brotherus in Malta, 1926) *nom. nud.*

Distr. Southern South America (Neger, 1899; Paris, 1905b; Malta, 1924, 1926; Brotherus, 1925; Herzog, 1938, 1954; Seki, 1974).

Z. caldensis Aongstr.

Zygodon reinwardtii

Z. coarctatus Schimp.

Zygodon reinwardtii

Z. coralensis Lor. ex C. Muell.

Zygodon coralensis

Z. coralensis Lor. (Lorentz, 1866).

- Syn. *Ceratodon coralensis* (Lor.) Jaeg. (Jaeger, 1873).
Zygodon coralensis Lor. ex C. Muell. (Lorentz ex Mueller, 1879a) err.

Distr. Juan Fernandez (Malta, 1926).

Southern South America (Lorentz, 1866; Jaeger, 1873, 1874; Paris, 1906; Malta, 1924, 1926; Brotherus, 1925; Herzog, 1938, 1954; Seki, 1974).

Z. curvicaulis Dus.

Zygodon intermedius

Z. cyathicarpus Mont.

Amphidium tortuosum

Z. cylindropyxis C. Muell.

Zygodon bartramoides

Z. denticulatus Tayl.

Zygodon reinwardtii

Z. erythrocarpus C. Muell.

Zygodon obtusifolius

Z. excelsus C. Muell.

Molendoa sendtneriana

Z. gracillimus Dus.

Zygodon papillatus

Z. hatcheri Dus.

Macromitrium sp.

Z. hyadesii Besch.

Zygodon reinwardtii var. *hyadesii*

var. *inermis* Dus.

Zygodon inermis

Z. hystrix Dus.

Zygodon pichinchensis

Z. inermis Malta (Malta, 1926).

Syn. *Zygodon hyadesii* Besch. var. *inermis* Dus. (Dusén *in* Malta, 1926) *nom. nud.*

Distr. Southern South America (Malta, 1926; Herzog, 1940; Seki, 1974).

Z. intermedius B.S.G. (Bruch and others, 1838).

Syn. *Zygodon kraussei* Lor. (Lorentz, 1864).

Zygodon curvicaulis Dus. (Dusén, 1903a).

Zygodon pseudoventricosus C. Muell. (Mueller *in* Malta, 1926) *nom. nud.*

Distr. Juan Fernandez (Malta, 1926; Thériot, 1927b; Bartram, 1959; Robinson, 1975).

Southern South America (Montagne, 1850; Lorentz, 1864, 1866; Jaeger, 1873, 1874; Dusén, 1903a; Paris, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Malta, 1926; Kühnemann, 1938; Herzog, 1954; Seki, 1974).

Z. jaffuelii Thér.

Zygodon liebmannii

Z. kraussei Lor.

Zygodon intermedius

Z. liebmannii Schimp. (Schimper *in* Mueller, 1849).

Syn. *Zygodon jaffuelii* Thér. (Thériot, 1921a).

Distr. Southern South America (Costes, 1921; Thériot, 1921a; Malta, 1926; Herzog, 1938, 1954; Seki, 1974).

Z. linearis C. Muell.

Anoectangium lineare

Z. magellanicus Dus. ex Malta (Dusén ex Malta, 1926).

Distr. Southern South America (Brotherus, 1925; Malta, 1926; Seki, 1974).

Z. mandonianus Schimp. ex C. Muell.

Anoectangium lineare

Z. menziesii (Schwaegr.) Arnott (Arnott, 1827).

Syn. *Codonoblepharon menziesii* Schwaegr. (Schwaegrichen, 1824).

Distr. Juan Fernandez (Brotherus, 1924b; Robinson, 1975).

Southern South America (Mueller, 1844b, 1849; Montagne, 1850; Mitten, 1869; Jaeger, 1874; Paris, 1906; Brotherus, 1925; Malta, 1926; Herzog, 1954; Seki, 1974).

Z. obovalis Mitt. (Mitten, 1885).

Distr. Juan Fernandez (Mitten, 1885; Johow, 1896; Paris, 1900a, 1906; Brotherus, 1924b, 1925; Malta, 1926).

Z. obtusifolius Hook. (Hooker, 1819).

Syn. *Zygodon erythrocarpus* C. Muell. (Mueller, 1879a).

Zygodon rufulus Dus. (Dusén *in* Malta, 1926) *nom. nud.*

Distr. Southern South America (Mueller, 1879a; Paris, 1906; Malta, 1924, 1926; Brotherus, 1925; Kühnemann, 1938).

var. antarcticus Malta (Malta, 1924).

Distr. Southern South America (Malta, 1924, 1926).

Z. ochraceus C. Muell. (Mueller, 1879a).

Syn. *Zygodon pygmaeus* C. Muell. (Mueller, 1879a).

Distr. Southern South America (Mueller, 1879a; Paris, 1906; Brotherus, 1925; Malta, 1926; Kühnemann, 1938).

Z. orthotrichoides (C. Muell.) Broth.

Leptodontium orthotrichoides

Z. palmarum C. Muell.

Leptodontium orthotrichoides

Z. papillatus Mont. (Montagne, 1845a).

Syn. *Codonoblepharon papillatum* (Mont.) Jaeg. (Jaeger, 1874).

Zygodon papillosum Mont. ex Broth. (Montagne *ex* Brotherus, 1902) *err.*

Zygodon gracillimus Dus. (Dusén, 1903a) *nom. nud.*

Distr. Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Mitten, 1869; Jaeger, 1874; Brotherus, 1902, 1925; Dusén, 1903a; Paris, 1906; Malta, 1926; Herzog, 1960; Seki, 1974).

Z. papillosum Mont. ex Broth.

Zygodon papillatus

***Z. pentastichus* (Mont.) C. Muell. (Mueller, 1849).**

Syn. *Aulacomnium pentastichum* Mont. (Montagne, 1845a).

Stenomitrium pentastichum (Mont.) Broth. (Brotherus, 1902).

Pentastichella pentasticha (Mont.) C. Muell. (Mueller in Thériot, 1918) *comb. inval.*

Zygodon uncinatus Mitt. (Mitten, 1869).

Codonoblepharon uncinatum (Mitt.) Jaeg. (Jaeger, 1874).

Macromitrium poeppigii Dub. (Duby, 1868).

Distr. Juan Fernandez (Skottsberg, 1914; Brotherus, 1924b; Malta, 1926; Espinosa B., 1941; Bartram, 1959; Robinson, 1975).

Southern South America (Montagne, 1845a, 1850, 1856; Mueller, 1849; Lorentz, 1866; Duby, 1868; Mitten, 1869; Jaeger, 1874; Neger, 1899; Brotherus, 1902, 1925; Paris, 1905a, b, 1906; Cardot, 1908a; Campo, 1915, 1921; Thériot, 1918; Cardot and Brotherus, 1923; Herzog, 1923, 1938, 1939, 1954, 1957, 1960; Malta, 1926; Reimers, 1926; Seki, 1974).

var. *crispatus* (Thér.) Malta (Malta, 1926).

Syn. *Zygodon porteri* Thér. var. *crispatus* Thér. (Thériot, 1926).

Distr. Southern South America (Malta, 1926; Thériot, 1926).

var. *strictifolius* Malta (Malta, 1926).

Syn. *Pentastichella pentasticha* (Mont.) C. Muell var. *strictifolia* Thér. (Thériot, 1924) *comb. inval.*

Distr. Southern South America (Thériot, 1924; Malta, 1926).

Z. perpusillus Thwait. et Mitt.

Rhachithecium perpusillum

***Z. pichinchensis* (Tayl.) Mitt. (Mitten, 1851).**

Syn. *Didymodon pichinchensis* Tayl. (Taylor, 1848b).

Zygodon hystrix Dus. (Dusén in Paris, 1906) *nom. nud.*

Distr. Southern South America (Paris, 1906; Brotherus, 1925; Malta, 1926; Seki, 1974).

***Z. podocarpi* C. Muell. ex Malta (Mueller ex Malta, 1926).**

Distr. Southern South America (Malta, 1926).

***Z. porteri* Thér. (Thériot, 1926).**

Syn. *Pentastichella jaffuelii* Thér. (Thériot, 1921a) *comb. inval.*

Distr. Southern South America. (Thériot, 1921a, 1926).

var. *crispatus* Thér.

Zygodon pentastichus* var. *crispatus

Z. pseudoventricosus C. Muell.

Zygodon intermedius

Z. pygmaeus C. Muell.

Zygodon ochraceus

***Z. reinwardtii* (Hornschr.) Braun (Braun in Bruch and others, 1838).**

Syn. *Syrrhopodon reinwardtii* Hornsch. (Hornschuch in Reinwardt and Hornschuch, 1826 [1829]).

Zygodon denticulatus Tayl. (Taylor, 1847).

Zygodon subdenticulatus Hamp. (Hampe, 1865).

Zygodon caldensis Aongstr. (Ångström, 1876a).

Zygodon coarctatus Schimp. (Schimper in Paris, 1898) *nom. nud.*

Distr. Southern South America (Mitten, 1869; Jaeger, 1874; Paris, 1898, 1906; Brotherus, 1925; Malta, 1926; Bartram, 1952).

var. *hyadesii* (Besch.) Malta (Malta, 1926).

Syn. *Zygodon hyadesii* Besch. (Bescherelle, 1885a).

Distr. Southern South America (Bescherelle, 1885a, 1889; Paris, 1906; Cardot, 1908a; Cardot and Brotherus, 1923; Brotherus, 1925; Malta, 1926; Kühnemann, 1938; Seki, 1974).

<i>Z. robustulus</i> Broth.	<i>Zygodon bartramoides</i>
<i>Z. rufulus</i> Dus.	<i>Zygodon obtusifolius</i>
<i>Z. subdenticulatus</i> Hamp.	<i>Zygodon reinwardtii</i>
<i>Z. tenerrimus</i> (C. Muell.) C. Muell.	<i>Gymnostomum tenerrimum</i>
<i>Z. theriotii</i> Herz. (Herzog, 1938).	
<i>Distr.</i> Southern South America (Herzog, 1938).	
<i>Z. tristis</i> Herz. (Herzog, 1960 [1961]).	
<i>Distr.</i> Southern South America (Herzog, 1960).	
<i>Z. uncinatus</i> Mitt.	<i>Zygodon pentastichus</i>
<i>Z. ventricosus</i> C. Muell.	<i>Ulota ventricosa</i>

III. ACKNOWLEDGEMENTS

It is a pleasure to acknowledge the support given by Sir Vivian Fuchs, FRS, who was Director of the British Antarctic Survey when this work was started. I am also grateful to Dr M. W. Holdgate, CB and Mr J. N. R. Jeffers, respectively, the former and current Directors of the Institute of Terrestrial Ecology, who facilitated its continuation to completion.

I wish to thank Miss P. I. Edwards, a former Head of the Library in the Department of Botany, British Museum (Nat. Hist.), who most generously spent many hours helping in the search for and verification of references, and Mr A. Eddy, of the same institution, who gave valuable expert advice on the presentation of data in *Sphagnum*. The help of Professor F. A. Stafleu and Dr W. D. Margadant, both of the University of Utrecht, is also gratefully acknowledged for assistance with dating queries. Staff in the Inter-Library Loans Service of the University of Birmingham (where the work was started), and of my present Institute, gave much essential assistance which facilitated my consultation of many works.

Of my colleagues, first Mrs J. Crawford and then Mrs A. Halcrow, gave continuous and unstinting support in the typing up of a voluminous manuscript, while Mr B. G. Bell is acknowledged for much helpful discussion throughout the work. A special word of thanks is reserved for Dr R. J. Adie, who advised on the presentation of the text and spent countless hours seeing it through the various stages of publication, and for Dr S. W. Greene, who suggested the project, helped in the search for literature and advised on the solution of many problems, and whose constant encouragement ensured its completion and final publication.

IV. REFERENCES

FOR each reference the date following the author's name is the bibliographic or title-page date. Where the date of issue, i.e. the taxonomically effective date, differs from the title-page date, the former is given in square brackets at the end of the reference. For further discussion see Introduction.

- AMANN, J. 1916. Étude sur la flore bryologique du Valais, 1912–1914. *Bull. Murithienne*, Fasc. 39 (for 1914–15), 350–55.
- AMANN, J. 1924. Le *Bryum schleicheri* Schwaegr. *Revue bryol.*, **51**, No. 2, 20–22.
- AMANN, J. 1928. Nouvelles additions et rectifications à la flore des mousses de la Suisse. *Mém. Soc. vaud. Sci. nat.*, Sér. 5, Mém. No. 16, 2, No. 9 [Mém. No. 17, 3, No. 2], 25–64.
- ANDO, H. 1972. Distribution and speciation in the genus *Hypnum* in the circum-Pacific region. *J. Hattori bot. Lab.*, No. 35, 68–98.
- ANDO, H. 1973. Studies on the genus *Hypnum* Hedw. II. *J. Sci. Hiroshima Univ.*, Ser. B, Div. 2 (Botany), **14**, No. 3, 165–207.
- ANDO, H. 1976. Studies on the genus *Hypnum* Hedw. III. *J. Sci. Hiroshima Univ.*, Ser. B, Div. 2 (Botany), **16**, No. 1, 1–46. [Effective date 20 November 1977, according to Ando personal communication.]
- ANDREWS, A. LE R. 1935. Bryaceae. (*In GROUT, A. J., ed. Moss flora of North America north of Mexico*. Newfane, Vermont, A. J. Grout, **2**, Pt. 3, 184–210.)
- ANDREWS, A. LE R. 1941. Notes on the Warnstorff *Sphagnum* herbarium. III. The subgenus *Inophloea* in South America. *Bryologist*, **44**, No. 6, 155–59.
- ANDREWS, A. LE R. 1947. Studies in the Warnstorff *Sphagnum* herbarium. IV. The group *Acutifolia* in South America. *Bryologist*, **50**, No. 2, 181–86.
- ANDREWS, A. LE R. 1949a. Taxonomic notes. VII. The genus *Acrocladium*. *Bryologist*, **52**, No. 2, 72–77.
- ANDREWS, A. LE R. 1949b. Studies in the Warnstorff *Sphagnum* herbarium. V. The group *Cuspidata* in South America. *Bryologist*, **52**, No. 3, 124–30.
- ANDREWS, A. LE R. 1951. Taxonomic notes. X. The family Leptostomaceae. *Bryologist*, **54**, No. 4, 217–23. [Effective date 16 February 1952; see **55**, No. 1, 84.]
- ÅNGSTRÖM, J. 1872. Förteckning och beskrifning öfver mossor, samlade af Professor N. J. Andersson under Fregatten Eugenies verldsomsegling åren 1851–53. *Övers. K. VetenskAkad. Förh.*, **29**, No. 4, 3–29.
- ÅNGSTRÖM, J. 1876a. Primae lineae muscorum cognoscendorum, qui ad Caldas Brasiliae sunt collecti. *Övers. K. VetenskAkad. Förh.*, **33**, No. 4, 3–50.
- ÅNGSTRÖM, J. 1876b. Rättelser och tillägg till förteckning och beskrifning öfver mossor, samlade af Professor N. J. Andersson under Fregatten Eugenies verldsomsegling 1851–1853, i Öfversigt af Kongl. VetenskapsAkademians Förhandlingar 1872, No. 4. *Övers. K. VetenskAkad. Förh.*, **33**, No. 4, 50–52.
- ARNELL, H. W. and C. JENSEN. 1910. Die Moose des Sarakgebietes, Abh. 2 und 3. *Naturw. Unters. sarekgebirg*, **3**, Botanik, Lief. 3, 144–268.
- ARNOTT, G. A. W. 1824a. Notice of a "Journal of a voyage from Rio de Janeiro to the coast of Peru, by Mr. William Jameson, surgeon, corresponding member of the Wernerian Natural History Society". *Mem. Wernerian nat. Hist. Soc.*, **5**, Pt. 1, 187–205.
- ARNOTT, [G.A.] W. 1824b. Notice sur quelques mousses de Rio-Janeiro. *Mém. Soc. Hist. nat. Paris*, **1**, Pt. 2, 346–52.
- ARNOTT, G. A. W. 1827. Nouvelle disposition méthodique. Des espèces de mousses exactement connues, par M. G.-A. Walker-Arnott, correspondent à Edimbourg; précédée d'un rapport fait à son sujet à la Société Linnéenne par M. B. Kittel, d.m.; suivie de notes critiques et 'additions qui complètent ce travail et le mettent à la hauteur des connaissances actuelles. *Mém. Soc. linn. Paris*, **5**, Pt. 2, 205–324.
- AUSTIN, C. F. 1877. Notes. *Bot. Gaz.*, **2**, No. 12, 142–43.
- AUSTIN, C. F. 1880. Bryological notes. *Bull. Torrey bot. Club*, **7**, No. 2, 15.
- BALANSA, B. and T. SAVÈS. 1887. *Musci exotici. Rev. bryol.*, **14**, No. 4, 56–57.
- BARTRAM, E. B. 1931. A review of the American species of *Daltonia*. *Bull. Torrey bot. Club*, **58**, No. 1, 31–48.
- BARTRAM, E. B. 1936. New mosses from Argentina. (*In BAUER, E., ed. Musci europaei et americani exsiccati. Appendix ad seriem 46. Stadt Olbersdorf, Ludwig Richter, 1–7.*)
- BARTRAM, E. B. 1938. The Second Byrd Antarctic Expedition — botany. III. Mosses. *Ann. Mo. bot. Gdn.*, **25**, No. 2, 719–24.
- BARTRAM, E. B. 1942. Additions to the moss flora of northern Chile. *Revue bryol. lichen.*, N. S., **13**, Fasc. 1, 130–34. [Issued as *Travaux bryologiques dédiés à le Mémoire de Pierre-Tranquille Husnot, Botaniste normand, Fondateur de la Revue Bryologique (1840–1929)*. Paris, Laboratoire de Cryptogamie, Muséum National d'Histoire Naturelle, 1942.]
- BARTRAM, E. B. 1943. Additions to the moss flora of northern Chile. *Farlowia*, **1**, No. 2, 191–94.
- BARTRAM, E. B. 1946. New mosses from Tierra del Fuego. *Farlowia*, **2**, No. 3, 309–19.
- BARTRAM, E. B. 1952. Mosses of Chile and Argentina mainly collected by R. Santesson. *Svensk bot. Tidskr.*, **46**, Ht. 2, 242–53.
- BARTRAM, E. B. 1957. Mosses from the United States Antarctic Service Expedition, 1940–41. *Bryologist*, **60**, No. 2, 139–43.
- BARTRAM, E. B. 1959. Mosses collected during Dr. and Mrs. C. Skottsberg's second expedition to the Juan Fernandez Islands, December 1954 to March 1955. *Ark. Bot.*, Ser. And., **4**, Häfte 1, 29–43.
- BARTRAM, E. B. 1965. New and noteworthy mosses from northern Argentina. *Revue bryol. lichen.*, N. S., **33** (for 1964–65), Fasc. 3–4, 323–27.
- BAUER, E. 1931. *Musci europaei et americani exsiccati. Schedae und Bemerkungen zur 43 serie*, Nos. 2101–50. [Prague, E. Bauer.]
- BAUER, E. 1934. *Musci europaei et americani exsiccati. Schedae und Bemerkungen zur 45 serie*. Nos. 2201–2250. Stadt Olbersdorf, Ludwig Richter.
- BÉHERÉ, J. B. J. 1826. *Muscologia Rothomagensis, ou tableau analytique des mousses découvertes jusqu'à ce jour aux environs de Rouen*. Rouen, F. Baudry.

- BELL, B. G. 1973a. Notes on Antarctic bryophytes: I. New records from the Antarctic botanical zone. *British Antarctic Survey Bulletin*, No. 36, 131–32. [Effective date 21 June 1974.]
- BELL, B. G. 1973b. A synoptic flora of South Georgian mosses: II. *Chorisodontium*, *Dicranoloma*, *Dicranum*, *Platyneurum* and *Conostomum*. *British Antarctic Survey Bulletin*, No. 37, 33–52. [Effective date 15 August 1974.]
- BELL, B. G. 1973c. Notes on Antarctic bryophytes: II. Records of Racomitria from the Antarctic botanical zone. *British Antarctic Survey Bulletin*, No. 37, 91–94. [Effective date 15 August 1974.]
- BELL, B. G. 1974. A synoptic flora of South Georgian mosses: V. *Willia* and *Racomitrium*. *British Antarctic Survey Bulletin*, No. 38, 73–101.
- BELL, B. G. 1976. Notes on Antarctic bryophytes: VI. The genus *Dicranoweisia* on Signy Island, South Orkney Islands. *British Antarctic Survey Bulletin*, No. 44, 97–100.
- BELL, B. G. 1977a. Notes on Antarctic bryophytes: VIII. Two species of *Campylopus* originally described from South Georgia. *British Antarctic Survey Bulletin*, No. 46, 136–37.
- BELL, B. G. 1977b. Notes on Antarctic bryophytes: IX. A previously unreported botanical collection made on South Georgia during the German International Polar-Year Expedition, 1882–83. *British Antarctic Survey Bulletin*, No. 46, 137–39.
- BERTSCH, K. 1949. *Moosflora*. Stuttgart, Verlag Eugen Ulmer.
- BESCHERELLE, E. 1871–72. *Prodromus bryologiae mexicanae ou énumération des mousses du Mexique avec description des espèces nouvelles*. *Mém. Soc. natn. Sci. nat. math. Cherbourg*, 16, [Sér. 2, 6], 145–256. [Effective date 1872, *fide* Crum, 1950.]
- BESCHERELLE, E. 1876. Florule bryologique des Antilles Françaises. *Annls Sci. nat.*, Sér. 6, 3, Botanique, 175–265.
- BESCHERELLE, E. 1877. Note sur les mousses de Paraguay. Récoltées par M. Balansa de 1874 à 1877. *Mém. Soc. natn. Sci. nat. math. Cherbourg*, 21, [Sér. 3, 1], 257–72.
- BESCHERELLE, E. 1885a. Mousses nouvelles de l'Amérique australe. *Bull. Soc. bot. Fr.*, 32, LIV–LXIX.
- BESCHERELLE, E. 1885b. Liste des mousses du Paraguay distribuées en 1884 par M. Balansa. *Revue bryol.*, 12, No. 2, 17–19.
- BESCHERELLE, E. 1889. Mousses. (*In* BESCHERELLE, E. and C. MASSALONGO. *Muscinées. Mission Scientifique du Cap Horn 1882–83*. Vol. 5. Botanique, 253–312.)
- BESCHERELLE, E. 1891. Selectio novorum muscorum. *J. Bot., Paris*, 5, No. 9, 142–48; No. 15, 252–55; No. 20, 342–50.
- BIZOT, M. 1967. Quelques mousses africaines et américaines. *Bull. Soc. bot. Fr.*, 114, No. 9, 423–27.
- BIZOT, M. and G. PIOVANO. 1953. Musci brasilienses. *Dusenia*, 5, Fasc. 5–6, 443–52.
- BLANDOW, O. 1809. *Uebersicht der Mecklenburgischen Moose nach alphabetischer Ordnung entworfen*. Neustrelitz, Kosten.
- BOULAY, J. N. 1884. *Muscinées de la France. Part 1. Mousses*. Paris, F. Savy.
- BRAITHWAITE, R. 1893. *The British moss flora*. Vol. 2. London, R. Braithwaite, Pt. 15, 183–218.
- BRAITHWAITE, R. 1895. *The British moss flora*. Vol. 2. London, R. Braithwaite, Pt. 16, 221–68.
- BRAITHWAITE, R. 1898. *The British moss flora*. Vol. 3. London, R. Braithwaite, Pt. 18, 37–64.
- BRIDEL, S. E. 1798. *Muscologia recentiorum seu analysis, historia, et descriptio methodica omnium muscorum frondosorum, hucusque cognitorum, ad norman Hedwigii*. Gothae, Carolum Guil. Ettinger; Paris, Barrois Junior, 2, Pt. 1, 1–222.
- BRIDEL, S. E. 1801a. *Animaduersiones in Muscologiae Recentiorum tomum secundum, ab ipso auctore propositae*. *J. Bot., Göttingen*, 1 (for 1800), Stück 2, 268–99.
- BRIDEL, S. E. 1801b. *Muscologia recentiorum seu analysis, historia et descriptio methodica omnium muscorum frondosorum hucusque cognitorum ad norman Hedwigii*. Gothae, Carol Guil. Ettinger; Paris, Barrois Junior, 2, Pt. 2, 1–198.
- BRIDEL, S. E. 1802. Fortsetzung der Nro. 13 abgebrochen Recension über: Sam. El. Bridel, *Muscologia etc. Bot. Ztg Erlangen*, 1, Ht. 3, No. 14, 209–18.
- BRIDEL, S. E. 1803. *Muscologia recentiorum seu analysis, historia et descriptio methodica omnium muscorum frondosorum hucusque cognitorum ad normam Hedwigii*. Gothae, Carol Guil. Ettinger; Paris, Barrois Junior, 2, Pt. 3, 1–182.
- BRIDEL, S. E. 1806. *Muscologiae recentiorum. Supplementum seu species muscorum, Pars I.* Gothae, Carol Guil. Ettingeriana.
- BRIDEL, S. E. 1812. *Muscologiae recentiorum. Supplementum seu species muscorum, Pars II.* Gothae, Libraria Ettingeriana.
- BRIDEL, S. E. 1817. *Muscologiae recentiorum. Supplementum seu species muscorum, Pars III.* Gothae, Libraria Ettingeriana.
- BRIDEL, S. E. 1819. *Muscologiae recentiorum supplementum Pars IV, seu mantissa generum specierumque muscorum frondosorum universa. (Methodus nova muscorum ad naturae normam melius instituta et muscologiae recentiorum accommodata)*. Gothae, A. Ukertum.
- BRIDEL, S. E. 1826. *Bryologia universa seu systematica ad novam methodum dispositio, historia et descripto omnium muscorum frondosorum hucusque cognitorum cum synonymia ex auctoribus probatissimis. Vol. I.* Lipsiae, Joan. Ambros. Barth. (1–746.)
- BRIDEL, S. E. 1827a. *Bryologia universa seu systematica ad novam methodum dispositio, historia et descripto omnium muscorum frondosorum hucusque cognitorum cum synonymia ex auctoribus probatissimis, supplementum ad volumen primum. Lipsiae, Joan. Ambros. Barth. (747–856.)*
- BRIDEL, S. E. 1827b. *Bryologia universa seu systematica ad novam methodum dispositio, historia et descripto omnium muscorum frondosorum hucusque cognitorum cum synonymia ex auctoribus probatissimis. Vol. II.* Lipsiae, Joan. Ambros. Barth. (1–848.)
- BRITTON, E. G. 1896. An enumeration of the plants collected by H. H. Rusby, in Bolivia, 1885–1886: II. Musci. *Bull. Torrey bot. Club*, 23, No. 12, 471–99.
- BRITTON, E. G. 1913. *Ditrichaceae. (In North America flora, Vol. 15, Pt. 1. New York, New York Botanical Garden, 55–67.)*
- BRITTON, E. G. 1914. Notes on nomenclature — XII. *Bryologist*, 17, No. 1, 7–10.
- BROTERO, F. A. 1804. Musci. (*In Flora Lusitanica, seu plantarum, quae in Lusitania vel sponte crescent, vel frequentius coluntur, ex florum praesertim sexibus systematicae distributarum*,

- synopsis. Pars 2. Olisipone, Typographia Regia, 403–25.)*
- BROTHERUS, V. F. 1891. Some new species of Australian mosses. *Öfvers. finska VetenskSoc. Forh.*, **33** (for 1890–91), 89–110.
- BROTHERUS, V. F. 1893. Contributions à la flore bryologique du Brésil. *Acta Soc. Sci. fenn.*, **19**, No. 5, 3–30. [Effective date 1891 according to *Revue bryol.*, **18**, No. 5, 80.]
- BROTHERUS, V. F. 1895. Nouvelles contributions à la flora bryologique du Brésil. *Bih. K. svenska VetenskAkad. Handl.*, **21**, Afd. 3, No. 3, 1–76.
- BROTHERUS, V. F. 1898. Some new species of Australian mosses. *Öfvers. finska VetenskSoc. Forh.*, **40** (for 1897–98), 159–93.
- BROTHERUS, V. F. 1900. Die Laubmoose der ersten regnellschen expedition. *Bih. K. svenska VetenskAkad. Handl.*, **26**, Afd. 3, No. 7, 1–65.
- BROTHERUS, V. F. 1901. Musci (Laubmoose). II Unterklasse Andreaeales, Andreaeaceae; III Unterklasse Bryales: II. Spezieller Teil, Gruppe I. Acrocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 1, Lief. 207, 265–68, 277–88; Lief. 208, 289–336; Lief. 212, 337–84.)
- BROTHERUS, V. F. 1902. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe I. Acrocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 1, Lief. 214, 385–432; Lief. 215, 433–80.)
- BROTHERUS, V. F. 1903. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe I. Acrocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 1, Lief. 216, 481–528; Lief. 218, 529–76.)
- BROTHERUS, V. F. 1904. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe I. Acrocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 1, Lief. 219, 577–624; Lief. 220, 625–72.)
- BROTHERUS, V. F. 1905a. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe I. Acrocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 1, Lief. 222, 673–700.)
- BROTHERUS, V. F. 1905b. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe II. Pleurocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 2, Lief. 222, 701–20; Lief. 223, 721–68.)
- BROTHERUS, V. F. 1905c. *Pleuroorthotrichum*, eine neue Laubmoosgattung aus Chile. *Öfvers. finska VetenskSoc. Förh.*, **47**, No. 15, 1–4.
- BROTHERUS, V. F. 1906a. Die Laubmoose der Deutschen Südpolar-Expedition 1901–1903. *Dt. Südpol.-Exped.*, **8**, Botanik, Ht. 1, 83–96.
- BROTHERUS, V. F. 1906b. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe II. Pleurocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 2, Lief. 224, 769–816; Lief. 226, 817–64.)
- BROTHERUS, V. F. 1907. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe II. Pleurocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 2, Lief. 227/28, 865–960; Lief. 229, 961–1008.)
- BROTHERUS, V. F. 1908. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe II. Pleurocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 2, Lief. 231, 1009–56; Lief. 232/33, 1057–152.)
- BROTHERUS, V. F. 1909. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe II. Pleurocarpi. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Teil 1, Abt. 3, Hälften 2, Lief. 234/35, 1153–246.)
- BROTHERUS, V. F. 1910a. Contributions to the bryological flora of the Philippines. III. *Philipp. J. Sci., Sect. C, Botany*, **5**, No. 2, 137–62.
- BROTHERUS, V. F. 1910b. *Schedae ad bryothecam fennicam. N: ri 1–100.* Helsingforsiae, J. Simelii.
- BROTHERUS, V. F. 1916. Descriptions of some new species of Australian, Tasmanian and New Zealand mosses. VI. *Proc. Linn. Soc. N.S.W.*, **41**, No. 163, 575–96.
- BROTHERUS, V. F. 1918. Contributions à la flore bryologique de l'Argentine. *Ark. Bot.*, **15**, Hälften 1–2, 1–15.
- BROTHERUS, V. F. 1920. Contributions à la flore bryologique de l'Ecuador. *Revue bryol.*, **47**, No. 1, 1–16.
- BROTHERUS, V. F. 1923. *Die Laubmoose Fennoskandias.* Helsingfors, Akademische Buchhandlung; Berlin, R. Friedländer & Sohn; London, Wheldon & Wesley Ltd. [Societas pro Fauna et Flora Fennica, Vol. I.]
- BROTHERUS, V. F. 1924a. Musci (Laubmoose). II Unterklasse Andreaeales, Andreaeaceae. III. Unterklasse Bryales: II. Spezieller Teil. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Aufl. 2, **10**, Hälften 1, 129–31, 143–478.)
- BROTHERUS, V. F. 1924b. The Musci of Juan Fernandez Islands. (*In SKOTTSBERG, C. [J. F.], ed. The natural history of Juan Fernandez and Easter Island.* Vol. 2, Botany, Pt. 3. Uppsala, Almqvist and Wiksell, 409–48.)
- BROTHERUS, V. F. 1924c. Musci nonnulli chilenses a C. Skottberg anno 1917 lectae. *Acta Horti gothoburg.*, **1**, 189–95.
- BROTHERUS, V. F. 1925. Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil. (*In ENGLER, A. and K. PRANTL. Die natürlichen Pflanzenfamilien.* Leipzig, Wilhelm Engelmann, Aufl. 2, **11**, Hälften 2, 1–542.)
- BROTHERUS, V. F. 1928. Musci. (*In IRMISCHER, E., [ed.] Beiträge zur Kenntnis der Flora von Borneo. Mitt. Inst. allg. Bot., Hamb.*, **7**, Ht. 2, 115–40.)
- BROTHERUS, V. F. 1931. Neue exotische Laubmoose. *Mitt. Inst. allg. Bot., Hamb.*, **8**, Ht. 2, 399–406.
- BROWN, R. 1811. Some observations on the parts of frutification in mosses, with characters and descriptions of two new genera of that order. *Trans. Linn. Soc. Lond.*, **10**, Pt. 2, 312–24.
- BROWN, R. 1902. On the Musci of the calcareous districts of New Zealand, with descriptions of new species. *Trans. Proc. N.Z. Inst.*, **35**, 323–42. [Effective date July 1903.]
- BRUCH, [P.] 1826. Bryologische Beobachtungen. *Flora, Jena*, **9**, No. 11, 161–66.
- BRUCH, P. 1832. Beschreibung einiger Laubmoose. *Abh. bayer. Akad. Wiss. Math.-phys. Kl.*, **1**, 277–86.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.* 1838. *Bryologia europaea seu genera muscorum*

- Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 3, [Fasc. 4, 33–41.], *Zygodon*, 1–9. Vol. 4, [Fasc. 5, 165–201.], *Mnium*, 1–37.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1839. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 4, [Fasc. 6–9, 71–152.], *Bryum*, 1–82.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1841. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 3, [Fasc. 11, 259–68], *Funaria*, 1–10.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1842. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 2, [Fasc. 13–15, 63–108.], *Barbula*, 1–46. Vol. 4, [Fasc. 12, 31–51.], *Bartramia*, 1–21.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1843. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 1, [Fasc. 17, 185–96.], *Fissidens*, 1–12. Vol. 2, [Fasc. 18–20, 29–41.], *Pottia*, 1–13.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1844. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 3, [Fasc. 23–24, 288–90.], *Tetraplodon*, 1–8. Vol. 4, [Fasc. 21–22, 245–54.], *Pogonatum*, 1–10; [Fasc. 23–24, 67–70.], *Orthodontium*, 1–4.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1845. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 3, [Fasc. 25–28, 93–102.], *Schistidium*, 1–10.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1846. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 1, [Fasc. 29–30, 173–78.], *Trematodon*, 1–6; [Fasc. 33–36, 61–71.], *Weisia*, 1–11; [Fasc. 33–36, 73–82.], *Gymnostomum*, 1–10; [Fasc. 33–36, 91–93.], *Eucladium*, 1–3; [Fasc. 33–36, 95–100.], *Rhabdoweisia*, 1–6. Vol. 2, [Fasc. 29–30, 135–41.], *Didymodon*, 1–7; [Fasc. 29–30, 153–57.], *Distichium*, 1–5; [Fasc. 33–36, 17–20.], *Blindia*, 1–4. Vol. 3, [Fasc. 29–30, 155–58.], *Hedwigidium*, 1–4.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1850. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 5, [Fasc. 44–45, 31–36.], *Cryphaea*, 1–6; [Fasc. 44–45, 53–55.], *Ormalia*, 1–3.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1852. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 5, [Fasc. 49–51, 157–67.], *Thuidium*, 1–11; [Fasc. 49–51, 197–208.], *Rhynchostegium*, 1–12.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1852. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 5, [Fasc. 48, 179–95.], *Plagiothecium*, 1–17.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1853. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 6, [Fasc. 52–54, 5–26.], *Brachythecium*, 1–22; [Fasc. 55–56, 45–61.], *Amblystegium*, 1–17.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1854. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 6, [Fasc. 57–61, 77–130.], *Hypnum*, 1–54.
- BRUCH, P., SCHIMPER, W. P. and W. T. GÜMBEL.*
 1855. *Bryologia europaea seu genera muscorum Europaeorum monographice illustrata*. Stuttgart, Librariae E. Schweizerbart, Vol. 6, [Fasc. 62–64, 131–56.], *Andraea*, 1–26.
- BRUGGEMAN-NANNENGA, M. A. 1973. The section *Pachylomidium* (Genus *Fissidens*). I. The species of tropical and subtropical South America. *Proc. K. ned. Akad. Wet., Series C, Biological and Medical Sciences*, **76**, No. 2, 172–89.
- BRYHN, N. 1902. *Sarconeurum*, genus muscorum novum. *Sarconeurum antarcticum* sp. nov. (*In WILLE*, N. Mittheilungen über einige von C. E. Borchgrevink auf dem antarctischen Festlande gesammelte Pflanzen. *Nyt. Mag. Naturvid.*, **40**, Ht. 3 and 4, 204–08.)
- BUCK, W. R. and L. R. LANDRUM. 1977. A new species of *Astomiopsis* (Ditrichaceae). *Bryologist*, **80**, No. 2, 359–62.
- BUSE, L. H. 1858. *Musci Neerlandici specimen exsiccati. 'Neerlands mossoorten in getroogde exemplaren*. Haarlem, Kruseman, Fasc. 3, No. 67.
- CAMPO, J. A. 1915. Mousses chiliens déterminés par V. F. Brotherus, et récoltées dans la province de Valdivia. *Revta chil. Hist. nat.* ..., **19**, Nos. 3–6, 74–75.
- CAMPO, J. A. 1921. Pequeñ catalogo briológico de Mariluan. *Revta chil. Hist. nat.*, **25**, 507–10.
- CARDOT, J. 1884. Notes sphagnologiques. *Revue bryol.*, **11**, No. 4, 54–56.
- CARDOT, J. 1887. Révision des Sphaignes de l'Amérique du Nord. *Bull. Soc. r. Bot. Belg.*, **26**, Pt. 1, 41–61.
- CARDOT, J. 1900. Note préliminaire sur les mousses recueillies par l'Expédition Antarctique Belge. *Revue bryol.*, **27**, No. 3, 38–46.
- CARDOT, J. 1901. Mousses et coup d'oeil sur la flore bryologique des Terres Magellaniques. *Résult. Voyage S. Y. Belgica*, **6**, Botanique, 44 pp.
- CARDOT, J. 1905a. Notice préliminaire sur les mousses recueillies par l'Expédition Antarctique Suédoise. I. Espèces de la région magellanique. *Bull. Herb. Boissier*, 2ème sér., **5**, No. 11, 997–1011.
- CARDOT, J. 1905b. Notice préliminaire sur les mousses recueillies par l'Expédition Antarctique Suédoise. Deux genres nouveaux de mousses acrocarpes. *Revue bryol.*, **32**, No. 3, 45–47.
- CARDOT, J. 1905c. Mousses de l'île Formose. *Beih. bot. Zbl.*, **19**, Abt. 2, 85–148.
- CARDOT, J. 1906a. Notice préliminaire sur les mousses recueillies par l'Expédition Antarctique Suédoise. *Bull. Herb. Boissier*, 2ème sér., **6**, No. 1, 1–17.
- CARDOT, J. 1906b. Les mousses de l'Expédition Charcot. *Revue bryol.*, **33**, No. 3, 33–35.
- CARDOT, J. 1907a. Musci. (*In National Antarctic Expedition, 1901–1904. Natural History, Vol. III. Zoology and Botany*. London, British Museum, 1–6.)

* Publication dates, fascicule numbers and figures for continuous pagination within square brackets from Stafleu (1967, 57–63).

- CARDOT, J. 1907b. Une mousse à péristome asymétrique. *Bull. Soc. Hist. nat. Ardennes*, **14**, 27–30.
- CARDOT, J. 1907c. *Mousses*. Paris, Masson et Cie. [Expédition Antarctique Française (1903–1905), Sciences naturelles: documents scientifiques, 1–9.]
- CARDOT, J. 1908a. La flore bryologique des Terres Magellaniques, de la Géorgie du Sud et de l'Antarctide. *Wiss. Ergebni. schwed. Südpolarexped.*, Bd. 4, Lief. 8, 298 pp.
- CARDOT, J. 1908b. Sur le genre *Hymenocleiston* Duby. *Annu. Conserv. Jard. bot. Genève*, **11**, and **12**, 165–69.
- CARDOT, J. 1910. Musci. Note sur les mousses rapportées par l'Expédition du "Nimrod". (*In British Antarctic Expedition, 1907–9. Reports on the Scientific Investigations. Biology*. Vol. I, Pt. 4, 77–79.)
- CARDOT, J. 1911a. Deux genres nouveaux de la région magellanique. *Revue bryol.*, **38**, No. 3, 49–52.
- CARDOT, J. 1911b. Sur les mousses rapportées par l'expédition antarctique du Pourquois-Pas? *C. r. Ass. fr. Avanc. Sci.*, **153**, No. 13, 599–603.
- CARDOT, J. 1911c. Les mousses de l'Expédition Nationale Antarctique Écossaise. *Trans. R. Soc. Edinb.*, **48**, Pt. 1, No. 3, 67–82.
- CARDOT, J. 1911d. Note sur les mousses rapportées par la seconde Expédition Antarctique Française, sous le commandement du Dr. Jean Charcot. *Revue bryol.*, **38**, No. 6, 124–27.
- CARDOT, J. 1912a. *Atrichopsis* Card., genre nouveau de la famille des Polytrichacées. *Revue bryol.*, **39**, No. 6, 95–96.
- CARDOT, J. 1912b. Les mousses de l'expédition national antarctique Écossaise. *Rep. scient. Results Scott. natn. antarct. Exped.*, **3**, Pt. 5, 55–57.
- CARDOT, J. 1913a. *Mousses*. Paris, Masson et Cie. [Deuxième Expédition Antarctique Française (1908–1910), Sciences naturelles: documents scientifiques, 1–30.]
- CARDOT, J. 1913b. Quelques observations sur la nomenclature bryologique. *Revue bryol.*, **40**, No. 2, 17–22.
- CARDOT, J. 1914a. *Acrocladiopsis* Card., genre nouveau de la tribu des Plagiothécées. *Revue bryol.*, **41**, No. 1, 9.
- CARDOT, J. 1914b. *Philibertiella* Card. Genre nouveau de la tribu des Ditrichées. *Revue bryol.*, **41**, No. 3, 37–38.
- CARDOT, J. and V. F. BROTHERUS. 1923. Les mousses. *K. svenska VetenskAkad. Handl.*, **63**, No. 10, 1–73.
- CHEN, P-C. 1941. Studien über die ostasiatischen Arten der Pottiaceae. I. *Hedwigia*, **80**, Ht. 1/2, 1–76; Ht. 3/4, 141–322.
- CHEVALLIER, F. F. 1827. Mousses. Musci. (*In Flore générale des environs de Paris, selon la méthode naturelle. Description de toutes les plantes agames, cryptogames et phanérogams qui y croissent spontanément; leurs propriétés, leur usage dans la médecine, les arts et l'économie domestique; avec une classification naturelle des agames et des cryptogames, basée sur l'organisation de ce végétaux*. Vol. 2. Paris, Chez Ferra Jeune, 26–99.)
- CLARKE, G. C. S. 1973a. A synoptic flora of South Georgian mosses: III. *Leptotheca*, *Philonotis*, *Mielichhoferia* and *Pohlia*. *British Antarctic Survey Bulletin*, No. 37, 53–79. [Effective date 15 August 1974.]
- CLARKE, G. C. S. 1973b. Notes on Antarctic bryophytes: III. The type specimen of *Pohlia inflexa* (C. Muell.) Wijk et Marg. *British Antarctic Survey Bulletin*, No. 37, 97–98. [Effective date 15 August 1974.]
- CLIFFORD, H. T. 1955. On the distribution of *Rhacomitrium crispulum* (H. f. & W.) H. f. & W. *Bryologist*, **58**, No. 4, 330–34.
- CLIFFORD, H. T. 1957. New records for Antarctic mosses. *Aust. J. Sci.*, **20**, No. 4, 115.
- CONARD, H. S. 1945. The Atracheata (Bryophyta) of Iowa. II. Iowa mosses in print: A critique of the species reported for the state. *Bryologist*, **48**, No. 3, 103–09.
- COSTES, N. 1921. Nomenclature de los principales musgos de la hoya de Marga-Marga (Quilpué-Provincia de Valparaíso). *Revta chil. Hist. nat.*, **25**, 130–32.
- CROME, G. E. W. 1805. *Sammlung deutscher Laubmoose. Nachlieferung I*. Schwerin, J. W. Barenspurungs Erben.
- CROSBY, M. R. 1975. Lectotypification of *Schizomitrion* B.S.G. (Musci, Hookeriaceae). *Taxon*, **24**, Nos. 2/3, 353–55.
- CROSBY, M. R. 1976. The correct name for the moss *Duseniella*. *Bryologist*, **79**, No. 2, 246–47.
- CROSBY, M. R. 1977a. Index muscorum. supplementum: 1974–1975. *Taxon*, **26**, Nos. 2–3, 285–307.
- CROSBY, M. R. 1977b. *Florschuetzia*, a new genus of Buxbaumiaceae (Musci) from southern Chile. *Bryologist*, **80**, No. 1, 149–52.
- CROSBY, M. R. 1978. *Muscoflorschuetzia* (Musci), a new name for *Florschuetzia* Crosby, nom. illeg. *Bryologist*, **81**, No. 2, 338.
- CRUM, H. A. 1950. Publication date of the prodromus bryologiae mexicanae. *Bryologist*, **53**, No. 1, 60–61.
- CRUM, H. A. 1952. Bibliographic notes on Palisot de Beauvois' prodrome. *Bull. Torrey bot. Club*, **79**, No. 5, 407–09.
- CRUM, H. A. 1957. A contribution to the moss flora of Ecuador. *Svensk bot. Tidskr.*, **51**, Ht. 1, 197–206.
- CRUM, H. 1972. A taxonomic account of the Eriodiaceae. *Nova Hedwigia*, **23**, Ht. 2–3, 201–24. [Effective date 8 January 1973.]
- CRUM, H. A. and W. C. STEERE. 1950. Additions to the moss flora of Panama. *Bryologist*, **53**, No. 2, 139–52.
- CRUNDWELL, A. C. and E. NYHOLM. 1963. Two new European species of *Bryum*. *Bot. Notiser*, **116**, Fasc. 1, 94–98.
- CRUNDWELL, A. C. and E. NYHOLM. 1964. The European species of the *Bryum erythrocarpum* complex. *Trans. Br. bryol. Soc.*, **4**, Pt. 4, 638–41.
- DALLA TORRE, K. W. and L. G. SARNTHEIN. 1904. *Die Moose (Bryophyta) von Tirol, Vorarlberg und Liechtenstein. Flora der Gefürsteten Grafschaft Tirol, des Landes Vorarlberg und des Fürstenthumes Liechtenstein. Band. 5*. Innsbruck, Verlag des Wagner'schen Universitäts-Buchhandlung.
- DARBISHIRE, O. V. 1923. Cryptogams from the Antarctic. *J. Bot., Lond.*, **61**, No. 724, 105–07.
- DE CANDOLLE, M. 1815. Famille des mousses. (*In Flore française, ou descriptions succinctes de toutes les plantes qui croissent naturellement en France, disposées selon une nouvelle méthode d'analyse, et précédées par un exposé des principes élémentaires de la botanique*. Vol. 5 (ou

- 6). 3rd edition. Paris, Chez Desray, 204–36.)
- DE LA PYLAIE, B. 1814. Suite de l'article sur la monographie des mousses rangées parmi les *Hypnum* de Linné, et que les auteurs modernes ou tantôt établies comme un genre particulier sous le nom de *Fissidens*, et tantôt rapportées au genre *Dicranum*. *J. Bot. Agric. Pharm. Med. Arts Paris*, Sér. 2, 4, Nos. 4–6, 145–69. [Effective date 1815, *fide* Sayre (1959, 47).]
- DELGADILLO, C. 1973. A new species, nomenclatural changes, and generic limits in *Aloina*, *Aloinella* and *Crossidium* (Musci). *Bryologist*, **76**, No. 2, 271–77.
- DELGADILLO, C. 1975. Taxonomic revision of *Aloina*, *Aloinella* and *Crossidium* (Musci). *Bryologist*, **78**, No. 3, 245–303.
- DE NOTARIS, J. 1838a. Specimen de *Tortulis italicis*. *Memorie Accad. Sci. Torino*, **40**, 281–325.
- DE NOTARIS, J. 1838b. *Syllabus muscorum in Italia et in insulis circumstantibus hucusque cognitorum*. Taurini, J. de Notaris.
- DE NOTARIS, J. 1859. Musci Napoani sive muscorum ad flumen Napo in Columbia a clar. Osculati lectorum. *Memorie Accad. Sci. Torino*. Ser. 2, **18**, 437–55.
- DE NOTARIS, G. 1867. Cronaca della Briologia Italiana. Part II. *Comment. Soc. crittogram. Ital.*, **2**, Fasc. 3, 269–312 [169–212].
- DE NOTARIS, G. 1869. Epilogo della briologia Italiana. *Atti Univ. Genova*, **1**, 1–781.
- DICKSON, J. 1801. *Fasciculus quartus plantarum cryptogamicarum Britanniae*. Londini, Auctor.
- DISMIER, G. 1910. Revision des *Philonotis* de l'Amérique. *Mém. Soc. bot. Fr.*, **2**, No. 17, 1–37.
- DIXON, H. N. 1896. *The student's handbook of British mosses, with illustrations, and keys to the genera and species by H. G. Jameson*. Eastbourne, V.T. Sumfield.
- DIXON, H. N. 1902. New varieties of British mosses. *J. Bot., Lond.*, **40**, No. 11, 374–80.
- DIXON, H. N. 1912. On some mosses of New Zealand. *J. Linn. Soc., Botany*, **40**, No. 277, 433–59.
- DIXON, H. N. 1913. Studies in the bryology of New Zealand, with special reference to the herbarium of Robert Brown of Christchurch, New Zealand. *Bull. N.Z. Inst.*, No. 3, Pt. 1, 3–29.
- DIXON, H. N. 1914. Studies in the bryology of New Zealand, with special reference to the herbarium of Robert Brown of Christchurch, New Zealand. *Bull. N.Z. Inst.*, No. 3, Pt. 2, 31–74.
- DIXON, H. N. 1920. Contributions to Antarctic bryology; mosses of Deception Island. *Bryologist*, **23**, No. 5, 65–71.
- DIXON, H. N. 1921. Miscellanea bryologica — VII. *J. Bot., Lond.*, **59**, No. 701, 132–39.
- DIXON, H. N. 1923. Studies in the bryology of New Zealand with special reference to the herbarium of Robert Brown, of Christchurch, New Zealand. *Bull. N.Z. Inst.*, No. 3, Pt. 3, 75–152.
- DIXON, H. N. 1924a. Miscellanea bryologica — IX. *J. Bot., Lond.*, **62**, No. 740, 228–36.
- DIXON, H. N. 1924b. *The student's handbook of British mosses, with illustrations, and keys to the genera and species by H. G. Jameson*. 3rd edition. Eastbourne, Sumfield and Day Ltd.
- DIXON, H. N. 1926. Studies in the bryology of New Zealand, with special reference to the herbarium of Robert Brown, of Christchurch, New Zealand. *Bull. N.Z. Inst.*, No. 3, Pt. 4, 153–238.
- DIXON, H. N. 1927. Studies in the bryology of New Zealand, with special reference to the herbarium of Robert Brown, of Christchurch, New Zealand. *Bull. N.Z. Inst.*, No. 3, Pt. 5, 239–98.
- DIXON, H. N. 1929a. Studies in the bryology of New Zealand, with special reference to the herbarium of Robert Brown of Christchurch, New Zealand. *Bull. N.Z. Inst.*, No. 3, Pt. 6, 299–372.
- DIXON, H. N. 1929b. Critical mosses. *Revue Bryol.*, **N.S.**, **2**, Fasc. 1, 21–29.
- DIXON, H. N. 1932. Some mosses from South Georgia. *K. norske Vidensk. Selsk. Forh.*, **4**, No. 47, 179–81.
- DIXON, H. N. 1933. The nomenclature of the *Species muscorum*. *Revue bryol. lichen.*, **N.S.**, **6**, Fasc. 1–4, 93–115. [Effective date 30 April 1934.]
- DIXON, H. N. 1934. More mosses from South Georgia. *K. norske Vidensk. Selsk. Forh.*, **7**, No. 1, 1–3.
- DIXON, H. N. 1935. The mosses. (*In* MARR, J. W. S. The South Orkney Islands. 'Discovery' Rep., **10**, 381–82.)
- DIXON, H. N. and M. FLEISCHER. 1937. *Bryophyta nova* (26–29). *Annls bryol.*, **9** (for 1936), 139–41.
- DIXON, H. N. and W. W. WATTS. 1918. Mosses. *Scient. Rep. Australas. Antarct. Exped.*, Ser. C (Zoology and Botany), Vol. 7, Pt. 1, 5–7.
- DONAT, A. 1936a. Resultados fitogeográficos de la expedición patagónica "Gaea" 1933. I. Algunas briófitas nuevas de la Cordillera Central Patagónica. *Revta sudam. Bot.*, **3**, No. 3, 61–68.
- DONAT, A. 1936b. Bericht über die pflanzengeographischen Ergebnisse der Inlandeisexpedition der Argentinischen Geographischen Gesellschaft "Gaea" 1933. *Ber. dt. bot. Ges.*, **44**, Ht. 1, 27–46.
- DOZY, F. and J. H. MOLKENBOER. 1845. *Musci frondosi inediti archipelagi indici, sive descriptio et adumbratio muscorum frondosorum in insulis Java, Borneo, Sumatra, Celebes, Amboina, nec non in Japonia nuper detectorum minusve cognitorum*. Lugduni-Batavorum, H. W. Hazenberg and Soc., Fasc. 1, 1–24.
- DU BUYSSEN, R. 1883. Essai analytique du genre *Amblystegium*. *Mém. Soc. natn. Agric. Sci. Angers*, **N.S.**, **25**, 161–79.
- DUBY, J. E. 1868. Choix de cryptogames exotiques nouvelles ou mal connues. *Mém. Soc. Phys. Hist. nat. Genève*, **19**, Pt. 2, 291–304.
- DUBY, J. E. 1870. Choix de cryptogames exotiques nouvelles ou mal connues. *Mém. Soc. Phys. Hist. nat. Genève*, **20**, Pt. 2, 351–64.
- DUBY, J. E. 1875a. Choix de mousses exotiques nouvelles ou mal connues. *Mém. Soc. Phys. Hist. nat. Genève*, **24**, Pt. 1, 361–74.
- DUBY, J. E. 1875b. Diagnoses muscorum quorum descriptiones et icones Soc. Phys. et Hist. nat. Genevensi primo Aprili communicavit. *Flora, Jena*, **58**, (N.R., **33**), No. 18, 282–85.
- DUBY, J. E. 1880a. Choix de mousses exotiques nouvelles ou mal connues. *Mém. Soc. Phys. Hist. nat. Genève*, **27**, Pt. 1, 1–10.
- DUBY, J. E. 1880b. Aliquot diagnoses muscorum novorum aut non cognitorum. *Flora, Jena*, **63**, (N.R., **38**), No. 11, 168–74.
- DUMONT-D'URVILLE, M. J. 1825. Flore des Malouines. *Mém. Soc. linn. Paris*, **4** (for 1826), Pt. 2, 573–621.
- DUSÉN, P. 1903a. Patagonian and Fuegian mosses. *Rep. Princeton Univ. Exped. Patagonia*, **8**, Botany, No. 1, Pt. 3, 63–126.
- DUSÉN, P. 1903b. Beiträge zur Bryologie der Magellansländer, von Westpatagonien und Südchile. Teil I. *Ark. Bot.*, **1**, Häfte 1–3, 441–65.
- DUSÉN, P. 1905a. Beiträge zur Bryologie der Magellansländer, von Westpatagonien und

- Südchile. Teil II. *Ark. Bot.*, 4, Hälften 1–3, 1–45.
- DUSÉN, P. 1905b. *Musci nonnulli novi e Fuegia et Patagonia reportati. Bot. Notiser*, No. 7, 299–310.
- DUSÉN, P. 1905c. Beiträge zur Bryologie der Magellansländer, von Westpatagonien und Südchile. Teil III. *Ark. Bot.*, 4, Hälften 4, 1–24.
- DUSÉN, P. 1905d. *Mousses et hépatiques*. (In AUTRAN, E. Énumération des plantes récoltées par Miles Stuart Pennington pendant son premier voyage à la Terre de Feu en 1903. *Revta Univ. B. Aires*, 4, 297–98.)
- DUSÉN, P. 1906. Beiträge zur Bryologie der Magellansländer, von Westpatagonien und Südchile. Teil IV. *Ark. Bot.*, 6, Hälften 1–2, 1–40.
- DUSÉN, P. 1907. Beiträge zur Bryologie der Magellansländer von Westpatagonien und Südchile. Teil V. *Ark. Bot.*, 6, Hälften 3–4, 1–32.
- EATON, D. C. 1892. List of mosses from Fuegia and Patagonia. (In List of plants collected by the U.S.S. *Albatross* in 1887–91 along the western coast of America. *Contr. U.S. natn. Herb.*, 1, No. 5, 135–42.)
- ESPINOSA B., M. R. 1924. Enumeración de plantas colectadas en "Los Bronces". *Revta chil. Hist. nat.*, 28, 88–97.
- ESPINOSA, B., M.R. 1941. Lista sistemática de musgos y liquenes chilenos, continentales e insulares enviados por el Dr. C. Skottsberg. *Boln Mus. nac. Hist. nat. Chile*, 19, 113–32.
- FAMILLER, I. 1913. Die Laubmose Bayerns. Eine Zusammenstellung der bisher bekannt gewordenen Standortsangaben. *Denkschr. K. bayer. bot. Ges. Regensb.*, 12, (N.F., 6), 1–174.
- FELIPPONE, F. 1909. *Contribution à la flore bryologique de l'Uruguay, 1er fascicule*. Buenos Aires, Jean-A. Alsina.
- FELIPPONE, F. 1912. *Contribution à la flore bryologique de l'Uruguay, 2 ème fascicule*. Montevideo, Ateliers Typographiques "Gimenez".
- FELIPPONE, F. 1917. *Contributriion à la flore bryologique de l'Uruguay, 3 ème fascicule*. Montevideo, Mosca Hnos.
- FELIPPONE, F. 1928. Contribución à la flora briológica del Uruguay. *Revta chil. Hist. nat.*, 32, 83–88. [Effective date February 1929.]
- FELIPPONE, F. 1929. Contribution à la flore bryologique de l'Uruguay. *Revue bryol.*, N.S., 2, Fasc. 3–4, 210–25. [Effective date 31 March 1930, *fide* title page.]
- FELIPPONE, F. 1930. Contribution à la flore bryologique de l'Uruguay. Corrigenda. *Revue bryol.*, N.S., 3, Fasc. 1–2, 98.
- FIEDLER, C. 1844. *Synopsis der Laubmose Mecklenburg's*. Schwerin, C. Kürschner.
- FILSON, R. B. 1966. The lichens and mosses of Mac. Robertson Land. A.N.A.R.E. Rep., Ser. B 2, (Botany), No. 82, 169 pp.
- FLEISCHER, M. 1904. *Die Musci der Flora von Buitenzorg (zugleich Laubmoosflora von Java)*. Enthaltend alle aus Java bekannt gewordenen Sphagnales und Bryales, nebst kritischen Bemerkungen über vieler Archipelarten, sowie indischer und australischer Arten. Leiden, E. J. Brill, Bd. 1, 1–379.
- FLEISCHER, M. 1906. *Der Musci der Flora von Buitenzorg. Zugleich Laubmoosflora von Java mit Berücksichtigung aller Familien und Gattungen der gesamten Laubmooswelt*. Enthaltend alle aus Java bekannt gewordenen Sphagnales und Bryales nebst kritischen Bemerkungen über viele Archipelarten, sowie indische und australische Arten. Leiden, E. J. Brill, Bd. 3, Bogen 42, 645–60. [Effective date 1908, *fide* Stafleu and Cowan (1976, 844).]
- FLEISCHER, M. 1908a. Grundlagen zu einer Monographie der Gattung *Stereohypnum* (Hpe.). *Hedwigia*, 47, Ht. 5/6, 271–99.
- FLEISCHER, M. 1908b. *Die Musci der Flora von Buitenzorg. Zugleich Laubmoosflora von Java mit Berücksichtigung aller Familien und Gattungen der gesamten Laubmooswelt*. Enthaltend alle aus Java bekannt gewordenen Sphagnales und Bryales nebst kritischen Bemerkungen über viele Archipelarten, sowie indische und australische Arten. Leiden, E. J. Brill, Bd. 3, Bogen 59, 917–32.
- FLEISCHER, M. 1914. Kritische Revision von Carl Müllerschen Laubmoosgattungen. I. *Hedwigia*, 55, Ht. 3, 280–85.
- FLEISCHER, M. 1917. Kritische Revision von Carl Müllerschen Laubmoosgattungen. II. *Hedwigia*, 59, Ht. 5, 212–19.
- FLEISCHER, M. 1920. Kritische Revision von Carl Müllerschen Laubmoosgattungen. III. *Hedwigia*, 61, Ht. 6, 402–08.
- FLEISCHER, M. 1922. Kritische Revision der Carl Müllerschen Laubmoosgattungen. IV. *Hedwigia*, 63, Ht. 3/4, 209–16.
- FLEISCHER, M. 1923. *Die Musci der Flora von Buitenzorg. Zugleich Laubmoosflora von Java mit Berücksichtigung aller Familien und Gattungen der gesamten Laubmooswelt*. Enthaltend alle aus Java bekanntgewordenen Sphagnales und Bryales nebst kritischen Bemerkungen über viele Archipelarten, sowie indische und australische Arten. Leiden, E. J. Brill, Bd. 4, 1105–729.
- FLOWERS, S. 1935. Bartramiaceae. (In GROUT, A. J., ed. *Moss flora of North America north of Mexico*. Newfane, Vermont, A. J. Grout, 2, Pt. 3, 152–83.)
- FLOWERS, S. 1952. Monograph of the genus *Anacolia*. *Bull. Torrey bot. Club*, 79, No. 2, 161–85.
- FRAHM, J. P. 1974. Zur Unterscheidung und Verbreitung von *Campylopus introflexus* (Hedw.) Brid. und *C. polytrichoides* De Not. *Revue bryol. lichen.*, N.S., 40, Fasc. 1, 33–44.
- FRAHM, J. P. 1975a. Conspectus der mittel- und südamerikanischen *Campylopus* — Arten (Dicranaceae). *Bryophytorum Bibliotheca*, 5, 1–143.
- FRAHM, J. P. 1975b. Taxonomische Notizen zur Gattung *Campylopus*. *Revue bryol. lichen.*, N.S., 41, Fasc. 3, 321–32.
- FRAHM, J. P. 1976a. Zur *Campylopus* — Flora von Chile. *Herzogia*, 4, Ht. 1–2, 141–60.
- FRAHM, J. P. 1976b. Taxonomische Notizen zur Gattung *Campylopus* II. *Revue bryol. lichen.*, N.S., 42, Fasc. 1, 603–16.
- FRIES, E. M. 1825. Musci. (In *Stirpes agri Fernionensis, quarum indicem cons. ampliss. phil. ord. Lundens. pp. J. G. Arrhenius, Vermelandus, in acad. Carol. D. XI Junii MDCCXXV. Continatio I. Londoni Gothorum, Officina Berliniana, 24–30.)*
- FROEHLICH, J. 1953. Zwei neue Laubmose aus Süd-Chile und Feuerland. *Annln naturh. Mus. Wien*, 59 (for 1952–53), 117–18.
- FRYE, T. C. and M. W. DUCKERING. 1948. *Atrichum polycarpum*. *Bryologist*, 51, No. 3, 170–74.
- FUNCK, H. C. 1802. Botanische Notizen. *Bot. Ztg. Erlangen*, 1, Ht. 2, No. 8, 120–24.
- FUNCK, H. C. 1820. *Deutschlands Moose. Ein*

- Taschenherbarium zum Gebrauch auf botanischen Excursionen.* Baiereuth, F. C. Birner.
- FÜRNROHR, A. E. 1827. Versuch einer näheren Beleuchtung der Disposition methodique des espèces de mousses par G. A. Walker-Arnott, M. Membre des Sociétés royale et Wernerienne d'Edinbourg etc., Paris, 1825 (4.72 seiten). *Flora, Jena*, **10**, (Bd. 2, Beil. 1), 22–86.
- FÜRNROHR, A. E. 1829. Sam. El. a Bridel-Brideri, a consil secret legation Saxo-Gothanae etc. *Bryologia universa seu systematica ad novam methodum dispositio, historia et descriptio omnium muscorum hucusque cognitorum cum synonymia ex auctoribus probatissimis.* Accedunt tabulae aeneae tredecim. Lipsiae sumtibus Joan. Ambros. Barth. 8 max. Vol. primum 1926, XLVI et 856 pag. Vol. secundum 1827, 848 pag. *Flora, Jena*, **12**, (Bd. 2, Ergänzungsblätter), 1–60.
- GAROVAGLIO, S. 1840. *Bryologia austriaca excursionia ta tamquam clavis analytica ad omnes in imperio austriaco hucusque inventos muscos facile et tuto determinandos.* Vindobonae, Friderici Volke.
- GÄRTNER, G., MEYER, B. and J. SCHERBIUS. 1802. Ordnung Musci, Moose. (*In Oekonomisch-technische Flora der Wetterau. Band 3, Abt. 2.* Frankfurt am Main, Philipp Heinrich Guilhauman, 68–139.)
- GAUDICHAUD, [M.] C. 1825. Rapport sur la flore des îles Malouines. *Annls sci. nat., Botanique*, **5**, 89–110.
- GAUDICHAUD, M. C. 1826. Botanique, Musci, Jussieu. (*In FREYCINET, M. L. DE. Voyage autour du monde, entrepris par ordre du roi, sous le ministère et conformément aux instructions de s. exc. M. le vicomte du Bouchage, secrétaire d'état au département de la marine, exécuté sur les corvettes de S. M. l'Uranie et la Physicienne, pendant les années 1817, 1818, 1819 et 1820.* Paris, Chez Pillot Aîné, Pt. 6, 221–30.) [Effective date February 1828, *fide* Stafleu and Cowan, 1976, 922.]
- GENTH, F. F. 1835. Flora des Herzogthums Nassau und der obern, so wie der untern Rheingegenden von speier bei Cöln Von F. F. Genth. I. Theil Cryptogamie. I. Abtheilung. Farmkräuter, Lebermoose, Moose und Flechten. Im Verlage von Florian Kupferberg. Mainz. 1836. 8. *LitBer. Flora allg. bot. Zeit.*, **5**, No. 12, 177–91.
- GEPP, A. 1902. Cryptogamia: Musci. (*In Report on the collections of natural history made in Antarctic regions during the voyage of the "Southern Cross".* Vol. 21. London, The Trustees, British Museum (Nat. Hist.), 319.)
- GIACOMINI, V. 1955. Sull'autonomia specifica e sul ciclo di forme di *Campylopus polytrichoides* De Not. *Atti Ist. bot. Univ. Lab. critogam.* Pavia, Ser. 5, **13**, Fasc. 1, 45–83.
- GIBERT, E. 1873. Musci. (*In Enumeratio plantarum sponte nascentium agro Montevidensi cum synonymis selectis.* Montevideo, La Asociación Rural del Uruguay, 129–32.)
- GODMAN, F. Du C. 1870. *Natural history of the Azores or western islands.* London, John von Voerst.
- GRADSTEIN, S. R. and H. J. M. SIPMAN. 1978. Taxonomy and world distribution of *Campylopus introflexus* and *C. pilifer* (—*C. polytrichoides*): a new synthesis. *Bryologist*, **81**, No. 1, 114–21.
- GRASSI, M. M. 1975a. Notas de clase Bryophyta. I. Parte general. *Fundación Miguel Lillo.*
- Miscelánea, No. 55, 1–38.
- GRASSI, M. M. 1975b. Notas de clase Bryophyta. (Atlas). *Fundación Miguel Lillo. Miscelánea*, No. 55, 1–57.
- GRASSI, M. M. 1976. Notas de clase Bryophyta. II. Parte especial. *Fundación Miguel Lillo. Miscelánea*, No. 55, 1–320.
- GRAY, S. F. 1821. *A natural arrangement of British plants, according to their relations to each other, as pointed out by Jussieu, de Candolle, Brown, etc. including those cultivated for use; with an introduction to botany, in which the terms newly introduced are explained; illustrated by figures.* Vol. 1. London, Baldwin, Cradock and Joy.
- GREENE, D. M. 1972. The herbarium of the British Antarctic Survey. *British Antarctic Survey Bulletin*, No. 31, 107–09.
- GREENE, S. W. 1964. Problems and progress in Antarctic bryology. (*In CARRICK, R., HOLDGATE, M. and J. PRÉVOST, ed. Biologie antarctique.* Paris, Hermann, 173–79.)
- GREENE, S. W. 1967. Bryophyte distribution. (*In BUSHNELL, V., ed. Terrestrial life in Antarctica. Antarct. Map Folio Ser., Folio 5*, 11–13.)
- GREENE, S. W. 1968a. Studies in Antarctic bryology: I. A basic check list for mosses. *Revue bryol. lichen.*, N.S., **36**, Fasc. 1–2, 132–38.
- GREENE, S. W. 1968b. Studies in Antarctic bryology: II. *Andreaea, Neuroloma.* *Revue bryol. lichen.*, N.S., **36**, Fasc. 1–2, 139–46.
- GREENE, S. W. 1972. A new combination in *Andreaea.* *British Antarctic Survey Bulletin*, No. 31, 109. [Effective date 17 May 1973.]
- GREENE, S. W. 1973. A synoptic flora of South Georgian mosses: I. *Dendroligotrichum, Polytrichum* and *Psilotilum.* *British Antarctic Survey Bulletin*, No. 36, 1–32. [Effective date 21 June 1974.]
- GREENE, S. W. 1975. The Antarctic moss *Sarconeurum glaciale* (C. Muell.) Card. et Bryhn in southern South America. *British Antarctic Survey Bulletin*, Nos. 41 and 42, 187–91. [Effective date 4 February 1976.]
- GREENE, S. W., GREENE, D. M., BROWN, P. D. and J. M. PACEY. 1970. Antarctic moss flora: I. The genera *Andreaea, Pohlia, Polytrichum, Psilotilum* and *Sarconeurum.* *British Antarctic Survey Scientific Reports*, No. 64, 118 pp.
- GREVILLE, R. K. 1824. *Flora Edinensis: or a description of plants growing near Edinburgh, arranged according to the Linnean system. With a concise introduction to the material orders of the class Cryptogamia, and illustrative plates.* Edinburgh, William Blackwood; London, T. Cadell.
- GREVILLE, R. K. 1826. Descriptions of two new species of Musci, belonging to the genera *Neckera* and *Hypnum.* *Mem. Wernerian nat. Hist. Soc.*, **5**, Pt. 2 (for 1824–25), 481–84.
- GREVILLE, R. K. 1830. Descriptions of some new mosses discovered in South America by Dr. Gillies; with a plate. *Edinb. J. nat. geogr. Sci.*, **2**, 1–4.
- GREVILLE, R. K. and G. A. W. Arnott. 1824. IV. *Tentamen methodi muscorum; or a new arrangement of the genera of mosses, with characters, and observations on their distribution, history and structure. Mem. Wernerian nat. Hist. Soc.*, **5**, Pt. 1 (for 1823–24), 42–89.
- GROUT, A. J. 1929. *Check list of the pleurocarpous mosses of North America, north of Mexico.* New Brighton, New York, A. J. Grout. [The only copy seen lacked a date, so that given in Index]

- muscorum*, 2, p. 172, under *Drepanocladus uncinatus* var. *symmetricus* has been accepted but not verified.]
- GROUT, A. J. 1939. Pottieae with *Aloina* by E. J. Craig; *Tortula* by W. C. Steere; *Pottia* and *Pterygoneurum* by W. T. Wareham. (*In* GROUT, A. J., ed. *Moss flora of North America north of Mexico*, Newfane, Vermont, A. J. Grout, 1, Pt. 4, 193–253.)
- GROUT, A. J. 1944. Preliminary synopsis of the North American Macromitriae. *Bryologist*, 47, No. 1, 1–22.
- GROUT, A. J. 1946. *North American flora*. New York, New York Botanical Garden, 15A, Pt. 1, 1–62.
- GUNCKEL, H. 1938. Los *Porothamnium* de la flora Corraleña. *Revta chil. Hist. nat.*, 42, 118–22. [Effective date 1939.]
- GUNCKEL, H. 1964. Un interesante musgo de la Selva Valdiviana: *Eriodon conostomus* Montagne. *Revta univ., Santiago*, 49 [An. Acad. chil. Cienc. nat., No. 27], 67–70. [Effective date 1965.]
- GUNCKEL, H. 1971. Las primeras plantas herborizadas en Chile en 1690. *An. Inst. Patagonia*, 2, Nos. 1–2, 134–41.
- HAGEN, I. 1915. Forarbejder til en Norsk Løvmosflora. XX. Dicranaceae. *K. norske Vidensk. Selsk. Skr.*, (for 1914), Nr. 1, 1–192.
- HAGEN, I. and H. PRINTZ. 1929. Forabeider til en Norsk Løvmosflora. XXI. Pottiaceae. *K. norske Vidensk. Selsk. Skr.*, (for 1928), Nr. 3, 1–96.
- HAMPE, E. 1837a. Musci frondosi Germaniae ad methodum naturalem dispositi. *Flora, Jena*, 20, No. 18, 273–87.
- HAMPE, E. 1837b. *Prodromus florae Hercyniae. Linnaea*, Halle, 11, Ht. 1, 17–105.
- HAMPE, E. 1837c. *Weberae novae chilensis descriptio. Linnaea*, Halle, 11, Ht. 3, 278.
- HAMPE, E. 1839. Relation über die von dem Reisenden C. Beyrich auf seiner letzten Reise in Nordamerika gesammelten Laubmose. *Linnaea*, Halle, 13, Ht. 1, 39–48.
- HAMPE, E. 1844. *Icones muscorum novorum vel minus cognitorum. Decades 1–3*. Bonnae, Henry et Cohen.
- HAMPE, E. 1846. Ueber *Hyophila javanica* Brid. *Bot. Ztg*, 4, No. 15, 266–69.
- HAMPE, E. 1847. Ein Referat über die Columbischen Moose, welche von Herrn Moritz gesammelt wurden. *Linnaea*, Halle, 20, [Beitr. *PflKunde*, 4], Ht. 1, 65–98.
- HAMPE, E. 1860. *Muscorum frondosorum florae Australasiae auctore Dr. F. Müller mox edendae species novas. Linnaea*, Halle, 30, [Beitr. *PflKunde*, 14], Ht. 5, 623–40; Ht. 6, 641–46.
- HAMPE, E. 1862a. Species novas muscorum ab Dr. A. Alexandro Lindigio in Nova-Granada mensibus Julio et Augusto a 1861 collectas. *Linnaea*, Halle, 31, [Beitr. *Pflkunde*, 15], Ht. 5, 518–32.
- HAMPE, E. 1862b. Beitrag zu einer Moosflora von Neu-Granada. *Flora, Jena*, 45, (N.R., 20), No. 29, 449–58.
- HAMPE, E. 1863. Species novas muscorum ab Alexandro Lindigio in Nova-Granada collectas. *Linnaea*, Halle, 32, [Beitr. *PflKunde*, 16], Ht. 1, 127–28; Ht. 2, 129–64.
- HAMPE, E. 1865. Musci. (*In* TRIANA, J. and J. E. PLANCHON. *Prodromus florae Novo-Granatensis ou énumération des plantes de la Nouvelle-Grenade avec description des espèces nouvelles. Annls Sci. nat.*, Sér. 5, 4, Botanique, 324–78.)
- HAMPE, E. 1866. Musci. (*In* TRIANA, J. and J. E. PLANCHON. *Prodromus florae Novo-Granatensis ou énumération des plantes de la Nouvelle-Grenade avec description des espèces nouvelles. Annls Sci. nat.*, Sér. 5, 5, Botanique, 300–42.)
- HAMPE, E. 1867a. *Bryologische Mittheilungen aus dem Herbarium. Flora, Jena*, 50, (N.R., 25), No. 5, 65–80.
- HAMPE, E. 1867b. *Ditrichum Timm*, statt *Leptotrichum* Hpe. *Flora, Jena*, 50, (N.R., 25), No. 12, 181–82.
- HAMPE, E. 1870. Musci frondosi. (*In* WARMING, E., ed. *Symbolae ad floram Brasiliæ centralis cognoscendam, Particula VIII. Vidensk. Meddr dansk naturh. Foren.*, Ser. 3, Nos. 18–20, 267–96.)
- HAMPE, E. 1872. Musci frondosi a clar. Dr. A. Glaziou in vicinia urbis Rio de Janeiro lecti. (*In* WARMING, E., ed. *Symbolae ad floram Brasiliæ centralis cognoscendam. Pt. 10. Vidensk. Meddr dansk naturh. Foren.*, Ser. 3, 4, Nr. 1–5, 36–59.)
- HAMPE, E. 1874. Musci frondosi a clar. Dr. A. Glaziou in vicinia urbis Rio de Janeiro lecti. (*In* WARMING, E., ed. *Symbolae ad floram Brasiliæ centralis cognoscendam. Particula 19. Vidensk. Meddr dansk naturh. Foren.*, Ser. 3, Nos. 9–11, 129–78.) [Effective date 1875, according to entry for *Rhodobryum gracilescens* in *Index muscorum*, 4, p. 148.]
- HAMPE, E. 1878. Musci frondosi, a clar. Dr. A. Glaziou in vicinia urbis Rio Janeiro lecti (continuatio). (*In* WARMING, E., ed. *Symbolae ad floram Brasiliæ centralis cognoscendam, Particula XXIV. Vidensk. Meddr dansk naturh. Foren.*, Ser. 3 (for 1877–78), 251–74.)
- HAMPE, E. 1879. *Enumeratio muscorum frondosorum Brasiliæ centralis, praecipue provinciarum Rio de Janeiro et S. Paulo, adhuc cognitorum. Vidensk. Meddr dansk naturh. Foren.*, Ser. 4 (for 1879–80), 73–164.
- HAMPE, E. 1880a. Choix de mousses exotiques, nouvelles ou mal connues, par J. E. Duby, ancien pasteur, Docteur de Sciences communiq. le 5 Febr. 1880. *Flora, Jena*, 63, (N.R., 38), No. 21, 332–36.
- HAMPE, E. 1880b. *Supplementum ad volumen undecimum fragmentorum phytographie Australiae, indices plantarum acotyledonearum complectens. III. Musci frondosi Australiae continentalis, praesertim e baronis de Mueller collectionibus. (In MUELLER, F. *Fragmenta phytographie Australiae*, Vol. XI. Melbourne, Joannis Ferres, 45–52.)*
- [HAMPE, E.] 1880c. Additmenta pro supplemento voluminis undecimi. (*In* MUELLER, F. *Fragmenta phytographie Australiae*, Vol. XI. Melbourne, Joannis Ferres, 105–21.) [Unsigned and presumed to be by Hampe who worked up the mosses.]
- HAMPE, E. and A. GEHEEB. 1881. Additmenta ad "Enumerationem muscorum hactenus in provinciis Brasiliensibus Rio de Janeiro et Sao Paulo detectorum". *Flora, Jena*, 64, (N.R., 39), No. 26, 401–16.
- HARIOT, P. 1891. Contribution à la flore cryptogamique de la Terre de Feu. *Bull. Soc. bot. Fr.*, 38, [Sér. 2, 13], 416–22.
- HARTMAN, C. J. 1838. Musci. (*In* Handbok i Skandinaviens Flora, innehållande Sveriges och Norriges vexter, till och med Mossorona; med ett utkast till botanologien eller vextlären i allmänhet. Vol. 2. 3rd edition. Stockholm, Zacharias

- Haeggström, 260–334.)
- HARTMAN, C. J. 1849. Musci. (*In Handbok i Skandinaviens Flora, innehållande Sveriges och Norriges vexter till och med Mossorna; ordnade efter Prof. Fries' system.* 5th edition. Stockholm, Zacharias Haeggström, 319–431.)
- HASSKARL, J. K. 1851. *Allgemeines Sach- und Namen-Register zu den ersten XXV Jahrgängen der Flora oder allgemeinen botanischen Zeitung von 1818–1842.* Regensburg, G. Joseph Manz. [Note this is general index to *Flora*, Vols 1–25, 1818–42.]
- HEDWIG, D. J. 1782. *Fundamentum historiae naturalis muscorum frondosorum concernens eorum flores, fructus, seminalem propagationem adiecta generum dispositione methodica, iconibus illustratis. Part 2.* Lipsiae, Siegfried Lebrecht Crusium.
- HEDWIG, J. 1801. *Species muscorum frondosorum. Descriptae et tabulis aeneis LXVII coloratis illustratae. Opus posthumum editum a Friderico Schwaegrichen.* Lipsiae, Joannis Ambrosii Barthii; Parisiis, Amand Koenig. [Reprinted: *Species muscorum frondosorum.* (*In CRAMER, J. and H. K. SWANN, ed. Historiae naturalis classica, 16,* 1960. Weinheim/Bergstr., H. R. Engelmann (J. Cramer); Codicote/Herts., Wheldon & Wesley, Ltd.; New York, Hafner Publishing Co.)]
- HERTER, [W.] G. 1927–28. Index montevidensis, plantae avasculares. (*In HERTER, [W.] G., ed. Estudios botánicos en la región Uruguaya.* No. 1. Montevideo, Asociación Rural del Uruguay, 36 pp.)
- HERTER, [W.] G. 1928. Las plantas uruguayas de Ernesto Gibert. Nomenclator Gibertianus. (*In HERTER, [W.] G., ed. Estudios botánicos en la región Uruguaya.* No. V. Montevideo, Universitatia rei Publicae, 60 pp.)
- HERTER, [W.] G. 1933a. *Florula uruguayensis, plantae avasculares.* (*In HERTER, [W.] G., ed. Ostenia, colección de trabajos botánicos dedicados a Don Cornelio Osten en ocasión del LXX aniversario de su nacimiento. Estudios botánicos en la región Uruguaya* No. XIII. Montevideo, [W. G. Herter], 7–84.) [Re-issued without change of pagination in separate covers entitled *Florula uruguayensis, plantae avasculares. Estudios botánicos en la región Uruguaya* No. III. Montevideo, W. G. Herter.]
- HERTER, [W.] G. 1933b. Las plantas uruguayas de Ernesto Gibert. Nomenclator Gibertianus 2. (*In HERTER, [W.] G., ed. Ostenia, colección de trabajos botánicos dedicados a Don Cornelio Osten en ocasión del LXX aniversario de su nacimiento. Estudios botánicos en la región Uruguaya* No. XIII. Montevideo, [W. G. Herter], 153–71.) [Re-issued without change of pagination in separate covers entitled *Las plantas uruguayas de Ernesto Gibert, segunda parte. Nomenclator Gibertianus II. Estudios botánicos en la región Uruguaya* No. Va.]
- HERTER, W. G. 1937. Plantae uruguayenses novae vel criticae I. *Revta sudam. Bot.*, 5, Nos. 1/2, 13–36. [Part of *Estudios botánicos en la región Uruguaya* No. XVI, *fide* Herter, 1953.]
- HERTER, W. G. 1939a. Zur Kenntnis der Flora von Osturuguay. *Beih. bot. Zbl.*, 59, Abt. B, Ht. 1, 245–86.
- HERTER, W. G. 1939b. Plantae uruguayenses novae vel criticae II. *Revta sudam. Bot.*, 6, Nos. 3/4, 69–107. [Effective date September 1941. Part of *Estudios botánicos en la región Uruguaya* No. XVI, *fide* Herter, 1953.]
- HERTER, W. G. 1943. Plantae uruguayenses novae vel criticae IV. *Revta sudam. Bot.*, 7, Nos. 6/8, 171–260. [Part of *Estudios botánicos en la región Uruguaya* No. XVI, *fide* Herter, 1953.]
- HERTER, [W.] G. 1953. Conspectus estudios botánicos en la región Uruguaya. *Revta sudam. Bot.*, 10, No. 3, 99–100.
- HERZOG, T. 1909. Beiträge zur Laubmoosflora von Bolivia. *Beih. bot. Zbl.*, 26 (for 1909–10), Abt. 2, Ht. 1, 45–102.
- HERZOG, T. 1911. Bemerkungen zu der neuren Laubmoos-Gattung *Wollyna*. *Beih. bot. Zbl.*, 28, Abt. 2, Ht. 2, 268–71.
- HERZOG, T. 1914. Zwei kleistokarpe Moose der bolivianischen Hochcordillere. *Flora, Jena, N.F.*, 7, Ht. 3, 317–26.
- HERZOG, T. 1916a. Die Bryophyten meiner zweiten Reise durch Bolivia. Lief. 1. *Biblthca bot.*, Ht. 87, 1–168.
- HERZOG, T. 1916b. Neue Laubmose aus Ostasien und Südamerika. *Hedwigia*, 57, Ht. 3, 233–50.
- HERZOG, T. 1920. Die Bryophyten meiner zweiten Reise durch Bolivia Nachtrag. *Biblthca bot.*, Ht. 88, 1–31.
- HERZOG, T. 1923. Beiträge zur Bryophytenflora von Chiles. *Hedwigia*, 64, Ht. 1/2, 1–18.
- HERZOG, T. 1926. Beiträge zur Moosflora Westpatagoniens. *Hedwigia*, 66, Ht. 2, 79–92.
- HERZOG, T. 1927. Zwei Bryophytensammlungen aus Südamerika. *Hedwigia*, 67, Ht. 6, 249–68.
- HERZOG, T. 1928. Drei neue Laubmoosgattungen. *Revue bryol.*, N.S., 1, Nos. 2–3, 98–108.
- HERZOG, T. 1937. Neue bryophyten von Ruwenzori und aus dem patagonischen Inlandesgebiet. *Reprium nov. spec. Regni veg.*, 41, Nos. 14–25, 285–92.
- HERZOG, T. 1938. Contribución al conocimiento de la flora Briofita del sur de Chile. a. Parte sistemática. *Archos Esc. Farm. Córdoba*, No. 7 (Sección científica), 3–56.
- HERZOG, T. 1939. I. Verzeichnis der gesammelten Bryophyten. (*In HERZOG, T., SCHWABE, G. H. and E. SCHWABE. Zur Bryophytenflora südchiles.* *Beih. bot. Zbl.*, 60, Abt. B, Ht. 1/2, 1–51.)
- HERZOG, T. 1940. Die Moose der Expedition Ljungner nach Patagonien, 1932/34. *Ark. Bot.*, 29A, Häfte 4, 1–17.
- HERZOG, T. 1952a. Beiträge zur Kenntnis der argentinischen Bryophytenflora. *Reprium nov. spec. Regni veg.*, 55, Ht. 1, 1–27.
- HERZOG, T. 1952b. Miscellanea bryologica. III. *Memo. Soc. Fauna Flora fenn.*, 27 (for 1950–51), 92–110. [Effective date 1952, according to footnotes on p. 80, 96 and 112.]
- HERZOG, T. 1952c. Eine neue Lebermoosgattung aus Westpatagonien, *Chondrophyllum* Herz. *Revue bryol. lichen.*, N.S., 21, Fasc. 1–2, 46–49.
- HERZOG, T. 1954. Zur Bryophytenflora chiles. *Revue bryol. lichen.*, N.S., 23, Fasc. 1–2, 27–99.
- HERZOG, T. 1957. Eine Briophytensammlung aus dem argentinischen Nationalpark Nahuelhuapi. *Darwiniana*, 11, No. 2, 207–22.
- HERZOG, T. 1960. Weitere Beiträge zur Bryophytenflora von Chile. *Revue bryol. lichen.*, N.S., 29, Fasc. 3–4, 183–206. [Effective date March 1961, *fide* Lamy, 1978.]
- HERZOG, T., THÉRIOT, I., BUCH, H., SCHIFFNER, V. and F. VERDOORN. 1933. Bryophyta nova (6–16). *Annls bryol.*, 6, 124–35.
- HILPERT, F. 1933. Studien zur Systematik der Trichostomaceen. *Beih. bot. Zbl.*, 50, Abt. 2, Ht. 2, 585–640; Ht. 3, 641–706.
- HOFFMAN, G. F. 1795. *Deutschlands Flora oder*

- botanisches Taschenbuch. Theil 2. Cryptogamie.* Erlangen, Johann Jacob Palm. [Effective date 1796, *fide* Stafleu and Cowan, 1979, 240.]
- HOLZINGER, J. M. 1911. Reviews of current literature. [Cardot, J. 1907. *Musci. (In National Antarctic Expedition, 1901–1904. Natural History, Vol. IV, Zoology and Botany. London, British Museum, 1–6.) Bryologist, 14, No. 1, 11–12.*]
- HOOKER, J. D. 1867. Order IV, *Musci. (In Handbook of the New Zealand flora: a systematic description of the native plants of New Zealand and the Chatham, Kermadec's, Lord Auckland's, Campbell's, and Macquarie's Islands. London, Reeve & Co., Pt. 2, 393–497.)*
- HOOKER, J. D. and W. WILSON. 1844. *Musci Antartici*, being characters with brief descriptions of the new species of mosses discovered during the voyage of H.M. Discovery Ships, *Erebus* and *Terror*, in the southern circumpolar regions, together with those of Tasmania and New Zealand. *Lond. J. Bot.*, 3, 533–56.
- HOOKER, W. J. 1816. *Plantae cryptogamicae, quas in plaga orbis novi aequinoctiali collegerunt Alexander de Humboldt et Amat. Bonpland, adjectis tabulis species quasdam novas minusve cognitas exhibentibus necnon Alexandri de Humboldt notationibus quibusdam plantarum geographiam spectantibus.* London, W. J. Hooker.
- HOOKER, W. J. 1818. *Musci exotici; containing figures and descriptions of new or little known foreign mosses and other cryptogamic subjects.* Vol. 1. London, Longman, Hurst, Rees, Orme and Brown.
- HOOKER, W. J. 1819. *Musci exotici; containing figures and descriptions of new or little known foreign mosses and other cryptogamic subjects.* Vol. 2. London, Longman, Hurst, Rees, Orme and Brown.
- HOOKER, W. J. 1820. *Musci exotici; containing figures and descriptions of new or little known foreign mosses and other cryptogamic subjects.* Vol. 2. London, Longman, Hurst, Rees, Orme and Brown.
- HOOKER, W. J. 1829. *Bryum gilliesii. Botanical miscellany containing figures and descriptions of such plants as recommend themselves by their novelty, rarity or history or by the uses to which they are applied in the arts, in medicine, and in domestic economy together with occasional botanical notices and information.* London, John Murray, 1 (for 1830), Pt. 1, 3–4, pl. 2.
- HOOKER, W. J. 1830. *Gymnostomum amblophyllum. Botanical miscellany containing figures and descriptions of such plants as recommend themselves by their novelty, rarity or history or by the uses to which they are applied in the arts, in medicine, and in domestic economy together with occasional botanical notices and information.* London, John Murray, 1 (for 1830), Pt. 3, 352–53, pl. 75.
- HOOKER, W. J. 1833. *Cryptogamia: Musci. (In SMITH, J. E. The English flora, Vol. 5 (or Vol. 2 of Dr. Hooker's British flora), Part 1, comprising the mosses, Hepaticae, lichens, Characeae and Algae.* London, Longman, Rees, Orme, Brown, Green and Longman, 1–96.]
- HOOKER, W. J. 1836. *Bryum (Pohlia) pellucens. Hooker's Icon. Pl.*, 1, Tab. 34.
- HOOKER, W. J. 1837. *Dicranum jamesoni. Hooker's Icon. Pl.*, 2, Tab. 179.
- HOOKER, W. J. 1839. *Andreaea subulata* Harv. mst. *Hooker's Icon. Pl.*, 3, Tab. 201.
- HOOKER, W. J. 1840. *Bruchia brevipes* Harv. *Hooker's Icon. Pl.*, 3, Tab. 231.
- HOOKER, W. J. and G. A. W. ARNOTT. 1832. Order LXXIV. *Musci. (In The botany of Captain Beechey's voyage comprising an account of the plants collected by Messrs Lay and Collie and other officers of the expedition, during the voyage to the Pacific and Bering's Strait, performed in His Majesty's ship Blossom, under the command of Captain F. W. Beechey, R. N., F. R., & A. S., in the years 1825, 26, 27 and 28. London, Henry G. Bohn, Pt. 2, 53–54.)*
- HOOKER, W. J. and R. K. GREVILLE. 1824a. Sketch of the characters of the species of mosses, belonging to the genera *Orthotrichum* (including *Schlottheimia*, *Micromitrium* and *Ulota*), *Glyphomitrium*, and *Zygodon*. *Edinb. J. Sci.*, 1, No. 1, 110–33.
- HOOKER, W. J. and R. K. GREVILLE. 1824b. On the genus *Tortula* of the order *Musci*. *Edinb. J. Sci.*, 1, No. 2, 287–302.
- HOOKER, W. J. and R. K. GREVILLE. 1825a. On the genus *Hookeria* of Smith, of the order *Musci*. *Edinb. J. Sci.*, 2, No. 2, 221–36.
- HOOKER, W. J. and R. K. GREVILLE. 1825b. On the genus *Calymperes* of Swartz and *Syrrhopodon* of Schwaegrichen of the order *Musci*. *Edinb. J. Sci.*, 3, No. 2, 218–27.
- HOOKER, W. J. and T. TAYLOR. 1818. *Muscologia Britannica; containing the mosses of Great Britain and Ireland systematically arranged and described; with plates illustrative of the characters of the genera and species.* London, Longman, Hurst, Rees, Orme and Brown.
- HOOKER, W. J. and W. WILSON. 1844. Enumeration of the mosses and Hepaticae, collected in Brazil by George Gardner, Esq. *Lond. J. Bot.*, 3, 149–67.
- HOPPE, D. H. 1801. Nachtrag zu Abhandlung über die in Deutschland nachsenden Arten des Widerthons (*Polytrichum*). *Bot. Taschenb.*, 1801, 196–99.
- HORIKAWA, Y. 1961. Mosses from the Ongul Islands and its neighbouring areas (preliminary report). *Antarctic Rec.*, 11, No. 1, 876–78.
- HORIKAWA, Y. and H. ANDO. 1961. Mosses of the Ongul Islands collected during the 1957–1960 Japanese Antarctic Research Expedition. *Hikobia*, 2, No. 3, 160–78.
- HORIKAWA, Y. and H. ANDO. 1963. A review of the Antarctic species of *Ceratodon* described by Cardot. *Hikobia*, 3, No. 4, 275–80.
- HORIKAWA, Y. and H. ANDO. 1967. The mosses of the Ongul Islands and adjoining coastal areas of the Antarctic continent. *JARE Sci. Rep., Special issue*, No. 1, 245–52.
- HORNSCHUCH, [C.] F. 1820a. *Musci frondosi exotici herbarii Willdenowiani tum capensis A. B. Bergio lecti tum alii quidam ex australasiae alliisque orbis terrauei plagis AL. a Chamissoe relati. Descripsit et iconibus novarum maximeque memorabilium specierum illustravit. (In NEES AB ESENBECK, C. G., ed. *Horae physicae Berolinenses collectae ex symbolis virorum doctorum*. Bonnae, Adolphi Marcus, 57–68.)*
- HORNSCHUCH, [C. F.] 1820b. Ueber die von Chamisso und Bergius gesammelten Moose. *Flora, Jena*, 3, No. 33, 511–22.
- HORNSCHUCH, C. F. 1840. *Musci. (In DE MARTIUS, C. F. P., ed. *Flora Brasiliensis, enumeratio plantarum in Brasilia hactenus detectarum quas**

- suis aliorumque botanicorum studiis descriptas et methodo naturali digestas partim iconē illustratas. Vol. I. Monachii, Lipsiae apud Frid. Fleischner, Pt. 2, 2–100.)*
- HORNSCHUCH, [C.] F. 1841. Muscorum frondosorum novorum, quos in Africa Australiori colegerunt Ecklon, drēge mundt et maire, descriptiones. *Linnaea*, Halle, 15, Ht. 1, 113–57.
- HORRELL, E. C. 1900. The European Sphagnaceae (after Warnstorff). *J. Bot., Lond.*, 38, No. 12, 469–80.
- HOSSEUS, C. C. 1934. Bemerkungen zum Vorkommen einiger Südamerikanischer Fabroniaceen. *Reprint nov. spec. Regni veg.*, 35, Nos. 8–17, 174–76.
- HOSSEUS, C. C. 1935a. Beiträge zur Kenntnis der Laubmoose der Provinz Cordoba (Argentinien). I. *Reprint nov. spec. Regni veg.*, 38, Nos. 6–12, 131–36.
- HOSSEUS, C. C. 1935b. Beiträge zur Kenntnis der Laubmoose der Provinz Cordoba (Argentinien). II. *Reprint nov. spec. Regni veg.*, 38, Nos. 13–25, 209–18.
- HOSSEUS, C. C. 1936. Beiträge zur Kenntnis der Laubmoose der Provinz Cordoba (Argentinien). III. *Reprint nov. spec. Regni veg.*, 40, Nos. 1–10, 135–48.
- HOSSEUS, C. C. 1937. Musgos de la República Argentina. Contribución al conocimiento de los Musgos de la Provincia de Córdoba (Argentina). I. *Archos Esc. Farm. Córdoba*, No. 5 (Sección científica), 1–42.
- HOSSEUS, C. C. 1938a. Beiträge zur Kenntnis der Laubmoose der Provinz Cordoba (Argentinien). IV. *Reprint nov. spec. Regni veg.*, 44, Nos. 5–10, 136–44.
- HOSSEUS, C. C. 1938b. Beiträge zur Kenntnis der Laubmoose der Provinz Cordoba (Argentinien). V. *Reprint nov. spec. Regni veg.*, 44, Nos. 11–20, 241–43.
- HOSSEUS, C. C. 1938c. Beiträge zur Kenntnis der Laubmoose der Provinz Cordoba (Argentinien). VI. *Reprint nov. spec. Regni veg.*, 45, Nos. 1–7, 20–43.
- HOSSEUS, C. C. 1938d. Musgos de la República Argentina. Contribución al conocimiento de los Musgos de la Provincia de Córdoba (Argentina). II. *Archos Esc. Farm. Córdoba*, No. 8 (Sección científica), 1–58.
- HOSSEUS, C. C. 1939. La flore briológica Argentina. Contribución a su conocimiento. *Physis, B. Aires*, 15, No. 47, 1–12.
- HOSSEUS, C. C. 1940. *Musgos del Territorio de Misiones, nuevos para la Argentina*. Cordoba, Imprenta de la Universidad.
- HÜBENER, J. W. P. 1833. *Muscologia Germanica, oder Beschreibung der deutschen Laubmoose*. Leipzig, Friedrich Hofmeister.
- HULT, R. 1886. Mossfloran i trakterna mellan Aavasaksa och Pallastunturit. *Acta Soc. Fauna Flora fenn.*, 3, No. 1, 1–111.
- HUSNOT, T. 1888. *Muscologia gallica. Descriptions et figures des mousses de France et des contrées voisines*. Orne, T. Husnot, A. Cahan, par Athis; Paris, F. Savy. Pt. 1, Livr. 7, 193–224.
- HUSNOT, T. 1894. *Muscologia gallica. Descriptions et figures des mousses de France et des contrées voisines*. Orne, T. Husnot, A. Cahan, par Athis, Pt. 2, Livr. 12, 349–80.
- JAEGER, A. 1869. Uebersicht über die cleistocarpischen Moose. *Ber. Tät. St Gall. naturw. Ges.*, 1868–69, 62–125.
- JAEGER, A. 1870. Adumbratio florae muscorum totius orbis terrarum. *Ber. Tät. St Gall. naturw. ges.*, 1869–70, 245–99.
- JAEGER, A. 1872. Genera et species muscorum systematicae disposita seu adumbratio florae muscorum totius orbis terrarum. *Ber. Tät. St Gall. naturw. Ges.*, 1870–71, 357–451.
- JAEGER, A. 1873. Genera et species muscorum systematicae disposita seu adumbratio florae muscorum totius orbis terrarum. *Ber. Tät. St Gall. naturw. Ges.*, 1871–72, 309–490.
- JAEGER, A. 1874. Genera et species muscorum systematicae disposita seu adumbratio florae muscorum totius orbis terrarum. *Ber. Tät. St Gall. naturw. Ges.*, 1872–73, 61–236.
- JAEGER, A. 1875. Genera et species muscorum systematicae disposita seu adumbratio florae muscorum totius orbis terrarum. *Ber. Tät. St Gall. naturw. Ges.*, 1873–74, 53–278.
- JAEGER, A. 1876. Genera et species muscorum systematicae disposita seu adumbratio florae muscorum totius orbis terrarum. *Ber. Tät. St Gall. naturw. Ges.*, 1874–75, 85–188.
- JAEGER, A. 1877. Genera et species muscorum systematicae disposita seu adumbratio florae muscorum totius orbis terrarum. *Ber. Tät. St Gall. naturw. Ges.*, 1875–76, 201–371.
- JAEGER, A. 1878. Genera et species muscorum systematicae disposita seu adumbratio florae muscorum totius orbis terrarum. *Ber. Tät. St Gall. naturw. Ges.*, 1876–77, 211–454.
- JAEGER, A. and F. SAUERBECK. 1879a. Genera et species muscorum systematicae disposita seu adumbratio florae muscorum totius orbis terrarum. *Ber. Tät. St Gall. naturw. Ges.*, 1877–78, 257–361.
- JAEGER, A. and F. SAUERBECK. 1879b. Supplementum ad adumbrationem muscorum et conspectus systematis. *Ber. Tät. St Gall. naturw. Ges.*, 1877–78, 362–514.
- JENNINGS, O. E. 1913. *A manual of the mosses of western Pennsylvania*. Pittsburgh, O. E. Jennings.
- JENSEN, C. 1890. De danske Sphagnum-arten. (*In Festskrift, udgivet af den Botaniske Forening i Kjøbenhavn i anledning af dens halvhundred-arsfest, den 12, April 1890*. Kjøbenhavn, Hoffensberg & Trap's Etabl., 52–116.)
- JENSEN, C. 1897. Mosser fra Øst-Grønland. *Meddr Grønland*, 15, Afd. 7 (for 1898), 365–443.
- JENSEN, C. 1923. *Danmarks Mosser eller Beskrivelse af de i Danmark med faærerne fundne Bryofyter II. Andreaeales og Bryales*. København, Kristiania, Glyndendalske Boghandel, Nordisk Forlag.
- JENSEN, C. 1939. *Skandinaviens Bladmossflora*. København, Ejnar Munksgaard.
- JENSEN, T. 1856. *Bryologica Danica eller de Danske bladmosser*. Kjøbenhavn, C. G. Iversens Boghandel Theiles Bogtrykkeri.
- JOHOW, F. 1896. Cryptogamae — Bryophyta. (*In Estudios sobre la flora de las Juan Fernandez*. Santiago de Chile, Imprenta Cervantes, 174–84.)
- [JURATZKA, J.] 1861. Verine, Gesellschaften, Anstalten. *Öst. bot. Z.*, 11, No. 12, 409–13. [Article signed J. J. Ascribed to J. Juratzka by Index muscorum, 3, p. 80 under *Hypnum heufleri*.]
- JURATZKA, J. 1882. *Die Laubmossflora von Oesterreich-Ungarn*. Wien, W. Braumüller.
- KABERSCH, W. 1937. Studien über die ostasiatischen Arten einiger Laubmoosfamilien:

- II. Rhizogoniaceae, Bartramiaceae,
Aulacomniaceae, Meeseaceae. *Hedwigia*, 77, Ht.
2/3, 71–136.
- KARCZMARZ, K. 1966. Taxonomic studies on the
genus *Acrocladium* Mitt. *Nova Hedwigia*, 11, Ht.
1–4, 499–505.
- KICKX, J. 1867. *Flore cryptogamique des Flandres.*
Vol. 1. Gand, Librairie Générale de H. Hoste;
Paris, Librairie de J. B. Bailliére et Fils.
- KINDBERG, N. C. 1883. Die arten der Laubmoose
(Bryineae) Schwedens und Norwegens. *Bih. K.*
svenska VetensAkad. Handl., 7, No. 9, 1–167.
- KINDBERG, N. C. 1885. Table analytique des
mousses pleurocarpes européennes. *Revue*
bryol., 12, No. 2, 24–31.
- KINDBERG, N. C. 1888. *Enumeratio bryinearum*
exoticarum quam alphabetice dispositi.
Linköping, Officina Corresp. Ostrogoth.
- KINDBERG, N. C. 1889. *Enumeratio bryinearum*
exoticarum quam alphabetice dispositi.
Supplementum primum, circiter 1000 acrocarps
complectens. Linköping, Officina Corresp.
Ostrogoth.
- KINDBERG, N. C. 1891. *Enumeratio bryinearum*
exoticarum quam alphabetice dispositi.
Supplementum secundum, circiter 970 species
complectens. Linköping, Officina Corresp.
Ostrogoth.
- KINDBERG, N. C. 1894. Check-list of European and
North American mosses (Bryineae). (Concluded.)
Can. Rec. Sci., 6, No. 2, 72–76.
- KINDBERG, N. C. 1897. *European and North*
American Bryineae (mosses) Parts I & II.
Linköping, Lithografiska Aktiebolag.
- KINDBERG, N. C. 1901. Grundzüge einer
Monographie über die Laubmoos-Familie
Hypopterygiaceae. *Hedwigia*, 40, Ht. 5, 275–
303.
- KINDBERG, N. C. 1902. Grundzüge einer
Monographie der Laubmoos-Gattung
Thamnium. *Hedwigia*, 41, Ht. 4, 203–24; Ht. 5,
225–68.
- KLINGGRÄFF, H. E. M. von. 1858. *Die höheren*
Cryptogamen Preussens. Ein Beitrag zur Flora
der Provinz. Königsberg, Verlag von Wilhelm
Koch.
- KOPONEN, T. 1968. Generic revision of Mniaceae
Mitt. (Bryophyta). *Ann. bot. fenn.*, 5, No. 2,
117–51.
- KOPONEN, T. 1969. The taxonomic status and
typification of *Mnium subpunctatum*. *Bryologist*,
72, No. 1, 61–62.
- KOPONEN, T. 1971. A monograph of *Plagiomnium*
sect. Rosulata (Mniaceae). *Ann. bot. fenn.*, 8, No.
4, 305–67.
- KRAUSS, F. 1846. Pflanzen des Cap- und Natal-
Landes. *Flora, Jena*, 29, (N.R., 4), No. 9, 129–38.
- KRIEGER, W. 1904. Ein Beitrag zur Kenntnis der
Moosflora von Uruguay. *Hedwigia*, 43, Ht. 5,
349–50.
- KUC, M. 1969. Some mosses from an Antarctic
oasis. *Revue bryol. lichen.*, N.S., 36, Fasc. 3–4,
655–72.
- KÜHNEMANN, O. 1938. Catalogo de los musgos
argentinos. *Lilloa*, 2, 37–183.
- KÜHNEMANN, O. 1944. Generos de briofitas de los
alrededores de Buenos Aires. *Lilloa*, 10, 5–232.
- KÜHNEMANN, O. and M. F. GONÇALVES. 1975.
Monografía preliminar para la flora criptogámica
fueguina; Los musgos de la familia
Ptychomniaceae (Bryophyta). *Darwiniana*, 19,
Nos. 2–4, 583–617.
- KÜHNEMANN, O. and M. F. GONÇALVES. 1976. Sobre
Cryphaeophilum molle (Dus.) Fleisch.
Meteoriaceae-Briofitas. *Boln Soc. argent. Bot.*,
17, Nos. 3–4, 247–51.
- KUNTH, C. S. 1822. Musci, Juss. (*In Synopsis*
plantarum, quasin itinere ad plagam
aequinoctialem Orbis Novi, collegunt A, de
Humboldt et A. Bonpland. Vol. 1. Paris, F. G.
Levraut atque Argentorati, 46–65.)
- KUNTZE, O. 1891. Cryptogamae cellulares Musci. (*In*
Revisio generum plantarum vascularium omnium
ataque cellularium multarum secundum leges
nomenclaturae internationales, cum
enumeratione plantarum exoticarum in itinere
mundi collectarum. Pt. 1. Leipzig, Arthur Felix;
London, Dulau & Co.; Milano, U. Hoepli; New
York, Gust E. Stechert; Paris, Charles
Klincksieck, 829–40.)
- KUNTZE, O. 1898. Cryptogamae cellulares Musci. (*In*
Revisio generum plantarum vascularium omnium
ataque cellularium multarum secundum leges
nomenclaturae internationales, cum
enumeratione plantarum exoticarum in itinere
mundi collectarum. Pt. 2. Leipzig, Arthur Felix;
London, Dulau & Co.; Melbourne, George
Robertson & Co.; Milano, U. Hoepli; New York,
Gust E. Stechert; Paris, Charles Klincksieck,
382–83.)
- KURTZ, F. 1904. Flora. (*In Río, M. E. and L. Achával.*
Geografía de la Provincia de Córdoba. Vol. I.
Buenos Aires, Compañía Sud-Americana de
Billets de Banco, 270–343.)
- LAMARK, M. DE and [A.P.] DE CANDOLLE. 1805.
Mousses. Musci. (*In Flore française, ou*
descriptions succinctes de toutes les plantes qui
croissent naturellement en France, disposées
selon une nouvelle méthode d'analyse, et
précédées par un exposé des principes
élémentaires de la botanique. Troisième édition.
Paris, Chez Desray, 438–545.)
- LAMY, D. 1978. Dates de parution des volumes de la
2e. série de la *Revue bryologique et*
lichénologique. Revue bryol. lichen., N.S., 44,
Fasc. 2, 203–04.
- LANGE, J. and C. JENSEN. 1887. Muscineae,
Cryptogamae cellulares. (*In* LANGE, J.
Conspectus florae Groenlandiae. Part 2. *Meddr*
Groenland, 1880–87, Ht. 3, Afdel. 2, 309–446.)
- LAURER, [J.F.] 1827. Beiträge zur kryptogamischen
Flora der Insel Rügen. *Flora, Jena*, 10, No. 19,
289–99.
- LAWTON, E. 1961. A revision of the genus
Rhabdoweisia. *Bryologist*, 64, Nos. 2–3, 140–56.
- LECHLER, W. 1857. *Berberides americae australis,*
accedit enumeratio plantarum, quas in america
australi auctor detexit. Stuttgart, E.
Schweizerbart, 49–59.
- LIMPRICHT, K. G. 1881. Zur Systematik der
Torfmoose. *Bot. Zbl.*, 7, No. 36, 311–19.
- LIMPRICHT, K. G. 1886. Die Laubmoose
Deutschlands, Oesterreichs und der Schweiz. (*In*
RABENHORST, L., ed. *Kryptogamen-Flora von*
Deutschland, Oesterreich und der Schweiz,
Auflage 2, Bd. 4. Leipzig, Eduard Kummer, Abt.
1, Lief. 4, 193–256.)
- LIMPRICHT, K. G. 1888. Die Laubmoose
Deutschlands, Oesterreichs und der Schweiz. (*In*
RABENHORST, L., ed. *Kryptogamen-Flora von*
Deutschland, Oesterreich und der Schweiz,
Auflage 2, Bd. 4. Leipzig, Eduard Kummer, Abt.
1, Lief. 10, 577–640.)
- LIMPRICHT, K. G. 1889. Die Laubmoose
Deutschlands, Oesterreichs und der Schweiz. (*In*

- RABENHORST, L., ed. *Kryptogamen-Flora von Deutschland, Oesterreich und der Schweiz, Auflage 2, Bd. 4.* Leipzig, Eduard Kummer, Abt. 1, Lief. 12, 705–68.)
- LIMPRICHT, K. G. 1892. Die Laubmoose Deutschlands, Oesterreichs und der Schweiz. (*In RABENHORST, L., ed. Kryptogamen-Flora von Deutschland, Oesterreich und der Schweiz, Auflage 2, Bd. 4.* Leipzig, Eduard Kummer, Abt. 2, Lief. 18, 257–320; Lief. 20, 385–448.)
- LINDBERG, S. O. 1861. Om en ny art af slägten *Hypnum*. *Övers. K. VetenskAkad. Förh.*, **18**, No. 8, 371–75.
- LINDBERG, S. O. 1862. Om ett nytt släkte *Epipterygium*, bland bladmossorna. *Övers. K. VetenskAkad. Förh.*, **19**, No. 10, 599–609. [Effective date 1863.]
- LINDBERG, S. O. 1863. Bidrag till mossornas synonymi. *Övers. K. VetenskAkad. Förh.*, **20**, No. 7, 383–418.
- LINDBERG, S. O. 1864a. De Tortulis et ceteris Trichostomeis europaeis. *Övers. K. VetenskAkad. Förh.*, **21**, No. 4, 213–54.
- LINDBERG, S. O. 1864b. Uppställning af familjen Funariaceae. *Övers. K. VetenskAkad. Förh.*, **21**, No. 10, 589–608. [Effective date 15 December 1865.]
- LINDBERG, S. O. 1866. Förteckning öfver mossor, insamlade under de svenska expeditinerna till Spitsbergen 1858 och 1861. *Övers. K. VetenskAkad. Förh.*, **23**, No. 10, 535–61. [Effective date 1867, according to *Index muscorum*, 3, 170.]
- LINDBERG, S. O. 1868a. Animadversiones de *Hypno elegante* Hook. et speciebus europaeis Plagiothecii. *Notis. Sällsk. Faun. Fl. fenn. Förh.*, **9**, [N.S., 6], 19–38.
- LINDBERG, S. O. 1868b. Observationes de Mniaceis europaeis. *Notis. Sällsk. Faun. Fl. fenn. Förh.*, **9**, [N.S., 6], 39–88.
- LINDBERG, S. O. 1868c. Observationes de formis praesertim europaeis Polytrichoidearum (Bryacearum nematodontearum). *Notis. Sällsk. Faun. Fl. fenn. Förh.*, **9**, [N.S., 6], 89–158.
- LINDBERG, S. O. 1868d. Musci novi scandinavici. *Notis. Sällsk. Faun. Fl. fenn. Förh.*, **9**, [N.S., 6], 253–99.
- LINDBERG, S. O. 1871. Revisio critica iconum in opere Flora Danica muscos illustrantium. *Acta Soc. Sci. fenn.*, **10** (for 1875), 3–118.
- LINDBERG, S. O. 1872. Contributio ad floram cryptogamam Asiae boreali-orientalis. *Acta Soc. Sci. fenn.*, **10** (for 1875), 221–80.
- LINDBERG, S. O. 1879. *Musci Scandinavici in systemate novo naturali dispositi.* Upsaliae, Officina lesiae Edquist.
- LINDBERG, S. O. 1883. *Kritisk granskning af mossorna uti Dillenii historia muscorum.* Program. Helsingfors, J. C. Frenckell and Son.
- LINDBERG, S. O. and H. W. ARNELL. 1890. Musci Asiae borealis. Beschreibung der von den Schwedischen Expeditionen nach Sibirien in den Jahren 1875 und 1876 gesammelten Moose mit Berücksichtigung aller Früheren bryologischen angaben für das Russische Nord-Asien. Zweiter Theil. Laubmose. *K. svenska VetenskAkad. Handl.*, **23**, No. 10, 163 pp.
- LINDSAY, W. L. 1866. On new or rare cryptogams from Otago, New Zealand. *Trans. Proc. bot. Soc. Edinb.*, **8**, 280–84.
- LOESKE, L. 1903. *Mossflora des Harzes. Hilfbuch für die bryologische Forschung im Harze und dessen Umgebund mit Verbreitungssangaben und Bestimmungstabellen.* Leipzig, Gebrüder Borntraeger.
- LOESKE, L. 1907. *Drepanocladus*, eine biologische Mischgattung. *Hedwigia*, **46** (for 1906–07), Ht. 5, 300–21.
- LOESEKE, L. 1910. *Studien zur vergleichenden Morphologie und phylogenetischen Systematik der Laubmose.* Berlin, Max Lande.
- LOESKE, L. 1911. Zur Moosflora von Füssen und Hohenschwangau. *Hedwigia*, **50**, Ht. 5, 210–48.
- LOESKE, L. 1913. *Die Laubmose Europas: I. Grimmiaceae.* Berlin, Schöneberg.
- LONGTON, R. E. and M. W. HOLDGATE. 1979. The South Sandwich Islands: IV. Botany. *British Antarctic Survey Scientific Reports*, No 94, 55 pp.
- LOOSER, G. 1932. Excursión botánica y zoológica a la alta Cordillera de las Condes. *An. Univ. Chile*, Ser. 3, **2**, No. 3, 275–301.
- LOOSER, G. 1952. El Botánico francés Thériot y su labor acerca de los musgos Chilenos. *Revta univ., Santiago*, **37**, No. 1, [An. Acad. chil. Cienc. nat., No. 17], 221–43.
- LORENTZ, P. G. 1864. III. *Pugillus specierum novarum e exoticarum.* (*In LORENTZ, P. G. Moosstudien.* Leipzig, Wilhelm Engelmann, 155–71.)
- LORENTZ, P. G. 1866. *Musci frondosi in Chile prope Valdiviam et prope Corral lecti per Dr. Krause.* *Bot. Ztg*, **24**, No. 24, 185–89.
- LORENTZ, P. G. 1868. *Musci frondosi a clarissimo H. Krause in Ecuador, Prov. Loja collecti.* *Bot. Ztg*, **26**, No. 47, 793–800; No. 48, 809–22.
- MACHADO, A. 1919. *Catálogo descritivo de Briologia Portuguesa.* Lisboa, Imprensa de Manuel Lucas Torres. [Title page lacks date, so that given in *Index muscorum*, 5, p. 144, under *Trichostomum humile* has been accepted but not verified.]
- MACOUN, J. and N. C. KINDBERG. 1892. *Catalogue of Canadian plants. Part VI. Musci.* Montreal, William Foster Brown & Co.
- MAHU, M. 1979. Familias y géneros de musgos Chilenos. *Bryologist*, **82**, No. 4, 513–24.
- MALTA, N. 1924. Studien über die Laubmoosgattung *Zygodon* Hook. et Tayl. (11 and 12). *Latv. Univ. Rak.*, **10**, 303–35.
- MALTA, N. 1926. Die Gattung *Zygodon* Hook. et Tayl. Eine monographische studie. *Latv. Univ. bot. dárza Darbi*, No. 1, 185 pp.
- MALTA, N. 1927. Die Ulota-Arten Süd-Amerikas. *Acta Horti bot. Univ. latv.*, **2s**, Nos. 2/3, 165–208.
- MARGADANT, W. D. 1959. Typification and conservation of generic names in Musci. *Acta Bot. Neerl.*, **8**, No. 3, 271–76.
- MARTIN, R. R. 1949. Consideraciones generales sobre la flora actual de los turberos de Tierra del Fuego. *Revta argent. Agron.*, **16**, No. 3, 149–55.
- MATHIEU, C. 1853. Mousses. (*In Flore générale de Belgique, contenant la description de toutes les plantes qui croissent dans ce pays. Ouvrage publié sous le patronage de sa Majesté Le Roi des Belges. Tome II. Cryptogamie.* Bruxelles, Leipzig, Gand, C. Muquardt, 20–73.)
- MATTERI, C. M. 1968. Las especies de "Philonotis (Bartramiaeae)" del sur de Argentina. *Revta Mus. argent. Cienc. nat. Bernardino Rivadavia, Ciencias botánica*, **3**, No. 4, 185–234.
- MATTERI, C. M. 1972. Las Hookeriaceae (Musci) Andino-Patagonicas II. *Revta Mus. argent. Cienc. nat. Bernardino Rivadavia, Ciencias botánica*, **4**, No. 2, 243–80.
- MATTERI, C. M. 1973a. El genero *Breutelia* (Bartramiaeae, Musci) en la region Andino-Patagonica. *Revta Mus. argent. Cienc. nat.*

- Bernardino Rivadavia, Ciencias botánica*, 4, No. 4, 321–60.
- MATTERI, C. M. 1973b. Revision de las Hypopterygiaceae (Musci) Austrosudamericanas. *Boln Soc. argent. Bot.*, 25, Nos. 2–3, 229–50.
- MATTERI, C. M. 1973c. Notas briológicas (Musci) I. *An. Inst. Patagonia*, 4, Nos. 1–3, 273.
- MATTERI, C. M. 1973d. Notas briológicas (Musci) II. *An. Inst. Patagonia*, 4, Nos. 1–3, 274.
- MATTERI, C. M. 1975a. Hookeriales. (*In GUARRERA, S. A., GAMUNDI DE AMOS, I. and D. RABINOVICH DE HALPERIN, ed. Flora criptogámica de Tierre del Fuego*. Buenos Aires, Fundación para la Educación, la Ciencia y la Culutra, 14, Fasc. 9, 7–60.)
- MATTERI, C. M. 1975b. Las Hookeriaceae (Musci) Andino-Patagonicas, I. *Nova Hedwigia*, 26, No. 4, 649–724. [Effective date 1976, personal communication from C. M. Matteri.]
- MATTERI, C. M. 1977a. A synoptic flora of South Georgian mosses: VII. *Pottia*. *British Antarctic Survey Bulletin*, No. 46, 23–28.
- MATTERI, C. M. 1977b. Notes on Antarctic bryophytes: X. The genus *Pottia* from the Antarctic botanical zone. *British Antarctic Survey Bulletin*, No., 46, 140–43.
- MATTERI, C. M. 1979. Notas briológicas (Musci) IV y V. *Bryologist*, 82, No. 1, 99–102.
- MEIJER, W. 1952. The genus *Orthodontium*. *Acta bot. neerl.*, 1, No. 1, 3–80.
- MENZIES, A. 1798. A new arrangement of the genus *Polytrichum*, with some emendations. *Trans. Linn. Soc. Lond.*, 4, 63–84.
- MEYLAN, C. 1905. Catalogue des mousses du Jura. *Bull. Soc. vaud. Sci. nat.*, 5 ème sér., 41, No. 152, 41–96; No. 153, 97–172.
- MIDDLETON, R. M. 1909. The first Fuegian collection. *J. Bot., Lond.*, 47, No. 6, 207–12.
- MILDE, J. 1864a. Ein Sommer in Süd-Tirol. *Bot. Ztg.*, 22, Nos. 17–19 (Beilag), 1–22.
- MILDE, J. 1864b. Zur Cryptogamen-flora Süd-Tirols. *Bot. Ztg.*, 22, Nos. 43 and 44 (Beilag), 1–18.
- MILDE, J. 1869. *Bryologia Silesia. Laubmoose Flora von Nord- und Mitte-Deutschland, unter besonderer Berücksichtigung Schlesiens und mit Hinzunahme der Floren von Jütland, Holland, der Rheinpfalz, von Baden, Franken, Böhmen, Mähren und der Umgegend von München*. Leipzig, Arthur Felix.
- MITTEN, W. 1851. Catalogue of cryptogamic plants collected by Professor W. Jameson in the vicinity of Quito. *Hooker's J. Bot. & Kew Gdns Miscell.*, 3, 49–57; 351–61.
- MITTEN, W. 1856. A list of the Musci and Hepaticae collected in Victoria, Australia, by Dr. F. Mueller. *Hooker's J. Bot. & Kew Gdns Miscell.*, 8, 257–66.
- MITTEN, W. 1859. *Musci Indiae Orientalis*; an enumeration of the mosses of the East Indies. *J. Proc. Linn. Soc.*, Supplement to Botany, 1, [supplement to Vol. III of main series], 1–171.
- MITTEN, W. 1860. Descriptions of some new species of Musci from New Zealand and other parts of the Southern Hemisphere, together with an enumeration of the species collected in Tasmania by William Archer, Esq.; arranged upon the plan proposed in the 'Musci Indiae Orientalis'. *J. Linn. Soc.*, Botany, 4, 64–100. [Read 17 March 1859.]
- MITTEN, W. 1863. On the Musci and Hepaticae from the Cameroons Mountain and from the River Niger. *J. Proc. Linn. Soc.*, Botany, 7 (for 1864), 147–69.
- MITTEN, W. 1864. The "Bryologia" of the survey of the 49th parallel of latitude. *J. Linn. Soc.*, Botany, 8 (for 1865), 12–55.
- MITTEN, W. 1868. A list of Musci collected by the Rev. Thomas Powell in the Samoa or Navigator's Islands. *J. Linn. Soc.*, Botany, 10 (for 1869), 166–95.
- MITTEN, W. 1869. Musci austro-americani. *J. Linn. Soc.*, Botany, 12, 1–659.
- MITTEN, W. 1873. New species of Musci collected in Ceylon by Dr Thwaites. *J. Linn. Soc.*, Botany, 13, 293–326.
- MITTEN, W. 1875. Musci (mosses). (*In MELLIS, J. C. St. Helena; physical, historical, and topographical description of the island, including its geology, fauna, flora and meteorology*. London, L. Reeve & Co., 357–66.)
- MITTEN, W. 1876. The Musci and Hepaticae collected by H.N. Moseley, M.A., naturalist to H.M.S. 'Challenger'. *J. Linn. Soc.*, Botany, 15, No. 82, 59–73.
- MITTEN, W. 1879. Musci. (*In An account of the petrological, botanical and zoological collections made in Kerguelen's Land and Rodriguez during the Transit of Venus Expeditions, carried out by order of Her Majesty's Government in the years 1874–75. Phil. Trans. R. Soc.*, 168 (extra volume), 24–39, 388–96.)
- MITTEN, W. 1883. Australian mosses. *Trans. R. Soc. Vict.*, 19, 49–96.
- MITTEN, W. 1885. Musci. (*In HEMSLEY, W. B. Report on the botany of Juan Fernandez, the south-eastern Moluccas, and the Admiralty Islands. Report on the Scientific Results of the Voyage of H.M.S Challenger during the years 1873–76, under the command of Captain George S. Nares, R.N., F.R.S., and the late Captain Frank Tourle Thomson, R.N. Botany*, Vol. 1, Pt. 3, 78–83; 258–62.)
- MITTEN, W. 1886. The mosses and Hepaticae collected in Central Africa by the late Right Rev. James Hannington, Bishop of Mombasa, F.L.S., F.G.S., &c., with some others, including those gathered by Mr. H. H. Johnson on Kilimanjaro. *J. Linn. Soc.*, Botany, 22, No. 146, 298–329.
- MORR, D. M. H. 1806. Observations on *Orthotrichum* and *Neckera*, together with some other genera of mosses. *Ann. Bot., Lond.*, 2, No. 6, 532–47.
- MOLENDO, I. 1875. Bayerns Laubmoose. Vorläufige Uebersicht mit besonderer Rücksicht auf Niederbayern. *Ber. naturw. Ver. Passau*, 1871–74, 1–278.
- MÖLLER, H. 1907. Ett par upplysningar angående den snart utkommande förteckningen öfver Skandinaviens mossor. *Bot. Notiser* for 1907, 141–45.
- MÖNKEMEYER, W. 1911. Untersuchungen über Cratoneura und Hygroamblystegia. *Hedwigia*, 50, (for 1910–11), Ht. 5/6, 263–78.
- MÖNKEMEYER, W. 1927. Die Laubmoose Europas. (*In RABENHORST, L., ed. Kryptogamenflora von Deutschland, Österreich und der Schweiz*, Bd. 4, Ergänzungsband. Leipzig, Akademische Verlagsgesellschaft, m.b.H., 960 pp.)
- MONTAGNE, C. 1835. Prodromus florae fernandesiana. Pars prima, sistens enumerationem plantarum cellularium quas in Insulā Juan Fernandez à Cl. Bertero collectas describi edique curavit. *Annls Sci. nat.*, Sér. 2, 4, Botanique, 86–99.
- MONTAGNE, C. 1837. Monographie du genre *Conomitrium*, de la famille des mousses. *Annls*

- Sci. nat.*, Sér. 2, **8**, Botanique, 239–52.
- MONTAGNE, C. 1838. Centurie de plantes cellulaires exotiques nouvelles, Musci. *Annls Sci. nat.*, Sér. 2, **9**, Botanique, 38–57.
- MONTAGNE, C. 1839a. Cryptogamae brasilienses seu plantae cellulares quas in itinere per Brasiliam à celeb. Auguste de Saint-Hilaire collectas recensuit observationibusque nonnullis illustravit. *Annls Sci. nat.*, Sér. 2, **12**, Botanique, 42–55.
- MONTAGNE, C. 1839b. *Sertum patagonicum*. Cryptogames de la Patagonie. (*In* D'ORBIGNY, A. *Voyage dans l'Amérique méridionale (le Brésil, la République orientale de l'Uruguay, la République Argentine, la Patagonie, la République du Chili, la République de Bolivie, la République du Pérou)*, exécuté pendant les années 1826, 1827, 1828, 1829, 1830, 1831, 1832 et 1833. Paris, Chez P. Bertrand; Strasbourg, Chez V^e Levrault, Vol. 7, Botanique, Cryptogamie, Pt. 1, 1–19.) [Effective date p. 1–16, 1839 (?); p. 17–19, 1840, *fide* Stafleu, 1967, 326.]
- MONTAGNE, C. 1839c. *Florula boliviensis*. Cryptogams de la Bolivia recueillies par Alcide d'Orbigny. (*In* D'ORBIGNY, A. *Voyage dans l'Amérique méridionale (le Brésil, la République orientale de l'Uruguay, la République Argentine, la Patagonie, la République du Chili, la République de Bolivie, la République du Pérou)*, exécuté pendant les années 1826, 1827, 1828, 1829, 1830, 1831, 1832 et 1833. Paris, Chez P. Bertrand; Strasbourg, Chez V^e Levrault, Vol. 7, Botanique, Cryptogamie, Pt. 2, 1–119.) [Effective date p. 1–104, 1839; p. 105–19, 1840, *fide* Stafleu, 1967, 326.]
- MONTAGNE, C. 1840a. Seconde centurie de plantes cellulaires exotiques nouvelles. Décades VI, VII et VIII. *Annls Sci. nat.*, Sér. 2, **14**, Botanique, 321–50.
- MONTAGNE, C. 1840b. Plantes cellulaires. D. C. Muscinae. (*In* BARKER-WEBB, P. and S. BERTHELOT, *Histoire naturelle des îles Canaries*, Vol. 3, Pt. 2. *Phytographia Canariensis, Sectio Ultima*. Paris, Béthune, 1–67.)
- MONTAGNE, C. 1841. Seconde centurie de plantes cellulaires exotiques nouvelles. Décade X (1) Cryptogamarum Brasiliensium continuatio. *Annals Sci. nat.*, Sér. 2, **16**, Botanique, 266–82.
- MONTAGNE, C. 1842. Botanique — plantes cellulaires: Musci. (*In* SAGRA, R. DE LA. *Histoire physique, politique et naturelle de l'île de Cuba*. Paris, Arthus Bertrand, 493–533.)
- MONTAGNE, C. 1843. Quatrième centurie de plantes cellulaires exotiques nouvelles, Musci. *Annls Sci. nat.*, Sér. 2, **19**, Botanique, 238–43.
- MONTAGNE, C. 1845a. Cinquième centurie de plantes cellulaires exotiques nouvelles. Décades I à VI. *Annls Sci. nat.*, Sér. 3, **4**, Botanique, 86–123.
- MONTAGNE, C. 1845b. Plantes cellulaires. (*In* HOMBROU, M. and C. JACQUINOT. *Voyage au Pole Sud et dans l'Océanie sur les corvettes l'Astrolabe et la Zélée, exécuté par ordre du roi pendant les années 1837–1838–1839–1840, sous le commandement de M. J. Dumont-D'Urville, Capitaine de vaisseau*. Botanique, Vol. 1, Paris, Gide et Cie, 281–335.)
- MONTAGNE, C. 1846. Cryptogames cellulaires, Musci. (*In* GAUDICHAUD, M. 1844–46. *Voyage autour du monde exécuté pendant les années 1836 et 1837 sur la corvette La Bonite commandée par M. Vaillant capitaine de Vaisseau publié par ordre du rois sous les auspices du département de la marine*.
- Botanique*, Vol. 1, Paris, Arthus Bertrand, 267–314.) [Date *fide* Stafleu, 1967, 166.]
- MONTAGNE, C. 1850. Flora chilena, plantas cellulaires. I. Musgos. (*In* GAY, C. *Historia física y política de Chile*. Vol. 7, *Botanica*. Paris, C. Gay; Chile, Museo de Historia natural de Santiago, 5–202.)
- MONTAGNE, C. xs 1856. *CLASSIS MUSCINAE. (In SYLLOGE GENERUM SPECIERUMQUE CRYPTOGAMARUM.* PARISIIS, J-B. BAILLIERE, LONDINI, H. BAILLIERE, 1–95.)
- MONTAGNE, C. 1888. *Pottia flavideta*. (*In* JACKSON, D. B., ed General index to the Journal (Botany, Vols. I–XX) and Proceedings, Nov. 1838 to June 1886, of the Linnean Society of London, 328.)
- MOORE, D. 1873. A synopsis of the mosses of Ireland. *Proc. R. Ir. Acad.*, Ser. 2, **1**, Science (for 1872–73), No. 8, 329–474.
- MORITZI, A. 1846. *Systematisches Verzeichniss der von H. Zollinger in den Jahren 1842–1844 auf Java gesammelten Pflanzen, nebst einer kurzen Beschreibung der neuen Gattungen und Arten*. Solothurn, A. Moritzi. [Date *fide* Stafleu, 1967, 329.]
- MUELLER, C. 1843a. De muscis frondosis nonnullis novis vel minus cognitis. *Linnaea*, Halle, **17**, [Beitr. *PflKunde*, 1], Ht. 5, 581–601.
- MUELLER, C. 1843b. *Aulacomnion chilense* nov. sp. *Bot. Ztg*, **1**, No. 38, 649–56.
- MUELLER, C. 1844a. Relation über die von Gardner in Brasilien gesammelten Laubmoose. *Bot. Ztg*, **2**, No. 42, 726–28.
- MUELLER, C. 1844b. Beiträge zu einer Flora der Aequinoctial-Gegenden. Laubmoose. *Linnaea*, Halle, **18**, [Beitr. *PflKunde*, 12], Ht. 6, 667–709. [Effective date 1845, according to note on p. 508.]
- MUELLER, C. 1845. Synopsis *Macromitriorum hactenus cognitorum*. *Bot. Ztg*, **3**, No. 33, 539–45.
- MUELLER, C. 1847a. Beiträge zu einer Flora der Aequinoctial-Gegenden der neuen Welt. Columbische Laubmoose. *Linnaea*, Halle, **19**, [Beitr. *PflKunde*, 3], Ht. 2, 193–220. [Effective date February 1846, *fide* Van Steenis-Kruseman, M. J. and W. T. Stearn, 1954, p. cxcviii.]
- MUELLER, C. 1847b. Systema muscorum ordinis "Cleistocarpi" Brid. *Bot. Ztg*, **5**, No. 7, 97–102.
- MUELLER, C. 1847c. De muscis nonnullis novis vel minus cognitis exoticis. *Bot. Ztg*, **5**, No. 46, 801–06; No. 47, 825–30.
- MUELLER, C. 1848a. *Synopsis muscorum frondosorum omnium hucusque cognitorum*. Vol. I. Berlin, Alb. Foerstner. [Reprinted 1973: *Synopsis muscorum frondosorum omnium hucusque cognitorum, with a supplement consisting of the articles published in the Botanische Zeitung, Vols. 6–9: Vol. I. Musci vegetationis acrocarpiae*. Amsterdam, A. Asher & Co. B. V., Fasc. 1, 1–160; Fasc. 2, 161–320; Fasc. 3, 321–480.]
- MUELLER, C. 1848b. Plantae Kegelianae surinamenses. Musci frondosi. *Linnaea*, Halle, **21**, [Beitr. *PflKunde*, 5], Ht. 2, 181–200.
- MUELLER, C. 1848c. Ueber die Laubmoose der, von den Herren Funck and Schlium in Columbien veranstalteten, Käuflichen Sammlung des Herren J. Linden in Luxemburg. *Bot. Ztg*, **6**, No. 44, 761–68.
- MUELLER, C. 1849. *Synopsis muscorum frondosorum omnium hucusque cognitorum*. Vol. I. Berlin, Alb. Foerstner, [Reprinted 1973: *Synopsis muscorum frondosorum omnium hucusque cognitorum, with a supplement*

- consisting of the articles published in the *Botanische Zeitung*, Vols. 6–9: Vol. I. *Musci vegetationis acrocarpiae*. Amsterdam, A. Asher & Co. B. V., Fasc. 4, 481–640; Fasc. 5, 641–812.]
- MUELLER, C. 1850. *Synopsis muscorum frondosorum omnium hucusque cognitorum*. Vol. II. Berlin, Alb. Foerstner. [Reprinted 1973: *Synopsis muscorum frondosorum omnium hucusque cognitorum, with a supplement consisting of the articles published in the Botanische Zeitung, Vols. 6–9: Vol. II. Musci vegetationis pleurocarpiae cum appendice*. Amsterdam, A. Asher & Co. B. V., Fasc. 6, 1–160.]
- MUELLER, C. 1851a. *Synopsis muscorum frondosorum omnium hucusque cognitorum*. Vol. II. Berlin, Alb. Foerstner. [Reprinted 1973: *Synopsis muscorum frondosorum omnium hucusque cognitorum, with a supplement consisting of the articles published in the Botanische Zeitung, Vols. 6–9: Vol. II. Musci vegetationis pleurocarpiae cum appendice*. Amsterdam, A. Asher & Co. B. V., Fasc. 7 and 8, 161–510; Fasc. 9 and 10, 511–722.]
- MUELLER, C. 1851b. Die, von Samuel Mossman im Jahre 1850 in Van Dieman's Land, Neuseeland und Neuholland gemachte Laubmoossammlung bestimmt. *Bot. Ztg*, **9**, No. 31, 545–52; No. 32, 561–57.
- MUELLER, C. 1855. De muscis novis, incomplete descriptis, neglectis criticisve. *Bot. Ztg*, **13**, No. 44, 761–69; No. 45, 782–89.
- MUELLER, C. 1856. Symbolae ad synopsin muscorum. *Bot. Ztg*, **14**, No. 24, 415–21; No. 26, 455–59.
- MUELLER, C. 1857. Beiträge zu einer Flora der Kryptogamen Brasiliens, insbesondere der Insel Santa Catharina. *Bot. Ztg*, **15**, No. 23, 378–87.
- MUELLER, C. 1858. Additamenta nova ad synopsin muscorum. *Bot. Ztg*, **16**, No. 22, 154–56; No. 24, 169–72.
- MUELLER, C. 1859. Supplementum novum ad synopsin muscorum. *Bot. Ztg*, **17**, No. 22, 197–98; No. 23, 205–07; No. 24, 214–15; No. 25, 219–21; No. 26, 229–31; No. 28, 246–48.
- MUELLER, C. 1862. Additamenta ad synopsin muscorum nova. *Bot. Ztg*, **20**, No. 40, 327–29; No. 41, 337–39; No. 42, 348–50; No. 43, 361–62; No. 44, 373–74.
- MUELLER, C. 1864. Manipulus muscorum novorum. *Bot. Ztg*, **22**, No. 45, 339–42; No. 49, 373.
- MUELLER, C. 1869. Mittheilungen über *Rhacomitrium lanuginosum* und die verwandten Arten. *Verh. zool.-bot. Ges. Wien*, **19**, Ht. 1, 223–24.
- MUELLER, C. 1872. Musci Australici praesertim Brisbanici novi. *Linnaea*, Berlin, **37** (for 1871–73), [Beitr. *PflKunde*, N.F., **3**], Ht. 2, 143–82.
- MUELLER, C. 1874. Novitates bryothecae müllerianae: II. Musci Novo-Granatenses. Wallisiani adjectis nonnullis aliis muscis novis andinis vel tropico-Americanis vel australasiacis. *Linnaea*, Berlin, **38**, [Beitr. *PflKunde*, N.F., **4**], Ht. 5, 572–620.
- MUELLER, C. 1875. Musci Schweinfurthianoi in itineribus duobus in Africam Centralem per annos 1868–1871 collecti, determinati ex expositi. *Linnaea*, Berlin, **39**, [Beitr. *PflKunde*, N.F., **5**], Ht. 4–6, 325–474.
- MUELLER, C. 1879a. Prodromus bryologiae Argentinicae. I. *Linnaea*, Berlin, **42** (for 1878–79), [Beitr. *PflKunde*, N.F., **8**], Ht. 3–5, 217–460.
- MUELLER, C. 1879b. Musci Fendlerianoi Venezuelenses. *Linnaea*, Berlin, **42**, [Beitr. *PflKunde*, N.F., **8**], Ht. 5–7, 461–502.
- MUELLER, C. 1882. Prodromus bryologiae Argentinicae, II. *Linnaea*, Berlin, **43** (for 1880–82), [Beitr. *PflKunde*, N.F., **9**], Ht. 5 and 6, 341–486.
- MUELLER, C. 1884. Die auf der Expedition SMS "Gazelle" von Dr. Naumann gesammelten Laubmoose. *Bot. Jb.*, **5**, Ht. 1, 76–88. [Effective date 31 December 1883.]
- MUELLER, C. 1885. Bryologia fuegiana. *Flora*, Jena, **68**, Nos. 21–23, 391–429.
- MUELLER, C. 1887a. Sphagnorum novorum descriptio. *Flora*, Jena, **70**, Nos. 26 and 27, 403–22.
- MUELLER, C. 1887b. Erpodiaceae quatuor novae. *Flora*, Jena, **70**, No. 28, 446–50.
- MUELLER, C. 1888. Musci cleistocarpici novi. *Flora*, Jena, **71**, No. 1, 1–13.
- MUELLER, C. 1889. Laubmoose (Musci frondosi). *Die Forschungsreise S.M.S. "Gazelle"*, Theil IV, Botanik. Berlin, Ernst Siegfried Mittler und Sohn.
- MUELLER, C. 1890a. Bryologia austro-georgiae. (*In NEUMAYER, G. Die Internationale Polarforschung 1882–83. Die Deutschen Expeditionen und ihre Ergebnisse*. Berlin, A. Asher and Co., Bd. 2, 279–322.)
- MUELLER, C. 1890b. Die Moose von vier Kilimandscharo Expeditionen. *Flora*, Jena, **73**, [N.R., **48**], No. 5, 465–99.
- MUELLER, C. 1896a. Bryologia provinciae Schensi Sinensis. *Nuovo G. bot. ital.*, N.S., **3**, Fasc. 1, 89–129.
- MUELLER, C. 1896b. Musci nonnulli novi Guianae Anglicae prope Georgetown ad cataractas "Marshall Falls" fluvii Mazarunii a Cl. J. Quelch collecti. *Malpighia*, **10**, Fasc. 11–12, 512–20.
- MUELLER, C. 1897a. Prodromus bryologiae Argentinicae atque regionum vicinarum. III. *Hedwigia*, **36**, Ht. 2, 84–128; Ht. 3, 129–44.
- MUELLER, C. 1897b. Prodromus bryologiae Boliviiana. *Nuovo G. bot. ital.*, N.S., **4**, Fasc. 1, 5–50; Fasc. 2, 113–72.
- MUELLER, C. 1897c. Bryologia Guatemalensis ex collectionibus domin Bernoulli & Cario (1866–1878). V. Türcheim et aliorum. *Bull. Herb. Boissier*, **5**, No. 3, 171–220.
- MUELLER, C. 1897d. Synopsis generis *Harrisonia*. *Öst. bot. Z.*, **47**, No. 11, 387–98; No. 12, 417–20.
- MUELLER, C. 1897e. *Triquetrella* genus muscorum novum conditum et descriptum. *Öst. bot. Z.*, **47**, No. 12, 420–24.
- MUELLER, C. 1898a. Symbolae ad bryologiam Australiae. II. *Hedwigia*, **37**, Ht. 2, 76–112; Ht. 3, 113–44; Ht. 4, 145–71.
- MUELLER, C. 1898b. Analecta bryographica Antillarum. *Hedwigia*, **37**, Ht. 4, 219–24; Ht. 5, 225–66.
- MUELLER, C. 1898c. Bryologia serrae itatiae (Minas Geraës Brasiliae) adjectis nonnullis speciebus affinibus regionum vicinarum. *Bull. Herb. Boissier*, **6**, No. 1, 18–48; No. 2, 89–126.
- MUELLER, C. 1900. Symbolae ad bryologiam Brasiliæ et regionum vicinarum. *Hedwigia*, **39**, Ht. 5, 235–89.
- MUELLER, C. 1901a. Genera muscorum frondosorum, classes *Schistocarporum*, *Cleistocarporum*, *Stegocarporum* complectentia, exceptis *Orthotrichaceis* et *Pleurocarpis*. Leipzig, Eduard Kummer. [Effective date 1900, fide *Index muscorum*, 1, p. 56 under *Andreaea liliputana*.]
- MUELLER, C. 1901b. Symbolae ad bryologiam Brasiliæ et regionum vicinarum. *Hedwigia*, **40**, Ht. 1, 55–80; Ht. 2, 81–99.
- MUELLER, C. and V. F. BROTHERUS. 1900. Ergebnisse

- einer Reise nach dem Pacific. (H. Schauimland, 1896–97.) Musci Schauinslandiani. Ein Beitrag zur Kenntnis der Moosflora der Pacificischen Inseln. *Abh. naturw. Ver. Bremen*, **16**, Ht. 3, 493–512.
- MUELLER, C. and E. HAMPE. 1855. Musci frondosi Australasiae ab Drc. Ferd. Müller lecti. *Linnaea*, **26** (for 1853), Ht. 4, 489–505.
- MUELLER, F. A. 1829. Erestes Verzeichnis sardinischer Laubmoose, wie auch derjenigen welche von meinem Freunde Herm Fleischer bei Smyrna aufgefunden worden sind, nebst Beschreibungen und Abbildungen einiger neuen Arten. *Flora, Jena*, **12**, No. 25, 385–400.
- MULLEN, D. and T. C. FRYE. 1947. Middle and South American species of *Oligotrichum*. *Bryologist*, **50**, No. 1, 67–79.
- NAVEAU, R. 1928. Mousses critiques. *Revue bryol.*, N.S., **1**, No. 1, 38–40.
- NEES VON ESENBECK, C. G., HORNSCHUCH, F. and J. STURM. 1823. *Bryologia Germanica, oder Beschreibung der in Deutschland und in der Schweiz wachsenden Laubmoose*, Vol. 1. Nürnberg, J. Sturm.
- NEGER, F. W. 1899. Informe sobre las observaciones botánicas efectuadas en la Cordillera de Villarrica en el verano 1896–97. *An. Univ. Chile*, **103**, No. 6, 903–67.
- NEWTON, M. E. 1974a. A synoptic flora of South Georgian mosses: IV. *Bartramia* and *Breutelia*. *British Antarctic Survey Bulletin*, No. 38, 59–71.
- NEWTON, M. E. 1974b. Taxonomic notes on *Cheilotrichia* (Lindb.) Broth. and *Dicranella* (C. Muell.) Schimp. *J. Bryol.*, **8**, Pt. 2, 265–68.
- NEWTON, M. E. 1974c. Notes on Antarctic bryophytes: IV. *Encalypta* Hedw. *British Antarctic Survey Bulletin*, No. 39, 1–6.
- NEWTON, M. E. 1974d. Notes on Antarctic bryophytes: V. *Brachythecium majusculum*. M. E. Newton, sp. nov. *British Antarctic Survey Bulletin*, No. 39, 45–48.
- NEWTON, M. E. 1977a. A synoptic flora of South Georgian mosses: VI. *Cheilotrichia*, *Dicranella*, *Distichium*, *Myurella*, and *Catagonium*. *British Antarctic Survey Bulletin*, No. 46, 1–21.
- NEWTON, M. E. 1977b. Notes on Antarctic bryophytes: VII. The occurrence of *Distichium* B.S.G. and *Dicranella* (C. Muell.) Schimp. in the Antarctic botanical zone. *British Antarctic Survey Bulletin*, No. 46, 131–35.
- NEWTON, M. E. 1979a. A taxonomic assessment of *Brachythecium* on South Georgia. *British Antarctic Survey Bulletin*, No. 48, 119–32.
- NEWTON, M. E. 1979b. A synoptic flora of South Georgian mosses: VIII. *Calliergon* and *Brachythecium*. *British Antarctic Survey Bulletin*, No. 48, 133–57.
- NIEUWLAND, J. A. 1917. Critical notes on new and old genera of plants. X. *Am. Midl. Nat.*, **5**, No. 2, 50–52.
- NOGUCHI, A. 1947. A review of the Leucodontinae and Neckerinae of Japan, Loo Choo and of Formosa I. *J. Hattori bot. Lab.*, No. 2, 27–79.
- NYLANDER, W. and T. SAELEN. 1859. *Herbarium musei fennici: förteckning öfver Finska musei växtsamling, utgifven af sällskapet pro fauna et flora fennica*. Helsingfors, Finska Litteratur-sällskapets Tryckeri.
- OCHI, H. 1967. Notes on moss flora. V. *Hikobia*, **5**, Nos. 1–2, 14–38.
- OCHI, H. 1969. Notes on moss flora. VI. *Hikobia*, **5**, Nos. 3–4, 153–71.
- OCHI, H. 1970. A revision of the sub-family Bryoideae in Australia, Tasmania, New Zealand and the adjacent islands. *J. Fac. Educ. Tottori Univ.*, Natural Science, **21**, No. 1, 7–64.
- OCHI, H. 1972. A revision of African Bryoideae, Musci (first part). *J. Fac. Educ. Tottori Univ.*, Natural Science, **23**, No. 1, 1–126.
- OCHI, H. 1976. On the taxonomic status of *Bryum inconnexum* Card. in Antarctica. *Misc. bryol. lichen.*, **7**, No. 6, 116–17.
- OCHI, H. 1977. Central and South American bryaceous mosses, new to science or of geographical significance. *J. Fac. Educ. Tottori Univ.*, Natural Science, **27**, No. 2, 33–41.
- OCHI, H. 1979. A taxonomic review of the genus *Bryum*, Musci, in Antarctica. (*In* MATSUDA, T. and T. HOSHIAI, ed. *Proceedings of the Symposium on Terrestrial Ecosystem in the Syowa Station Area. Mem. natn. Inst. Polar Res. Tokyo*, Special Issue, No. 11, 70–80.)
- PALISOT DE BEAUVOIS, A. M. F. J. 1805a. *Prodrome des cinquième et sixième familles de l'Aethéogamie, les mousses, les lycopodes*. Paris, l'Auteur et Fournier fils. [For combinations effective in 1804 see Crum, 1952.]
- PALISOT DE BEAUVOIS, A. M. F. J. 1805b. Bartrame. (*In Dictionnaire des Sciences Naturelles*, Vol. 4. Strasburg and Paris, F. G. Levrault, 87–88.) [Effective date 8 January 1806, *fide* Sayre, 1959, 70.]
- PALISOT DE BEAUVOIS, [A. M. F. J.] 1822. Muscologie, ou traité sur les mousses. *Mém. Soc. linn. Paris*, **1**, Pt. 2, 388–472.
- PARIS, E. G. 1894. *Index bryologicus sive enumeratio muscorum hucusque cognitorum adjunctis synonymia distributioneque geographicā locupletissimis*. Pt. 1. Pariis, Paul Klincksieck, 1–324. [Date of issue *fide* Paris, 1900b.]
- PARIS, E. G. 1895. *Index bryologicus sive enumeratio muscorum hucusque cognitorum adjunctis synonymia distributioneque geographicā locupletissimis*. Pt. 2. Pariis, Paul Klincksieck, 325–644. [Date of issue *fide* Paris, 1900b.]
- PARIS, E. G. 1896. *Index bryologicus sive enumeratio muscorum hucusque cognitorum adjunctis synonymia distributioneque geographicā locupletissimis*. Pt. 3. Pariis, Paul Klincksieck, 645–964. [Date of issue *fide* Paris, 1900b.]
- PARIS, E. G. 1897. *Index bryologicus sive enumeratio muscorum hucusque cognitorum adjunctis synonymia distributioneque geographicā locupletissimis*. Pariis, Paul Klincksieck, Pt. 4, 965–1284. [Date of issue *fide* Paris, 1900b.]
- PARIS, E. G. 1898. *Index bryologicus sive enumeratio muscorum hucusque cognitorum adjunctis synonymia distributioneque geographicā locupletissimis*. Pariis, Paul Klincksieck, Pt. 5, 1285–380. [Date of issue *fide* Paris, 1900b.]
- PARIS, E. G. 1900a. *Index bryologicus. Supplement*. Genève et Bale, Georg and Cie.
- PARIS, E. G. 1900b. *Musci Novo-Zelandici Browniana. Revue bryol.*, **27**, No. 4, 49–51.
- PARIS, E. G. 1903. *Index bryologicus sive enumeratio muscorum ad diem ultimam anni 1900 cognitorum adjunctis synonymia distributioneque geographicā locupletissimis*. 2nd edition. Paris, Librairie Scientifique A. Hermann, 1, Fasc. 1, 1–64.

- PARIS, E. G. 1904a. *Index bryologicus sive enumeratio muscorum ad diem ultimam anni 1900 cognitorum adjunctis synonymia distributioneque geographicā locupletissimis.* 2nd edition. Paris, Librairie Scientifique A. Hermann, 1, Fasc. 2–6, 65–384.
- PARIS, E. G. 1904b. *Index bryologicus sive enumeratio muscorum ad diem ultimam anni 1900 cognitorum adjunctis synonymia distributioneque geographicā locupletissimis.* 2nd edition. Paris, Librairie Scientifique A. Hermann, 2, Fasc. 1–6 [7–12], 1–375.
- PARIS, E. G. 1904c. *Index bryologicus sive enumeratio muscorum ad diem ultimam anni 1900 cognitorum adjunctis synonymia distributioneque geographicā locupletissimis.* 2nd edition. Paris, Librairie Scientifique A. Hermann, 3, Fasc. 13, 1–72.
- PARIS, E. G. 1905a. *Index bryologicus sive enumeratio muscorum ad diem ultimam anni 1900 cognitorum adjunctis synonymia distributioneque geographicā locupletissimis.* 2nd edition. Paris, Librairie Scientifique A. Hermann, 3, Fasc. 14–18, 73–400.
- PARIS, E. G. 1905b. *Index bryologicus sive enumeratio muscorum ad diem ultimam anni 1900 cognitorum adjunctis synonymia distributioneque geographicā locupletissimis.* 2nd edition. Paris, Librairie Scientifique A. Hermann, 4, Fasc. 19–24, 1–368.
- PARIS, E. G. 1906. *Index bryologicus sive enumeratio muscorum ad diem ultimam anni 1900 cognitorum adjunctis synonymia distributioneque geographicā locupletissimis.* 2nd edition. Paris, Librairie Scientifique A. Hermann, 5, Fasc. 25–27, 1–160.
- PARIS, E. G. 1909. *Collatio nominum brotherianorum et indicis bryologici.* Paris, Libraria J.-B. Bailliére et filii.
- PIOVANO, G. 1954. Alcuni muschi nuovi per l'Argentina. *Nuovo G. bot. ital.*, N.S., 61, Fasc. 2–3, 425–28. [Effective date 10 June 1955.]
- PODPĚRA, J. 1911. Ein Beitrag zu der Kryptogamenflora der bulgarischen Hochgebirge. *Beih. bot. Zbl.*, 28, Abt. 2, Ht. 2, 173–224.
- PODPĚRA, J. 1921. Ad Bryophytorum Cisuralensium cognitionem additementum. *Spisy vydáv. prřír. Fak. Masaryk. Univ.*, Čis. 5, 1–42.
- POIRET, J. L., ed. 1808. *Encyclopédie méthodique botanique, par M. Lamarck, de l'Institut National de France.* Vol. 8. Paris, H. Agasse.
- PORTER, C. E. 1900. Los musgos colectados por la expedición Antártica Belga en el estrecho de Magallanes i Tierra del Fuego estractados de la "Revue Bryologique". *Revta chil. Hist. nat.*, 4, No. 7, 102–06.
- RAAB [W.] 1819. Excursion in die Gegend von Muggendorf von Herrn Präsidenten und Professor Nees von Esenbeck und seinen Zuhörern aus Erlangen. *Flora, Jena*, 2, No. 19, 289–304.
- RABENHORST, [G.] L. 1848. Leber-, Laubmose und Farrn. (*In RABENHORST, L. Deutschlands Kryptogamen-Flora. Bd. 2.* Leipzig, Eduard Kummer, Abt. 3, 1–352.)
- RABENHORST, [G.] L. 1863. *Kryptogamen-flora, Sachsen, der Ober-Lausik, Thüringen und Nordböhmen mit Berücksichtigung der benachbarten Länder.* Abt. 1. Leipzig, Eduard Kummer.
- RADDI, G. 1823. Crittigame brasiliiane. *Memorie Mat. Fis. Soc. Ital. Sci.*, 19, Fasc. 1, 27–57.
- RAYNER, G. W. 1940. MacRobertson Land and Kemp Land, 1936. *Discovery' Rep.*, 19, 165–79.
- REESE, W. R. 1977. The genus *Syrhopodon* in the Americas. I. The elimbate species. *Bryologist*, 80, No. 1, 2–31.
- REICHARDT, H. W. 1868. Diagnosen der neuen Arten von Laubmoosen welche die Novara-Expedition mitbrachte. *Verh. zool.-bot. Ges. Wien*, 18, Ht. 1 and 2, 193–98.
- REICHARDT, H. W. 1870. Fungi, Hepaticae et Musci frondosi. (*In GRUNOW, A., KREMPELHUBER, J., REICHARDT, H. W., METTENIUS, G. and J. MILDE. Reise de österreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859 unter den Befehlen des Commodore B. von Wüllerstorffurbair.* Botanischer theil, Bd. 1. Wien, Kaiserlich-Königlichen Hof- und Staatsdruckerei, 166–96.)
- REID, C. and H. N. RIDLEY. 1888. Fossil Arctic plants from the lacustrine deposit at Hoxne, in Suffolk. *Geol. Mag.*, N.S., 5, Dec. 3, No. 10, 441–44.
- REIMERS, H. 1926. Beiträge zur Bryophytenflora Südamerikas. I. Die von Pater A. Hollermayer in Chile 1918/1923 gesammelten Leber- und Laubmose. *Hedwigia*, 66, Ht. 1, 27–75.
- REIMERS, H. 1936. Über die Laubmoosgattungen *Bryopoteria* Thér., *Camptodontium* Dus., *Hymenoloma* Dus., *Verrucidens* Card. und ihre systematische Stellung. *Notizbl. bot. Gart. Mus. Berl.*, 13, No. 116, 36–52.
- REIMERS, H. 1937. Die europäischen *Haplocladion*—Arten mit besonderer Berücksichtigung ihrer aussereuropäischen Verbreitung und ihrer Verwandtschaft. *Hedwigia*, 76 (for 1936–37), Ht. 5/6, 191–298.
- REINWARDT, [G. C. C.] and [C. F.] HORNSCHUCH. 1826. Musci frondosi javanici reddidi coniunctis studiis et opera. *Nova Acta physico-med. Acad. Caesar. Leop. Carol.*, 14, Pt. 2 (for 1829), 697–732.
- RELHAN, R. 1802. Cryptogamia, Musci. (*In Flora Cantabrigiensis exhibens plantas agri Cantabrigiensis indigenas, secundum systema sexuale digestas.* Ed altera [2nd edition]. *Cantabrigiae, J. Burges*, 412–41.)
- RENAULD, F. 1901. Nouvelle classification des *Leucoloma*. *Revue bryol.*, 28, No. 4, 66–70.
- RENAULD, F. 1909. *Essai sur les Leucoloma et supplément au prodrome de la flore bryologique de Madagascar des Mascareignes et des comores publiés par ordre de S.A.S. Le Prince Albert 1^{er}.* Monaco, Institut de France.
- RENAULD, F. and J. CARDOT. 1888. Notice sur quelques mousses de l'Amérique du Nord. *Revue bryol.*, 15, No. 5, 69–72.
- RENAULD, F. and J. CARDOT. 1889. Mousses nouvelles de l'Amérique du Nord. *Mém. Soc. r. bot. Belg.*, 28, Pt. 1, 121–34.
- RENAULD, F. and J. CARDOT. 1892. Musci. (*In DURAND, T. and H. PITIER, ed. *Primitiae florae Costaricensis*, Fasc. 2. *Bull. Soc. r. Bot. Belg.*, 31, Pt. 1 (Mémoires), 143–74.*) [Effective date 1893, *fide Index muscorum*, 1, 6, under *Acidodontium megalocarpum*.]
- RENAULD, F. and J. CARDOT. 1893a. Pleurocarpae, Hypnaceae. (*In Nordamerikanische Laubmose Torfmoose und Lebermoose gesammelt von Dr. Julius Röll in Darmstadt.* *Hedwigia*, 32, Ht. 5, 334–71.)
- RENAULD, F. and J. CARDOT. 1893b. Musci Americae Septentrionalis, ex operibus novissimis recensiti et methodici dispositi (suite). *Revue bryol.*, 20, Nos. 1–2, 1–32.
- RICHARDS, P. W. 1937. A collection of bryophytes

- from the Azores. *Annals bryol.*, **9** (for 1936), 131–38.
- RICHARDS, P. W. and E. C. WALLACE. 1950. An annotated list of British mosses. *Trans. Br. bryol. Soc.*, **1**, Pt. 4, i–xxi.
- ROBINSON, H. [E.] 1964. Five bryophytes of interest from Chile. *Bryologist*, **67**, No. 1, 53–55.
- ROBINSON, H. [E.] 1970a. A revision of the genus *Trichostomopsis*. *Phytologia*, **20**, No. 3, 184–91.
- ROBINSON, H. [E.] 1970b. Notes on the moss genera, *Camptochaete*, *Physcomitrium* and *Ptychomnion*, in Chile. *Phytologia*, **20**, No. 6, 329–31.
- ROBINSON, H. E. 1972. Observations on the origin and taxonomy of the Antarctic moss flora. (*In* LLANO, G. A., ed. *Antarctic terrestrial biology*. Washington, D.C., American Geophysical Union, 163–77.) [Antarctic Research Series, Vol. 20.]
- ROBINSON, H. [E.] 1974. Notes on the mosses of Juan Fernandez and southern South America. *Phytologia*, **29**, No. 2, 116–19.
- ROBINSON, H. [E.] 1975. The mosses of Juan Fernandez Islands. *Smithson. Contrib. Bot.*, No. 27, 88 pp.
- RÖHLING, I. C. 1809. Kritische Bemerkungen über meine Laubmoose. Ein Beitrag zur richtigen Bestimmung der Moosarten. *Annals wetterauisch. Ges.*, **1**, Ht. 2, 178–201.
- RÖHLING, I. C. 1812. Kritische Bemerkungen über meine Laubmoose. *Annals wetterauisch. Ges.*, **3**, Ht. 1, 79–122.
- RÖHLING, I. C. 1813. Moose (*In Deutschlands Flora. Ein botanisches Taschenbuch. 2nd edition. Vol. 3. Kryptogamische Gewächse*. Frankfurt am Main, Friedrich Wilmans, Abt. 1, 32–134.)
- RÖHLING, I. C. 1814. Kritische Bemerkungen über meine Laubmoose. *Annals wetterauisch. Ges.*, **3**, Ht. 2, 193–235.
- ROI VAINEN, H. 1934. Musci nonnulli ex Chile austro-centrali. *Suomal. eläin- ja kasvit. Seur. van. kasvit. Julk.*, **5**, No. 10, 30–36.
- ROI VAINEN, H. 1954. Studien über die Moose Feuerlands. *Suomal. eläin- ja kasvit. Seur. van. kasvit. Julk.*, **28**, No. 2, 1–205.
- ROI VAINEN, H. 1955a. Contribution to the Fuegian species of the genus *Rhacomitrium* Brid. *Suomal. eläin- ja kasvit. Seur. van. Tiedon.*, **9**, No. 2, 85–98.
- ROI VAINEN, H. 1955b. *Bucklandia bartramii*, a new genus and species of leaf moss from Tierra del Fuego. *Suomal. eläin- ja kasvit. Seur. van. Tiedon.*, **9**, No. 2, 98–100.
- ROI VAINEN, H. 1972. *Bucklandiella* Roiv., nomen novum (Musci). *Ann. bot. fenn.*, **9**, No. 2, 116.
- ROI VAINEN, H. and E. B. BARTRAM. 1937. Bryological investigations in Tierra del Fuego. I. Sphagnaceae — Dicranaceae. *Suomal. eläin- ja kasvit. Seur. van. kasvit. Julk.*, **9**, No. 2, 1–58.
- RÖLL, [J.] 1886. Zur Systematik der Torfmoose. *Flora, Jena*, **69**, No. 6, 89–94; No. 15, 227–42; Nos. 27–28, 419–27.
- RÖLL, J. 1915. Die Thüringer Torfmoose und Laubmoose. *Hedwigia*, **56**, Ht. 1–2, 1–176.
- ROTH, A. W. 1802. *Neue Beyträge zur Botanik. Theil 1*. Frankfurt am Main, Friedrich Wilmans.
- ROTH, G. 1904. *Die europäischen Laubmoose*, Bd. II. Leipzig, Wilhelm Engelmann, Lief. 10, 513–640.
- ROTH, G. 1908. Übersicht über die europäischen *Drepanocladus*, einschliesslich der neueren Formen. *Hedwigia*, **48**, Ht. 3, 152–77.
- ROTH, G. 1910. *Die aussereuropäischen Laubmoose*. Bd. I. Dresden, C. Heinrich, Lief. 1, 1–96.
- ROTH, G. 1911. *Die aussereuropäischen Laubmoose*. Bd. I. Dresden, C. Heinrich, Lief. 2, 97–192; Lief. 3, 193–272; Lief. 4, 273–331.
- ROTH, G. 1913. Nachtrag I zu Band I der aussereuropäischen Laubmoose von 1910–11. *Hedwigia*, **53**, Ht. 2, 81–98.
- ROTH, G. 1914. Nachtrag II zu Band I der aussereuropäischen Laubmoose von 1910–11. *Hedwigia*, **55**, Ht. 6, 267–74.
- ROWLEE, W. W. 1916. Plants from southern Patagonia collected by Charles Wellington Furlong. *Bull. Torrey bot. Club*, **43**, No. 6, 305–21.
- RUSOW, E. 1867. Beiträge zur Kenntniss der Torfmoose. *Arch. Naturk. Liv- Est- u. Kurlands*, Ser. 2, Biologische Naturkunde, **7**, Lief. 1, 83–162.
- SAINS BURY, G. O. K. 1931. The moss *Dicranoloma setosum* (H. f. & W.) Par. *Bryologist*, **34**, No. 5, 72–74. [Effective date 8 February 1932.]
- SAINS BURY, G. O. K. 1952. Critical New Zealand mosses. *Revue bryol. lichen.*, N.S., **21**, Fasc. 3–4, 213–25.
- SAINS BURY, G. O. K. 1955. A handbook of the New Zealand mosses. *Bull. R. Soc. N.Z.*, No. 5, 490 pp.
- SALMON, E. S. 1903. Bryological notes. (30) *Calyptopogon mnioides* (Schwaegr.) Broth. *J. Bot., Lond.*, **41**, No. 1, 1–8.
- SANIONE, C. 1883. Additamentum secundum in harpidiorum cognitionem. *Bot. Zbl.*, **13**, No. 13, 425–40.
- SAVICH-LJUBITSKAYA, L. I. 1978. Listostebel'nye mkhi vostochnoi Antarktidy [Leafy mosses of eastern Antarctic region]. *Bot. Zh. SSSR*, **63**, No. 2, 272–90.
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1959. Novyi vid roda *Bryum* Hedw. iz oazisa Bangera [A new species of the genus *Bryum* Hedw. from Bunger's Oasis]. *Inf. Byull. sov. antarkt. Eksped.*, No. 7, 34–39. [English translation: Vol. 1, 1964, 308–13.]
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1960. Novaya raznovidnost *Bryum korotkeviciae* Sav.-Ljub. et Z. Smirn. [New variety of *Bryum korotkeviciae* Sav.-Ljub. et Z. Smirn.]. *Inf. Byull. sov. antarkt. Eksped.*, No. 17, 25–27. [English translation: Vol. 2, 1964, 206–08.]
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1962. Endem Antarktidy — mokh *Sarconeurum glaciale* (Hook. fil. et Wils.) Card. et Bryhn [The endemic moss of the Antarctic continent — *Sarconeurum glaciale* (Hook. fil. et Wils.) Card. et Bryhn]. (*In Rezul'taty biologicheskikh issledovaniy Sovetskoy Antarkticheskoi Ekspeditsii, 1955–58 [Biological Results of Soviet Antarctic Expedition, 1955–58]*. Moscow and Leningrad, Akademii Nauk SSSR, **1**, 295–309.)
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1963a. O predstaviteleyakh roda *Pottia* Fuernr. v Antarktide. De specibus *Pottiae* Fuernr. in Antarctica inventis. (*In Botanicheskie materialy Otdela sporovyleh rastenii. Akademiya Nauk SSSR Botanicheskii Institut im V.L. Komarova*, **16**, 188–95.)
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1963b. K biologii i geografii *Bryoerythrophyllum recurvirostre* (Hedw.) Chen — novogo vida dlya brioflory Antarktidy [A contribution to the biology and geography of *Bryoerythrophyllum recurvirostre* (Hedw.) Chen — a new species in the bryoflora of Antarctica]. *Bot. Zh. SSSR*, **48**,

- No. 3, 350–61.
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1964a. Glubokovodnyi predstavite' roda *Plagiothecium* Br. et Sch. v Antarktide [A deep-water member of the genus *Plagiothecium* Br. et Sch. in Antarctica]. *Inf. Byull. sov. antarkt. Eksped.*, No. 49, 33–39. [English translation: Vol. 5, No. 4, 1965, 240–43.]
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1964b. Zametka o *Bryum argenteum* Hedw. iz Antarktidy. Notula de *Bryo argenteno* Hedw. ex Antartida. *Nov. sist. Nizshikh Rast.*, (for 1964), 292–301.
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1965a. O *Grimmia antarctici* Card. iz Vostochnoy Antarktidy. De *Grimmia antarctici* Card. ex Antartide Orientalis. *Nov. sist. Nizshikh Rast.*, (for 1965), 251–58.
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1965b. Dopolneniya k state o predstaviteleyakh roda *Pottia* Fuernr. v Antarktide. De specibus *Pottiae* Fuernr. in Antartide inventis supplementa. *Nov. sist. Nizshikh Rast.*, (for 1965), 158–59.
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1969. *Grimmia doniana* Sm. iz Vostochnoy Antarktidy. *Grimmia doniana* Sm. ex Antartida orientali. *Nov. sist. Nizshikh Rast.*, 6, 249–53.
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1970. *Grimmia plagiopoda* Hedw. iz Vostochnoy Antarktidy [*Grimmia plagiopoda* Hedw. from East Antarctica]. *Inf. Byull. sov. antarkt. Eksped.*, No. 80, 94–97. [English translation: Vol. 8, No. 2, 1972, 97–98.]
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1971. Mokh *Pohlia nutans* (Hedw.) Lindb. v Vostochnoy Antarktide [Moss *Pohlia nutans* (Hedw.) Lindb. in East Antarctica]. *Inf. Byull. sov. antarkt. Eksped.*, No. 82, 72–76. [English translation: Vol. 8, No. 4, 1973, 222–24.]
- SAVICH-LJUBITSKAYA, L. I. and Z. N. SMIRNOVA. 1972. *Bryum algens* Card.—naibolee pasprostranennyy mokh Vostochnoy Antarktidy. *Bryum algens* Card. is the most common moss in East Antarctica. (*In Trudy Sovetskoy Antarkticheskoy Ekspeditsii. Nauchnye rezul'taty rabot Sovetskikh Antarkticheskikh Ekspeditsii*, 60, 328–45.)
- SAYRE, G. 1959. *Dates of publications describing Musci 1801–1821*. Troy, New York, G. Sayre.
- SAYRE, G. 1969. Cryptogamae exsiccatae—an annotated bibliography of published exsiccatae of Algae, Lichenes, Hepaticae and Musci. Introduction, I. General cryptogams, II. Algae, III. Lichenes. *Mem. N.Y. bot. Gdn.*, 19, No. 1, 1–174.
- SAYRE, G. 1971. Cryptogamae exsiccatae—an annotated bibliography of published exsiccatae of Algae, Lichenes, Hepaticae and Musci. IV. Bryophyta. *Mem. N.Y. bot. Gdn.*, 19, No. 2, 175–276.
- SAYRE, G. 1975. Cryptogamae exsiccatae—an annotated bibliography of published exsiccatae of Algae, Lichenes, Hepaticae and Musci. V. Unpublished exsiccatae. *Mem. N.Y. bot. Gdn.*, 19, No. 3, 277–423.
- SCHIMPER, W. P. 1836. Muscorum chilensem species novas descriptis. *Annls Sci. nat.*, Sér. 2, 6, Botanique, 145–49.
- SCHIMPER, W. P. 1838. Ueber *Octodiceras julianum*. *Flora, Jena*, 21, No. 17, 267–72.
- SCHIMPER, W. P. 1855. *Corollarium bryologiae europaeae, conspectum diagnosticum familiarum, generum et specierum, adnotaciones novas atque emendationes complectens*. Stuttgariae, E. Schweizerbart. [Effective date 1856, according to Stafleu and Cowan (1976, 379).]
- SCHIMPER, W. P. 1858. Mémoire pour servir à l'histoire naturelle des Sphaignes (*Sphagnum* L.). *Mém. prés. div. Sav. Acad. Sci. Inst. Fr., Science mathématiques et physiques*, 15, 1–97. [Effective date 1857, according to *Index muscorum*, 4, 63, under *Sphagnum acutifolium* var. *purpureum*. According to the author catalogue of the British Museum (Nat. Hist.) preprints were circulated in 1857.]
- SCHIMPER, W. P. 1860. *Synopsis muscorum europaeorum praemissa introductione de elementis bryologicis tractante*. Stuttgariae, E. Schweizerbart.
- SCHIMPER, W. P. 1876. *Synopsis muscorum europaeorum praemissa introductione de elementis bryologicis tractante. Vol. II. Specierum descriptio. Editio secunda*. Stuttgariae, E. Schweizerbart (E. Koch).
- SCHLEICHER, J. C. 1821. *Catalogus hucusque absolutus omnium plantarum in Helvetia cis et transalpina sponte nascentium*. 4th edition. Camberii, Gorrii et Routin.
- SCHLIEPHACKE, K. 1884. *Pottia güßfeldtii*, ein neues Laubmoos. *Ber. dt. bot. Ges.*, 2, Ht. 10, 461–62. [Effective date 20 January 1885.]
- SCHRADER, H. A. 1794. *Spicilegium florae germanicae*. Pars. I. Hannoverae, Ritscher.
- SCHRADER, H. A. 1803a. *Plantae cryptogamicae nouae, rariores aut minus cognitae*. J. Bot., Göttingen, 1 (for 1801), Stück 1, 66–80.
- SCHRADER, M. 1803b. [Review of:] *Muscologia recentiorum seu analysis, historia et descriptio methodica omnium muscorum hucusque cognitorum ad Norman Hedwigii. A. Sam. El. Bridel Gotha et Parisiis T. I. (xxiv und 179s) 4. 1797; Tom. II Pars I (x und 222s) cum tab. aen sex 1798; Pars 2 (xiv und 182s) cum tab. aen sex 1801. J. Bot., Göttingen, 1 (for 1801), Stück 1, 188–205.*
- SCHRANK, F. von PAULA. 1789. *Baiersche Flora*. Vol. II. München, Strobl.
- SCHREBIUS, J. 1802. Recensionen. Leipzig bei Barth: Jo. Hedwig, Prof. Botan. in acad. Lips. Species muscorum frondosorum descriptae et tabulis aeneis LXXVII coloratis illustratae. Opus posthumum editum a Fr. Schwägrichen, 1801, 352 S. in 4. *Bot. Ztg Erlangen*, 1, Ht. 1, No. 5, 65–80.
- SCHULTZ, C. F. 1823. Recensio generum *Barbulae* et *Syntrichiae*. *Nova Acta physico-med. Acad. Caesar. Leop. Carol.*, 11, Pt. 1, 191–232.
- SCHULTZ, C. F. 1828. Observationes bryologicae. (*In HORNSCHUCH, C. F. Sylloge plantarum novarum itemque minus cognitarum a praestantissimus botanicis adhuc viventibus collecta et a societate regia botanica ratisbonensi edita, tomus secundus. Ratisbonae*, C. E. Brenck, 119–54, 177–217.) [Editorship ascribed to Hornschuch following Stafleu, 1967, 220.]
- SCHULTZE-MOTEL, W. 1970. Monographie de Laubmoosgattung *Andreaea*. I. Die costaten arten. *Wildenowia*, 6, No. 1, 25–110.
- SCHWAEGRICHEN, A. D. 1804. Musci novi detecti et descripti. (*In SCHULTE, J. A. Reise auf dem Glockner, Theil 2*. Wien, J. V. Degen, Appendix, 362–66.)
- SCHWAEGRICHEN, F. 1811. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum*.

- Supplementum primum, Sectio primum.* Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz.
- SCHWAEGRICHEN, F. 1816. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum primum, Sectio posterior [=Section 2]. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz.
- SCHWAEGRICHEN, F. 1823. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum secundum, Sectio prior. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz; Londini, Treuttel et Würtz, Treuttel filium et Richter.
- SCHWAEGRICHEN, F. 1824. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum secundum, Sectio posterior. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz; Londini, Treuttel et Würtz, Treuttel filium et Richter.
- SCHWAEGRICHEN, F. 1826. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum secundum, Volumen secundum, Sectio prior. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz; Londini, Treuttel et Würtz, Treuttel filium et Richter.
- SCHWAEGRICHEN, F. 1827a. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum secundum, Volumen secundum, Sectio secundae. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz; Londini, Treuttel et Würtz, Treuttel filium et Richter.
- SCHWAEGRICHEN, F. 1827b. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum tertium, Volumen primum, Sectio prima. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz; Londini, Treuttel et Würtz, Treuttel filium et Richter.
- SCHWAEGRICHEN, F. 1828. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum tertium, Volumen primum, Sectio secunda. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz; Londini, Treuttel et Würtz, Treuttel filium et Richter.
- SCHWAEGRICHEN, F. 1829. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum tertium, Volumen secundum, Sectio prima. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz; Londini, Treuttel et Würtz, Treuttel filium et Richter.
- SCHWAEGRICHEN, F. 1830. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum tertium, Volumen secundum, Sectio secunda. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz; Londini, Treuttel et Würtz, Treuttel filium et Richter.
- SCHWAEGRICHEN, F. 1842. *Joannis Hedwig. Species muscorum frondosorum. Descriptae et tabulis aeneis coloratis illustratae. Opus postumum.* Supplementum quartum, Sectio prima. Lipsiae, Joannis Ambrosii Barth; Parisiis, Treuttel et Würtz; Londini, Longman, Brown, Rees and Longmans.
- SCHWAEGRICHEN, C. F. 1844. Ueber einige Moose. *Linnaea*, Halle, 18, [Beitr. PflKunde, 2], Ht. 5, 557–62. [Effective date 1845, according to note on p. 508.]
- SEEMANN, B. C. 1873. *Flora vitiensis: a description of the plants of the Fiji or Viti Islands, with an account of their history, uses and properties.* London, L. Reeve and Co., Pt. 10, 325–434. [Dates of whole work 1865–73.]
- SEKI, T. 1974. A moss flora of Provincia de Aisén, Chile. Results of the second scientific expedition to Patagonia by Hokkaido and Hiroshima Universities, 1967. *J. Sci. Hiroshima Univ., Ser. B, Div. 2 (Botany)*, 15, No. 1, 9–101. [Effective date 31 March 1976; personal communication from author.]
- SENDTNER, O. 1848. Beobachtungen über die klimatische Verbreitung der Laubmose durch das österreichische Küstenland und Dalmatien. *Flora, Jena*, 31, (N.R., 6), No. 14, 229–40.
- SKOTTSBERG, C. [J.F.] 1914. Botanische Ergebnisse der Schwedischen Expedition nach Patagonien und dem Feuerlande 1907–1909. IV. Studien über die Vegetation der Juan Fernandez-Inseln. *K. svenska VetenskAkad. Handl.*, 51, No. 9, 1–73.
- SKOTTSBERG, C. [J.F.] 1950. Apuntes sobre la flore y vegetación de Frai Jorge (Coquimbo, Chile). *Acta Horti gothoburg.*, 18 (for 1948), 91–184.
- SMITH, G. L. 1969a. On *Atrichopsis*, with notes on some austral *Psilopilum* species (Polytrichaceae). *Bull. Torrey bot. Club*, 96, No. 1, 60–69.
- SMITH, G. L. 1969b. New combinations in Polytrichaceae. *Phytologia*, 18, No. 7, 403.
- SMITH, G. L. 1971. A conspectus of the genera of Polytrichaceae. *Mem. N.Y. bot. Gdn*, 21, No. 3, 1–83.
- SMITH, J. E. 1803. *Trichostomum capillaceum. (In English botany; or coloured figures of British plants, with their essential characters, synonyms, and places of growth: to which will be added occasional remarks. The figures by James Sowerby.* London, J. E. Smith, 16, Tab. 1152.)
- SMITH, J. E. 1804a. *Grimmia donniana. (In English botany; or coloured figures of British plants, with their essential characters, synonyms, and places of growth: to which will be added occasional remarks. The figures by James Sowerby.* London, J. E. Smith, 18, Tab. 1259).
- SMITH, J. E. 1804b. *Flora britannica*, Vol. 3. London, J. Taylor.
- SMITH, J. E. 1807. *Grimmia pulvinata. (In English botany; or coloured figures of British plants, with their essential characters, synonyms, and places of growth: to which will be added occasional remarks. The figures by James Sowerby.* London, J. E. Smith, 24, Tab. 1728.)
- SMITH, J. E. 1809. *Grimmia atro-virens. (In English botany; or coloured figures of British plants, with their essential characters, synonyms, and places of growth: to which will be added occasional remarks. The figures by James Sowerby.* London, J. E. Smith, 28, Tab. 2015.)
- SMITH, J. E. 1811. *Gymnostomum imberbe. (In English botany; or coloured figures of British plants, with their essential characters, synonyms, and places of growth: to which will be added occasional remarks. The figures by James Sowerby.* London, J. E. Smith, 32, Tab. 2237.)
- SMITH, R. I. L. 1972. Vegetation of the South Orkney Islands with particular reference to Signy Island. *British Antarctic Survey Scientific Reports*, No. 68, 124 pp.

- SPEGAZZINI, C. 1922. Resultados de la primera expedición a Tierra del Fuego (1921) enviada por la Facultad de Ciencias Exactas, Físicas y Naturales de la Universidad Nacional de Buenos Aires. *Cryptogamae nonnullae fuegianae. An. Soc. cient. argent.*, **94**, Entr. 1–2, 59–85.
- SPRENGEL, C. 1827. *Cryptogamia. Sect. II. Musci frondosi et Hepatici. (In LINNAEI, C. Systema vegetabilium. Editio decima sexta. Vol. IV. Pars 1. Gottingae, Librariae Dieterichianaes, 135–236.)*
- SPRUCE, R. 1845. On several mosses, new to the British flora. *Lond. J. Bot.*, **4**, 169–95.
- SPRUCE, R. 1847. *Musci pyrenaici, quos in Pyrenaeis centralibus occidentalibusque nec non in agro syrtico, AD MDCCCXLV-VI, Fasc. 1, Nos. 1–160. Londoni.*
- SPRUCE, R. 1849. The Musci and Hepaticae of the Pyrenees. *Ann. Mag. nat. Hist.*, Ser. 2, **3**, No. 16, 269–93.
- SPRUCE, R. 1867. *Catalogus muscorum fere omnium quos in terris Amazonicis et Andinis per annos 1849–1860. Londini, E. Newman.*
- STAFLEU, F. A. 1967. Taxonomic literature. A selective guide to botanical publications with dates, commentaries and types. *Regnum veg.*, **52**, 556 pp.
- STAFLEU, F. A., ED. 1978. *International code of botanical nomenclature, adopted by the twelfth International Botanical Congress, Leningrad, July 1975. Regnum veg.*, **97**, 457 pp.
- STAFLEU, F. A. and R. S. COWAN. 1976. Taxonomic literature, a selective guide to botanical publications and collections with dates, commentaries and types. Edition 2. Vol. 1, A–G. *Regnum veg.*, **94**, 1136 pp.
- STAFLEU, F. A. and R. S. COWAN. 1979. Taxonomic literature, a selective guide to botanical publications and collections with dates, commentaries and types. Edition 2. Vol. 2, H–Le. *Regnum veg.*, **98**, 991 pp.
- STAFLEU, F. A. and R. S. COWAN. 1983. Taxonomic literature, a selective guide to botanical publications and collections with dates, commentaries and types. Edition 2. Vol. 4. *Regnum veg.*, **110**, 1214 pp.
- STEERE, W. C. 1948. Contribution to the bryogeography of Ecuador. 1. A review of the species of musci previously reported. *Bryologist*, **51**, No. 3, 65–167.
- STEERE, W. C. 1961a. A preliminary review of the bryophytes of Antarctica. (In *Science in Antarctica. Part I. The life sciences in Antarctica*. Washington, D.C., National Academy of Sciences—National Research Council, 20–33.) [National Academy of Sciences—National Research Council Publication 839.]
- STEERE, W. C. 1961b. The bryophytes of South Georgia. (In *Science in Antarctica. Part I. The life sciences in Antarctica*. Washington, D.C., National Academy of Sciences—National Research Council, 34–48.) [National Academy of Sciences—National Research Council Publication 839.]
- STEDEL, E. [G.] 1824. *Nomenclator botanicus enumerans ordine alphabetico nomina atque synomina tum genericum tum specifica et a Linnaeo et recentioribus de re botanica scriptoribus plantis cryptogamicis imposita. Part 2. Stuttgardiae et Tubingae, J. G. Cottae.*
- STURM, J. 1813. *Deutschlands Flora in Abbildungen nach der Natur mit Beschreibungen. Abt. II. Cryptogamia. Ht. 14. Nürnberg, J. Sturm.*
- SULLIVANT, W. S. 1846. *Musci Alleghanienses, sive enumeratio muscorum atque hepaticarum quos in itinere a Marylandia usque ad Georgiam per tractus montium. Columbus, Ohio, W. S. Sullivant.*
- SULLIVANT, W. S. 1849. Contributions to the bryology and hepaticology of North America. Pt. II. *Mem. Am. Acad. Arts Sci.*, N.S. **4**, Pt. 1, 169–76.
- SULLIVANT, W. S. 1850. Notices of several new species of mosses and Hepaticae from Tierra del Fuego. *Hooker's J. Bot. & Kew Gdns Miscell.*, **2**, 315–18.
- SULLIVANT, W. S. 1857. Notices of new species of mosses, from the Pacific islands. *Proc. Am. Acad. Arts Sci.*, **3** (for 1852–57), 73–81. [Effective date 1854, according to entry under *Cryphaea cuspidata* in *Index muscorum*, 1, 514.]
- SULLIVANT, W. S. 1859. Musci. (In *United States Exploring Expedition during the years 1838, 1839, 1840, 1841, 1842 under the command of Charles Wilkes, U.S.N.* Vol. XVII. *Botany*. Philadelphia, C. Sherman and Son, 3–32.)
- SULLIVANT, W. S. and [C.] L. LESQUEREAUX. 1856. *Musci boreali-Americanorum specimina exsiccata. Columbi Ohioensium, Auctorum.*
- SULLIVANT, W. S. and L. LESQUEREAUX, ED. 1865. *Musci boreali-Americanorum specimina exsiccata. Editio secunda. Columbi Ohioensium, W. S. Sullivant and L. Lesquereux.*
- [SWARTZ, O.] 1801. *Dispositio systematica muscorum frondosorum Sueciae, adjectis descriptionibus et iconibus novarum specierum. 12 m. pp. 112. Erlangae, apud Jo. Jac. Palm.* 1799. *Mon. Rev. Lond.*, Ser. 2, **34**, Appendix, 535–38.
- SWARTZ, O. 1803. *Cinclidium, eine neue Moos-gattung. J. Bot., Göttingen*, **1** (for 1801), Stück 1, 25–31.
- SWARTZ, O. 1806a. *Conostomum, eine neue Moos-gattung. Neues J. Bot.*, **1**, Stück 3, 24–36.
- SWARTZ, O. 1806b. *Cryptogamia. In Flora Indiae occidentalis aucta atque illustrata sive descriptiones plantarum in Prodromo recensitarum. Vol. 3. Erlangen, Jo. Jacobi Palmii.* 1569–940.
- SYDOW, P. 1910. Moose. *Just's bot. Jber.*, **36** (for 1908), Abt. 1, 30–90.
- TAYLOR, N. 1914. Plants collected on the South Georgian Expedition. (In MURPHY, R. C. A report on the South Georgia Expedition. *Sci. Bull. Brooklyn Inst.*, **2**, No. 4, 60–63.)
- TAYLOR, T. 1846. The distinctive characters of some new species of Musci, collected by Professor William Jameson, in the vicinity of Quito, and by Mr. James Drummond at Swan River. *Lond. J. Bot.*, **5**, 41–67.
- TAYLOR, T. 1847. Description of new Musci and Hepaticae collected by Prof. W. Jameson on Pichincha, near Quito by Thomas Taylor M.D. *Lond. J. Bot.*, **6**, 328–42.
- TAYLOR, T. 1848a. On some new Musci, collected by Professor W. Jameson on Pichincha. *Lond. J. Bot.*, **7**, 187–99.
- TAYLOR, T. 1848b. On the specific characters of certain new cryptogamic plants, lately received from, and collected by, Professor William Jameson, on Pichincha, near Quito. *Lond. J. Bot.*, **7**, 278–85.
- THÉRIOT, I. 1911. *Breutelia (Eubreutelia) subplicata* Broth. sp. nov. ined. *Revta chil. Hist. nat.*, **15**, No. 1, 38–39.
- THÉRIOT, I. 1915. Sur quelques mousses recueillies au Chili par le Prof. Carlos E. Porter. *Revta chil. Hist. nat.*, **19**, Nos. 1–2, 30–37.

- THÉRIOT, I. 1916. Une rectification. *Bryologist*, **19**, No. 1, 10.
- THÉRIOT, I. 1917a. Note sur une mousse du Chili, *Barbula flagellaris* Sch. *Recl Publs Soc. Havr. Étud. Diverses*, **84**, 55–62.
- THÉRIOT, I. 1917b. Contribution à la flore bryologique du Chili, II. *Revta chil. Hist. nat.*, **21**, Nos. 1–2, 6–37.
- THÉRIOT, I. 1918. Contribution à la flore bryologique du Chili, III. *Revta chil. Hist. nat.*, **22**, Nos. 2–3, 79–94.
- THÉRIOT, I. 1919. Notes bryologiques, [I et II]. *Bull. Soc. bot. Genève*, 2ème sér., **11**, Nos. 1–4, 24–29.
- THÉRIOT, I. 1921a. Contribution à la flore bryologique du Chili, IV. *Revta chil. Hist. nat.*, **25**, 289–312.
- THÉRIOT, I. 1921b. Courte historie de rescapées. *Recl Publs Soc. Havr. Étud. Diverses*, **88**, 29–36.
- THÉRIOT, I. 1922. Deuxième contribution à la flore bryologique de Madagascar. *Recl Publs Soc. Havr. Étud. Diverses*, **89**, 111–32.
- THÉRIOT, I. 1923a. Notes bryologiques [III, IV et V]. *Bull. Soc. bot. Genève*, 2ème sér., **14**, Nos. 1–10 (for 1922), 44–49. [Effective date 1919, according to *Index muscorum*, 5, 72, *Tortula amblyophylla* yet on p. 44 of article "Communiqué en séance du 12 mars 1919. Par suite d'une regrettable omission, les présentes notes n'ont pas été publiées dans le vol. XII du Bulletin, auquel elles étaient destinées: toutefois le tirage à part a été distribué par l'auteur dès 1921 [Réd.]".]
- THÉRIOT, I. 1923b. Contribution à la flore bryologique du Chili, V. *Revta chil. Hist. nat.*, **27**, 9–15.
- THÉRIOT, I. 1924. Contribution à la flore bryologique du Chili, VI. *Revta chil. Hist. nat.*, **28**, 129–39.
- THÉRIOT, I. 1925a. Contribution à la flore bryologique du Chili, VII. *Revta chil. Hist. nat.*, **29**, 287–92.
- THÉRIOT, I. 1925b. Notes bryologiques, [VI et VII]. *Bull. Soc. bot. Genève*, 2ème sér., **17**, 252–57. [Effective date 10 April 1926.]
- THÉRIOT, I. 1926. Contribution à la flore bryologique du Chili, VIII. *Revta chil. Hist. nat.*, **30**, 341–61.
- THÉRIOT, I. 1927a. Le genre *Funaria* au Chili. *Revta chil. Hist. nat.*, **31**, 30–37.
- THÉRIOT, I. 1927b. Mousses récoltées dans l'île Más a Tierra (Juan Fernández) en 1927, par M. Gualterio Looser. *Revta chil. Hist. nat.*, **31**, 256–59.
- THÉRIOT, I. 1928. Contribution à la flore bryologique du Chili, IX. *Revta chil. Hist. nat.*, **32**, 252–55. [Effective date February 1929.]
- THÉRIOT, I. 1929a. La genre *Leptodontium* au Chili. *Revta chil. Hist. nat.*, **33**, 515–17.
- THÉRIOT, I. 1929b. Mousses du Chili austral récoltées par le Prof. Rol. Thaxter, de Cambridge, Mass. *Revue bryol.*, N.S., **2**, Fasc. 3–4, 165–66. [31 March 1930, *fide* title page.]
- THÉRIOT, I. 1929c. Mousses critiques. *Revue bryol.*, N.S., **2**, Fasc. 3–4, 231–33. [31 March 1930, *fide* title page.]
- THÉRIOT, I. 1930. Contribution à la flore bryologique du Chili, X. *Revta chil. Hist. nat.*, **34**, 258–62.
- THÉRIOT, I. 1931. Liste et correction des fautes orthographiques ou autres erreurs contenues dans la 2e édition des *Musci* de Brotherus, in Engler-Prantl, *Die natürlichen Pflanzenfamilien*. *Revue bryol.*, N.S., **4**, Fasc. 4, 170–85. [Effective date September 1932.]
- THÉRIOT, I. 1932. *Barbula ventanica* ou *Barbula ventanae*? *Revue bryol. lichen.*, N.S., **5**, Fasc. 2–3, 140. [Effective date 10 April 1933.]
- THÉRIOT, I. 1933. *Bryoporteria* nouveau genre de la famille Seligeriaceae. *Revta chil. Hist. nat.*, **37**, 23–25.
- THÉRIOT, I. 1934a. Quelques mousses du Chili austral récoltées à freire par Mr. Raul Gillet L., en février 1934. *Revta chil. Hist. nat.*, **38**, 83–85.
- THÉRIOT, I. 1934b. Contribucion à la flore bryologique du Chili, 11e article. *Revue bryol. lichen.*, N.S., **7**, Fasc. 3–4, 167–183. [Effective date 10 April 1935.]
- THÉRIOT, I. 1935a. Contribucion à la flore bryologique du Chili, XII. *Revta chil. Hist. nat.*, **39**, 16–21.
- THÉRIOT, I. 1935b. Quelques nouveautés pour la flore bryologique du Sud-Amérique. *Annls Cryptog. exot.*, **8**, Fasc. 3–4, 135–37. [Effective date November 1936.]
- THÉRIOT, I. 1936. Quatre formes nouvelles de mousses de l'Argentine. (*In* BAUER, E., ed. *Musci europaei et americanii exsiccati. Schedae und Bemerkungen zur 46, Ser. 1936*. [Stadt Obersdorf, Ludwig Richter].)
- THÉRIOT, I. 1939. Bryales uruguayenses. *Revta sudam. Bot.*, **6**, Nos. 3–4, 108–09. [Effective date September 1941.]
- TIXIER, P. 1977. Clastobryoidées et taxa apparentés. *Revue bryol. lichen.*, **43**, Fasc. 4, 397–464.
- TOUW, A. 1971. A taxonomic revision of the Hypnodendraceae (Musci). *Blumea*, **19**, No. 2, 211–354.
- TURNER, D. 1804a. *Muscologiae hiberniae spicilegium*. London, J. White.
- TURNER, D. 1804b. Remarks upon the Dillenian Herbarium. *Trans. Linn. Soc. Lond.*, **7**, 101–15.
- TURNER, D. 1805. Remarks upon some parts of the Hedwigian system of mosses, with a monograph of the genus *Bartramia*. *Ann. Bot., Lond.*, **1**, No. 3, 517–30.
- ULE, E. 1899. *Bryotheca brasiliensis*, herausgegeben von Dr. C. H. Brotherus in Helsingfors, bestimmt von Carl Müller, C. H. Brotherus und C. Warnstorff. *Hedwigia, Beibl.*, **38**, No. 1, 57–59.
- VAN DEN BOSCH, R. B. and C. M. VAN DER SANDE LACOSTE. 1860. *Bryologia javanica seu descriptio muscorum frondosorum Archipelagi indici iconibus illustrata*. Vol. 1. Lugduni-Batavorum, E. J. Brill, Fasc. 24, 141–48.
- VAN DER WIJK, R. and W. D. MARGADANT. 1958. New combinations in mosses. I. *Taxon*, **7**, No. 9, 287–90.
- VAN DER WIJK, R. and W. D. MARGADANT. 1959. New combinations in mosses. II. *Taxon*, **8**, No. 2, 70–75.
- VAN DER WIJK, R. and W. D. MARGADANT. 1960a. NEW COMBINATIONS IN MOSSES. IV. *TAXON*, **9**, No. 2, 50–52.
- VAN DER WIJK, R. and W. D. MARGADANT. 1960b. New combinations in mosses. V. *Taxon*, **9**, No. 6, 189–91.
- VAN DER WIJK, R. and W. D. MARGADANT. 1961. New combinations in mosses. VI. *Taxon*, **10**, No. 1, 24–26.
- VAN DER WIJK, R. and W. D. MARGADANT. 1962. New combinations in mosses. VII. *Taxon*, **11**, No. 7, 221–23.
- VAN DER WIJK, R. and W. D. MARGADANT. 1965. New combinations in mosses. VIII. *Taxon*, **14**, No. 6, 196–98.
- VAN DER WIJK, R., MARGADANT, W. D. and P. A. FLORSCHÜTZ. 1959. Index muscorum. Vol. 1 (A–

- C). *Regnum veg.*, **17**, 548 pp.
- VAN DER WIJK, R., MARGADANT, W. D. and P. A. FLORSCHÜTZ. 1962. Index muscorum. Vol. 2 (D–*Hypno*). *Regnum veg.*, **26**, 535 pp.
- VAN DER WIJK, R., MARGADANT, W. D. and P. A. FLORSCHÜTZ. 1964. Index muscorum. Vol. 3 (*Hypnum*–O). *Regnum veg.*, **33**, 529 pp.
- VAN DER WIJK, R., MARGADANT, W. D. and P. A. FLORSCHÜTZ. 1967. Index muscorum. Vol. 4 (P–S). *Regnum veg.*, **48**, 604 pp.
- VAN DER WIJK, R., MARGADANT, W. D. and P. A. FLORSCHÜTZ. 1969. Index muscorum. Vol. 5 (T–Z, Appendix). *Regnum veg.*, **65**, 922 pp.
- VAN STEENIS-KRUSEMAN, M. J. and W. T. STEARN. 1954. Dates of publication. (*In* VAN STEENIS-KRUSEMAN, C. G. G. J., ed. *Flora malesiana, being an illustrated systematic account of Malaysian flora including keys for determination, diagnostic descriptions, references to the literature, synonymy, and distribution, and notes on the ecology of its wild and commonly cultivated plants*. Djakarta, Noordhoff-Kolff N. V. Ser. 1, 4, No. 5, CLXIII–CCXIX.)
- VENTURI, G. 1872. Muschi raccolti dal Signor Odoardo Beccari nella Terra dei Bogos in Abissinia. *Nuovo G. bot. Ital.*, **4**, Fasc. 1, 7–22.
- VENTURI, G. and A. BOTTINI. 1884. Enumerazione critica del Muschi Italiani. *Atti Soc. crittogram. Ital.*, Ser. 2, **3**, Dispensa 3a, 153–227.
- VERDOORN, F. 1938. Hepaticae selectae et criticae Series IX (1936) et Series X (1937) and Musci selecti et critici Series III (1936) et Series IV (1937). *Annls bryol.*, **10** (for 1937), 124–30.
- VITT, D. H. 1973. A revisionary study of the genus *Macrocoma*. *Revue bryol. lichen.*, N.S., **39**, Fasc. 2, 205–20.
- VITT, D. H. 1976. A monograph of the genus *Muelleriella* Dusén. *J. Hattori bot. Lab.*, No. 40, 91–113.
- VITT, D. H. and M. R. CROSBY. 1972. *Achrophyllum* — a new name for a genus of mosses. *Bryologist*, **75**, No. 2, 174–75.
- WAHLENBERG, G. 1812. Musci. (*In Flora Lapponica exhibens plantas geographicae et botanice consideratas, in Lapponiis Suecicis scilicet Umensi, Pitensi, Lulensi, Tornensi et Kemensi nec non Lapponiis Norvegicis scilicet Nordlandia et Finmarkia utraque indigenas, et itineribus annorum 1800, 1802, 1807 et 1810 denuo investigatas*. Berolini, Taberna libraria scholae realis, 300–400.)
- WAHLENBERG, G. 1826. Musci. (*In Flora Suecica enumerans plantas sueciae indigenas cum synopsi classium ordinumque, characteribus generum, differentiis specierum, synonymis citationibusque selectis, locis regionibusque natalibus, descriptionibus habitualibus nomina incolarum et qualitates plantarum illustrantibus. Post Linnaeum edita. Part 2*. Upsaliae, Palmblad & Co., 697–795.)
- WALTHER, A. and L. MOLENDÖ. 1868. *Die Laubmoose Oberfrankens. Beiträge zur Pflanzengeographie und Systematik und zur Theorie vom Ursprunge der Arten*. Leipzig, Wilhelm Engelmann.
- WALTHER, K. 1934. Untersuchungen über die Variabilität innerhalb des Formenkreises von *Polytrichum juniperinum* Willd. *Annls bryol.*, **7**, 121–56.
- WALTHER, K. 1940. Revision des Verwandtschaftskreises von *Polytrichum piliferum* Schreb. *Reprium nov. Spec. Regni veg.*, **48**, Nos. 12–20, 251–55.
- WARBURG, E. F. 1962. New combinations in British mosses. *Trans. Br. bryol. Soc.*, **4**, Pt. 2, 247–48.
- WARNSTORF, C. 1881a. *Die europäischen Torfmoose. Eine Kritik und Beschreibung derselben*. Berlin, Theobald Grießen.
- WARNSTORF, C. 1881b. *Thuidium delicatulum* (Hedw.) Lindberg, in Steiermark und wahrscheinlich auch in Deutschland verbreitet. *Bot. Zbl.*, **5**, No. 6, 183–85.
- WARNSTORF, C. 1883. Die Torfmoose des v. Flotow'schen Herbarium in königl bot. Museum in Berlin. *Flora, Jena*, **66**, (N.R., **41**), No. 24, 371–80.
- WARNSTORF, C. 1884a. Neue europäische Sphagnumformen. *Hedwigia*, **23**, Ht. 8, 116–25.
- WARNSTORF, C. 1884b. Spagnologische Rückblicke, (Schluss). Verzeichniss der mir ois jetzt aus Europa bekannt gewordenen Sphagnumformen. *Flora, Jena*, **67**, (N.R., **42**), No. 31, 597–611.
- WARNSTORF, C. 1888. *Europäische Torfmoose Serie 1. Flora v. Deutschland (Brandenburg)*. Nearuppin, Waldumpf bei Stendenitz.
- WARNSTORF, C. 1890a. Beiträge zur Kenntniss exotischer Sphagna. *Hedwigia*, **29**, Ht. 4, 179–211; Ht. 5, 213–58.
- WARNSTORF, C. 1890b. Contributions to the knowledge of North American Sphagna. *IV. Bot. Gaz.*, **15**, No. 10, 242–55.
- WARNSTORF, C. 1894. Characteristik und Uebersicht der nord-, mittel- und südamerikanischen Torfmoose nach dem heutigen Standpunkte der Sphagnologie (1893). *Hedwigia*, **33**, Ht. 6, 307–37.
- WARNSTORF, C. 1895. Beiträge zur Kenntnis exotischer Sphagna. *Allg. bot. Z.*, **1**, No. 10, 187–89.
- WARNSTORF, C. 1898. Beiträge zur Kenntniss exotischer und europäischer Torfmoose. *Bot. Zbl.*, **76**, No. 12, 385–90; No. 13, 417–23.
- WARNSTORF, C. 1903. Die europäischen Harpidien. *Beih. bot. Zbl.*, **13**, 388–430.
- WARNSTORF, C. 1904. Laubmoose. (*In Kryptogamenflora der Mark Brandenburg und angrenzender Gebiete herausgegeben von dem Botanischen Verein der Provinz Brandenburg. Bd. 2*. Leipzig, Gebrüder Borntraeger, Ht. 1, 1–240; Ht. 2, 241–432.)
- WARNSTORF, C. 1906. Laubmoose. (*In Kryptogamenflora der Mark Brandenburg und angrenzender Gebiete herausgegeben von dem Botanischen Verein der Provinz Brandenburg. Bd. 2*. Leipzig, Gebrüder Borntraeger, Ht. 5, 833–1160.)
- WARNSTORF, C. 1907. Neue europäische und ausereuropäische Torfmoose. *Hedwigia*, **47**, Ht. 1/2, 76–112.
- WARNSTORF, C. 1911. Sphagnales-Sphagnaceae (Sphagnologia universalis). (*In ENGLER, A. Das Pflanzenreich, Regni vegetabilis conspectus im Auftrage der Königl. preuss. Akademie der Wissenschaften*, Ht. 51. Leipzig, Wilhelm Engelmann.)
- WARNSTORF, C. 1915. Bryophyta nova europaea et exotica. III. Musci. *Hedwigia*, **57**, Ht. 1, 78–80; Ht. 2/3, 81–131.
- WARNSTORF, C. 1916. Pottia-Studien als Vorarbeiten zu einer Monographie des Genus "Pottia Ehrh." sens. str. *Hedwigia*, **58**, Ht. 1/2, 35–80; Ht. 3/4, 81–152.
- WATTS, W. W. and T. WHITELEGGE. 1902. Census muscorum australiensium. A classified catalogue

- of the frondose mosses of Australia and Tasmania, collated from available publications and herbaria records. Part 1, *Proc. Linn. Soc. N.S.W.*, Ser. 2, **27**, Pt. 3, No. 107, Supplement, 1–90.
- [WEBER, F. and D. M. H. MOHR]. 1803. *Index musei plantarum cryptogamarum*. Kiloniae, 1–6. [Authorship of this work, seen in the Linnean Society of London, follows the ascription by Sayre, 1959, 17.]
- WEBER, F. and D. M. H. MOHR. 1807. *Botanisches Taschenbuch auf das Jahr 1807. Deutschlands kryptogamische Gewächse. Abt. 1. Filices, Musci frondosi et Hepatici*. Kiel, Akademischen Buchhandlung.
- WEIMARCK, H. 1937. *Förteckning över Skandinaviens Växter utgiven av Lunds Botaniska Förening. Pt. 2. Mossor*. Lund, Carl Bloms.
- WELCH, W. 1969. Hookeriaceae species and distribution in South America. *Proc. Indiana Acad. Sci.*, **78** (for 1968), 396–405.
- WHELDON, J. A. 1921. Key to the Harpidoid Hypna. *Naturalist, Hull*, (for 1921), No. 774, [No. 548 of current series], 245–48.
- WILLIAMS, R. S. 1901. Contributions to the botany of the Yukon Territory. 2. An enumeration of the mosses collected. *Bull. N.Y. bot. Gdn*, **2**, No. 6, 105–48.
- WILLIAMS, R. S. 1903a. Bolivian mosses. Part I. *Bull. N.Y. bot. Gdn*, **3**, No. 9, 104–34.
- WILLIAMS, R. S. 1903b. Additional mosses of the Upper Yukon River. *Bryologist*, **6**, No. 4, 61–62.
- WILLIAMS, R. S. 1915. Mosses from the west coast of South America. *Bull. Torrey bot. Club*, **42**, No. 7, 393–404.
- WILLIAMS, R. S. 1916. Peruvian mosses. *Bull. Torrey bot. Club*, **43**, No. 6, 323–34.
- WILLIAMS, R. S. 1924. *Orthorrhynchium chilense* sp. nov. *Bryologist*, **27**, No. 6, 87–88. [Effective date 15 January 1925, *fide* title page of Vol. 28, No. 1.]
- WILLIAMS, R. S. 1930. Mosses collected in Brazil and Argentina by J. N. Rose in 1915. *J. Wash. Acad. Sci.*, **20**, No. 19, 465–75.
- WILSON, W. 1844. *Grimmia orbicularis*. (*In HOOKER, W. J. Supplement to the English botany of the late Sir J. E. Smith and Mr. Sowerby*. London, J. Sowerby, J. W. Salter and Mrs. Sowerby, Vol. 4, Tab. 2888.)
- WILSON, W. 1846. Remarks on the new species of Musci from Quito and Swan River, indicated by Dr. Taylor in the London Journal of Botany, Vol. V, p. 41. *Lond. J. Bot.*, **5**, 447–55.
- WILSON, W. 1847. Brief characters of some new mosses, collected in New Granada by Mr. W. Purdie. *Lond. J. Bot.*, **6**, 289–92.
- WILSON, W. 1848. Brief characters of *Aulacopilum*, a new genus of mosses, from New Zealand. *Lond. J. Bot.*, **7**, 90–91.
- WILSON, W. 1854. Musci, Juss. (*In HOOKER, J. D. The botany of the Antarctic voyage of H.M. Discovery ships Erebus and Terror in the years 1839–1843, under the command of Captain Sir James Clark Ross, Kt., R.N., F.R.S., & L.S. etc. Vol. II. Flora Novae-Zelandiae. Pt. 2. Flowerless plants*. London, Lovell Reeve, No. 5, 57–80; No. 6, 81–125.)
- WILSON, W. 1855. *Bryologia Britannica; containing the mosses of Great Britain and Ireland, systematically arranged and described according to the method of Bruch and Schimper, with illustrative plates*. London, Longman, Brown, Green and Longmans.
- WILSON, W. 1859. Musci, Juss. (*In HOOKER, J. D. The botany of the Antarctic voyage of H.M. Discovery ships Erebus and Terror, in the years 1839–1843, under the command of Captain Sir James Clark Ross, Kt., R.N., F.R.S. & L.S., etc. Vol. III. Flora Tasmaniae. Pt. 2. Monocotyledones and Acotyledones*. London, Lovell Reeve, No. 8, 161–221.)
- WILSON, W. and J. D. HOOKER. 1845. Musci. (*In HOOKER, J. D. The botany of the Antarctic voyage of H.M. Discovery ships Erebus and Terror in the years 1839–1843, under the command of Captain Sir James Clark Ross, Kt., R.N., F.R.S., &c. Vol. I. Flora Antarctica. Pt. I. Botany of Lord Auckland's group and Campbell's Island*. London, Reeve Brothers. [Reprinted 1963, Weinheim, J. Cramer], No. 8, 117–43.)
- WILSON, W. and J. D. HOOKER. 1847. Musci. (*In HOOKER, J. D. The botany of the Antarctic voyage of H.M. Discovery ships Erebus and Terror, in the years 1839–43, under the command of Captain Sir James Clark Ross, Kt., R.N., F.R.S., &c. Vol. I. Flora Antarctica. Pt. 2. Botany of Fuegia, The Falklands, Kerguelen's Land, etc.* London, Reeve Brothers. [Reprinted 1963. Weinheim, J. Cramer], No. 22, 395–96; No. 23, 397–423.)
- WITHERING, W. 1801. *A systematic arrangement of British plants; with an easy introduction to the study of botany. Vol. III. 4th edition*. London, T. Cadell, Jun. and W. Davies, G. and J. Robinson, Wynne and Scholey, J. Walker, J. Cuthell, and James Wallis.
- WRIGHT, C. H. 1905. The mosses of the South Orkneys. (*In BROWN, R. N. R., WRIGHT, C. H. and O. V. DARBSHIRE. The botany of the South Orkneys. Trans. Proc. bot. Soc. Edinb.*, **23**, Pt. 1, 103–04.)
- ZANDER, R. H. 1972. Revision of the genus *Leptodontium* (Musci) in the New World. *Bryologist*, **75**, No. 3, 213–80.
- ZANDER, R. H. 1977. The tribe Pleuroweisiae (Pottiaceae, Musci) in middle America. *Bryologist*, **80**, No. 2, 233–69.
- ZANDER, R. H. 1978. New combinations in *Didymodon* (Musci) and a key to the taxa in North America north of Mexico. *Phytologia*, **41**, No. 1, 11–32.
- ZETTERSTEDT, J. E. 1876. *Musci et Hepaticae Finmarkiae circa sinum altensem crescentes. K. svenska VetenskAkad. Handl.*, **13**, No. 13, 42 pp.