

Guide to ECOLOGICAL SITES OF THE KAZAN UPLAND SUBREGION



ECOLOGICAL SITES OF THE KAZAN UPLAND SUBREGION

First approximation

This is the first approximation of the Kazan Upland subregion.

2017

Prepared by:

**Michael G. Willoughby,
Margaret Meijer
and
Dave J. Downing**

Cover Photo Credits: (Natural Regions and Subregions of Alberta 2006, A. Landals, D. Vujnovic, L. Allen)

Ecological Site Photo Credits: Allen et al. (2002,2004)

Environment and Parks
Land Policy
Policy and Planning
Edmonton

ISBN No.: 978-1-4601-3321-7 (Printed Edition)

ISBN No.: 978-1-4601-3320-0 (PDF)

For more information regarding this guide contact:

Michael G. Willoughby
Range Management
Alberta Environment and Parks
4th Flr. Great West Life Building
Edmonton, Alta.
T5K 2M4
email: mike.willoughby@gov.ab.ca

This publication is the property of the Government of Alberta and is available under the Alberta Open Government Licence (<http://open.alberta.ca/licence>) and the publication is available online at: <http://open.alberta.ca/publications/9781460133200>

© Her Majesty the Queen in Right of Alberta, as represented by the Minister of Environment and Parks, 2017

Table of Contents

ECOLOGICAL SITES OF THE KAZAN UPLAND SUBREGION	i
Table of Contents.....	iii
Executive Summary	vi
Acknowledgements.....	vii
Introduction and Background	1
Physiography, Climate and Soils.....	2
Approach and Methods of Classification	3
Correlation of Soils and Ecological Sites	6
Guidelines for Determining Ecological Sites	7
How to use the Guide.....	9
Results.....	11
General Ecological Descriptions.....	12
Plant Community Keys.....	15
Plant Community Tables	17
KU Kazan Upland (n=73).....	20
a lichen stone fields (n=1).....	21
a1 lichen stonefields (n=1).....	22
KUA1 Lichen stonefield (n=1).....	23
aa Sand heather/moss (xeric/poor) (n=0).....	24
aa1 grass/Sand heather (n=0).....	25
b bearberry/lichen(subxeric/poor) (n=18).....	26
b1 bearberry/lichen - Pj (n=15).....	27
KUE1 Pj/Lichen (n=9).....	28
KUE2 Pj/Bearberry (n=6).....	29
b2 bearberry/lichen - shrub (n=3).....	30
KUA2 Bearberry/Indian ricegrass (n=2).....	31
KUB1 Juniper/Lichen(Pj) (n=1).....	32
c blueberry(submesic/medium) (n=12).....	33
c1 blueberry - Pj-Aw(Bw) (n=7).....	34
KUD3 Pj-Aw(Bw)/Bearberry-Bog cranberry (n=4).....	35
KUE3 Pj/Blueberry (n=3).....	36
c2 blueberry - Aw(Bw) (n=1).....	37
KUC1 Aw/Rose/Spreading dogbane (n=1).....	38
c6 blueberry grassland (n=4).....	39
KUA3 Hay sedge-Slender wheatgrass (n=4).....	40
c7 tame/disturbed (n=0).....	41
d Labrador tea-mesic(mesic/poor) (n=10).....	42
d1 Labrador tea-mesic Pj-Sb (n=10).....	43
KUC4 Aw-Pj-Sb/Labrador tea (n=1).....	44
KUD1 Sb-Aw/Moss (n=3).....	45
KUE11 Pj/Bog cranberry/lichen (n=1).....	46
KUE4 Pj- Sb/Bog cranberry/lichen (n=5).....	47
e buffaloberry/alder(mesic/medium) (n=16).....	48
e1 buffaloberry-alder/ Aw (n=8).....	49
KUC2 Aw(Bw)/Alder (n=3).....	50
KUC3 Aw/Buffaloberry (n=4).....	51
KUC7 Aw-Pb/Saskatoon-Red Osier Dogwood/Spreading dogbane (n=1).....	52
e2 buffaloberry-alder/ Pj-Sw-Sb-Aw-Bw (n=5).....	53
KUD2 Aw-Sw/Buffaloberry (n=3).....	54
KUD4 Aw- Pj/Alder (n=2).....	55

e3	buffaloberry-alder shrubland (n=0)	56
e4	buffaloberry-alder/ Pj-Sw-Sb (n=3)	57
	KUE5 Pj/Alder (n=1)	58
	KUE6 Sb-Sw/Moss (n=2)	59
f	red osier dogwood/horsetail (subhygric/rich) (n=2)	60
f1	red osier dogwood/horsetail Sw (n=0)	61
f2	red osier dogwood/horsetail Pb-Sw (n=0)	62
f3	red osier dogwood/horsetail - Pb-Aw(Ba) (n=2)	63
	KUC5 Pb/Alder-Red osier dogwood/Horsetail (n=1)	64
	KUC6 Bw-Aw-Pb/Willow (n=1)	65
f4	shrub (n=0)	66
f5	tame/disturbed (n=0)	67
g	Labrador tea-hygric(hygric/medium) (n=2)	68
g1	Labrador tea-hygric Sb-Pj (n=2)	69
	KUE10 Lt-Sb/River alder/Sedge (n=1)	70
	KUE7 Sb/Alder/Tufted moss (n=1)	71
h	bog (subhygric/very poor) (n=3)	72
h1	bog - treed (n=1)	73
	KUE8 Sb/Labrador tea/peat moss (n=1)	74
h2	bog - shrubby (n=2)	75
	KUB2 Labrador tea/Peat moss (Sb) (n=2)	76
h3	bog -graminoid (n=0)	77
i	poor fen (subhygric/medium) (n=3)	78
i1	poor fen - treed (n=1)	79
	KUE9 Sb-Lt/Labrador tea/Golden moss (n=1)	80
i2	poor fen - shrubby (n=0)	81
i3	poor fen - graminoid (n=2)	82
	KUA4 Shooting star/Mat muhly (saline) (n=1)	83
	KUA5 Sedge/Brown moss (saline) (n=1)	84
j	rich fen (subhygric/rich) (n=3)	85
j1	rich fen - treed (n=0)	86
j2	rich fen - shrubby (n=3)	87
	KUB3 Willow/Marsh reed grass (n=1)	88
	KUB4 Willow/Sedge (n=1)	89
	KUC8 Willow/Brown moss/ Aw(Bw) (n=1)	90
j3	rich fen - graminoid (n=0)	91
k	marsh (hygric/rich) (n=3)	92
k1	marsh (n=3)	93
	KUA6 Water parsnip/Northern manna grass (n=1)	94
	KUA7 Northern quillwort (n=1)	95
	KUA8 Water arum/Sedge (n=1)	96
	Literature Cited	97

List of Figures

Figure 1. Edatope grid and ecological sites for the Kazan Upland subregion.	14
---	----

List of Tables

Table 1. Kazan Upland Communities..... 17

Executive Summary

The Kazan Upland Natural Subregion occurs in the far northeastern corner of Alberta and occupies 1.5% of the province (Natural Regions Committee 2006). The main area lies north of Lake Athabasca. It is bordered on the east and north by the Alberta–Saskatchewan and Alberta–Northwest Territories boundaries, respectively, on the west by the Slave River, and on the south by Lake Athabasca. There is a small outlier east of the Athabasca River between Fort McMurray and Fort Chipewyan embedded within the Athabasca Plain Natural Subregion. Elevations range from about 150 m to over 400 m. Approximately 60% of the landscape of the Kazan Upland subregion is exposed precambrian bedrock (Natural Regions Committee 2006).

This guide represents the analysis of 73 plots described in the Kazan Upland subregion. The 73 plots represent 34 community types. These types are split into: A. Native grasslands 8 community types B. Native shrublands 4 community types C. Deciduous types 8 community types D. Mixedwood types 4 community types E. Conifer types 10 community types. The dominant plant species, canopy cover, soil and environmental conditions are outlined for each type.

Acknowledgements

Landscape classification is the process of breaking the landscape into definable and manageable pieces through a hierarchical classification. In the early 1990's the forested landscape of Alberta was classified using a well organized hierarchical system (Archibald/ Beckingham / Klappstein). Unfortunately this left about 50% of the remaining natural landscapes of the province unclassified. Starting in the late 1990's rangelands undertook efforts to classify the rangelands of Alberta. A need for consistency across the province was recognized. Therefore a hierarchical classification that built on the forested classification was used for all forest dominated subregions in the province.

These classifications involve taking large amounts of vegetation, soils and landscape inventory data taken from the ECOSYS database and compiling the data into succinct descriptive summaries for each ecological site, ecological site phase and plant community within a subregion.

In 2010 funding was provided by Policy and Planning Division, Alberta Environment and Parks in order to produce hard copy pdf documents from the ECOSYS website.

Introduction and Background

The province of Alberta is covered by a broad spectrum of vegetation regions from prairie in the South, to alpine vegetation in the mountains and dense forests and extensive wetlands in the Central and Northern parts of the province. These broad vegetation regions have been classified into 6 natural regions and 21 subregions for the province (Natural Regions and Subregions of Alberta 2006). Each of the regions consists of groups of plant communities which are influenced by environmental conditions and human impacts. Intensive management of these regions requires the ability to recognize the vegetative communities that have similar productivities and respond to disturbance in the same way. These ecological classification guides are highly regarded by most resource managers as a tool which may assist in decision making processes related to livestock grazing, prime habitat for wildlife, productive watersheds and recreational areas, addressing biodiversity and conservation matters and numerous other potential related topics.

The purpose of this guide was to develop a framework that would easily group the ecological sites and vegetative community types in the Kazan Upland Natural Subregion of the province. Ecological site classification helps to organize our current understanding about ecosystem function. This organization is achieved by grouping research plots into similar and functional units that respond to disturbance in a similar and predictable manner.

The ecological site classification system outlined in this document organizes ecological information into a format that facilitates understanding and provides a structure for ecologically based management. The system has been developed primarily as a field tool to complement the user's knowledge about ecological site classification, soil description, and plant identification. The objectives of the ecological site classification are to provide a consistent overview of the common vegetation communities and site characteristics in order to:

1. to facilitate the application of ecological information to decisions on a wide variety of activities within the realm of land resource management
2. to facilitate the collection and organization of information to expedite the development of resource management applications and decision support systems
3. to promote communication among resource managers and between managers and the public
4. to provide a common basis for integrated planning, and
5. to reduce resource management costs by integrating ecological information into the decision-making process.

This guide builds on the work outlined in the Field guide to Ecosites of Northern Alberta (Beckingham and Archibald 1996) for the Canadian Shield Natural Region. In 2006 (Natural Regions and Subregions of Alberta 2006) the original Canadian shield region was split into the Athabasca Plain and Kazan Upland subregions. This guide outlines the analysis of 73 plots described in the Kazan Upland subregion.

Physiography, Climate and Soils

Please note this summary of Natural Subregion characteristics is extracted directly from the Natural Subregions guide (Natural Regions Committee 2006) and is presented here for the reader's convenience.

The Kazan Upland Natural Subregion is 9719 km², covering only 1.5% of the province. It features short, warm summers where July is the warmest month, but possesses the coldest winters of any Natural Subregion in Alberta due to continental polar and continental arctic weather systems. July is also the month of maximum precipitation. Winter snowfall accounts for 40% of the annual total (Natural Regions Committee 2006). The Alberta Climate Model indicates the Kazan Upland Natural Subregion receives about the same annual precipitation as the Dry Mixedgrass Natural Subregion, and has a potential summer moisture deficit (summer moisture index) comparable to that of the Northern Fescue Natural Subregion. Figure 4-6.1 and Table 3-2 in Natural Regions and Subregions guide of Alberta provides monthly and annual climate data summaries. The prevalence of rock barrens and well to rapidly drained glacial deposits, together with low rainfalls, produce conditions that are favorable for non-vascular and vascular plants that are adapted to dry conditions.

Extensive outcrops of Precambrian bedrock, composed of Archean and Proterozoic granitoids define the Kazan Upland subregion. The westernmost edge of the Canadian Shield, defines the limits of the Kazan Upland; approximately, 60 percent of the landscape is exposed bedrock. The surficial landscape is comprised of about 45 percent glacially scoured rock, about 40 percent sandy and gravelly ice-contact glacial drift, and about 15 percent organic accumulations in low-lying areas. Parent materials are ice-scoured bedrock and coarse textured glacial.

The terrain is hummocky to rolling with relief up to 50 m. Nonsoils occur across about half the area. Surficial deposits between bedrock outcrops are mainly coarse and acidic. Soils on these materials are predominantly eluviated Dystric Brunisols with associated Orthic, Gleyed and Lithic Subgroups, the latter on very thin deposits over bedrock. Wetlands are a complex of Typic and Fibric Mesisols, many with Terric Subgroups, Peaty Gleysols also occur (Natural Regions Committee 2006).

Approach and Methods of Classification

Approach:

Ecological classification hierarchy and terminology

The system of classification in this guide was initially based on the community type approach of Mueggler (1988). Mueggler's system was chosen over the habitat type approach (Daubenmire 1952) or ecosystem association approach (Corns and Annas 1986) because it could classify plant communities irregardless of their successional status. However, as the philosophy of proper functioning condition of a site evolved, it became apparent (through data analysis) that there was a need to also organize the various plant communities based on their response to disturbance (i.e. disturbance vs. natural succession) within an area under similar environmental influences.

It was determined that the ecosystem classification system developed by Corns and Annas (1986) and Beckingham et al. (1996) could accommodate this additional requirement. Thus, this classification system is a combination of Mueggler (1988) and Beckingham et al. (1996). Consequently, this guide adopts a similar ecological unit classification hierarchy (ecodistrict, ecosection, ecological site, ecological site phase, plant community). The ecological classification system is nested within Alberta's geographically based natural region and subregion classification system (Natural Regions Committee 2006).

Ecodistrict

The ecodistrict level is a unique pattern of slope, landform, soils and vegetation. Mapping of this unit is usually done at a scale of 1:1,000,000 to 1: 250,000 within the whole province (Strong and Anderson 1980). This level of the classification hierarchy is spatially defined and may or may not be unique to a subregion.

Ecosection

The natural subregion used by the Alberta Government is equivalent to the ecoregion defined by the Canada Committee on Ecological Land Classification (CCELC) as part of a multi-level national mapping system for Canada and that was used for integrated resource planning in Alberta (Marshall et al. 1996). Similarly, the ecodistrict as presently used and its associated scale of mapping is equivalent to the ecodistrict defined by the CCELC. However, the ecosection has a somewhat different meaning in the current context than it did in the national system or than it did when it was applied to integrated planning maps in Alberta in the 1980's and 1990's. For those mapping projects, the ecosection was a subdivision of the ecodistrict and was mapped at 1:20 000 to 1:50 000 as a more specific delineation of recurring landform and vegetation patterns, usually with reference to major community type groups or soil subgroups. In the current scheme, the ecosection is a term used to define one ecodistrict or an aggregation of ecodistricts that represent one or more climatic variants within a natural subregion; therefore, its mapping scale is flexible. This level of the classification system is not spatially defined. The ecosection is a unique pattern of slope, landform, soils and vegetation and may also represent a slight change in the climate of a subregion. Mapping of this unit is usually done at a scale of 1:1,000,000 to 1:100,000 and can be a grouping of ecodistricts or at smaller scales outliers in a subregion. For example the Lower Boreal Highlands subregion is split into the foothills and boreal ecosections which are influenced by their proximity and location within the Boreal and Foothills Natural Regions. Spatially these two ecosections are split by grouping ecodistricts. In contrast an example of a smaller scale ecosection (1:100,000) is the Cypress Hills outlier of the Montane subregion. Subregion ecosections have a characteristic sequence of ecological sites according to soil moisture regime (SMR) and, to a lesser degree, soil nutrient regime (SNR). Currently there is no ecosection described for this subregion.

Ecological Site

Ecological sites are ecological units that develop under similar environmental influences (climate, moisture, nutrient regime). They are groups of one or more ecological site phases that occur within the same portion of

the edatope (moisture/nutrient grid). Each ecological site is designated with a small letter. These letters range from "a" the driest ecological site and the last letter being the wettest. Each ecological site has been given a name that conveys some information about the ecology of the unit. Ecological sites are typically named after plant species that are common or typical of the site (eg. e low-bush cranberry). The plant that the ecological site is named after, however, may not be present in every plot or stand belonging to the site. Ecological site in this classification system, is a functional unit defined by moisture and nutrients. It is based on the combined interaction of biophysical factors which together dictate the availability of moisture and nutrients for plant growth. Thus, different ecological sites vary in their moisture and nutrient regime and have similar characteristic plants and soils.

Ecological site phase

An ecological site phase is a subdivision of the ecological site based on the dominant species in the canopy. On lowland, meadow or grassland sites where tree canopy is not present the tallest structural vegetation layer with greater than 5% cover determines the ecological site phase. Generally, ecological site phases are mappable units and spatial ecological site phase land cover datasets have been developed from AVI (Alberta Vegetation Inventory) (Derived Ecosite Phase (DEP)) and PLVI (Primary Land Vegetation Inventory). Ecological site phases are identified by the ecological site letter "a" along with a number "a1" representing the phase within the ecological site. Ecological site phases have a distinct range in canopy composition, lower strata plant species and pedogenic processes. The ecological site phase has a strong ecological basis and correlates well with forest cover on forest inventory maps.

Plant community type

Ecological site phases may be subdivided into plant community types, which are the lowest taxonomic unit in the classification system. While plant community types of the same ecological site phase share vegetational similarities they differ in their understory species composition and abundance. Generally the plant community types are named by combining the name of the dominant plant species in each structural layer (eg. White spruce/Horsetail/Moss)

Methods:

Plant community classification

Data used to create this guide were collected from field plots within the Kazan Upland subregion. Seventy three plots were used to create the classification for this subregion. Field inventory for these plots generally followed the Ecological Land Survey Site Description Manual (2003) and uses various site, vegetation and soils forms. Plot data was analyzed using the multivariate analysis techniques of classification and ordination. Classification is the assignment of plots to classes or groups based on the similarity of species within each plot. A polythetic agglomerative approach was used to group the samples. This technique assigns each plot to a cluster which has a single measure. It then agglomerates these clusters into a hierarchy of larger and larger clusters until finally a single cluster contains all the plots (Gauch 1982). The cluster analysis was performed in SAS with Euclidean distance used as the Cluster Distance Measure and Ward's method was used in the Group Linkage Method. The groupings generated in cluster analysis were overlain on the site ordination to determine final groupings.

Ordination was used to find relationships among species, communities and environmental variables. Ordination reduces the dimensionality of the data to 1-3 most important axes to which environmental gradients can be assigned. The ordination technique used in the analysis of the data was DECORANA (Detrended Correspondence Analysis). Once final groupings were determined on the ordination specific environmental variables can be assigned to the variation outlined on the ordination axes.

Plant community summaries were generated by averaging plant species composition, range in composition, and percent constancy of occurrence, among vegetation inventory plots which were part of a community type. Environmental data was sorted into the same plant community groupings to create the plant community descriptions outlined in this guide. The number of sample plots on which the description was based is also

provided (e.g. n=16).

Ecological Health and Ecological Status Score

Ecological health is determined by comparing the functioning of ecological processes on an area (e.g. plant community polygon) of to a standard (i.e. Reference Plant Community) described within an ecological site description. An ecological site is defined by the Task Group on Unity and Concepts (1995) as, “a distinctive kind of land with specific physical characteristics that differs from other kinds of land in its ability to produce a distinctive kind and amount of vegetation”. This guide can be used to determine the appropriate reference plant community, within an ecological site, for a health assessment. We use health terminology (healthy, healthy with problems, or unhealthy), to rank the ability of the land to perform certain ecological functions. These functions include: net primary production, maintenance of soil/site stability, capture and beneficial release of water, nutrient and energy cycling and plant species functional diversity. For a detailed description on how to assess health for various plant communities please refer to “Rangeland Health Assessment for Grassland, Forest and Tame Pasture” (Adams et al. 2009).

An ecological status score (i.e. the integrity of the plant community composition compared to the reference plant community) has been added to each community type description. These values are based on what is currently known about how a reference plant community (RPC) responds to various kinds and levels of disturbance or successional processes. The values indicate how a particular plant community fits in the state and transition model relative to the RPC. If an experienced observer wishes to estimate the health of a plant community without completing a health form, (e.g. a small riparian area), these values can be used as a guide. Occasionally there are 2 options provided for the ecological status score. This was done for two reasons: 1) to express the range of divergence from the RPC possible for a particular plant community; or 2) to allow for different health forms to be used in communities with variable shrub or tree cover (e.g. on sites with high woody cover and/or an obvious LFH layer use the forest rangeland health form and the corresponding ecological status score; on sites dominated by herbaceous cover and/or an obvious herbaceous litter layer use the native grassland form). Late seral plant communities tend to be superior in the efficient capture of solar energy, in cycling of organic matter and nutrients, in retaining moisture, in supporting wildlife habitat values and in providing the highest potential productivity for the site (Adams et al. 2009). In contrast, early seral disturbed stages ability to produce a distinctive kind and amount of vegetation” represent plant communities with diminished ecological processes, which are less stable and more vulnerable to erosion and invasion by weeds and non-native species. In most cases these late seral plant communities are used as the RPC, but sometimes management goals influence the choice of RPC (e.g. a cut block to be maintained as untimbered rangeland).

Correlation of Soils and Ecological Sites

Please note this summary of Natural Subregion characteristics is extracted directly from the Natural Subregions guide (Natural Regions Committee 2006) and is presented here for the reader's convenience.

The acidic characteristics of granitic bedrock exposures, the well to rapidly drained glacial drift, and frequent fires significantly impact the vegetation in this subregion. Communities are associated with dry rock barrens are widespread, and although species diversity is low in any given locale, the variety of habitat results in high species variety across the barrens. Various lichen communities occupy south-facing and steep rock faces and slopes--many found only in the Kazan Upland (Natural Regions Committee 2006). "Pocket" communities grow in rock crevices and sheltered locations where mineral soil has accumulated and moisture conditions are right. The species associated with these types of areas are stunted jack pine and Alaska birch forming open stands, with a sparse understory of bearberry, ground juniper, bog cranberry, and a variety of drought-tolerant ferns and other herbs, mosses, and lichens (Natural Regions Committee 2006).

Places with coarse textured, rapidly drained and dry sandy or gravelly soils support more vigorous pine growth. The driest sites are vegetated by open jack pine stands with a patchy carpet of lichens below. Moister sites support more diverse understories of green alder, common blueberry, bearberry, common Labrador tea, Canada buffaloberry, bunchberry, and other herbs and feathermosses. Aspen, Alaska birch, and black spruce are locally common in places.

Small lakes occupy about 10% of the Kazan Upland Natural Subregion, and wetlands are common in the lower relief western half, covering approximately 20%. Adjacent to these areas, communities of moister adapted species can develop. These include aspen, balsam poplar, Alaska birch, white spruce, and a diverse shrub and forb understory. Bog communities are the dominant wetland type. Black spruce forms open-to-dense stands with an understory of common Labrador tea, leatherleaf, bog cranberry, cloudberry, and peat moss on Organic soils. Permafrost is discontinuous but widespread. Nutrient rich wetlands typically have open forests of tamarack, willow, dwarf birch, sedges, and rich-site mosses.

Nonsoils (e.g., bedrock outcrops) occur across about half the area. Surficial deposits between bedrock outcrops are mainly coarse and acidic. Soils on these materials are predominantly eluviated Dystric Brunisols with associated Orthic, Gleyed and Lithic Subgroups, the latter on very thin deposits over bedrock. Wetlands are a complex of Typic and Fibric Mesisols, many with Terric Subgroups. Peaty Gleysols also occur. Appendix 7 summarizes the proportional occurrence of soil types in the Kazan Upland Natural Subregion (Natural Regions Committee 2006).

Guidelines for Determining Ecological Sites

Alberta currently uses two ecological classification methods to determine ecological sites. In the agricultural settlement area of the Province, resource managers can determine site soil conditions using AGRASID (Agricultural Region of Alberta Soil Inventory Database). In the Rocky Mountain, Foothills and Boreal Natural Regions, the Ecological Landscape Classification approach incorporates both vegetation and site conditions (climate, soils and geology) into a hierarchical ecological unit classification (e.g. subregion, ecodistrict, ecosection, ecological site, ecological site phase, plant community) (Strong and Thompson 1995). Ecological sites are areas of similar climate, moisture and nutrient regimes. The combination of moisture and nutrient regimes can be represented on a two-dimensional grid called the edatope grid. The edatope grid is a two-dimensional table with soil moisture regime increasing from bottom to top along the vertical axis and soil nutrient regime increasing from left to right on the horizontal axis. Soil moisture regime (SMR) is defined as the average amount of soil water available annually for evapotranspiration by vascular plants (Meidinger and Pojar 1991). The SMR uses nine classes to define the available soil moisture, which range from the driest (very xeric) to the wettest (hydric). Soil nutrient regime (SNR) is defined as the amount of essential soil nutrients that are available to vascular plants over a period of several years (Meidinger and Pojar 1991). SNR is broken down into five classes that range from A (very poor) to E (very rich). Generally ecological sites are named from low moisture/low nutrient to high moisture/high nutrient. Ecological sites within a Natural subregion are defined unique combinations of soil moisture and nutrients. These conditions, in addition to climate, terrain, and elevations create conditions favourable to specific suite of plants referred to as Indicator species. For example a site with a subxeric moisture regime and poor nutrient regime site is characterized by the "a" [bearberry (subxeric/poor)] ecological site. A resource manager can review the indicator plant species of the ecological site, plant community types, soils and site conditions to see if the plant community in question fits the general descriptions. The following steps provide a framework for determining ecological sites.

Step 1 Review background information and pre-stratify the area to be classified

Review information about the area of interest to learn what you can about the landscape and ecology. Consult the natural subregions and Derived Ecosite Phase (DEP) or Primary Land Vegetation Inventory (PLVI) maps to ensure you are using the correct subregion guide. DEP and PLVI classification will also give you the common ecological site phase for a particular forest polygon.

Step 2 Carry out a quick reconnaissance of the site to be classified

Take note of the variability and relationship between topography and position on the landscape and the general plant species distribution including trees and understory. Check the DEP and PLVI ecological site phase maps.

Step 3 Choose a location that appears to be representative of the area to be classified

Locate an area for your assessment that appears to be representative of the site to be classified, and is homogeneous in slope, plant cover, and overstory canopy conditions as possible. Avoid locating the sample in areas that have received significant natural or artificial disturbance. Also avoid ecotone areas or relatively small areas that are transitional between homogenous ecological units such as slope breaks.

Step 4 Determine the plant species composition and abundance

Determine the plant species composition and abundance within a 10x10 m plot. Also record any species that appear to be representative of the ecological unit but occur outside the plot within the same slope position and on the same parent materials. Abundance is estimated by determining the amount of ground area that is covered by the plant species when its canopy is projected onto the ground surface (Ecological Land Survey Site Description Manual 2003).

Step 5 Determine the important soil properties

To collect soils data, a soil pit must be dug or augered. In most cases a soil pit 60 cm deep will be adequate. A deeper pit is required when the soil has a coarse to moderately coarse texture. In these cases the pit is dug deeper to see if there are finer-textured layers that are influencing ecological function below the 60 cm of coarse material. A deeper pit is also required when the plant community on the site cannot be explained by the site conditions and soil conditions above 60 cm. The minimum soils data that should be collected within a plot to classify it correctly are organic matter thickness, humus form, Ah horizon thickness, surface texture, effective texture, presence of seepage, depth to mottles, depth to gley, coarse fragment content, parent material/landform and drainage.

Step 6 Determine important site properties

Important site variables that should be collected include topographic position, slope, aspect, moisture regime, and nutrient regime.

Step 7 Determine the natural subregion, ecological site, ecological site phase and plant community type.

There are several ways to determine the ecological site, ecological site phase and plant community type. The first way is to assign an ecological classification to a site is to use the field data collected and go through the various subregion guides to identify the ecological site. You can also use the dichotomous keys to ecological site and ecological site phase. Once you find a potentially correct plant community type, check the soil, site and vegetational characteristics of your site to make sure it matches the ecological site, ecological site phase and plant community type on the various fact sheets. To consider all ecological site choices, you must compare the characteristics of your site, with the descriptions on the fact sheets for all ellipses that overlap the moisture and nutrient classes of your site on the edatope grid for the subregion and adjacent subregions within the area (Ecological Land Survey Site Description Manual 2003).

How to use the Guide

Organization of the guide

This guide is an expansion of the Ecosites of Northern Alberta guide (Beckingham and Archibald 1996). It contains new information and it is recommended that the reader has access to relevant information from both guides. The community types in this guide are closely related to the ecosites and ecosite phases outlined in Ecosites of Northern Alberta (Beckingham and Archibald 1996), and are similarly arranged (e.g. Table 1). Table 1 is a reproduction of Figure 20 in Ecosites of Northern Alberta with community types in this guide further separated into reference range plant communities, successional communities and harvesting and fire communities. The "Successional community types" or "Harvesting and Fire succession" categories outline the successional sequence the community types undergo with heavy grazing pressure, harvesting or fire disturbance.

The bulk of this guide consists of ecological sites, ecological site phase and vegetation community descriptions which include information on the dominant plant species, canopy cover and environmental conditions. Where available, plant community successional information (ie response to grazing) has been included to help you determine rangeland health and the successional relationships of an ecological site.

Ecological units within a subregion are classified by their position on the edatopic grid.

The information in this guide is presented and named by:

1. Subregion/Ecological area
 - a. Kazan Upland (KU)
2. Dominant cover type
 - a. A-grassland
 - b. B-shrubland
 - c. C-deciduous
 - d. D-mixedwood
 - e. E-conifer
3. A number- Generally, communities are named/numbered from low moisture /nutrient status to high moisture/nutrient status. For example, KUE1 Pj/Lichen community on the "b" subxeric/poor ecological site, while KUE10 Lt/Alder/Sedge community type on the "j" subhydric/rich ecological site.

Eg. KUE1 Pj/Lichen

NOTE: As additional information is collected and new ecological units are identified and described, an attempt is made to fit them into the pre-existing ones.

How to read the fact sheets

The field guide contains 4 types of fact sheets: One for ecosection, one for ecological site, one for ecological site phase and one for plant community type.

Ecosection

There is an identification code at the top of the ecosection fact sheet and a name followed by the number of sample sites (pg 20). Each ecosection has been given a name that conveys information about the location of the unit and are frequently named after a general location within the subregion (Ecosection: Cypress Hills (Mc) of the Montane subregion). A short text description of the site is given under the General Description (pg 20), this is followed by a picture or a cross section diagram and map of the ecosection(pg 20). The section on successional relationships gives a brief note about the spatial locations and differences in ecosections (pg

20). This is followed by a list of environmental variables (elevation), ecodistricts and ecological sites associated with the ecosection (pg 20).

Ecological site

There is an identification letter at the top of the ecological site fact sheet and a name, moisture and nutrients followed by the number of sample sites (pg 21). Each ecological site has been given a name that conveys information about the ecology of the unit and are frequently named after a common plant species. A short text description of the site is given under the General Description (pg 21), this is followed by a picture or a cross section diagram of the ecological site (pg 21). The section on successional relationships gives a brief note about the temporal development of the ecological site (pg 21). It generally describes the successional relationships among the ecological site phases and plant community types. Plant species that are indicators of the ecological conditions on the site are listed (pg 21). Site index at 50 years of age at breast height (1.3 m) is presented next. The mean site index is presented in meters followed by the standard error and the number of trees used to calculate the mean (pg 21). Environment and soil variables are then listed and represent a roll-up from the plant community and ecological site phase descriptions (pg 21). Variables that represent environment and soils have a number (1) that indicates the number of the samples in which each variable class occurred. Data has been collected and analyzed from many sources over 40 years and data gaps may exist for many variables. The frequency of occurrence value indicates the number of sampled plots for which data was collected for that variable at the Ecological site, Ecological site phase and plant community fact sheets. Optional variables such as soil exposure, LFH thickness, forage production and stocking rate for livestock may also be listed and represent a roll-up for the plant community and ecological site phase.

Ecological site phase

There is an identification code at the top of the ecological site phase fact sheet and a name followed by the number of sample sites (pg 22). Each ecological site phase has been given a name that conveys information about the dominant tree species or lifeform (shrubland, grassland, tame/disturbance) of the unit and are frequently named after a common plant species. A short text description of the site and successional information maybe given under the General Description or Successional relationships (pg 22) if it provides more detail than is available on the ecological site fact sheet. Plant species that are indicators of the ecological conditions on the site are listed with the average cover summarized from the various plant communities (pg 22). Indicator species for the ecological site phase are identified with an asterix "*" and are rolled-up to develop the indicator species list for the ecological site fact sheet. Environment and soil variables are then listed and represent a roll-up from the plant community (pg 22). Optional variables such as soil exposure, LFH thickness, forage production and stocking rate for livestock may also be listed and represent a roll-up for the plant communities.

Plant community

There is an identification code at the top of the plant community fact sheet and a name followed by the number of sample sites (pg 23). The name of the plant community is generally the common name of the indicator plant species within the various lifeform layers (tree, shrub, forb, grass, lichen, moss). This is followed by the latin name of each indicator species and a general description of the community type describing its unique ecology. Plant species that are indicators of the ecological conditions on the site are listed with the mean cover summarized, range in cover and overall constancy (frequency of plots that the species was described (pg 23)). Environment and soil variables are then listed and represent a roll-up from the various plots and assessments (pg 23). Optional variables such as soil exposure, LFH thickness, forage production and stocking rate for livestock may also be listed and represent a roll-up for various plots.

Results

This guide represents the analysis of 73 grassland/lichen, shrubland, conifer and deciduous field plots described from the Kazan Upland subregion. These plots are distributed between 34 community types:

- A. Native grasslands 8 types
- B. Shrubland 4 types
- C. Deciduous 8 types
- D. Mixedwood 4 types
- E. Conifer 10 types

The dominant plant species, canopy cover and environmental conditions are outlined for each community type.

General Ecological Descriptions

Please note this summary of Natural Subregion characteristics is largely extracted directly from the Natural Subregions guide (Natural Regions Committee 2006) and is presented here for the reader's convenience.

Forest Communities

Vegetation in the Kazan Upland Natural Subregion is strongly influenced by the distribution and acidic characteristics of granitic bedrock exposures and well to rapidly drained glacial drift, and by frequent fires. Beckingham and Archibald (1996) produced a general description of vegetation communities based on limited plot data. Coarse textured, rapidly drained and dry sandy or gravelly soils support vigorous jack pine growth. The driest sites are vegetated by open jack pine stands with a patchy carpet of lichens on the forest floor. Moister sites with submesic-mesic moisture regimes support more diverse understories of green alder, common blueberry, bearberry, common Labrador tea, Canada buffaloberry, bunchberry, other herbs and feathermosses with an overstory of jack pine, white spruce or aspen. Aspen, Alaska birch and black spruce are locally common in places. Brunisols are common soils on these upland forested community types. On moister sites with subhygric to hygric moisture regimes communities of aspen, balsam poplar, Alaska birch, white spruce and a diverse and lush shrub (red osier dogwood and willow on rich sites and Labrador tea on poorer sites) and forb (horsetail) understories develop in bands adjacent to wetlands and along lakes.

Wetlands

The Alberta Wetland Classification System (2015) recognizes the hydrological, biogeochemical and biotic processes that affect differing characteristics that can be used to define a wetland. The AWCS recognizes five classes of wetlands in Alberta: bogs, fens, marshes, shallow open water and swamps. Wetlands can be divided into two broad groups: **peatlands** and **mineral wetlands**. In general the AWCS considers bogs and fens to be peatlands and all other wetland classes (i.e. swamps, marshes and shallow open waters) are considered to be mineral wetlands. For the most part the ecological sites align with AWCS five classes of wetlands (Table A), however some willow, bog birch, sedge, marsh reedgrass and tufted hairgrass dominated ecological sites because of their moisture regime and species composition are classified as meadows and fens and have mineral soils but in the AWCS classification these sites are mineral wetlands which are considered marshes or swamps.

Table A. Cross walk of broad AWCS classes to general Ecological site

AWCS	Ecological Sites
Bog	Bog
Fen	Poor fen (Organic soils)
Fen	Rich fen (Organic soils)
Marsh	Poor fen, Rich fen (Mineral soils)
Marsh	Meadows
Marsh	Marsh
Swamp	Red osier dogwood, Honeysuckle, Horsetail and wetter Labrador tea, black spruce dominated ecological sites.

Dozens of small lakes occupy about 10 percent of the Kazan Upland Natural Subregion, the largest being Cornwall, Colin, Charles, Wylie and Andrew Lakes. A few small streams drain into the Slave River. Wetlands are most common in the lower relief western half of the Natural Subregion, and cover about 20 percent of the total area. Treed poor fens and bogs on Organic soils are the most common wetland type. Shrubby fens and marshes on Organic and wet mineral soils also occur.

Bog communities are the dominant wetland type. Black spruce forms open-to-dense stands with an understory of common Labrador tea, leatherleaf, bog cranberry, cloudberry and peat moss on Organic soils. Permafrost is discontinuous but widespread. Nutrient-rich wetlands typically have open forests of tamarack, willow, dwarf

birch, sedges and rich-site mosses. Marshes can be locally extensive in sheltered lake bays or along creek channels, and are dominated by water and small bottle sedge, bulrushes, and in deeper water, pondweeds.


Grasslands

Upland dry grasslands are not common in the Kazan Upland subregion. Allen et al. 2004 did observe a hay sedge dominated community on upper to mid slope positions on steep south and west facing slopes with sandy or gravelly glacial deposits in the Colin-Cornwall Lakes Wildland Provincial Park. Saline meadow complexes were described within the La Butte Creek Wildland Provincial Park (Allen et al. 2002) at two sites. Site 1 which is described was slightly drier than site 2 and was dominated by mat muhly, shooting star and seaside arrow grass. The other site which was slightly wetter was dominated by brown moss, awned sedge and seaside arrowgrass on the drier margins. It was felt that the mat muhly dominated community was subjected to higher levels of salt through evaporation. Raup (1935) described similar saline complexes in Wood Buffalo National Park and Bailey et al. (1992) described saline complexes in the Yukon Territory. The wettest edges were dominated by samphire sea blight, the Nuttall's saltgrass, foxtail barley and reedgrass and willow communities on the drier edges.

Lowland sites dominated by water sedge and marsh reedgrass were also observed in the Colin-Cornwall Lakes Wildland Park (Allen et al. 2004). The marsh reedgrass community was observed in patches or in narrow bands besides lakes, rivers and streams. They also observed fairly extensive meadows of marsh reedgrass near ponds and lakes. Water sedge and beaked sedge communities were observed along the edges of small ponds or a linear community along drainages where moisture was sufficient. Marshes dominated by cattails tend to occur in small patches along creeks (Allen et al. 2004).

Shrublands

Upland shrub types develop when soil conditions are too poor for tree growth or tree canopy has been removed mechanically or by fire. On eroded glaciofluvial kames a Bearberry/Indian ricegrass community was described (Allen et al. 2002). Allen et al. (2004) found this community type graded into open jack pine stands in La Butte Creek Wildland Provincial Park. Upland forest sites disturbed by fire will often have a early successional shrub phase. Labrador tea, lichen and feather moss tend to dominate the poorer sites and rose, Saskatoon, green alder and Bebb's willow the more mesic sites. Shrub communities dominated by willow, water birch and bog birch dominate the moister and richer sites. Sedges and reed grasses (*Calamagrostis* spp) dominate the drier edges of marshes or areas that have standing water only during spring runoff (i.e. fens). Willow will invade into these fens to form the Willow/Sedge and Willow/Reed grass plant communities and bog birch and Labrador tea tend to dominate the poorer nutrient sites.



Ecological sites

a=lichen stone fields
xeric/very poor
aa=Sand heather/moss
xeric/poor
b=bearberry/lichen
subxeric/poor
c=blueberry
submesic/medium
d=Labrador tea-mesic
mesic/poor
e= buffaloberry/alder
mesic/medium

g=red osier dogwood/horsetail
subhygric/rich
g=Labrador tea - hygric
hygric/medium
h=bog
subhydric/very poor
j=poor fen
subhydric/medium
j=rich fen
subhydric/rich
k=marsh
hydric/rich

Figure 1. Edatope grid and ecological sites for the Kazan Upland subregion.

Plant Community Keys

1. Kazan Upland.....	2
2. a. (xeric/very poor) exposed bedrock with crustose lichens (ecosite a lichen stonefield).....	3
aa. (xeric/poor) open sand dunes with sand heather, Plains wormwood or bearberry (ecosite aa Sand heather/moss)	5
b. (subxeric/poor) coarse textured soils dominated by jack pine, bearberry, lichen or hair cap moss (ecosite b bearberry/lichen)	6
c. (submesic/medium) coarse textured soils with aspen, jackpine and blueberry understory (ecosite c blueberry)	9
d. (mesic/poor) mesic sites with poor nutrient regimes dominated by black spruce, jack pine and labrador tea and aspen mixedwoods (ecosite d Labrador tea - mesic)	13
e. (mesic/medium) mesic site with medium nutrient regimes dominated by aspen, jackpine, white spruce and understory of buffaloberry and alder (ecosite e buffaloberry/alder)	15
f. (subhygric/rich) moist sites with aspen, balsam poplar, white spruce or balsam fir with an understory of red osier dogwood or horsetail (ecosite f red osier dogwood/horsetail)	19
g. (hygric/medium) very moist sites with poor nutrient regime dominated by black spruce and jack pine with labrador tea in the understory (ecosite g Labrador tea - hygric)	21
h. (subhydric/very poor) bog dominated by black spruce, sphagnum, and labrador tea (ecosite h bog)	23
i. (subhydric/medium) poor fen site codominated by black spruce and larch, understory has bog birch with some sphagnum and some golden moss (ecosite i poor fen).....	26
j. (subhydric/rich) rich fen dominated by larch, with willows and bog birch, little sphagnum (ecosite j fen)	29
k. (hydric/rich) aquatic sites with open standing water, dominated by cattails, northern quillwort and rush species (ecosite k marsh)	32
3. exposed bedrock with crustose lichens (ecosite phase a1 lichen stonefields).....	4
4. Lichen stonefield (KUA1)	p 23
5. low shrub and grass dominated community types (ecosite phase aa1 grass/sand heather)	
6. jack pine dominated phases (ecosite phase b1 bearberry/lichen - Pj)	7
low shrub dominated phase (ecosite phase b2 bearberry/lichen - shrub)	8
7. Pj/Lichen (KUE1)	p 28
Pj/Bearberry (KUE2).....	p 29
8. Bearberry/Indian ricegrass (KUA2)	p 31
Juniper/Lichen(Pj) (KUB1)	p 32
9. jack pine and aspen dominated phase (ecosite phase c1 blueberry - Pj-Aw(Bw))	10
deciduous dominated phase little jack pine (ecosite phase c2 blueberry - Aw(Bw)).....	11
site dominated by grasses (hay sedge, northern ricegrass) (ecosite phase c6 - grassland)	12
site dominated by tame forage species (brome, creeping red fescue, timothy) (ecosite phase c7 tame/disturbed)	
10. Pj-Aw(Bw)/Bearberry-Bog cranberry (KUD3)	p 35
Pj/Blueberry (KUE3)	p 36
11. Aw/Rose/Spreading dogbane (KUC1).....	p 38
12. Hay sedge-Slender wheatgrass (KUA3)	p 40
13. predominantly fine textured soils with forests dominated by black spruce, aspen and jack pine (ecosite phase d1 Labrador tea - mesic Pj-Sb)	14
14. Aw-Pj-Sb/Labrador tea (KUC4).....	p 44
Sb-Aw/Moss (KUD1)	p 45
Pj/Bog cranberry/lichen (KUE11)	p 46
Pj- Sb/Bog cranberry/lichen (KUE4).....	p 47
15. mesic site dominated by aspen (ecosite phase e1 buffaloberry/alder - Aw).....	16
site dominated by a mixture of deciduous and conifer species (ecosite phase e2 buffaloberry/alder - Aw-Sw-Pj)	17
site dominated by conifer trees (jack pine, white spruce) (ecosite phase e4 buffaloberry/alder - Pj-Sw).....	18
mesic site dominated by shrubs (ecosite phase e3 buffaloberry/alder - shrubland)	
16. Aw(Bw)/Alder (KUC2).....	p 50
Aw/Buffaloberry (KUC3)	p 51
Aw-Pb/Saskatoon-Red Osier Dogwood/Spreading dogbane (KUC7)	p 52
17. Aw-Sw/Buffaloberry (KUD2)	p 54
Aw- Pj/Alder (KUD4).....	p 55
18. Pj/Alder (KUE5)	p 58
Sb-Sw/Moss (KUE6).....	p 59
19. moist site dominated by deciduous species (aspen, balsam poplar) (ecosite phase f3 fern/horsetail - Pb-Aw).....	20
moist sites dominated by conifer trees (white spruce, balsam fir) (ecosite phase f1 fern/horsetail - Sw(Fb))	
moist site dominated by mixture of conifer and deciduous species (ecosite phase f2 fern/horsetail - Pb-Sw)	

moist site dominated by shrub species (red osier dogwood) (ecosite phase f4 shrub)
 site dominated by tame species (brome, creeping red fescue,timothy) (ecosite phase f5 tame/disturbed)

20. Pb/Alder-Red osier dogwood/Horsetail (KUC5)	p 64
Bw-Aw-Pb/Willow (KUC6).....	p 65
21. conifer dominated site (black spruce and jack pine)(ecosite phase g1 Labrador tea - hygric Sb-Pj)	22
22. Sb/Alder/Tufted moss (KUE7).....	p 71
23. site dominated by trees (black spruce) (ecosite phase h1 bog - treed).....	24
site dominated by shrub species (labrador tea) (ecosite phase h2 - bog shrubby)	25
site dominated by graminoid and sphagnum species (mud sedge) (ecosite phase h3 - graminoid bog)	
24. Sb/Labrador tea/peat moss (KUE8)	p 74
25. Labrador tea/Peat moss (Sb) (KUB2)	p 76
26. site dominated by tree species (larch, black spruce) (ecosite phase i1 poor fen - treed)	27
site dominated by graminoid species (sedges) (ecosite phase i3 poor fen - graminoid)	28
site dominated by shrub species (bog birch) (ecosite phase i2 poor fen - shrubby)	
27. Sb-Lt/Labrador tea/Golden moss (KUE9)	p 80
28. Shooting star/Mat muhly (saline) (KUA4)	p 83
Sedge/Brown moss (saline) (KUA5)	p 84
29. site dominated by tree species (larch) (ecosite phase j1 rich fen - treed)	30
site dominated by shrub species (willow) (ecosite phase j2 rich fen - shrubby)	31
site dominated by graminoid species (sedge) (ecosite phase j3 rich fen - graminoid)	
30. Lt-Sb/River alder/Sedge (KUE10)	p 70
31. Willow/Marsh reed grass (KUB3)	p 88
Willow/Sedge (KUB4)	p 89
Willow/Brown moss/ Aw(Bw) (KUC8).....	p 90
32. marsh sites with standing water (ecosite phase k1 marsh).....	33
33. Water parsnip/Northern manna grass (KUA6).....	p 94
Northern quillwort (KUA7).....	p 95
Water arum/Sedge (KUA8).....	p 96

Plant Community Tables

Table 1. Kazan Upland Communities

Ecological Site / Range Site	Ecosite Phase / Ecological Range Site	Reference Plant Community	Grazing Succession	Modified Plant Community	Harvesting Succession
a lichen stone fields	a1 lichen stonefields	KUA1 Lichen stonefield			
aa Sand heather/moss (xeric/poor)	aa1 grass/Sand heather				
b bearberry/lichen(subxeric/poor)	b1 bearberry/lichen - Pj	KUE1 Pj/Lichen			
		KUE2 Pj/Bearberry			
	b2 bearberry/lichen - shrub	KUA2 Bearberry/Indian ricegrass			
		KUB1 Juniper/Lichen(Pj)			
c blueberry(submesic/medium)	c1 blueberry - Pj-Aw(Bw)	KUD3 Pj-Aw(Bw)/Bearberry-Bog cranberry			
		KUE3 Pj/Blueberry			
	c2 blueberry - Aw(Bw)	KUC1 Aw/Rose/Spreading dogbane			
	c6 blueberry grassland	KUA3 Hay sedge-Slender wheatgrass			
	c7 tame/disturbed				
d Labrador tea-mesic(mesic/poor)	d1 Labrador tea-mesic Pj-Sb	KUC4 Aw-Pj-Sb/Labrador tea			
		KUD1 Sb-Aw/Moss			
		KUE11 Pj/Bog cranberry/lichen			
		KUE4 Pj- Sb/Bog cranberry/lichen			
e buffaloberry/alder(mesic/medium)	e1 buffaloberry-alder/ Aw	KUC2 Aw(Bw)/Alder			
		KUC3 Aw/Buffaloberry			
		KUC7 Aw-Pb/Saskatoon-Red Osier Dogwood/Spreading dogbane			
	e2 buffaloberry-alder/ Pj-Sw-Sb-Aw-	KUD2 Aw-Sw/Buffaloberry			

Ecological Site / Range Site	Ecosite Phase / Ecological Range Site	Reference Plant Community	Grazing Succession	Modified Plant Community	Harvesting Succession
	Bw	KUD4 Aw- Pj/Alder			
	e3 buffaloberry-alder shrubland				
	e4 buffaloberry-alder/ Pj-Sw-Sb	KUE5 Pj/Alder			
		KUE6 Sb-Sw/Moss			
f red osier dogwood/horsetail (subhygric/rich)	f1 red osier dogwood/horsetail Sw				
	f2 red osier dogwood/horsetail Pb-Sw				
	f3 red osier dogwood/horsetail - Pb-Aw(Ba)	KUC5 Pb/Alder-Red osier dogwood/Horsetail			
		KUC6 Bw-Aw-Pb/Willow			
	f4 shrub				
	f5 tame/disturbed				
g Labrador tea-hygric(hygric/medium)	g1 Labrador tea-hygric Sb-Pj	KUE10 Lt-Sb/River alder/Sedge			
		KUE7 Sb/Alder/Tufted moss			
h bog (subhydric/very poor)	h1 bog - treed	KUE8 Sb/Labrador tea/peat moss			
	h2 bog - shrubby	KUB2 Labrador tea/Peat moss (Sb)			
	h3 bog -graminoid				
i poor fen (subhydric/medium)	i1 poor fen - treed	KUE9 Sb-Lt/Labrador tea/Golden moss			
	i2 poor fen - shrubby				
	i3 poor fen - graminoid	KUA4 Shooting star/Mat muhly (saline)			
		KUA5 Sedge/Brown moss (saline)			
j rich fen (subhydric/rich)	j1 rich fen - treed				
	j2 rich fen - shrubby	KUB3 Willow/Marsh reed grass			


Ecological Site / Range Site	Ecosite Phase / Ecological Range Site	Reference Plant Community	Grazing Succession	Modified Plant Community	Harvesting Succession
		KUB4 Willow/Sedge			
		KUC8 Willow/Brown moss/ Aw(Bw)			
	j3 rich fen - graminoid				
k marsh (hydric/rich)	k1 marsh	KUA6 Water parsnip/Northern manna grass			
		KUA7 Northern quillwort			
		KUA8 Water arum/Sedge			

KU Kazan Upland (n=73)

Natural Subregion: Kazan Upland

General Description

This subregion has not been split into ecosections. It includes the Uranium City and Hart Lake Plain ecodistricts. The subregion is located north of Lake Athabasca in the Northeastern part of the province with a small outlier at the headwaters of the Marguerite River. It is bordered by Athabasca Plain in the south and Northern Mixedwood and Peace Athabasca Delta subregions on its eastern boundary.



Environmental Variables

Elevation (range): 248.97 (180-330) M

Ecological Sites

Site Count

a	lichen stone fields	1
b	bearberry/lichen(subxeric/poor)	18
c	blueberry(submesic/medium)	12
d	Labrador tea-mesic(mesic/poor)	10
e	buffaloberry/alder(mesic/medium)	16
f	red osier dogwood/horsetail (subhygric/rich)	2
g	Labrador tea-hygric(hygric/medium)	2
h	bog (subhydric/very poor)	3
i	poor fen (subhydric/medium)	3
j	rich fen (subhydric/rich)	3
k	marsh (hydric/rich)	3

a lichen stone fields (n=1)

Natural Subregion: Kazan Upland

General Description

This ecosite occurs on xeric rocky outcrops throughout the Kazan Upland subregion. Moisture and nutrients are virtually nonexistent on these rock outcrops, where nearly 60% of the landscape in the subregion is exposed bedrock. Where there is some moisture accumulation in the rock crevices *Cladina* species and some grasses, sedges and xeric forb species are found growing. These sites also have high potential for extreme exposure to wind and insolation. There is very little soil or organic material present; therefore, the lichens have to rely on the rock substrate and the air for their moisture and nutrient requirements. Only lichen appears to tolerate the existent conditions on these rocky outcrops.



Ecosection: KU Kazan Upland

Environmental Variables

Moisture Regime: Very Xeric (very dry) (0)
Nutrient Regime: Oligotrophic (very poor) (0)
Elevation (range): 295 (295-295) M
Slope (%): very gentle slope (1)
Aspect: Southerly (1)
Topographic Position: Crest (1)

Soil Variables

Soil Drainage: Rapidly drained (1)
Soil Subgroup:
Surface Texture:
Effective Texture:
Depth to Mottles/Gley: Not Applicable (0)
Organic Thickness: 0 - 5 cm (1)
Parent Material: Rock (1)
Humus Form

LFH Thickness	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

Successional Relationships

This ecosite is characteristic of weakly or noncalcareous sites and is successional mature.

Indicator Species

Lichen

N/A
Umbilicaria muehlenbergii
N/A
Arctoparmelia centrifuga
REINDEER LICHEN
Cladina mitis

a1 lichen stonefields (n=1)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: a lichen stone fields

Characteristic Species

Shrub

- [2.0] COMMON BEARBERRY
Arctostaphylos uva-ursi
- [1.0] GROUND JUNIPER
Juniperus communis

Lichen

- [15.0] REINDEER LICHEN*
Cladonia mitis
- [8.0] N/A*
Umbilicaria muehlenbergii
- [5.0] N/A*
Arctoparmelia centrifuga

Moss and Liverwort

- [1.0] AWNEED HAIR-CAP
Polytrichum piliferum

Environmental Variables

Moisture Regime: Very Xeric (very dry) (0)
 Nutrient Regime: Oligotrophic (very poor) (0)
 Elevation (range): 295 (295-295) M
 Slope (%): very gentle slope (1)
 Aspect: Southerly (1)
 Topographic Position: Crest (1)

Soil Variables

Soil Drainage: Rapidly drained (1)
 Soil Subgroup:
 Surface Texture:
 Effective Texture:
 Depth to Mottles/Gley: Not Applicable (0)
 Organic Thickness: 0 - 5 cm (1)
 Parent Material: Rock (1)
 Humus Form

LFH Thickness	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

KUA1 Lichen stonefield (n=1)

(*Umblicaria spp.*)

This community type represents the lichen cover on rock outcrops throughout the subregion. These Precambrian rock outcrops represent 60% of the landscape in the subregion (Natural Regions Committee 2006). There are no soil or organic material associated with the exposed bedrock which favours the growth of crustose lichens. In the rock crevices where some organic matter and moisture accumulates reindeer lichen is common.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: a lichen stone fields
Ecosite Phase: a1 lichen stonefields

Plant Composition	Canopy Cover (%)			Const.	Environmental Variables				
	Mean	Range							
Medium Shrub (0.5 to 2 m)					Ecological Status Score: 40-40				
GROUND JUNIPER (<i>Juniperus communis</i>)	1.0	1.0-1.0		100	Moisture Regime: Very Xeric (very dry) (0)				
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	2.0	2.0-2.0		100	Nutrient Regime: Oligotrophic (very poor) (0)				
Low Forb (< 30 cm)					Elevation (range): 295 (295-295) M				
HAREBELL (<i>Campanula rotundifolia</i>)	1.0	1.0-1.0		100	Slope (%): 2.5 - 5.99 (1)				
THREE-TOOTHED CINQUEFOIL (<i>Potentilla tridentata</i>)	1.0	1.0-1.0		100	Aspect: Southerly (1)				
THREE-TOOTHED SAXIFRAGE (<i>Saxifraga tricuspidata</i>)	1.0	1.0-1.0		100	Topographic Position: Crest (1)				
RUSTY WOODSIA (<i>Woodsia ilvensis</i>)	1.0	1.0-1.0		100	Soil Variables				
Graminoid					Soil Drainage: Rapidly drained (1)				
SILVERY-FLOWERED SEDGE (<i>Carex aenea</i>)	1.0	1.0-1.0		100	Soil Subgroup:				
UMBELLATE SEDGE (<i>Carex umbellata</i>)	1.0	1.0-1.0		100	Surface Texture:				
SMALL BOTTLE SEDGE (<i>Carex utriculata</i>)	1.0	1.0-1.0		100	Effective Texture:				
ROCKY MOUNTAIN FESCUE (<i>Festuca saximontana</i>)	1.0	1.0-1.0		100	Depth to Mottles/Gley: Not Applicable (0)				
NORTHERN RICE GRASS (<i>Oryzopsis pungens</i>)	1.0	1.0-1.0		100	Organic Thickness: 0 - 5 cm (1)				
TIMBERLINE BLUEGRASS (<i>Poa glauca</i>)	1.0	1.0-1.0		100	Parent Material: Rock (1)				
Moss					Humus Form				
AWNED HAIR-CAP (<i>Polytrichum piliferum</i>)	1.0	1.0-1.0		100	LFH Thickness				
Lichen					Mean	Min	Max	Count	
N/A (<i>Arctoparmelia centrifuga</i>)	5.0	5.0-5.0		100	0.00	0.00	0.00	0	
N/A (<i>Umblicaria muehlenbergii</i>)	8.0	8.0-8.0		100					
REINDEER LICHEN (<i>Cladina mitis</i>)	15.0	15.0-15.0		100					

aa Sand heather/moss (xeric/poor) (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

This ecological site represents the sand of paleodunes of various types, including parabolic and longitudinal dunes and is more common in the southern outlier of the Kazan Upland and Athabasca Plain subregions. This ecological site is primarily made up of large, distinctive dune ridges that rise well above the surrounding wetlands or sand plains. The dunes characteristically have steep slopes, dune crests and often include areas of active sand (blowouts). Tops of stabilized dune ridges have mature jack pine / lichen communities. Blowouts are generally unvegetated, with areas of sand heather / lichen crust beginning the stabilization process. Jack pine / green alder communities can be found in moister locations (e.g. bases of sand ridges). This ecological site is not common in the Kazan Upland subregion and currently no data has been collected.



Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

LFH Thickness

cm:

Mean

Min

Max

Count

0.00

0.00

0.00

0

Successional Relationships

Due to the dry nature of the site often only sand heather and lichen will dominate the site. Jack pine and alder will invade in the moister areas or on northerly aspects where the dunes have stabilized. See the Athabasca Plain subregion guide for a complete description of this ecological site.

Indicator Species

aa1 grass/Sand heather (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: aa Sand heather/moss (xeric/poor)

General Description

A number of ecological site phases currently have no data. These ecological site phases have been created as place holders because they were described in adjacent subregions (Athabasca Plain). This ecosite phase is more common in the southern outlier of the Kazan Upland subregion. See the Athabasca Plain subregion guide for a description of the plant communities.

Characteristic Species

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

b bearberry/lichen(subxeric/poor) (n=18)

Natural Subregion: Kazan Upland

General Description

This ecosite has dry conditions with rapidly drained acidic soils and poor nutrient status due to the coarse-textured glacial parent materials. These sites commonly have bedrock within 1m of the surface. Plants that are indicative of the nutrient-poor substrate include bearberry, lichen, bog cranberry, and blueberry. Open-canopied jack pine stands dominate this ecosite that commonly has a carpet of lichens covering the forest floor. The picture of this ecosite is for the bearberry/lichen shrub phase (b2) of this ecological site.



Successional Relationships

Due to the dry nature of this ecosite, succession to a black spruce canopy is slower than the fire return interval. Pine is maintained for relatively long periods from colonization of the site after fire to the climax stages. The pine phase of this ecosite can be considered a fire edaphic climax.

Indicator Species

Tree

JACK PINE
Pinus banksiana

Shrub

COMMON BEARBERRY
Arctostaphylos uva-ursi
GROUND JUNIPER
Juniperus communis

Lichen

REINDEER LICHEN
Cladonia mitis

Moss and Liverwort

AWNED HAIR-CAP
Polytrichum piliferum

Graminoid

NORTHERN RICE GRASS
Oryzopsis pungens

Ecosection: KU Kazan Upland

Environmental Variables

Moisture Regime: Xeric (dry) (1), Submesic (moderately fresh) (7), Subxeric (moderately dry) (9)

Nutrient Regime: Mesotrophic (medium) (1), Submesotrophic (poor) (16)

Elevation (range): 248.25 (200-330) M

Slope (%): strong slope (1), nearly level (2), moderate slope (3), very gentle slope (4), gentle slope (6)

Aspect: Westerly (3), Easterly (4), Level (4), Southerly (5)

Topographic Position: Level (2), Lower Slope (3), Midslope (3), Crest (4), Upper Slope (5)

Soil Variables

Soil Drainage: Very poorly drained (1), Well drained (7), Rapidly drained (10)

Soil Subgroup: REGOSOL (1), DYSTRIC BRUNISOL (2), EUTRIC BRUNISOL (12)

Surface Texture: Sandy loam (0), Silt (1), Loamy fine sand (1), Sandy clay (1), Loamy sand (3), Sand (9)

Effective Texture: Sandy loam (0), Loamy fine sand (1), Sand (14)

Depth to Mottles/Gley: Not Applicable (0), 51 - 100 (1)

Organic Thickness: 0 - 5 cm (16)

Parent Material: Glaciolacustrine (2), Morainal (2), Residual (2), Rock (7), Glaciofluvial (8)

Humus Form FIBRIMOR (1)

LFH Thickness	Mean	Min	Max	Count
cm:	3.00	2.00	5.00	15

b1 bearberry/lichen - Pj (n=15)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: b bearberry/lichen(subxeric/poor)

Characteristic Species

Tree

- [30.0] JACK PINE*
Pinus banksiana

Shrub

- [2.0] COMMON BEARBERRY
Arctostaphylos uva-ursi
- [2.0] SAND HEATHER
Hudsonia tomentosa
- [1.0] COMMON BLUEBERRY
Vaccinium myrtilloides
- [1.0] BOG CRANBERRY
Vaccinium vitis-idaea

Forb

- [1.0] WILD LILY-OF-THE-VALLEY
Maianthemum canadense

Lichen

- [20.0] REINDEER LICHEN*
Cladina mitis
- [1.0] N/A
Peltigera malacea

Moss and Liverwort

- [3.0] AWNED HAIR-CAP*
Polytrichum piliferum

Environmental Variables

Moisture Regime: Submesic (moderately fresh) (7), Subxeric (moderately dry) (8)

Nutrient Regime: Mesotrophic (medium) (1), Submesotrophic (poor) (14)

Elevation (range): 254 (200-330) M

Slope (%): nearly level (1), strong slope (1), moderate slope (2), very gentle slope (3), gentle slope (6)

Aspect: Westerly (2), Southerly (3), Easterly (4), Level (4)

Topographic Position: Level (2), Crest (2), Lower Slope (3), Midslope (3), Upper Slope (4)

Soil Variables

Soil Drainage: Very poorly drained (1), Well drained (7), Rapidly drained (7)

Soil Subgroup: REGOSOL (1), DYSTRIC BRUNISOL (2), EUTRIC BRUNISOL (12)

Surface Texture: Silt (1), Sandy clay (1), Loamy fine sand (1), Loamy sand (3), Sand (9)

Effective Texture: Loamy fine sand (1), Sand (14)

Depth to Mottles/Gley: 51 - 100 (1)

Organic Thickness: 0 - 5 cm (15)

Parent Material: Glaciolacustrine (2), Residual (2), Morainal (2), Rock (6), Glaciofluvial (8)

Humus Form FIBRIMOR (1)

LFH Thickness

	Mean	Min	Max	Count
cm:	3.00	2.00	5.00	15

KUE1 Pj/Lichen (n=9)

(*Pinus banksiana*/*Cladonia spp.*)

This community type is found on very dry sandy or gravelly sites that are rapidly drained. Stunted jack pine and Alaska birch form open pocket communities with a sparse understory of bearberry, juniper and lichen in rock crevices or sheltered locations where mineral soil has accumulated (Natural Regions Committee 2006). Where the soil is better developed vigorous jack pine growth is evident. As the moisture increases the understory often becomes dominated by blueberry.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: b bearberry/lichen(subxeric/poor)
Ecosite Phase: b1 bearberry/lichen - Pj

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
JACK PINE (<i>Pinus banksiana</i>)	12.8	0.0-42.0	78	Moisture Regime: Submesic (moderately fresh) (3), Subxeric (moderately dry) (6)
Understory Tree				Nutrient Regime: Mesotrophic (medium) (1), Submesotrophic (poor) (8)
JACK PINE (<i>Pinus banksiana</i>)	8.8	0.0-30.0	67	Elevation (range): 268 (235-330) M
Tall Shrub (2 to 5m)				Slope (%): 0.5 - 2.49 (1), 2.5 - 5.99 (1), 10 - 15.99 (1), 16 - 30.99 (1), 6 - 9.99 (4)
JACK PINE (<i>Pinus banksiana</i>)	8.8	0.0-30.0	78	Aspect: Southerly (1), Westerly (2), Level (2), Easterly (3)
Medium Shrub (0.5 to 2 m)				Topographic Position: Crest (1), Level (2), Midslope (2), Lower Slope (3)
GREEN ALDER (<i>Alnus crispa</i>)	2.7	0.0-5.0	67	Soil Variables
JACK PINE (<i>Pinus banksiana</i>)	3.3	0.0-20.0	56	Soil Drainage: Very poorly drained (1), Rapidly drained (4), Well drained (4)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	4.0	0.0-18.0	56	Soil Subgroup: DYSTRIC BRUNISOL ELUVIATED (1), REGOSOL ORTHIC (1), EUTRIC BRUNISOL ELUVIATED (7)
COMMON BLUEBERRY (<i>Vaccinium myrtilloides</i>)	5.5	0.0-12.0	89	Surface Texture: Loamy fine sand (1), Sandy clay (1), Loamy sand (3), Sand (4)
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	9.5	0.0-30.0	89	Effective Texture: Loamy fine sand (1), Sand (8)
Moss				Depth to Mottles/Gley:
AWNED HAIR-CAP (<i>Polytrichum piliferum</i>)	2.2	0.0-10.0	22	Organic Thickness: 0 - 5 cm (9)
Lichen				Parent Material: Glaciolacustrine (1), Morainal (1), Residual (1), Rock (5), Glaciofluvial (5)
N/A (<i>Stereocaulon tomentosum</i>)	1.6	0.0-10.0	22	Humus Form FIBRIMOR (1)
REINDEER LICHEN (<i>Cladina rangiferina</i>)	2.0	0.0-18.0	11	
REINDEER LICHEN (<i>Cladina mitis</i>)	51.8	30.0-90.0	100	
				LFH Thickness
				Mean
				Min
				Max
				Count
				cm: 3.00 2.00 4.00 9

KUE2 Pj/Bearberry (n=6)

(*Pinus banksiana*/*Arctostaphylos uva-ursi*)

This community type is very similar to the jack pine/lichen community type that was previously described but appears to be slightly moister and more diverse. The understory of this community type is dominated by bearberry and there is an increase in cover of more mesic plant species like alder, rose, bedstraw and wild lily of the valley. However despite the increase in moisture this community type still occupies very coarse textured sites that are well to rapidly drained.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: b bearberry/lichen(subxeric/poor)

Ecosite Phase: b1 bearberry/lichen - Pj

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
JACK PINE (<i>Pinus banksiana</i>)	29.8	15.0-50.0	100	Moisture Regime: Subxeric (moderately dry) (2), Submesic (moderately fresh) (4)
Understory Tree				Nutrient Regime: Submesotrophic (poor) (6)
JACK PINE (<i>Pinus banksiana</i>)	5.8	0.0-20.0	50	Elevation (range): 240 (200-300) M
Tall Shrub (2 to 5m)				Slope (%): 10 - 15.99 (1), 6 - 9.99 (2), 2.5 - 5.99 (2)
JACK PINE (<i>Pinus banksiana</i>)	1.3	0.0-4.0	33	Aspect: Easterly (1), Level (2), Southerly (2)
GREEN ALDER (<i>Alnus crispa</i>)	3.3	0.0-8.0	50	Topographic Position: Crest (1), Midslope (1), Upper Slope (4)
Medium Shrub (0.5 to 2 m)				Soil Variables
PRICKLY ROSE (<i>Rosa acicularis</i>)	1.0	0.0-3.0	50	Soil Drainage: Rapidly drained (3), Well drained (3)
COMMON BLUEBERRY (<i>Vaccinium myrtilloides</i>)	3.6	2.0-8.0	100	Soil Subgroup: DYSTRIC BRUNISOL ELUVIATED (1), EUTRIC BRUNISOL GLEYED ELUVIATED (1), EUTRIC BRUNISOL ELUVIATED (4)
TWINFLOWER (<i>Linnaea borealis</i>)	6.0	0.0-20.0	83	Surface Texture: Silt (1), Sand (5)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	10.6	0.0-20.0	83	Effective Texture: Sand (6)
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	31.5	15.0-60.0	100	Depth to Mottles/Gley: 51 - 100 (1)
Graminoid				Organic Thickness: 0 - 5 cm (6)
NORTHERN RICE GRASS (<i>Oryzopsis pungens</i>)	1.3	0.0-2.0	83	Parent Material: Glaciolacustrine (1), Morainal (1), Rock (1), Residual (1), Glaciofluvial (3)
Lichen				Humus Form
REINDEER LICHEN (<i>Cladina mitis</i>)	22.5	3.0-42.0	100	LFH Thickness
				Mean
				Min
				Max
				Count
				cm:
				3.00
				2.00
				5.00
				6

b2 bearberry/lichen - shrub (n=3)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: b bearberry/lichen(subxeric/poor)

Characteristic Species

Tree

- [6.0] JACK PINE
Pinus banksiana

Shrub

- [32.5] COMMON BEARBERRY*
Arctostaphylos uva-ursi
- [4.0] GROUND JUNIPER*
Juniperus communis
- [1.0] COMMON BLUEBERRY
Vaccinium myrtilloides

Forb

- [1.5] HAREBELL
Campanula rotundifolia
- [1.2] WILD STRAWBERRY
Fragaria virginiana
- [1.0] THREE-TOOTHED SAXIFRAGE
Saxifraga tricuspidata
- [1.0] THREE-TOOTHED CINQUEFOIL
Potentilla tridentata

Lichen

- [20.0] REINDEER LICHEN*
Cladina mitis
- [2.5] N/A
Stereocaulon tomentosum

Graminoid

- [5.2] NORTHERN RICE GRASS*
Oryzopsis pungens
- [1.7] SLENDER WHEAT GRASS
Agropyron trachycaulum
- [1.5] ROUGH HAIR GRASS
Agrostis scabra

Environmental Variables

Moisture Regime: Xeric (dry) (1), Subxeric (moderately dry) (1)
 Nutrient Regime: Submesotrophic (poor) (2)
 Elevation (range): 242.5 (240-250) M
 Slope (%): moderate slope (1), nearly level (1), very gentle slope (1)
 Aspect: Westerly (1), Southerly (2)
 Topographic Position: Upper Slope (1), Crest (2)

Soil Variables

Soil Drainage: Rapidly drained (3)
 Soil Subgroup:
 Surface Texture: Sandy loam (0)
 Effective Texture: Sand (0), Sandy loam (0)
 Depth to Mottles/Gley: Not Applicable (0)
 Organic Thickness: 0 - 5 cm (1)
 Parent Material: Glaciofluvial (0), Rock (1)
 Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

KUA2 Bearberry/Indian ricegrass (n=2)

(*Arctostaphylos uva-ursi*/*Oryzopsis pungens*)

This community type represents open areas on steep south and west facing slopes with coarse textured soils. One site was described on the top of an eroded glaciofluvial kame and graded into an open jack pine/buffaloberry community type. The other site was described in an opening amongst jack pine and bog cranberry and was relatively flat with bedrock at or near the surface. The first site had better developed soils and a higher cover of slender wheatgrass. In contrast the second site had a predominance of lichen and Indian ricegrass (Allen et al. 2002).

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: b bearberry/lichen(subxeric/poor)

Ecosite Phase: b2 bearberry/lichen - shrub

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Medium Shrub (0.5 to 2 m)				Ecological Status Score: 40-40
GROUND JUNIPER (<i>Juniperus communis</i>)	1.0	0.0-2.0	50	Moisture Regime: Xeric (dry) (1)
Low Shrub (< 0.5m)				Nutrient Regime: Submesotrophic (poor) (1)
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	65.0	50.0-80.0	100	Elevation (range): 245 (240-250) M
Tall Forb (>= 30 cm)				Slope (%): 0.5 - 2.49 (1), 2.5 - 5.99 (1)
LONG-LEAVED ARNICA (<i>Arnica longifolia</i>)	1.0	1.0-1.0	100	Aspect: Southerly (1), Westerly (1)
Low Forb (< 30 cm)				Topographic Position: Crest (1), Upper Slope (1)
PRAIRIE CROCUS (<i>Anemone patens</i>)	1.0	1.0-1.0	100	Soil Variables
HAREBELL (<i>Campanula rotundifolia</i>)	1.0	1.0-1.0	100	Soil Drainage: Rapidly drained (2)
WILD LILY-OF-THE-VALLEY (<i>Maianthemum canadense</i>)	1.0	0.0-2.0	50	Soil Subgroup:
MOUNTAIN GOLDENROD (<i>Solidago spathulata</i>)	1.0	1.0-1.0	100	Surface Texture: Sandy loam (0)
NORTHERN BEDSTRAW (<i>Galium boreale</i>)	2.0	1.0-3.0	100	Effective Texture: Sandy loam (0)
WILD STRAWBERRY (<i>Fragaria virginiana</i>)	2.5	2.0-3.0	100	Depth to Mottles/Gley: Not Applicable (0)
Graminoid				Organic Thickness:
PURPLE REED GRASS (<i>Calamagrostis purpurascens</i>)	1.0	1.0-1.0	100	Parent Material: Glaciofluvial (0)
SLENDER WHEAT GRASS (<i>Agropyron trachycaulum</i>)	3.5	0.0-7.0	50	Humus Form
NORTHERN RICE GRASS (<i>Oryzopsis pungens</i>)	10.5	1.0-20.0	100	LFH Thickness
Lichen				Mean Min Max Count
REINDEER LICHEN (<i>Cladina mitis</i>)	10.0	0.0-20.0	50	cm: 0.00 0.00 0.00 0

KUB1 Juniper/Lichen(Pj) (n=1)

(*Juniperus communis*/*Cladonia spp.*(*Pinus banksiana*))

This community type is very similar to the jack pine/lichen community types that were previously described but appears to be early seral. Fire is extensive throughout the Kazan Upland and jack pine dominated stands will burn frequently. In the absence of disturbance these sites will slowly regrow jack pine.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: b bearberry/lichen(subxeric/poor)
Ecosite Phase: b2 bearberry/lichen - shrub

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
JACK PINE (<i>Pinus banksiana</i>)	2.0	2.0-2.0	100	Moisture Regime: Subxeric (moderately dry) (1)
Tall Shrub (2 to 5m)				Nutrient Regime: Submesotrophic (poor) (1)
JACK PINE (<i>Pinus banksiana</i>)	10.0	10.0-10.0	100	Elevation (range): 240 (240-240) M
Medium Shrub (0.5 to 2 m)				Slope (%): 10 - 15.99 (1)
PIN CHERRY (<i>Prunus pensylvanica</i>)	1.0	1.0-1.0	100	Aspect: Southerly (1)
WILD RED RASPBERRY (<i>Rubus idaeus</i>)	1.0	1.0-1.0	100	Topographic Position: Crest (1)
COMMON BLUEBERRY (<i>Vaccinium myrtilloides</i>)	2.0	2.0-2.0	100	
GROUND JUNIPER (<i>Juniperus communis</i>)	7.0	7.0-7.0	100	Soil Variables
Tall Forb (>= 30 cm)				Soil Drainage: Rapidly drained (1)
SHOWY GOLDENROD (<i>Solidago nemoralis</i>)	1.0	1.0-1.0	100	Soil Subgroup:
Low Forb (< 30 cm)				Surface Texture:
RUSTY WOODSIA (<i>Woodsia ilvensis</i>)	1.0	1.0-1.0	100	Effective Texture: Sand (0)
HAREBELL (<i>Campanula rotundifolia</i>)	2.0	2.0-2.0	100	Depth to Mottles/Gley: Not Applicable (0)
PARSLEY FERN (<i>Cryptogramma acrostichoides</i>)	2.0	2.0-2.0	100	Organic Thickness: 0 - 5 cm (1)
THREE-TOOTHED CINQUEFOIL (<i>Potentilla tridentata</i>)	2.0	2.0-2.0	100	Parent Material: Rock (1)
THREE-TOOTHED SAXIFRAGE (<i>Saxifraga tricuspidata</i>)	2.0	2.0-2.0	100	Humus Form
Graminoid				
ROCKY MOUNTAIN FESCUE (<i>Festuca saximontana</i>)	1.0	1.0-1.0	100	
ROUGH HAIR GRASS (<i>Agrostis scabra</i>)	3.0	3.0-3.0	100	
Lichen				
N/A (<i>Stereocaulon tomentosum</i>)	5.0	5.0-5.0	100	
REINDEER LICHEN (<i>Cladina mitis</i>)	30.0	30.0-30.0	100	

LFH Thickness	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

c blueberry(submesic/medium) (n=12)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

This ecosite tends to be subxeric to submesic partly due to coarse-textured parent materials. It is intermediate in both moisture and nutrient regime between the bearberry ecosite (b) and the buffaloberry/alder ecosite (e) as there is some moisture at depth. As such it has species characteristic of the bearberry ecosite such as jack pine, blueberry, bearberry, bog cranberry, Labrador tea, and reindeer lichen and also species characteristic of the buffaloberry ecosite such as aspen, buffaloberry, bunchberry, alder, and fireweed.



Successional Relationships

The pine, aspen, and white birch-dominated phases of this ecosite may, in some cases, succeed to white spruce, however the process is expected to be relatively slow due to the dry nature of these sites.

Indicator Species

Tree

ASPEN
Populus tremuloides
JACK PINE
Pinus banksiana

Shrub

PRICKLY ROSE
Rosa acicularis
GREEN ALDER
Alnus crispa
COMMON BEARBERRY
Arctostaphylos uva-ursi
COMMON BLUEBERRY
Vaccinium myrtilloides
BOG CRANBERRY
Vaccinium vitis-idaea

Forb

PLAINS WORMWOOD
Artemisia campestris
SPREADING DOGBANE
Apocynum androsaemifolium

Lichen

REINDEER LICHEN
Cladina mitis

Graminoid

HAY SEDGE
Carex siccata

Site Index at 50 Years	Height (m)	Variation (m)	Count
ASPEN (<i>Populus tremuloides</i>)	17.20	0.40	0

Environmental Variables

Moisture Regime: Subxeric (moderately dry) (1), Mesic (fresh) (3), Submesic (moderately fresh) (4), Xeric (dry) (4)

Nutrient Regime: Mesotrophic (medium) (3), Submesotrophic (poor) (9)

Elevation (range): 277.75 (240-330) M

Slope (%): moderate slope (1), nearly level (1), strong slope (2), gentle slope (2), very strong slope (2), very gentle slope (3)

Aspect: Northerly (1), Easterly (2), Level (2), Southerly (6)

Topographic Position: Level (1), Midslope (3), Upper Slope (8)

Soil Variables

Soil Drainage: Rapidly drained (6), Well drained (6)

Soil Subgroup: DYSTRIC BRUNISOL (2), EUTRIC BRUNISOL (5)

Surface Texture: Sandy loam (0), Sand (7)

Effective Texture: Sandy clay (1), Sand (6)

Depth to Mottles/Gley: Not Applicable (0)

Organic Thickness: 0 - 5 cm (7)

Parent Material: Fluvial (0), Glaciolacustrine (1), Lacustrine (1), Rock (1), Morainal (2), Glaciofluvial (4)

Humus Form

LFH Thickness	Mean	Min	Max	Count
cm:	3.50	2.00	6.00	7

c1 blueberry - Pj-Aw(Bw) (n=7)

Natural Subregion: Kazan Upland

Ecosite: c blueberry(submesic/medium)

Ecosection: KU Kazan Upland

Characteristic Species

Tree

- [32.5] JACK PINE*
Pinus banksiana
- [7.5] WHITE BIRCH
Betula papyrifera
- [3.1] ASPEN*
Populus tremuloides

Shrub

- [14.8] GREEN ALDER*
Alnus crispa
- [12.2] BOG CRANBERRY*
Vaccinium vitis-idaea
- [11.2] COMMON BEARBERRY*
Arctostaphylos uva-ursi
- [10.5] COMMON BLUEBERRY*
Vaccinium myrtilloides
- [4.0] COMMON BEARBERRY
Arctostaphylos uva-ursi
- [3.5] BOG CRANBERRY
Vaccinium vitis-idaea
- [3.0] TWINFLOWER
Linnaea borealis
- [2.0] PRICKLY ROSE
Rosa acicularis
- [1.2] LOW-BUSH CRANBERRY
Viburnum edule

Forb

- [1.6] WILD SARSAPARILLA
Aralia nudicaulis
- [1.1] NORTHERN BASTARD TOADFLAX
Geocaulon lividum
- [1.0] WILD LILY-OF-THE-VALLEY
Maianthemum canadense

Lichen

- [8.5] REINDEER LICHEN*
Cladina mitis

Moss and Liverwort

- [8.5] SCHREBER'S MOSS
Pleurozium schreberi
- [2.7] WAVY DICRANUM
Dicranum polysetum

Environmental Variables

Moisture Regime: Subxeric (moderately dry) (1), Mesic (fresh) (2), Submesic (moderately fresh) (4)

Nutrient Regime: Mesotrophic (medium) (2), Submesotrophic (poor) (5)

Elevation (range): 287.5 (240-330) M

Slope (%): moderate slope (1), nearly level (1), very gentle slope (2), gentle slope (2)

Aspect: Northerly (1), Southerly (1), Easterly (2), Level (2)

Topographic Position: Level (1), Midslope (2), Upper Slope (4)

Soil Variables

Soil Drainage: Rapidly drained (2), Well drained (5)

Soil Subgroup: DYSTRIC BRUNISOL (2), EUTRIC BRUNISOL (5)

Surface Texture: Sand (7)

Effective Texture: Sandy clay (1), Sand (6)

Depth to Mottles/Gley:

Organic Thickness: 0 - 5 cm (7)

Parent Material: Lacustrine (1), Rock (1), Glaciolacustrine (1), Morainal (2), Glaciofluvial (4)

Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	3.50	2.00	6.00	7

KUD3 Pj-Aw(Bw)/Bearberry-Bog cranberry (n=4)

(*Pinus banksiana*-*Populus tremuloides*(*Betula papyrifera*)/*Arctostaphylos uva-ursi*-*Vaccinium vitis-idaea*)

Community types KUE3 and KUD3 are variants occurring modal conditions for this ecological site differing mainly in their initial secondary successional pathways. This community is very similar to the jack pine/blueberry community type but appears to be slightly moister. The understory of this community type is dominated by blueberry and bearberry and there is an increase in cover of more mesic plant species like alder, rose, bedstraw and wild sarsaparilla. However despite the increase in moisture this community type still occupies very dry coarse textured sites that are rapidly drained at the soil surface.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: c blueberry(submesic/medium)
Ecosite Phase: c1 blueberry - Pj-Aw(Bw)

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
ASPEN (<i>Populus tremuloides</i>)	6.2	0.0-20.0	50	Moisture Regime: Submesic (moderately fresh) (2), Mesic (fresh) (2)
WHITE BIRCH (<i>Betula papyrifera</i>)	15.0	0.0-42.0	50	Nutrient Regime: Submesotrophic (poor) (2), Mesotrophic (medium) (2)
JACK PINE (<i>Pinus banksiana</i>)	32.0	20.0-50.0	100	Elevation (range): 290 (240-330) M
Tall Shrub (2 to 5m)				Slope (%): 10 - 15.99 (1), 6 - 9.99 (2)
GREEN ALDER (<i>Alnus crispa</i>)	27.7	0.0-63.0	75	Aspect: Level (1), Easterly (2)
Medium Shrub (0.5 to 2 m)				Topographic Position: Level (1), Upper Slope (3)
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	2.5	0.0-8.0	75	Soil Variables
PRICKLY ROSE (<i>Rosa acicularis</i>)	4.0	0.0-15.0	50	Soil Drainage: Rapidly drained (2), Well drained (2)
TWINFLOWER (<i>Linnaea borealis</i>)	6.0	2.0-10.0	100	Soil Subgroup: DYSTRIC BRUNISOL ELUVIATED (2), EUTRIC BRUNISOL ELUVIATED (2)
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	22.5	0.0-42.0	75	Surface Texture: Sand (4)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	24.5	0.0-40.0	75	Effective Texture: Sand (4)
Tall Forb (>= 30 cm)				Depth to Mottles/Gley:
WILD SARSAPARILLA (<i>Aralia nudicaulis</i>)	3.2	1.0-10.0	100	Organic Thickness: 0 - 5 cm (4)
Low Forb (< 30 cm)				Parent Material: Glaciolacustrine (1), Lacustrine (1), Morainal (1), Rock (1), Glaciofluvial (2)
NORTHERN BASTARD TOADFLAX (<i>Geocaulon lividum</i>)	2.2	0.0-8.0	50	Humus Form
Graminoid				LFH Thickness
NORTHERN RICE GRASS (<i>Oryzopsis pungens</i>)	1.0	0.0-2.0	75	Mean
Moss				Min
WAVY DICRANUM (<i>Dicranum polysetum</i>)	5.5	0.0-18.0	50	Max
SCHREBER'S MOSS (<i>Pleurozium schreberi</i>)	10.0	0.0-18.0	75	Count
				cm:
				4.00
				3.00
				6.00
				4

KUE3 Pj/Blueberry (n=3)

(*Pinus banksiana/Vaccinium myrtilloides*)

Community types KUE3 and KUD3 are variants occurring modal conditions for this ecological site differing mainly in their initial secondary successional pathways. This community is very similar to the jack pine/bearberry community type but appears to be slightly moister. The understory of this community type is dominated by blueberry and there is an increase in cover of more mesic plant species like alder, rose, bedstraw and wild lily of the valley. However despite the increase in moisture this community type still occupies very dry coarse textured sites that are rapidly drained at the soil surface.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: c blueberry(submesic/medium)

Ecosite Phase: c1 blueberry - Pj-Aw(Bw)

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
JACK PINE (<i>Pinus banksiana</i>)	33.3	25.0-45.0	100	Moisture Regime: Subxeric (moderately dry) (1), Submesic (moderately fresh) (2)
Understory Tree				Nutrient Regime: Submesotrophic (poor) (3)
WHITE BIRCH (<i>Betula papyrifera</i>)	3.3	0.0-10.0	33	Elevation (range): 285 (260-330) M
JACK PINE (<i>Pinus banksiana</i>)	6.6	0.0-10.0	67	Slope (%): 0.5 - 2.49 (1), 2.5 - 5.99 (2)
Tall Shrub (2 to 5m)				Aspect: Level (1), Northerly (1), Southerly (1)
JACK PINE (<i>Pinus banksiana</i>)	3.3	0.0-6.0	67	Topographic Position: Upper Slope (1), Midslope (2)
GREEN ALDER (<i>Alnus crispa</i>)	4.3	3.0-7.0	100	
Medium Shrub (0.5 to 2 m)				Soil Variables
JACK PINE (<i>Pinus banksiana</i>)	1.0	0.0-2.0	67	Soil Drainage: Well drained (3)
COMMON LABRADOR TEA (<i>Ledum groenlandicum</i>)	1.3	0.0-4.0	33	Soil Subgroup: EUTRIC BRUNISOL ELUVIATED (3)
TWINFLOWER (<i>Linnaea borealis</i>)	2.0	0.0-5.0	67	Surface Texture: Sand (3)
GREEN ALDER (<i>Alnus crispa</i>)	2.3	0.0-4.0	67	Effective Texture: Sandy clay (1), Sand (2)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	9.6	7.0-12.0	100	Depth to Mottles/Gley:
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	11.6	5.0-15.0	100	Organic Thickness: 0 - 5 cm (3)
COMMON BLUEBERRY (<i>Vaccinium myrtilloides</i>)	15.6	10.0-25.0	100	Parent Material: Morainal (1), Glaciofluvial (2)
Low Forb (< 30 cm)				Humus Form
GREENISH-FLOWERED WINTERGREEN (<i>Pyrola chlorantha</i>)	1.0	1.0-1.0	100	
Lichen				
REINDEER LICHEN (<i>Cladina mitis</i>)	21.6	0.0-35.0	67	
				LFH Thickness
				Mean
				Min
				Max
				Count
				cm:
				3.00
				2.00
				5.00
				3

c2 blueberry - Aw(Bw) (n=1)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: c blueberry(submesic/medium)

Characteristic Species

Tree

- [70.0] ASPEN
Populus tremuloides

Shrub

- [15.0] PRICKLY ROSE*
Rosa acicularis
- [5.0] COMMON BEARBERRY
Arctostaphylos uva-ursi
- [5.0] BUNCHBERRY
Cornus canadensis
- [2.0] BOG CRANBERRY
Vaccinium vitis-idaea
- [1.0] TWINFLOWER
Linnaea borealis

Forb

- [20.0] SPREADING DOGBANE*
Apocynum androsaemifolium
- [1.0] WILD LILY-OF-THE-VALLEY
Maianthemum canadense

Graminoid

- [5.0] HAIRY WILD RYE
Elymus innovatus

Environmental Variables

Moisture Regime: Mesic (fresh) (1)
 Nutrient Regime: Mesotrophic (medium) (1)
 Elevation (range): 245 (245-245) M
 Slope (%): very gentle slope (1)
 Aspect: Southerly (1)
 Topographic Position: Midslope (1)

Soil Variables

Soil Drainage: Well drained (1)
 Soil Subgroup:
 Surface Texture: Sand (0)
 Effective Texture: Sand (0)
 Depth to Mottles/Gley:
 Organic Thickness:
 Parent Material: Fluvial (0)
 Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

KUC1 Aw/Rose/Spreading dogbane (n=1)

(*Populus tremuloides*/*Rosa acicularis*/*Androsace androsaefolium*)

In La Butte Creek Wildland Provincial Park, Allen et al. (2002) described this community type on gently sloping fluvial, sandy terraces. This community type also appears to be transitional to the Peace-Athabasca Delta subregion and is associated with the large river systems described in these two subregions. This community type has species characteristic of mesic sites rose, bunchberry and peavine, but it also has species characteristic of drier sites bearberry, spreading dogbane and hairy wildrye. It is intermediate in moisture and nutrients between the blueberry and the buffaloberry dominated ecological sites.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: c blueberry(submesic/medium)

Ecosite Phase: c2 blueberry - Aw(Bw)

Plant Composition	Canopy Cover (%)		
	Mean	Range	Const.
Overstory Tree			
ASPEN (<i>Populus tremuloides</i>)	70.0	70.0-70.0	100
Understory Tree			
ASPEN (<i>Populus tremuloides</i>)	1.0	1.0-1.0	100
Tall Shrub (2 to 5m)			
BEAKED WILLOW (<i>Salix bebbiana</i>)	1.0	1.0-1.0	100
Medium Shrub (0.5 to 2 m)			
TWINFLOWER (<i>Linnaea borealis</i>)	1.0	1.0-1.0	100
SNOWBERRY (<i>Symphoricarpos albus</i>)	1.0	1.0-1.0	100
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	1.0	1.0-1.0	100
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	2.0	2.0-2.0	100
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	5.0	5.0-5.0	100
PRICKLY ROSE (<i>Rosa acicularis</i>)	15.0	15.0-15.0	100
Tall Forb (>= 30 cm)			
CREAM-COLORED VETCHLING (<i>Lathyrus ochroleucus</i>)	3.0	3.0-3.0	100
SPREADING DOGBANE (<i>Apocynum androsaemifolium</i>)	20.0	20.0-20.0	100
Low Forb (< 30 cm)			
NORTHERN BEDSTRAW (<i>Galium boreale</i>)	1.0	1.0-1.0	100
WILD LILY-OF-THE-VALLEY (<i>Maianthemum canadense</i>)	1.0	1.0-1.0	100
ONE-SIDED WINTERGREEN (<i>Orthilia secunda</i>)	1.0	1.0-1.0	100
BUNCHBERRY (<i>Cornus canadensis</i>)	5.0	5.0-5.0	100
Graminoid			
HAIRY WILD RYE (<i>Elymus innovatus</i>)	5.0	5.0-5.0	100

Environmental Variables

Ecological Status Score: 25-25
 Moisture Regime: Mesic (fresh) (1)
 Nutrient Regime: Mesotrophic (medium) (1)
 Elevation (range): 245 (245-245) M
 Slope (%): 2.5 - 5.99 (1)
 Aspect: Southerly (1)
 Topographic Position: Midslope (1)

Soil Variables

Soil Drainage: Well drained (1)
 Soil Subgroup:
 Surface Texture: Sand (0)
 Effective Texture: Sand (0)
 Depth to Mottles/Gley:
 Organic Thickness:
 Parent Material: Fluvial (0)
 Humus Form

LFH Thickness	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

c6 blueberry grassland (n=4)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: c blueberry(submesic/medium)

Characteristic Species

Shrub

- [1.0] GROUND JUNIPER
Juniperus communis
- [1.0] SASKATOON
Amelanchier alnifolia

Forb

- [1.0] PLAINS WORMWOOD*
Artemisia campestris

Lichen

- [15.0] REINDEER LICHEN
Cladina mitis

Graminoid

- [15.0] HAY SEDGE*
Carex siccata
- [2.0] SLENDER WHEAT GRASS*
Agropyron trachycaulum
- [2.0] ROCKY MOUNTAIN FESCUE
Festuca saximontana

Environmental Variables

Moisture Regime: Xeric (dry) (4)
 Nutrient Regime: Submesotrophic (poor) (4)
 Elevation (range): 291 (290-292) M
 Slope (%): very strong slope (2), strong slope (2)
 Aspect: Southerly (4)
 Topographic Position: Upper Slope (4)

Soil Variables

Soil Drainage: Rapidly drained (4)
 Soil Subgroup:
 Surface Texture: Sandy loam (0)
 Effective Texture: Sand (0)
 Depth to Mottles/Gley: Not Applicable (0)
 Organic Thickness: 0 - 5 cm (0)
 Parent Material: Glaciolacustrine (0)
 Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

KUA3 Hay sedge-Slender wheatgrass (n=4)

(*Carex siccata*-*Agropyron trachycaulum*)

This community type is found on coarse textured, sandy soils. with southerly facing slope. It is generally found on hilltops and steep (>20%) south-facing slopes throughout the Colin Cornwall Lakes Wildland Provincial Park (Allen et al. 2004). This community type generally has better soil conditions than bearberry dominated communities previously described in the bearberry ecological site. This community type was also described on benches and south facing slopes in the Athabasca Plain subregion.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: c blueberry(submesic/medium)

Ecosite Phase: c6 blueberry grassland

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Medium Shrub (0.5 to 2 m)				Ecological Status Score: 40-40
GROUND JUNIPER (<i>Juniperus communis</i>)	1.3	0.0-5.0	50	Moisture Regime: Xeric (dry) (4)
Tall Forb (>= 30 cm)				Nutrient Regime: Submesotrophic (poor) (4)
COMMON BLUE-EYED GRASS (<i>Sisyrinchium montanum</i>)	0.1	0.0-0.5	25	Elevation (range): 291 (290-292) M
PLAINS WORMWOOD (<i>Artemisia campestris</i>)	1.2	0.0-3.0	75	Slope (%): 31 - 45.99 (2), 16 - 30.99 (2)
Low Forb (< 30 cm)				Aspect: Southerly (4)
ROCK LITTLE CLUB-MOSS (<i>Selaginella rupestris</i>)	0.1	0.0-0.5	25	Topographic Position: Upper Slope (4)
RUSTY WOODSIA (<i>Woodsia ilvensis</i>)	0.1	0.0-0.5	25	
THREE-TOOTHED SAXIFRAGE (<i>Saxifraga tricuspidata</i>)	0.2	0.0-0.5	50	
Graminoid				Soil Variables
SLENDER WHEAT GRASS (<i>Agropyron trachycaulum</i>)	1.8	0.5-3.0	100	Soil Drainage: Rapidly drained (4)
ROCKY MOUNTAIN FESCUE (<i>Festuca saximontana</i>)	2.3	0.0-6.0	75	Soil Subgroup:
HAY SEDGE (<i>Carex siccata</i>)	14.5	10.0-25.0	100	Surface Texture: Sandy loam (0)
Moss				Effective Texture: Sand (0)
HAIRY SCREW MOSS (<i>Tortula ruralis</i>)	0.2	0.0-1.0	25	Depth to Mottles/Gley: Not Applicable (0)
Lichen				Organic Thickness: 0 - 5 cm (0)
REINDEER LICHEN (<i>Cladina mitis</i>)	15.2	1.0-50.0	100	Parent Material: Glaciolacustrine (0)
				Humus Form
				LFH Thickness
				Mean
				Min
				Max
				Count
				cm:
				0.00
				0.00
				0.00
				0

c7 tame/disturbed (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: c blueberry(submesic/medium)

General Description

A number of ecological site phases currently have no data. These ecological site phases have been created as place holders because they were described in adjacent subregions (Central Mixedwood).

Characteristic Species

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

d Labrador tea-mesic(mesic/poor) (n=10)

Natural Subregion: Kazan Upland

General Description

This ecosite generally occurs in submesic to subhygric nutrient-poor substrates. Labrador tea and bog cranberry are indicative of the relatively acidic surface soil conditions. It occurs in upland (midslope, upper slope, and crest) or level topographic positions dominantly on till or fluvial over till parent materials. There is commonly a two-tiered even-aged canopy with faster growing jack pine in the higher layer and slower growing black spruce as a secondary canopy. While the Labrador tea-mesic ecosite has plant community types similar to the Labrador tea-hygric ecosite (g), the mesic ecosite tends to occur in upper topographic positions and the presence of mottles in the upper 50 cm. is less common (Beckingham and Archibald 1996).



Successional Relationships

Successionally mature stands that develop on these ecosites may be dominated by black spruce. Residual pine occurring in the climax community are generally very old. The successional mature stage is rare due to high fire frequency.

Indicator Species

Tree

ASPEN
Populus tremuloides
BLACK SPRUCE
Picea mariana
JACK PINE
Pinus banksiana

Shrub

COMMON LABRADOR TEA
Ledum groenlandicum
BOG CRANBERRY
Vaccinium vitis-idaea

Lichen

REINDEER LICHEN
Cladonia mitis

Moss and Liverwort

STAIR-STEP MOSS
Hylocomium splendens

Ecosection: KU Kazan Upland

Environmental Variables

Moisture Regime: Mesic (fresh) (5), Subhygric (moderately moist) (5)
Nutrient Regime: Permesotrophic (rich) (2), Submesotrophic (poor) (4), Mesotrophic (medium) (4)
Elevation (range): 246.5 (180-300) M
Slope (%): gentle slope (2), nearly level (3), very gentle slope (5)
Aspect: Level (1), Northerly (1), Westerly (2), Easterly (3), Southerly (3)
Topographic Position: Level (1), Midslope (2), Upper Slope (3), Lower Slope (4)

Soil Variables

Soil Drainage: Rapidly drained (1), Well drained (2), Moderately well drained (3), Imperfectly drained (4)
Soil Subgroup: GLEYSOL (1), LUVIC GLEYSOL (1), DYSTRIC BRUNISOL (2), EUTRIC BRUNISOL (2), GRAY LUVISOL (4)
Surface Texture: Sandy loam (1), Silty clay loam (1), Clay loam (2), Sand (3), Silt (3)
Effective Texture: Heavy clay (1), Loam (1), Silt (1), Silt loam (1), Sand (6)
Depth to Mottles/Gley:
Organic Thickness: 0 - 5 cm (10)
Parent Material: Glaciofluvial (2), Lacustrine (2), Morainal (3), Glaciolacustrine (3)
Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	4.75	2.00	11.00	10

d1 Labrador tea-mesic Pj-Sb (n=10)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: d Labrador tea-mesic(mesic/poor)

Characteristic Species

Tree

- [20.6] JACK PINE*
Pinus banksiana
- [12.4] ASPEN*
Populus tremuloides
- [19.1] BLACK SPRUCE*
Picea mariana
- [3.3] WHITE BIRCH
Betula papyrifera

Shrub

- [6.0] BOG CRANBERRY*
Vaccinium vitis-idaea
- [5.0] COMMON LABRADOR TEA*
Ledum groenlandicum
- [4.3] BUNCHBERRY
Cornus canadensis
- [2.5] LOW-BUSH CRANBERRY
Viburnum edule
- [1.7] TWINFLOWER
Linnaea borealis
- [1.2] CANADA BUFFALOBERRY
Shepherdia canadensis

Lichen

- [7.9] REINDEER LICHEN*
Cladonia mitis

Moss and Liverwort

- [13.7] STAIR-STEP MOSS*
Hylocomium splendens
- [13.4] SCHREBER'S MOSS
Pleurozium schreberi

Environmental Variables

Moisture Regime: Mesic (fresh) (5), Subhygric (moderately moist) (5)

Nutrient Regime: Permesotrophic (rich) (2), Mesotrophic (medium) (4), Submesotrophic (poor) (4)

Elevation (range): 246.5 (180-300) M

Slope (%): gentle slope (2), nearly level (3), very gentle slope (5)

Aspect: Level (1), Northerly (1), Westerly (2), Easterly (3), Southerly (3)

Topographic Position: Level (1), Midslope (2), Upper Slope (3), Lower Slope (4)

Soil Variables

Soil Drainage: Rapidly drained (1), Well drained (2), Moderately well drained (3), Imperfectly drained (4)

Soil Subgroup: GLEYSOL (1), LUVIC GLEYSOL (1), EUTRIC BRUNISOL (2), DYSTRIC BRUNISOL (2), GRAY LUVISOL (4)

Surface Texture: Silty clay loam (1), Sandy loam (1), Clay loam (2), Silt (3), Sand (3)

Effective Texture: Silt (1), Silt loam (1), Heavy clay (1), Loam (1), Sand (6)

Depth to Mottles/Gley:

Organic Thickness: 0 - 5 cm (10)

Parent Material: Glaciofluvial (2), Lacustrine (2), Morainal (3), Glaciolacustrine (3)

Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	4.75	2.00	11.00	10

KUC4 Aw-Pj-Sb/Labrador tea (n=1)

(*Populus tremuloides*-*Picea mariana*-*Pinus banksiana*/*Ledum groenlandicum*)

This community type is a moisture variant of the ecological with a slightly richer nutrient regime which favours the growth of aspen and more medium nutrient species like wild sarsaparilla. Successionally mature sites are often dominated by black spruce within this ecological site. The presence of a high cover of aspen and alder indicate the slightly better nutrients. This community type is rapidly drained and has a sandy soil texture at the surface. In the absence of disturbance this community type will likely succeed to a black spruce dominated community.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: d Labrador tea-mesic(mesic/poor)

Ecosite Phase: d1 Labrador tea-mesic Pj-Sb

Plant Composition	Canopy Cover (%)			Const.	Environmental Variables				
	Mean	Range							
Overstory Tree					Ecological Status Score: 25-25				
JACK PINE (<i>Pinus banksiana</i>)	10.0	10.0-10.0		100	Moisture Regime: Subhygric (moderately moist) (1)				
ASPEN (<i>Populus tremuloides</i>)	35.0	35.0-35.0		100	Nutrient Regime: Permesotrophic (rich) (1)				
Understory Tree					Elevation (range): 230 (230-230) M				
BLACK SPRUCE (<i>Picea mariana</i>)	8.0	8.0-8.0		100	Slope (%): 6 - 9.99 (1)				
ASPEN (<i>Populus tremuloides</i>)	8.0	8.0-8.0		100	Aspect: Southerly (1)				
Tall Shrub (2 to 5m)					Topographic Position: Lower Slope (1)				
GREEN ALDER (<i>Alnus crispa</i>)	4.0	4.0-4.0		100	Soil Variables				
Medium Shrub (0.5 to 2 m)					Soil Drainage: Imperfectly drained (1)				
TWINFLOWER (<i>Linnaea borealis</i>)	2.0	2.0-2.0		100	Soil Subgroup: GRAY LUVISOL GLEYED (1)				
PRICKLY ROSE (<i>Rosa acicularis</i>)	2.0	2.0-2.0		100	Surface Texture: Sandy loam (1)				
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	2.0	2.0-2.0		100	Effective Texture: Sand (1)				
BLACK SPRUCE (<i>Picea mariana</i>)	4.0	4.0-4.0		100	Depth to Mottles/Gley:				
COMMON LABRADOR TEA (<i>Ledum groenlandicum</i>)	15.0	15.0-15.0		100	Organic Thickness: 0 - 5 cm (1)				
Tall Forb (>= 30 cm)					Parent Material: Morainal (1)				
COMMON FIREWEED (<i>Epilobium angustifolium</i>)	1.0	1.0-1.0		100	Humus Form				
WOODLAND HORSETAIL (<i>Equisetum sylvaticum</i>)	1.0	1.0-1.0		100	LFH Thickness				
WILD SARSAPARILLA (<i>Aralia nudicaulis</i>)	2.0	2.0-2.0		100	cm:	Mean	Min	Max	Count
Low Forb (< 30 cm)						6.00	6.00	6.00	1
WILD LILY-OF-THE-VALLEY (<i>Maianthemum canadense</i>)	1.0	1.0-1.0		100					
ONE-SIDED WINTERGREEN (<i>Orthilia secunda</i>)	1.0	1.0-1.0		100					
BUNCHBERRY (<i>Cornus canadensis</i>)	8.0	8.0-8.0		100					
Moss									
STAIR-STEP MOSS (<i>Hylocomium splendens</i>)	2.0	2.0-2.0		100					
SCHREBER'S MOSS (<i>Pleurozium schreberi</i>)	2.0	2.0-2.0		100					

KUD1 Sb-Aw/Moss (n=3)

(*Picea mariana*-*Populus tremuloides*/Stair step moss)

This community type is a moist variant with a slightly richer nutrient regime of the Labrador tea-mesic ecological site. Successionally mature sites are often dominated by black spruce within this ecological site. The presence of a high cover of aspen and stair step moss indicate the slightly better nutrients. This community type is well drained and has a sandy soil texture. In the absence of disturbance this community type will likely succeed to a black spruce dominated community.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: d Labrador tea-mesic(mesic/poor)
Ecosite Phase: d1 Labrador tea-mesic Pj-Sb

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
ASPEN (<i>Populus tremuloides</i>)	5.0	0.0-10.0	67	Moisture Regime: Mesic (fresh) (1), Subhygric (moderately moist) (2)
WHITE BIRCH (<i>Betula papyrifera</i>)	13.3	0.0-35.0	67	Nutrient Regime: Submesotrophic (poor) (1), Mesotrophic (medium) (1), Permesotrophic (rich) (1)
BLACK SPRUCE (<i>Picea mariana</i>)	23.3	20.0-30.0	100	Elevation (range): 222 (180-255) M
Understory Tree				Slope (%): 0.5 - 2.49 (1), 2.5 - 5.99 (2)
ASPEN (<i>Populus tremuloides</i>)	1.6	0.0-5.0	33	Aspect: Westerly (1), Southerly (2)
BLACK SPRUCE (<i>Picea mariana</i>)	6.6	0.0-20.0	33	Topographic Position: Lower Slope (1), Midslope (1), Upper Slope (1)
Tall Shrub (2 to 5m)				Soil Variables
BEAKED WILLOW (<i>Salix bebbiana</i>)	2.3	0.0-4.0	67	Soil Drainage: Imperfectly drained (1), Moderately well drained (2)
Medium Shrub (0.5 to 2 m)				Soil Subgroup: GRAY LUVISOL GLEYED (1), GRAY LUVISOL ORTHIC (1), LUVIC GLEYSOL ORTHIC (1)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	0.6	0.0-2.0	33	Surface Texture: Silt (1), Clay loam (2)
CANADA BUFFALOBERRY (<i>Shepherdia canadensis</i>)	5.0	0.0-15.0	33	Effective Texture: Heavy clay (1), Silt (1), Silt loam (1)
Low Forb (< 30 cm)				Depth to Mottles/Gley:
NORTHERN BASTARD TOADFLAX (<i>Geocaulon lividum</i>)	3.3	0.0-10.0	33	Organic Thickness: 0 - 5 cm (3)
Graminoid				Parent Material: Glaciolacustrine (1), Lacustrine (2)
BLUEJOINT (<i>Calamagrostis canadensis</i>)	5.0	0.0-15.0	33	Humus Form
Moss				LFH Thickness
SCHREBER'S MOSS (<i>Pleurozium schreberi</i>)	13.3	5.0-30.0	100	Mean
STAIR-STEP MOSS (<i>Hylocomium splendens</i>)	35.0	15.0-50.0	100	Min
Lichen				Max
REINDEER LICHEN (<i>Cladina mitis</i>)	2.6	2.0-3.0	100	Count
				cm: 7.00 5.00 11.00 3

KUE11 Pj/Bog cranberry/lichen (n=1)

(*Pinus banksiana/Vaccinium vitis idea/Cladina spp.*)

This community type is a drier variant of the Labrador tea-mesic ecological site. Successionally mature sites are often dominated by black spruce within this ecological site. This community type is rapidly drained and has a coarse texture.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: d Labrador tea-mesic(mesic/poor)

Ecosite Phase: d1 Labrador tea-mesic Pj-Sb

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
JACK PINE (<i>Pinus banksiana</i>)	42.0	42.0-42.0	100	Moisture Regime: Mesic (fresh) (1)
Understory Tree				Nutrient Regime: Submesotrophic (poor) (1)
JACK PINE (<i>Pinus banksiana</i>)	1.0	1.0-1.0	100	Elevation (range): 300 (300-300) M
Medium Shrub (0.5 to 2 m)				Slope (%): 2.5 - 5.99 (1)
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	1.0	1.0-1.0	100	Aspect: Easterly (1)
TWINFLOWER (<i>Linnaea borealis</i>)	2.0	2.0-2.0	100	Topographic Position: Upper Slope (1)
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	8.0	8.0-8.0	100	
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	18.0	18.0-18.0	100	
Low Forb (< 30 cm)				Soil Variables
ONE-SIDED WINTERGREEN (<i>Orthilia secunda</i>)	1.0	1.0-1.0	100	Soil Drainage: Well drained (1)
NORTHERN BASTARD TOADFLAX (<i>Geocaulon lividum</i>)	2.0	2.0-2.0	100	Soil Subgroup: DYSTRIC BRUNISOL ELUVIATED (1)
BUNCHBERRY (<i>Cornus canadensis</i>)	8.0	8.0-8.0	100	Surface Texture: Silt (1)
				Effective Texture: Sand (1)
				Depth to Mottles/Gley:
				Organic Thickness: 0 - 5 cm (1)
				Parent Material: Glaciolacustrine (1)
				Humus Form
Moss				LFH Thickness
STAIR-STEP MOSS (<i>Hylocomium splendens</i>)	18.0	18.0-18.0	100	Mean
SCHREBER'S MOSS (<i>Pleurozium schreberi</i>)	29.0	29.0-29.0	100	Min
				Max
				Count
Lichen				cm:
N/A (<i>Cladonia multiformis</i>)	1.0	1.0-1.0	100	3.00
N/A (<i>Cladonia uncialis</i>)	1.0	1.0-1.0	100	3.00
REINDEER LICHEN (<i>Cladina mitis</i>)	18.0	18.0-18.0	100	3.00
				1

KUE4 Pj- Sb/Bog cranberry/lichen (n=5)

(*Pinus banksiana*-*Picea mariana*/*Vaccinium vitis idea*/*Cladina spp.*)

This community type represents a successionaly immature stand of this ecological site. Successionaly mature sites are often dominated by black spruce within this ecological site. This community type is rapidly drained and has a coarse texture.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: d Labrador tea-mesic(mesic/poor)

Ecosite Phase: d1 Labrador tea-mesic Pj-Sb

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
JACK PINE (<i>Pinus banksiana</i>)	20.2	15.0-25.0	100	Moisture Regime: Subhygric (moderately moist) (2), Mesic (fresh) (3)
Understory Tree				Nutrient Regime: Submesotrophic (poor) (2), Mesotrophic (medium) (3)
BLACK SPRUCE (<i>Picea mariana</i>)	6.4	0.0-15.0	80	Elevation (range): 234 (230-239) M
JACK PINE (<i>Pinus banksiana</i>)	9.6	5.0-15.0	100	Slope (%): 6 - 9.99 (1), 2.5 - 5.99 (2), 0.5 - 2.49 (2)
Tall Shrub (2 to 5m)				Aspect: Level (1), Northerly (1), Westerly (1), Easterly (2)
JACK PINE (<i>Pinus banksiana</i>)	2.4	0.0-4.0	80	Topographic Position: Level (1), Midslope (1), Upper Slope (1), Lower Slope (2)
BLACK SPRUCE (<i>Picea mariana</i>)	15.4	5.0-40.0	100	
Medium Shrub (0.5 to 2 m)				Soil Variables
TWINFLOWER (<i>Linnaea borealis</i>)	2.8	1.0-5.0	100	Soil Drainage: Rapidly drained (1), Well drained (1), Moderately well drained (1), Imperfectly drained (2)
COMMON BLUEBERRY (<i>Vaccinium myrtilloides</i>)	3.8	0.0-12.0	60	Soil Subgroup: DYSTRIC BRUNISOL GLEYED ELUVIATED (1), EUTRIC BRUNISOL ELUVIATED (1), EUTRIC BRUNISOL GLEYED ELUVIATED (1), GLEYSOL ORTHIC (1), GRAY LUVISOL ORTHIC (1)
COMMON LABRADOR TEA (<i>Ledum groenlandicum</i>)	5.0	0.0-10.0	80	Surface Texture: Silt (1), Silty clay loam (1), Sand (3)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	5.6	0.0-10.0	80	Effective Texture: Loam (1), Sand (4)
BLACK SPRUCE (<i>Picea mariana</i>)	12.6	2.0-40.0	100	Depth to Mottles/Gley:
Low Forb (< 30 cm)				Organic Thickness: 0 - 5 cm (5)
BUNCHBERRY (<i>Cornus canadensis</i>)	1.2	0.0-3.0	60	Parent Material: Glaciolacustrine (1), Morainal (2), Glaciofluvial (2)
Moss				Humus Form
JUNIPER HAIR-CAP (<i>Polytrichum juniperinum</i>)	2.0	0.0-10.0	20	LFH Thickness
SCHREBER'S MOSS (<i>Pleurozium schreberi</i>)	9.4	0.0-45.0	40	Mean
Lichen				Min
REINDEER LICHEN (<i>Cladina mitis</i>)	11.2	0.0-16.0	80	Max
				Count
				cm:
				3.00
				2.00
				5.00
				5

e buffaloberry/alder(mesic/medium) (n=16)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

This is the reference ecosite for the Kazan Upland subregion because it has a mesic moisture regime and a medium nutrient regime. However, the conventional reference site concept of deep, medium textured, well drained and associated vegetation does not generally fit for most of this subregion. The majority of the characteristic sites in this subregion are rocky exposures or dry, rapidly drained coarse glacial deposits. Generally, this ecosite has fine to medium textured glaciolacustrine and glacialfluvial parent materials. This ecological site is located at a mid to lower slope positions where some moisture accumulates. These sites are often dominated by aspen, green alder and buffaloberry and have a high diversity of shrubs and forbs in the understory.



Successional Relationships

Pioneer jack pine and deciduous tree species such as aspen, balsam poplar, and white birch are replaced by white spruce, black spruce and balsam fir as the sites develop successionaly. Along with a change in canopy composition is a change in understory structure and understory species composition and abundance. Generally, as a stand successionaly matures, the coniferous canopy cover increases, and understory species structure and diversity declines. This results in stands with low cover of shrub, forb, and grass species and high moss cover.

Indicator Species

Tree

ASPEN
Populus tremuloides
WHITE SPRUCE
Picea glauca
JACK PINE
Pinus banksiana

Shrub

CANADA BUFFALOBERRY
Shepherdia canadensis
GREEN ALDER
Alnus crispa

Moss and Liverwort

STAIR-STEP MOSS
Hylocomium splendens
SCHREBER'S MOSS
Pleurozium schreberi

Graminoid

HAIRY WILD RYE
Elymus innovatus

Site Index at 50 Years

Site Index at 50 Years	Height (m)	Variation (m)	Count
WHITE SPRUCE <i>(Picea glauca)</i>	12.00	0.00	0
WHITE BIRCH <i>(Betula papyrifera)</i>	10.40	0.60	0

Environmental Variables

Moisture Regime: Subhygric (moderately moist) (2), Submesic (moderately fresh) (3), Mesic (fresh) (10)

Nutrient Regime: Permesotrophic (rich) (2), Submesotrophic (poor) (6), Mesotrophic (medium) (7)

Elevation (range): 243.86 (180-330) M

Slope (%): extreme slope (1), nearly level (1), strong slope (1), level (2), gentle slope (3), very gentle slope (7)

Aspect: Level (2), Southerly (2), Westerly (3), Easterly (3), Northerly (4)

Topographic Position: Depression (1), Upper Slope (2), Level (3), Lower Slope (3), Midslope (7)

Soil Variables

Soil Drainage: Rapidly drained (1), Moderately well drained (6), Well drained (9)

Soil Subgroup: DYSTRIC BRUNISOL (3), EUTRIC BRUNISOL (5), GRAY LUVISOL (7)

Surface Texture: Sandy clay loam (1), Sandy loam (1), Silt (1), Loam (1), Silty clay (1), Silty clay loam (1), Silt loam (2), Sand (6)

Effective Texture: Clay (1), Silt loam (1), Silty clay loam (1), Sandy clay loam (2), Silty clay (3), Sand (6)

Depth to Mottles/Gley:

Organic Thickness: 0 - 5 cm (15)

Parent Material: Colluvial (1), Morainal (1), Rock (2), Fluvial (2), Lacustrine (3), Glaciolacustrine (5), Glaciofluvial (7)

Humus Form

LFH Thickness	Mean	Min	Max	Count
cm:	5.50	3.00	11.00	14

e1 buffaloberry-alder/ Aw (n=8)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Characteristic Species

Tree

- [67.4] ASPEN*
Populus tremuloides

Shrub

- [16.4] BOG CRANBERRY
Vaccinium vitis-idaea
- [11.6] CANADA BUFFALOBERRY*
Shepherdia canadensis
- [7.1] PRICKLY ROSE
Rosa acicularis
- [5.5] GREEN ALDER*
Alnus crispa
- [5.0] SASKATOON
Amelanchier alnifolia
- [3.7] LOW-BUSH CRANBERRY
Viburnum edule
- [3.4] BUNCHBERRY
Cornus canadensis
- [3.3] RED-OSIER DOGWOOD
Cornus stolonifera
- [3.0] TWINFLOWER
Linnaea borealis
- [2.3] BEAKED WILLOW
Salix bebbiana
- [2.0] DEWBERRY
Rubus pubescens

Forb

- [5.0] SPREADING DOGBANE
Apocynum androsaemifolium
- [3.8] COMMON FIREWEED
Epilobium angustifolium
- [3.3] CREAM-COLORED VETCHLING
Lathyrus ochroleucus
- [2.1] WILD SARSAPARILLA
Aralia nudicaulis

Graminoid

- [6.8] HAIRY WILD RYE*
Elymus innovatus

Environmental Variables

Moisture Regime: Submesic (moderately fresh) (1), Mesic (fresh) (6)

Nutrient Regime: Submesotrophic (poor) (2), Mesotrophic (medium) (5)

Elevation (range): 231.33 (200-260) M

Slope (%): extreme slope (1), gentle slope (1), level (1), nearly level (1), strong slope (1), very gentle slope (3)

Aspect: Easterly (1), Level (1), Southerly (1), Westerly (2), Northerly (2)

Topographic Position: Lower Slope (1), Upper Slope (2), Level (2), Midslope (3)

Soil Variables

Soil Drainage: Moderately well drained (3), Well drained (5)

Soil Subgroup: DYSTRIC BRUNISOL (1), EUTRIC BRUNISOL (2), GRAY LUVISOL (4)

Surface Texture: Sandy loam (1), Silt (1), Silty clay loam (1), Sand (2), Silt loam (2)

Effective Texture: Silty clay loam (1), Silt loam (1), Sandy clay loam (1), Sand (2), Silty clay (2)

Depth to Mottles/Gley:

Organic Thickness: 0 - 5 cm (7)

Parent Material: Fluvial (0), Morainal (1), Glaciofluvial (1), Rock (2), Lacustrine (2), Glaciolacustrine (3)

Humus Form

LFH Thickness

cm:

Mean	Min	Max	Count
6.50	4.00	9.00	7

KUC2 Aw(Bw)/Alder (n=3)

(*Populus tremuloides* (*Betula papyrifera*))/*Alnus crispa*)

This reference plant community represents the deciduous phase of the buffaloberry-alder ecological site in the Kazan upland subregion. Moisture and nutrients are intermediate between the drier blueberry ecosite and the moister fern/horsetail ecological site. Succession in the absence of disturbance is to a white spruce climax community.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)
Ecosite Phase: e1 buffaloberry-alder/ Aw

Plant Composition	Canopy Cover (%)			Const.	Environmental Variables
	Mean	Range			
Overstory Tree					Ecological Status Score: 25-25 Moisture Regime: Submesic (moderately fresh) (1), Mesic (fresh) (2) Nutrient Regime: Mesotrophic (medium) (1), Submesotrophic (poor) (2) Elevation (range): 237 (200-260) M Slope (%): 16 - 30.99 (1), 2.5 - 5.99 (2) Aspect: Northerly (1), Southerly (1), Westerly (1) Topographic Position: Lower Slope (1), Upper Slope (2)
JACK PINE (<i>Pinus banksiana</i>)	2.0	1.0-3.0	100		
ASPEN (<i>Populus tremuloides</i>)	45.0	35.0-60.0	100		
Understory Tree					
ASPEN (<i>Populus tremuloides</i>)	15.0	10.0-20.0	100		
Tall Shrub (2 to 5m)					
GREEN ALDER (<i>Alnus crispa</i>)	16.6	15.0-20.0	100		
Medium Shrub (0.5 to 2 m)					
TWINFLOWER (<i>Linnaea borealis</i>)	1.6	0.0-3.0	67		
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	1.6	1.0-2.0	100		
PRICKLY ROSE (<i>Rosa acicularis</i>)	2.0	1.0-4.0	100		
GREEN ALDER (<i>Alnus crispa</i>)	5.0	0.0-15.0	33		
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	13.0	2.0-30.0	100		
Low Shrub (< 0.5m)					
DEWBERRY (<i>Rubus pubescens</i>)	3.3	0.0-10.0	33		
Tall Forb (>= 30 cm)					
COMMON FIREWEED (<i>Epilobium angustifolium</i>)	3.0	0.0-8.0	67		
WILD SARSAPARILLA (<i>Aralia nudicaulis</i>)	6.3	0.0-15.0	67		
Low Forb (< 30 cm)					
PALMATE-LEAVED COLTSFOOT (<i>Petasites palmatus</i>)	1.0	0.0-3.0	33		
NORTHERN BASTARD TOADFLAX (<i>Geocaulon lividum</i>)	5.6	0.0-15.0	67		
BUNCHBERRY (<i>Cornus canadensis</i>)	8.3	6.0-12.0	100		
Moss					
SCHREBER'S MOSS (<i>Pleurozium schreberi</i>)	1.6	0.0-4.0	67		
Lichen					
STUDED LEATHER LICHEN (<i>Peltigera aphthosa</i>)	1.6	0.0-5.0	33		

Soil Variables

Soil Drainage: Moderately well drained (1), Well drained (2)
 Soil Subgroup: DYSTRIC BRUNISOL ELUVIATED (1), EUTRIC BRUNISOL ELUVIATED (1), GRAY LUVISOL ORTHIC (1)
 Surface Texture: Sand (1), Silt loam (1), Sandy loam (1)
 Effective Texture: Sandy clay loam (1), Sand (2)
 Depth to Mottles/Gley:
 Organic Thickness: 0 - 5 cm (3)
 Parent Material: Glaciofluvial (1), Lacustrine (1), Morainal (1), Rock (2)
 Humus Form

LFH Thickness	Mean	Min	Max	Count
cm:	6.00	4.00	8.00	3

KUC3 Aw/Buffaloberry (n=4)

(*Populus tremuloides*/ *Shepherdia canadensis*)

This community type was found on mesic sites, with shallow slopes and had medium nutrient regimes. Beckingham (1993) felt the Aw/Buffaloberry type was slightly drier and had a slightly poorer nutrient regime than the more modal Aw/Alder or Aw/Low bush cranberry dominated community types. A similar buffaloberry dominated community was described at upper elevations in the Birch and Saddle Hills of the Boreal Mixedwood Natural Region (Moisey et al. 2012).

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Ecosite Phase: e1 buffaloberry-alder/ Aw

Plant Composition	Canopy Cover (%)			Const.	Environmental Variables
	Mean	Range			
Overstory Tree					Ecological Status Score: 25-25
JACK PINE (<i>Pinus banksiana</i>)	1.7	0.0-5.0	50		Moisture Regime: Mesic (fresh) (4)
ASPEN (<i>Populus tremuloides</i>)	46.2	40.0-55.0	100		Nutrient Regime: Mesotrophic (medium) (4)
Understory Tree					Elevation (range): 212 (205-225) M
ASPEN (<i>Populus tremuloides</i>)	15.0	0.0-30.0	75		Slope (%): 0 - 0.49 (1), 0.5 - 2.49 (1), 6 - 9.99 (1), 2.5 - 5.99 (1)
Tall Shrub (2 to 5m)					Aspect: Level (1), Easterly (1), Westerly (1)
ASPEN (<i>Populus tremuloides</i>)	1.2	0.0-4.0	50		Topographic Position: Level (2), Midslope (2)
BEAKED WILLOW (<i>Salix bebbiana</i>)	2.0	1.0-3.0	100		
Medium Shrub (0.5 to 2 m)					Soil Variables
ASPEN (<i>Populus tremuloides</i>)	1.2	0.0-2.0	75		Soil Drainage: Well drained (2), Moderately well drained (2)
TWINFLOWER (<i>Linnaea borealis</i>)	2.5	1.0-5.0	100		Soil Subgroup: EUTRIC BRUNISOL ELUVIATED (1), GRAY LUVISOL BRUNISOLIC (1), GRAY LUVISOL ORTHIC (2)
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	2.7	0.0-5.0	75		Surface Texture: Sand (1), Silt (1), Silty clay loam (1), Silt loam (1)
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	4.5	0.0-15.0	50		Effective Texture: Silty clay loam (1), Silt loam (1), Silty clay (2)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	6.2	0.0-15.0	50		Depth to Mottles/Gley:
PRICKLY ROSE (<i>Rosa acicularis</i>)	9.5	3.0-20.0	100		Organic Thickness: 0 - 5 cm (4)
CANADA BUFFALOBERRY (<i>Shepherdia canadensis</i>)	35.0	10.0-60.0	100		Parent Material: Lacustrine (1), Glaciolacustrine (3)
Low Shrub (< 0.5m)					Humus Form
DEWBERRY (<i>Rubus pubescens</i>)	2.7	0.0-6.0	50		
Tall Forb (>= 30 cm)					LFH Thickness
COMMON FIREWEED (<i>Epilobium angustifolium</i>)	3.5	0.0-6.0	75		Mean
Low Forb (< 30 cm)					Min
NORTHERN BEDSTRAW (<i>Galium boreale</i>)	1.0	0.0-2.0	75		Max
PALMATE-LEAVED COLTSFOOT (<i>Petasites palmatus</i>)	1.0	0.0-3.0	50		Count
BUNCHBERRY (<i>Cornus canadensis</i>)	2.0	2.0-2.0	100		cm:
Graminoid					7.00
BLUEJOINT (<i>Calamagrostis canadensis</i>)	4.0	0.0-10.0	75		4.00
HAIRY WILD RYE (<i>Elymus innovatus</i>)	10.5	1.0-20.0	100		9.00
					4

KUC7 Aw-Pb/Saskatoon-Red Osier Dogwood/Spreading dogbane (n=1)

(*Populus tremuloides*-*Populus balsamifera*/ *Amelanchier alnifolia*-*Cornus stolonifera*/*Apocynum androsaefolium*)

This community type was found on mesic to subhygric sites, with shallow slopes and had medium nutrient regimes. The presence of Balsam poplar and red osier dogwood indicate the transition to the moisture and richer ecological site. A similar spreading dogbane community was described in the La Butte Creek Wildland Park but the other site was described on slightly drier sites. Spreading dogbane dominated community types appears to be a provincially rare plant community type and should be recognized as a special feature of provincial significance (Allen et al. 2002).

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Ecosite Phase: e1 buffaloberry-alder/ Aw

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
BALSAM POPLAR (<i>Populus balsamifera</i>)	5.0	5.0-5.0	100	Moisture Regime:
ASPEN (<i>Populus tremuloides</i>)	75.0	75.0-75.0	100	Nutrient Regime:
Tall Shrub (2 to 5m)				Elevation (range): 245 (245-245) M
ASPEN (<i>Populus tremuloides</i>)	5.0	5.0-5.0	100	Slope (%): > 100.99 (1)
BEAKED WILLOW (<i>Salix bebbiana</i>)	5.0	5.0-5.0	100	Aspect: Northerly (1)
Medium Shrub (0.5 to 2 m)				Topographic Position: Midslope (1)
TWINING HONEYSUCKLE (<i>Lonicera dioica</i>)	2.0	2.0-2.0	100	Soil Variables
TWINFLOWER (<i>Linnaea borealis</i>)	5.0	5.0-5.0	100	Soil Drainage: Well drained (1)
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	5.0	5.0-5.0	100	Soil Subgroup:
RED-OSIER DOGWOOD (<i>Cornus stolonifera</i>)	10.0	10.0-10.0	100	Surface Texture: Sand (0)
PRICKLY ROSE (<i>Rosa acicularis</i>)	10.0	10.0-10.0	100	Effective Texture:
SASKATOON (<i>Amelanchier alnifolia</i>)	15.0	15.0-15.0	100	Depth to Mottles/Gley:
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	30.0	30.0-30.0	100	Organic Thickness:
Tall Forb (>= 30 cm)				Parent Material: Fluvial (0)
COMMON FIREWEED (<i>Epilobium angustifolium</i>)	5.0	5.0-5.0	100	Humus Form
CREAM-COLORED VETCHLING (<i>Lathyrus ochroleucus</i>)	10.0	10.0-10.0	100	LFH Thickness
SPREADING DOGBANE (<i>Apocynum androsaemifolium</i>)	15.0	15.0-15.0	100	cm:
Low Forb (< 30 cm)				Mean
NORTHERN BEDSTRAW (<i>Galium boreale</i>)	5.0	5.0-5.0	100	Min
WILD LILY-OF-THE-VALLEY (<i>Maianthemum canadense</i>)	5.0	5.0-5.0	100	Max
Graminoid				Count
HAIRY WILD RYE (<i>Elymus innovatus</i>)	10.0	10.0-10.0	100	0

e2 buffaloberry-alder/ Pj-Sw-Sb-Aw-Bw (n=5)

Natural Subregion: Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Ecosection: KU Kazan Upland

Characteristic Species

Tree

- [28.2] ASPEN
Populus tremuloides
- [20.0] JACK PINE*
Pinus banksiana
- [6.3] WHITE SPRUCE*
Picea glauca
- [1.3] BLACK SPRUCE
Picea mariana

Shrub

- [15.0] CANADA BUFFALOBERRY
Shepherdia canadensis
- [13.2] BOG CRANBERRY
Vaccinium vitis-idaea
- [8.7] GREEN ALDER
Alnus crispa
- [6.6] BUNCHBERRY
Cornus canadensis
- [5.5] COMMON BEARBERRY
Arctostaphylos uva-ursi
- [4.9] TWINFLOWER
Linnaea borealis
- [2.0] DEWBERRY
Rubus pubescens
- [1.6] PRICKLY ROSE
Rosa acicularis

Lichen

- [6.2] REINDEER LICHEN
Cladonia mitis

Moss and Liverwort

- [8.3] STAIR-STEP MOSS*
Hylocomium splendens

Environmental Variables

Moisture Regime: Submesic (moderately fresh) (1), Mesic (fresh) (4)
 Nutrient Regime: Mesotrophic (medium) (1), Submesotrophic (poor) (2),
 Permesotrophic (rich) (2)
 Elevation (range): 216.5 (180-240) M
 Slope (%): level (1), very gentle slope (2), gentle slope (2)
 Aspect: Easterly (1), Level (1), Southerly (1), Northerly (2)
 Topographic Position: Lower Slope (1), Midslope (4)

Soil Variables

Soil Drainage: Rapidly drained (1), Well drained (2), Moderately well drained (2)
 Soil Subgroup: EUTRIC BRUNISOL (2), GRAY LUVISOL (3)
 Surface Texture: Sand (1), Silty clay (1), Sandy clay loam (1), Loam (1)
 Effective Texture: Silty clay (1), Sandy clay loam (1), Sand (1), Clay (1)
 Depth to Mottles/Gley:
 Organic Thickness: 0 - 5 cm (5)
 Parent Material: Lacustrine (1), Colluvial (1), Fluvial (2), Glaciofluvial (5)
 Humus Form

LFH Thickness	Mean	Min	Max	Count
cm:	4.50	4.00	6.00	4

KUD2 Aw-Sw/Buffaloberry (n=3)

(*Populus tremuloides*-*Picea glauca*/ *Shepherdia canadensis*)

This community type was found on mesic sites, with shallow slopes and had medium nutrient regimes. Beckingham (1993) felt the Aw/Buffaloberry type was slightly drier and had a slightly poorer nutrient regime than the more modal Aw/Alder or Aw/Low bush cranberry dominated community types. In the absence of disturbance Spruce will dominate this plant community type.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Ecosite Phase: e2 buffaloberry-alder/ Pj-Sw-Sb-Aw-Bw

Plant Composition	Canopy Cover (%)			Const.	Environmental Variables
	Mean	Range			
Overstory Tree					Ecological Status Score: 25-25
BLACK SPRUCE (<i>Picea mariana</i>)	6.0	5.0-8.0	100		Moisture Regime: Mesic (fresh) (3)
WHITE SPRUCE (<i>Picea glauca</i>)	6.6	0.0-15.0	67		Nutrient Regime: Mesotrophic (medium) (1), Permesotrophic (rich) (2)
ASPEN (<i>Populus tremuloides</i>)	18.3	5.0-35.0	100		Elevation (range): 193 (180-210) M
Understory Tree					Slope (%): 0 - 0.49 (1), 2.5 - 5.99 (1), 6 - 9.99 (1)
WHITE SPRUCE (<i>Picea glauca</i>)	6.0	2.0-10.0	100		Aspect: Level (1), Northerly (2)
BLACK SPRUCE (<i>Picea mariana</i>)	6.6	2.0-10.0	100		Topographic Position: Lower Slope (1), Midslope (2)
ASPEN (<i>Populus tremuloides</i>)	18.3	5.0-40.0	100		Soil Variables
Tall Shrub (2 to 5m)					Soil Drainage: Well drained (1), Moderately well drained (2)
BLACK SPRUCE (<i>Picea mariana</i>)	2.6	0.0-6.0	67		Soil Subgroup: GRAY LUVISOL BRUNISOLIC (1), GRAY LUVISOL ORTHIC (2)
WHITE SPRUCE (<i>Picea glauca</i>)	3.3	2.0-5.0	100		Surface Texture: Loam (1), Sandy clay loam (1), Silty clay (1)
Medium Shrub (0.5 to 2 m)					Effective Texture: Clay (1), Sandy clay loam (1), Silty clay (1)
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	2.0	2.0-2.0	100		Depth to Mottles/Gley:
PRICKLY ROSE (<i>Rosa acicularis</i>)	2.3	0.0-4.0	67		Organic Thickness: 0 - 5 cm (3)
BLACK SPRUCE (<i>Picea mariana</i>)	3.0	1.0-5.0	100		Parent Material: Colluvial (1), Lacustrine (1), Fluvial (2)
TWINFLOWER (<i>Linnaea borealis</i>)	3.3	0.0-5.0	67		Humus Form
WHITE SPRUCE (<i>Picea glauca</i>)	5.0	2.0-10.0	100		LFH Thickness
GREEN ALDER (<i>Alnus crispa</i>)	9.0	5.0-12.0	100		Mean
CANADA BUFFALOBERRY (<i>Shepherdia canadensis</i>)	30.0	20.0-50.0	100		Min
Low Shrub (< 0.5m)					Max
DEWBERRY (<i>Rubus pubescens</i>)	4.0	2.0-8.0	100		Count
Tall Forb (>= 30 cm)					cm:
COMMON FIREWEED (<i>Epilobium angustifolium</i>)	1.6	1.0-3.0	100		5.00
Low Forb (< 30 cm)					5.00
BUNCHBERRY (<i>Cornus canadensis</i>)	12.3	7.0-20.0	100		6.00
Moss					3
STAIR-STEP MOSS (<i>Hylocomium splendens</i>)	16.6	15.0-20.0	100		

KUD4 Aw- Pj/Alder (n=2)

(*Populus tremuloides*-*Pinus banksiana*/*Alnus crispa*)

A better water supply in the rooting zone due to aspect, finer soil textures at depth or local groundwater flow contributes to better understory growth of alder and feather mosses. Typically these sites are often dominated by a mixture of aspen and jackpine.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Ecosite Phase: e2 buffaloberry-alder/ Pj-Sw-Sb-Aw-Bw

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
ASPEN (<i>Populus tremuloides</i>)	15.0	0.0-30.0	50	Moisture Regime: Submesic (moderately fresh) (1), Mesic (fresh) (1)
JACK PINE (<i>Pinus banksiana</i>)	25.0	20.0-30.0	100	Nutrient Regime: Submesotrophic (poor) (2)
Understory Tree				Elevation (range): 240 (240-240) M
ASPEN (<i>Populus tremuloides</i>)	5.0	0.0-10.0	50	Slope (%): 2.5 - 5.99 (1), 6 - 9.99 (1)
JACK PINE (<i>Pinus banksiana</i>)	15.0	10.0-20.0	100	Aspect: Easterly (1), Southerly (1)
Tall Shrub (2 to 5m)				Topographic Position: Midslope (2)
GREEN ALDER (<i>Alnus crispa</i>)	17.5	5.0-30.0	100	Soil Variables
Medium Shrub (0.5 to 2 m)				Soil Drainage: Rapidly drained (1), Well drained (1)
PRICKLY ROSE (<i>Rosa acicularis</i>)	1.0	0.0-2.0	50	Soil Subgroup: EUTRIC BRUNISOL ELUVIATED (2)
COMMON BLUEBERRY (<i>Vaccinium myrtilloides</i>)	2.5	2.0-3.0	100	Surface Texture: Sand (1)
TWINFLOWER (<i>Linnaea borealis</i>)	6.5	3.0-10.0	100	Effective Texture: Sand (1)
COMMON BEARBERRY (<i>Arctostaphylos uva-ursi</i>)	11.0	2.0-20.0	100	Depth to Mottles/Gley:
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	26.5	3.0-50.0	100	Organic Thickness: 0 - 5 cm (2)
Low Forb (< 30 cm)				Parent Material: Glaciofluvial (5)
BUNCHBERRY (<i>Cornus canadensis</i>)	1.0	0.0-2.0	50	Humus Form
WILD LILY-OF-THE-VALLEY (<i>Maianthemum canadense</i>)	1.0	1.0-1.0	100	LFH Thickness
ONE-SIDED WINTERGREEN (<i>Orthilia secunda</i>)	1.0	1.0-1.0	100	Mean
Moss				Min
JUNIPER HAIR-CAP (<i>Polytrichum juniperinum</i>)	2.5	0.0-5.0	50	Max
Lichen				Count
REINDEER LICHEN (<i>Cladina mitis</i>)	12.5	0.0-25.0	50	cm:

e3 buffaloberry-alder shrubland (n=0)

Natural Subregion: Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Ecosection: KU Kazan Upland

General Description

A number of ecological site phases currently have no data. These ecological site phases have been created as place holders because they were described in adjacent subregions (Athabasca Plain and Northern Mixedwood).

Characteristic Species

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

e4 buffaloberry-alder/ Pj-Sw-Sb (n=3)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Characteristic Species

Tree

- [30.0] JACK PINE
Pinus banksiana
- [22.2] BLACK SPRUCE
Picea mariana
- [10.5] WHITE SPRUCE
Picea glauca
- [7.5] ASPEN
Populus tremuloides

Shrub

- [20.0] GREEN ALDER
Alnus crispa
- [1.5] TWINFLOWER
Linnaea borealis
- [1.0] BEAKED WILLOW
Salix bebbiana

Forb

- [3.5] NORTHERN BASTARD TOADFLAX
Geocaulon lividum

Moss and Liverwort

- [38.2] SCHREBER'S MOSS*
Pleurozium schreberi
- [7.0] STAIR-STEP MOSS
Hylocomium splendens
- [5.0] WAVY DICRANUM
Dicranum polysetum

Environmental Variables

Moisture Regime: Submesic (moderately fresh) (1), Subhygric (moderately moist) (2)

Nutrient Regime: Mesotrophic (medium) (1), Submesotrophic (poor) (2)

Elevation (range): 290 (250-330) M

Slope (%): very gentle slope (2)

Aspect: Easterly (1), Westerly (1)

Topographic Position: Depression (1), Level (1), Lower Slope (1)

Soil Variables

Soil Drainage: Moderately well drained (1), Well drained (2)

Soil Subgroup: EUTRIC BRUNISOL (1), DYSTRIC BRUNISOL (2)

Surface Texture: Sand (3)

Effective Texture: Sand (3)

Depth to Mottles/Gley:

Organic Thickness: 0 - 5 cm (3)

Parent Material: Glaciofluvial (1), Glaciolacustrine (2)

Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	5.50	3.00	11.00	3

KUE5 Pj/Alder (n=1)

(*Pinus banksiana*/*Alnus crispa*)

The increase in moisture favours the growth of alder, aspen and feather moss in the understory of this community type. This community type occupies lower slope positions that accumulate some moisture. Typically these sites are often dominated by a mixture of aspen and jackpine.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Ecosite Phase: e4 buffaloberry-alder/ Pj-Sw-Sb

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
JACK PINE (<i>Pinus banksiana</i>)	50.0	50.0-50.0	100	Moisture Regime: Submesic (moderately fresh) (1)
Understory Tree				Nutrient Regime: Mesotrophic (medium) (1)
ASPEN (<i>Populus tremuloides</i>)	5.0	5.0-5.0	100	Elevation (range): 250 (250-250) M
JACK PINE (<i>Pinus banksiana</i>)	10.0	10.0-10.0	100	Slope (%): 2.5 - 5.99 (1)
Tall Shrub (2 to 5m)				Aspect: Westerly (1)
BLACK SPRUCE (<i>Picea mariana</i>)	1.0	1.0-1.0	100	Topographic Position: Lower Slope (1)
BEAKED WILLOW (<i>Salix bebbiana</i>)	2.0	2.0-2.0	100	
ASPEN (<i>Populus tremuloides</i>)	10.0	10.0-10.0	100	Soil Variables
GREEN ALDER (<i>Alnus crispa</i>)	40.0	40.0-40.0	100	Soil Drainage: Moderately well drained (1)
Medium Shrub (0.5 to 2 m)				Soil Subgroup: EUTRIC BRUNISOL ELUVIATED (1)
BLACK SPRUCE (<i>Picea mariana</i>)	2.0	2.0-2.0	100	Surface Texture: Sand (1)
TWINFLOWER (<i>Linnaea borealis</i>)	3.0	3.0-3.0	100	Effective Texture: Sand (1)
Low Forb (< 30 cm)				Depth to Mottles/Gley:
NORTHERN BASTARD TOADFLAX (<i>Geocaulon lividum</i>)	1.0	1.0-1.0	100	Organic Thickness: 0 - 5 cm (1)
GROUND-PINE (<i>Lycopodium obscurum</i>)	1.0	1.0-1.0	100	Parent Material: Glaciofluvial (1)
ONE-SIDED WINTERGREEN (<i>Orthilia secunda</i>)	1.0	1.0-1.0	100	Humus Form
Moss				LFH Thickness
STAIR-STEP MOSS (<i>Hylocomium splendens</i>)	1.0	1.0-1.0	100	Mean
SCHREBER'S MOSS (<i>Pleurozium schreberi</i>)	1.0	1.0-1.0	100	Min
Lichen				Max
REINDEER LICHEN (<i>Cladina arbuscula</i>)	2.0	2.0-2.0	100	Count
				cm: 3.00 3.00 3.00 1

KUE6 Sb-Sw/Moss (n=2)

(*Picea mariana*-*Picea glauca*/Moss)

This community type represents the climax community type for the buffaloberry-alder ecological site in the Kazan Upland subregion. Pine, aspen and white birch dominated phases of this ecological site are seral to the white spruce dominated climax community. Black spruce once established may occupy a significant portion of the canopy in the climax community (Beckingham and Archibald 1996). On wetter sites in the absence of disturbance white and black spruce dominated communities are also found, but the understory species usually are dominated by a presence of horsetail species.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: e buffaloberry/alder(mesic/medium)

Ecosite Phase: e4 buffaloberry-alder/ Pj-Sw-Sb

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
WHITE SPRUCE (<i>Picea glauca</i>)	21.0	0.0-42.0	50	Moisture Regime: Subhygric (moderately moist) (2)
BLACK SPRUCE (<i>Picea mariana</i>)	31.5	0.0-63.0	50	Nutrient Regime: Submesotrophic (poor) (2)
Understory Tree				Elevation (range): 330 (330-330) M
WHITE SPRUCE (<i>Picea glauca</i>)	0.5	0.0-1.0	50	Slope (%): 2.5 - 5.99 (1)
BLACK SPRUCE (<i>Picea mariana</i>)	13.0	8.0-18.0	100	Aspect: Easterly (1)
Tall Shrub (2 to 5m)				Topographic Position: Level (1), Depression (1)
BLACK SPRUCE (<i>Picea mariana</i>)	18.0	18.0-18.0	100	Soil Variables
Medium Shrub (0.5 to 2 m)				Soil Drainage: Well drained (2)
PRICKLY ROSE (<i>Rosa acicularis</i>)	1.0	1.0-1.0	100	Soil Subgroup: DYSTRIC BRUNISOL ELUVIATED (2)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	1.5	1.0-2.0	100	Surface Texture: Sand (2)
Low Forb (< 30 cm)				Effective Texture: Sand (2)
NORTHERN BASTARD TOADFLAX (<i>Geocaulon lividum</i>)	6.0	4.0-8.0	100	Depth to Mottles/Gley:
Epiphyte				Organic Thickness: 0 - 5 cm (2)
OLD MAN'S BEARD (<i>Usnea hirta</i>)	1.0	1.0-1.0	100	Parent Material: Glaciolacustrine (2)
Moss				Humus Form
PIPECLEANER MOSS (<i>Rhytidium rugosum</i>)	4.0	0.0-8.0	50	LFH Thickness
WAVY DICRANUM (<i>Dicranum polysetum</i>)	10.0	2.0-18.0	100	Mean
STAIR-STEP MOSS (<i>Hylocomium splendens</i>)	13.0	8.0-18.0	100	Min
SCHREBER'S MOSS (<i>Pleurozium schreberi</i>)	75.5	63.0-88.0	100	Max
				Count
				cm:
				8.00
				6.00
				11.00
				2

f red osier dogwood/horsetail (subhygric/rich) (n=2)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

The red osier dogwood/horsetail ecosite is subhygric and nutrient rich. These sites are commonly found on level or depressional areas near the base of slopes or near watercourses where they receive nutrient-rich seepage or flood waters for a portion of the growing season. Upland tree dominated ecological sites in the Peace-Athabasca Delta subregion are often this ecological site. Fine-textured till and lacustrine parent materials are common. Horsetails commonly form a blanket over the forest floor.



Environmental Variables

Moisture Regime: Subhygric (moderately moist) (2)

Nutrient Regime: Permesotrophic (rich) (2)

Elevation (range): 217.5 (200-235) M

Slope (%): level (1), nearly level (1)

Aspect: Level (1), Westerly (1)

Topographic Position: Level (1), Upper Slope (1)

Soil Variables

Soil Drainage: Imperfectly drained (1), Moderately well drained (1)

Soil Subgroup: LUVIC GLEYSOL (1)

Surface Texture: Silty clay (1)

Effective Texture: Silt loam (1)

Depth to Mottles/Gley: 26 - 50 (1)

Organic Thickness: 0 - 5 cm (1)

Parent Material: Lacustrine (1)

Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	5.00	5.00	5.00	1

Successional Relationships

Succession on these rich sites proceeds slowly after disturbance due to the proliferation of grass, forb and shrub cover. This explosion of vegetational cover can make tree establishment (especially coniferous) difficult and can reduce early growth rates. Once white spruce becomes established, high growth rates can be expected.

Indicator Species

Tree

BALSAM POPLAR

Populus balsamifera

ASPEN

Populus tremuloides

Shrub

BEAKED WILLOW

Salix bebbiana

RIVER ALDER

Alnus tenuifolia

RED-OSIER DOGWOOD

Cornus stolonifera

Forb

MEADOW HORSETAIL

Equisetum pratense

f1 red osier dogwood/horsetail Sw (n=0)

Natural Subregion: Kazan Upland

Ecosite: f red osier dogwood/horsetail (subhygric/rich)

Ecosection: KU Kazan Upland

General Description

A number of ecological site phases currently have no data. These ecological site phases have been created as place holders because they were described in adjacent subregions (Athabasca Plain and Northern Mixedwood).

Characteristic Species

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

LFH Thickness

cm:

Mean

Min

Max

Count

0.00

0.00

0.00

0

f2 red osier dogwood/horsetail Pb-Sw (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: f red osier dogwood/horsetail (subhygric/rich)

General Description

A number of ecological site phases currently have no data. These ecological site phases have been created as place holders because they were described in adjacent subregions (Northern Mixedwood and Athabasca Plain).

Characteristic Species

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

LFH Thickness

cm:

Mean

Min

Max

Count

0.00

0.00

0.00

0

f3 red osier dogwood/horsetail - Pb-Aw(Ba) (n=2)

Natural Subregion: Kazan Upland

Ecosite: f red osier dogwood/horsetail (subhygric/rich)

Ecosection: KU Kazan Upland

Characteristic Species

Tree

- [32.0] BALSAM POPLAR*
Populus balsamifera
- [20.0] WHITE BIRCH
Betula papyrifera
- [7.5] ASPEN*
Populus tremuloides

Shrub

- [11.0] RIVER ALDER*
Alnus tenuifolia
- [10.5] RED-OSIER DOGWOOD*
Cornus stolonifera
- [7.5] BEAKED WILLOW*
Salix bebbiana
- [5.0] TWINFLOWER
Linnaea borealis
- [5.0] DEWBERRY
Rubus pubescens
- [4.0] BEAKED WILLOW
Salix bebbiana
- [3.5] PRICKLY ROSE
Rosa acicularis
- [2.0] LOW-BUSH CRANBERRY
Viburnum edule

Forb

- [5.0] COMMON PINK WINTERGREEN
Pyrola asarifolia
- [2.5] MEADOW HORSETAIL*
Equisetum pratense
- [1.5] COMMON FIREWEED
Epilobium angustifolium

Environmental Variables

Moisture Regime: Subhygric (moderately moist) (2)
 Nutrient Regime: Permesotrophic (rich) (2)
 Elevation (range): 217.5 (200-235) M
 Slope (%): level (1), nearly level (1)
 Aspect: Level (1), Westerly (1)
 Topographic Position: Level (1), Upper Slope (1)

Soil Variables

Soil Drainage: Imperfectly drained (1), Moderately well drained (1)
 Soil Subgroup: LUVIC GLEYSOL (1)
 Surface Texture: Silty clay (1)
 Effective Texture: Silt loam (1)
 Depth to Mottles/Gley: 26 - 50 (1)
 Organic Thickness: 0 - 5 cm (1)
 Parent Material: Lacustrine (1)
 Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	5.00	5.00	5.00	1

KUC5 Pb/Alder-Red osier dogwood/Horsetail (n=1)

(*Populus balsamifera*/*Alnus tenuifolia*-*Cornus stolonifera*/*Equisetum pratense*)

This plant community is transitional to the Peace-Athabasca subregion and is found in lower slope topographic positions near water courses where it receives nutrient-rich seepage or flood waters for a portion of the growing season. Undisturbed, this plant community tends to have dense shrub cover. Light is limited for lower herbaceous layers and succession in the absence of disturbance will be to white spruce.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: f red osier dogwood/horsetail (subhygric/rich)
Ecosite Phase: f3 red osier dogwood/horsetail - Pb-Aw(Ba)

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
BALSAM POPLAR (<i>Populus balsamifera</i>)	50.0	50.0-50.0	100	Moisture Regime: Subhygric (moderately moist) (1)
Understory Tree				Nutrient Regime: Permesotrophic (rich) (1)
BALSAM POPLAR (<i>Populus balsamifera</i>)	1.0	1.0-1.0	100	Elevation (range): 200 (200-200) M
Tall Shrub (2 to 5m)				Slope (%): 0 - 0.49 (1)
RIVER ALDER (<i>Alnus tenuifolia</i>)	20.0	20.0-20.0	100	Aspect: Level (1)
Medium Shrub (0.5 to 2 m)				Topographic Position: Level (1)
WILD RED RASPBERRY (<i>Rubus idaeus</i>)	1.0	1.0-1.0	100	Soil Variables
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	2.0	2.0-2.0	100	Soil Drainage: Moderately well drained (1)
PRICKLY ROSE (<i>Rosa acicularis</i>)	5.0	5.0-5.0	100	Soil Subgroup:
RED-OSIER DOGWOOD (<i>Cornus stolonifera</i>)	20.0	20.0-20.0	100	Surface Texture:
Low Shrub (< 0.5m)				Effective Texture:
DEWBERRY (<i>Rubus pubescens</i>)	10.0	10.0-10.0	100	Depth to Mottles/Gley:
Tall Forb (>= 30 cm)				Organic Thickness:
COMMON FIREWEED (<i>Epilobium angustifolium</i>)	1.0	1.0-1.0	100	Parent Material:
MEADOW HORSETAIL (<i>Equisetum pratense</i>)	5.0	5.0-5.0	100	Humus Form
Low Forb (< 30 cm)				LFH Thickness
COMMON PINK WINTERGREEN (<i>Pyrola asarifolia</i>)	10.0	10.0-10.0	100	Mean
Lichen				Min
N/A (<i>Melanelia albertana</i>)	1.0	1.0-1.0	100	Max
				Count
				cm:
				0.00
				0.00
				0.00
				0

KUC6 Bw-Aw-Pb/Willow (n=1)

(*Betula papyrifera*-*Populus tremuloides*-*Populus balsamifera*/*Salix spp.*)

This community type occupies moist rich seepage areas in lower slope positions. Succession in the absence of disturbance will likely be to White spruce.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: f red osier dogwood/horsetail (subhygric/rich)
Ecosite Phase: f3 red osier dogwood/horsetail - Pb-Aw(Ba)

Plant Composition	Canopy Cover (%)			cm	Environmental Variables
	Mean	Range	Const.		
Overstory Tree					
BALSAM POPLAR (<i>Populus balsamifera</i>)	8.0	8.0-8.0	100		Ecological Status Score: 25-25
ASPEN (<i>Populus tremuloides</i>)	10.0	10.0-10.0	100		Moisture Regime: Subhygric (moderately moist) (1)
WHITE BIRCH (<i>Betula papyrifera</i>)	40.0	40.0-40.0	100		Nutrient Regime: Permesotrophic (rich) (1)
Understory Tree					
BALSAM POPLAR (<i>Populus balsamifera</i>)	5.0	5.0-5.0	100		Elevation (range): 235 (235-235) M
ASPEN (<i>Populus tremuloides</i>)	5.0	5.0-5.0	100		Slope (%): 0.5 - 2.49 (1)
WHITE BIRCH (<i>Betula papyrifera</i>)	20.0	20.0-20.0	100		Aspect: Westerly (1)
Tall Shrub (2 to 5m)					
RIVER ALDER (<i>Alnus tenuifolia</i>)	2.0	2.0-2.0	100		Topographic Position: Upper Slope (1)
BEAKED WILLOW (<i>Salix bebbiana</i>)	15.0	15.0-15.0	100		
Medium Shrub (0.5 to 2 m)					
RED-OSIER DOGWOOD (<i>Cornus stolonifera</i>)	1.0	1.0-1.0	100		Soil Variables
TWINING HONEYSUCKLE (<i>Lonicera dioica</i>)	1.0	1.0-1.0	100		Soil Drainage: Imperfectly drained (1)
PRICKLY ROSE (<i>Rosa acicularis</i>)	2.0	2.0-2.0	100		Soil Subgroup: LUVIC GLEYSOL ORTHIC (1)
LOW-BUSH CRANBERRY (<i>Viburnum edule</i>)	2.0	2.0-2.0	100		Surface Texture: Silty clay (1)
BEAKED WILLOW (<i>Salix bebbiana</i>)	8.0	8.0-8.0	100		Effective Texture: Silt loam (1)
TWINFLOWER (<i>Linnaea borealis</i>)	10.0	10.0-10.0	100		Depth to Mottles/Gley: 26 - 50 (1)
Tall Forb (>= 30 cm)					
LINDLEY'S ASTER (<i>Aster ciliolatus</i>)	1.0	1.0-1.0	100		Organic Thickness: 0 - 5 cm (1)
COMMON FIREWEED (<i>Epilobium angustifolium</i>)	2.0	2.0-2.0	100		Parent Material: Lacustrine (1)
Low Forb (< 30 cm)					
BUNCHBERRY (<i>Cornus canadensis</i>)	2.0	2.0-2.0	100		Humus Form

LFH Thickness	Mean	Min	Max	Count
cm:	5.00	5.00	5.00	1

f4 shrub (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

A number of ecological site phases currently have no data. These ecological site phases have been created as place holders because they were described in adjacent subregions (Northern Mixedwood and Athabasca Plain).

Characteristic Species

Ecosite: f red osier dogwood/horsetail (subhygric/rich)

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

f5 tame/disturbed (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

A number of ecological site phases currently have no data. These ecological site phases have been created as place holders because they were described in adjacent subregions (Central Mixedwood).

Characteristic Species

Ecosite: f red osier dogwood/horsetail (subhygric/rich)

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	0.00	0.00	0.00	0

g Labrador tea-hygric(hygric/medium) (n=2)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

The Labrador tea-hygric ecosite has a nutrient poor substrate with imperfectly to very poorly drained soils. Labrador tea and bog cranberry are indicative of the relatively acidic surface soil conditions. The Labrador tea - hygric ecosite has plant community types similar to the Labrador tea - mesic ecosite the hygric ecosite tends to be in mid to lower slope positions, has distinct mottling within the top 50cm of soil and tends to be dominated by black spruce rather than pine.



Environmental Variables

Moisture Regime: Hygric (moist) (1), Subhygric (moderately wet) (1)
Nutrient Regime: Mesotrophic (medium) (1), Permesotrophic (rich) (1)
Elevation (range): 245 (240-250) M
Slope (%): level (1), nearly level (1)
Aspect: Level (2)
Topographic Position: Depression (1), Midslope (1)

Soil Variables

Soil Drainage: Imperfectly drained (1), Very poorly drained (1)
Soil Subgroup: GLEYSOL (1), MELANIC BRUNISOL (1)
Surface Texture: Sand (2)
Effective Texture: Sand (2)
Depth to Mottles/Gley: 26 - 50 (1)
Organic Thickness: 0 - 5 cm (2)
Parent Material: Glaciofluvial (1), Lacustrine (1)
Humus Form

LFH Thickness

	Mean	Min	Max	Count
cm:	6.50	3.00	10.00	2

Successional Relationships

Young and mature stands developing in this ecosite often have a component of black spruce. The black spruce is often the same age as the pine but forms a secondary canopy due to slower growth rates. Successionally mature stands are dominated by black spruce with a small component of old residual pine.

Indicator Species

Tree

BLACK SPRUCE
Picea mariana
TAMARACK
Larix laricina

Shrub

SMALL BOG CRANBERRY
Oxycoccus microcarpus
GREEN ALDER
Alnus crispa
RIVER ALDER
Alnus tenuifolia

Moss and Liverwort

TUFTED MOSS
Aulacomnium palustre
PEAT MOSS
Sphagnum warnstorffii
GOLDEN MOSS
Tomenthypnum nitens

Graminoid

WATER SEDGE
Carex aquatilis

g1 Labrador tea-hygric Sb-Pj (n=2)

Natural Subregion: Kazan Upland
 Ecoregion: KU Kazan Upland

Ecosite: g Labrador tea-hygric(hygric/medium)

Characteristic Species

Tree

- [32.5] BLACK SPRUCE*
Picea mariana
- [3.5] TAMARACK
Larix laricina

Shrub

- [10.0] RIVER ALDER*
Alnus tenuifolia
- [10.0] GREEN ALDER*
Alnus crispa
- [7.5] RIVER ALDER
Alnus tenuifolia
- [5.0] LEATHERLEAF
Chamaedaphne calyculata
- [2.5] BEAKED WILLOW
Salix bebbiana
- [2.5] MYRTLE-LEAVED WILLOW
Salix myrtillifolia
- [2.5] DWARF BIRCH
Betula pumila
- [2.5] BOG ROSEMARY
Andromeda polifolia
- [1.0] SMALL BOG CRANBERRY*
Oxycoccus microcarpus

Moss and Liverwort

- [25.0] PEAT MOSS*
Sphagnum warnstorffii
- [17.5] TUFTED MOSS*
Aulacomnium palustre
- [10.0] GOLDEN MOSS*
Tomenthypnum nitens
- [10.0] STAIR-STEP MOSS
Hylocomium splendens
- [2.5] JUNIPER HAIR-CAP
Polytrichum juniperinum

Graminoid

- [7.5] WATER SEDGE*
Carex aquatilis

Environmental Variables

Moisture Regime: Hygric (moist) (1), Subhydryc (moderately wet) (1)
 Nutrient Regime: Permesotrophic (rich) (1), Mesotrophic (medium) (1)
 Elevation (range): 245 (240-250) M
 Slope (%): level (1), nearly level (1)
 Aspect: Level (2)
 Topographic Position: Depression (1), Midslope (1)

Soil Variables

Soil Drainage: Imperfectly drained (1), Very poorly drained (1)
 Soil Subgroup: GLEYSOL (1), MELANIC BRUNISOL (1)
 Surface Texture: Sand (2)
 Effective Texture: Sand (2)
 Depth to Mottles/Gley: 26 - 50 (1)
 Organic Thickness: 0 - 5 cm (2)
 Parent Material: Lacustrine (1), Glaciofluvial (1)
 Humus Form

LFH Thickness	Mean	Min	Max	Count
cm:	6.50	3.00	10.00	2

KUE10 Lt-Sb/River alder/Sedge (n=1)

(*Larix laricina*-*Picea mariana*/*Alnus tenuifolia*/*Carex spp*)

Fen ecosites are not common in the Canadian Shield natural region. Most wetlands are situated in the bog or poor fen ecosites because of the acidic nature of the soils in the region. Micro-topography allows trees to grow on slightly higher ground and contributes to species richness. Increased flooding and prolonged water-logging may result in the disappearance of trees and a transition to a willow/sedge fen.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: g Labrador tea-hygric(hygric/medium)
Ecosite Phase: g1 Labrador tea-hygric Sb-Pj

Plant Composition	Canopy Cover (%)			cm	Environmental Variables
	Mean	Range	Const.		
Overstory Tree					Ecological Status Score: 25-25
TAMARACK (<i>Larix laricina</i>)	2.0	2.0-2.0	100		Moisture Regime: Subhydryc (moderately wet) (1)
Tall Shrub (2 to 5m)					Nutrient Regime: Mesotrophic (medium) (1)
TAMARACK (<i>Larix laricina</i>)	5.0	5.0-5.0	100		Elevation (range): 240 (240-240) M
BLACK SPRUCE (<i>Picea mariana</i>)	5.0	5.0-5.0	100		Slope (%): 0 - 0.49 (1)
RIVER ALDER (<i>Alnus tenuifolia</i>)	10.0	10.0-10.0	100		Aspect: Level (1)
Medium Shrub (0.5 to 2 m)					Topographic Position: Depression (1)
TAMARACK (<i>Larix laricina</i>)	2.0	2.0-2.0	100		Soil Variables
SMALL BOG CRANBERRY (<i>Oxycoccus microcarpus</i>)	2.0	2.0-2.0	100		Soil Drainage: Very poorly drained (1)
BOG ROSEMARY (<i>Andromeda polifolia</i>)	5.0	5.0-5.0	100		Soil Subgroup: GLEYSOL REGO (1)
DWARF BIRCH (<i>Betula pumila</i>)	5.0	5.0-5.0	100		Surface Texture: Sand (1)
MYRTLE-LEAVED WILLOW (<i>Salix myrtillifolia</i>)	5.0	5.0-5.0	100		Effective Texture: Sand (1)
LEATHERLEAF (<i>Chamaedaphne calyculata</i>)	10.0	10.0-10.0	100		Depth to Mottles/Gley:
RIVER ALDER (<i>Alnus tenuifolia</i>)	15.0	15.0-15.0	100		Organic Thickness: 0 - 5 cm (1)
Tall Forb (>= 30 cm)					Parent Material: Lacustrine (1)
MARSH CINQUEFOIL (<i>Potentilla palustris</i>)	2.0	2.0-2.0	100		Humus Form
Graminoid					
WATER SEDGE (<i>Carex aquatilis</i>)	15.0	15.0-15.0	100		
Moss					
TUFTED MOSS (<i>Aulacomnium palustre</i>)	5.0	5.0-5.0	100		
GOLDEN MOSS (<i>Tomenthypnum nitens</i>)	20.0	20.0-20.0	100		
PEAT MOSS (<i>Sphagnum warnstorffii</i>)	50.0	50.0-50.0	100		
					LFH Thickness
					Mean
					Min
					Max
					Count
				cm:	3.00
					3.00
					3.00
					1

KUE7 Sb/Alder/Tufted moss (n=1)

(*Picea mariana*/*Alnus crispa*/*Aulacomnium palustre*)

This community type represents the transition between a treed bog and the upland jack pine/black spruce dominated community types. Alder and stair step-moss indicate the medium nutrient status of this community type.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: g Labrador tea-hygric(hygric/medium)

Ecosite Phase: g1 Labrador tea-hygric Sb-Pj

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
WHITE SPRUCE (<i>Picea glauca</i>)	1.0	1.0-1.0	100	Moisture Regime: Hygric (moist) (1)
BLACK SPRUCE (<i>Picea mariana</i>)	2.0	2.0-2.0	100	Nutrient Regime: Permesotrophic (rich) (1)
JACK PINE (<i>Pinus banksiana</i>)	2.0	2.0-2.0	100	Elevation (range): 250 (250-250) M
Tall Shrub (2 to 5m)				Slope (%): 0.5 - 2.49 (1)
BEAKED WILLOW (<i>Salix bebbiana</i>)	5.0	5.0-5.0	100	Aspect: Level (1)
RIVER ALDER (<i>Alnus tenuifolia</i>)	10.0	10.0-10.0	100	Topographic Position: Midslope (1)
GREEN ALDER (<i>Alnus crispa</i>)	20.0	20.0-20.0	100	Soil Variables
BLACK SPRUCE (<i>Picea mariana</i>)	40.0	40.0-40.0	100	Soil Drainage: Imperfectly drained (1)
Medium Shrub (0.5 to 2 m)				Soil Subgroup: MELANIC BRUNISOL GLEYED (1)
BEAKED WILLOW (<i>Salix bebbiana</i>)	2.0	2.0-2.0	100	Surface Texture: Sand (1)
BLACK SPRUCE (<i>Picea mariana</i>)	20.0	20.0-20.0	100	Effective Texture: Sand (1)
Low Forb (< 30 cm)				Depth to Mottles/Gley: 26 - 50 (1)
BUNCHBERRY (<i>Cornus canadensis</i>)	1.0	1.0-1.0	100	Organic Thickness: 0 - 5 cm (1)
VINE-LEAVED COLTSFOOT (<i>Petasites vitifolius</i>)	1.0	1.0-1.0	100	Parent Material: Glaciofluvial (1)
THREE-LEAVED SOLOMON'S-SEAL (<i>Smilacina trifolia</i>)	2.0	2.0-2.0	100	Humus Form
Graminoid				LFH Thickness
NORTHERN REED GRASS (<i>Calamagrostis inexpansa</i>)	1.0	1.0-1.0	100	Mean
Moss				Min
JUNIPER HAIR-CAP (<i>Polytrichum juniperinum</i>)	5.0	5.0-5.0	100	Max
STAIR-STEP MOSS (<i>Hylocomium splendens</i>)	20.0	20.0-20.0	100	Count
TUFTED MOSS (<i>Aulacomnium palustre</i>)	30.0	30.0-30.0	100	cm:
Lichen				10.00
N/A (<i>Cladonia gracilis</i>)	3.0	3.0-3.0	100	10.00
				10.00
				1

h bog (subhydric/very poor) (n=3)

Natural Subregion: Kazan Upland

General Description

Bog communities are the dominant wetland type in the Kazan Upland subregion. The bog ecosite has a very poor to poor nutrient regime and is poor to very poorly drained. Organic soils consisting of slowly decomposing peat moss are common. This ecosite occupies level and depressional areas where water tends to be stagnant and impeded drainage or high water tables enhance the accumulation of organic matter. Stunted black spruce forms a sparse canopy on the treed phase (h1) of the bog ecosite. Leatherleaf (*Chamaedaphne calyculata*) is very common in the bog ecosite.



Ecosection: KU Kazan Upland

Environmental Variables

Moisture Regime: Subhydric (moderately wet) (3)

Nutrient Regime: Oligotrophic (very poor) (1), Submesotrophic (poor) (2)

Elevation (range): 267.5 (230-300) M

Slope (%): level (1), very gentle slope (1)

Aspect: Level (1), Northerly (1)

Topographic Position: Lower Slope (1), Depression (2)

Soil Variables

Soil Drainage: Poorly drained (1), Very poorly drained (2)

Soil Subgroup: FIBRISOL (1), GLEYSOL (1), ORGANIC CRYOSOL (1)

Surface Texture: Fibric (2)

Effective Texture: Mesic (1), Sand (1)

Depth to Mottles/Gley:

Organic Thickness: 26 - 39 cm (1), 60 - 79 cm (1)

Parent Material: Glaciofluvial (1), Undifferentiated Organic (2)

Humus Form

Successional Relationships

The bog ecosite is an edaphic climax that is maintained by water tables. The hydrarch succession that leads to the bog ecosite is extremely slow.

Indicator Species

Tree

BLACK SPRUCE

Picea mariana

Shrub

COMMON LABRADOR TEA

Ledum groenlandicum

SMALL BOG CRANBERRY

Oxycoccus microcarpus

LEATHERLEAF

Chamaedaphne calyculata

NORTHERN LAUREL

Kalmia polifolia

Moss and Liverwort

N/A

Sphagnum nemoreum

PEAT MOSS

Sphagnum angustifolium

h1 bog - treed (n=1)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: h bog (subhydric/very poor)

Characteristic Species

Tree

- [99.0] BLACK SPRUCE
Picea mariana

Shrub

- [42.0] COMMON LABRADOR TEA*
Ledum groenlandicum
- [29.0] CLOUDBERRY
Rubus chamaemorus
- [8.0] LEATHERLEAF*
Chamaedaphne calyculata
- [8.0] BOG CRANBERRY
Vaccinium vitis-idaea

Moss and Liverwort

- [42.0] PEAT MOSS*
Sphagnum angustifolium
- [42.0] N/A*
Sphagnum nemoreum
- [18.0] SCHREBER'S MOSS
Pleurozium schreberi

Environmental Variables

Moisture Regime: Subhydric (moderately wet) (1)

Nutrient Regime: Oligotrophic (very poor) (1)

Elevation (range): 300 (300-300) M

Slope (%):

Aspect:

Topographic Position: Depression (1)

Soil Variables

Soil Drainage: Poorly drained (1)

Soil Subgroup: ORGANIC CRYOSOL (1)

Surface Texture: Fibric (1)

Effective Texture: Mesic (1)

Depth to Mottles/Gley:

Organic Thickness: 60 - 79 cm (1)

Parent Material:

Humus Form

KUE8 Sb/Labrador tea/peat moss (n=1)

(*Picea mariana*/*Ledum groenlandicum*/*Sphagnum spp.*)

This community type is typical if a treed bog ecological site phase. Rich fens tend to be dominated by larch, willow and golden moss, whereas, bogs are dominated by black spruce, Labrador tea and peat moss.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: h bog (subhydric/very poor)

Ecosite Phase: h1 bog - treed

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
BLACK SPRUCE (<i>Picea mariana</i>)	29.0	29.0-29.0	100	Moisture Regime: Subhydric (moderately wet) (1)
Understory Tree				Nutrient Regime: Oligotrophic (very poor) (1)
BLACK SPRUCE (<i>Picea mariana</i>)	42.0	42.0-42.0	100	Elevation (range): 300 (300-300) M
Tall Shrub (2 to 5m)				Slope (%):
BLACK SPRUCE (<i>Picea mariana</i>)	29.0	29.0-29.0	100	Aspect:
Medium Shrub (0.5 to 2 m)				Topographic Position: Depression (1)
LEATHERLEAF (<i>Chamaedaphne calyculata</i>)	8.0	8.0-8.0	100	Soil Variables
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	8.0	8.0-8.0	100	Soil Drainage: Poorly drained (1)
COMMON LABRADOR TEA (<i>Ledum groenlandicum</i>)	42.0	42.0-42.0	100	Soil Subgroup: ORGANIC CRYOSOL FIBRIC (1)
Low Shrub (< 0.5m)				Surface Texture: Fibric (1)
CLODBERRY (<i>Rubus chamaemorus</i>)	29.0	29.0-29.0	100	Effective Texture: Mesic (1)
Moss				Depth to Mottles/Gley:
SCHREBER'S MOSS (<i>Pleurozium schreberi</i>)	18.0	18.0-18.0	100	Organic Thickness: 60 - 79 cm (1)
N/A (<i>Sphagnum nemoreum</i>)	42.0	42.0-42.0	100	Parent Material:
PEAT MOSS (<i>Sphagnum angustifolium</i>)	42.0	42.0-42.0	100	Humus Form

h2 bog - shrubby (n=2)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: h bog (subhydric/very poor)

Characteristic Species

Tree

- [15.0] BLACK SPRUCE
Picea mariana

Shrub

- [40.0] LEATHERLEAF
Chamaedaphne calyculata
- [35.5] COMMON LABRADOR TEA
Ledum groenlandicum
- [10.0] NORTHERN LAUREL*
Kalmia polifolia
- [7.5] BOG CRANBERRY
Vaccinium vitis-idaea
- [5.0] SMALL BOG CRANBERRY*
Oxycoccus microcarpus
- [2.0] CLOUDBERRY
Rubus chamaemorus

Forb

- [4.0] THREE-LEAVED SOLOMON'S-SEAL
Smilacina trifolia

Lichen

- [12.5] REINDEER LICHEN
Cladina mitis

Moss and Liverwort

- [65.0] RUSTY PEAT MOSS
Sphagnum fuscum
- [17.5] MIDWAY PEAT MOSS
Sphagnum magellanicum
- [5.0] N/A
Sphagnum nemoreum
- [2.0] TUFTED MOSS
Aulacomnium palustre

Graminoid

- [4.0] SHEATHED COTTON GRASS
Eriophorum vaginatum
- [3.0] WATER SEDGE
Carex aquatilis

Environmental Variables

Moisture Regime: Subhydric (moderately wet) (2)

Nutrient Regime: Submesotrophic (poor) (2)

Elevation (range): 235 (230-240) M

Slope (%): level (1), very gentle slope (1)

Aspect: Level (1), Northerly (1)

Topographic Position: Depression (1), Lower Slope (1)

Soil Variables

Soil Drainage: Very poorly drained (2)

Soil Subgroup: FIBRISOL (1), GLEYSOL (1)

Surface Texture: Fibric (1)

Effective Texture: Sand (1)

Depth to Mottles/Gley:

Organic Thickness: 26 - 39 cm (1)

Parent Material: Glaciofluvial (1), Undifferentiated Organic (2)

Humus Form

KUB2 Labrador tea/Peat moss (Sb) (n=2)

(*Ledum groelandicum*/*Sphagnum* spp (*Picea mariana*))

This plant community represents an early seral community of the treed bog ecosite phase. The bog ecosite commonly has organic soils consisting of slowly decomposing peat moss. In the absence of disturbance this community type will become dominated by black spruce. The soils for one plot in this community type was described as peaty phase of a Rego Gleysol as the soil was described at the outer edge of the bog.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: h bog (subhydric/very poor)
Ecosite Phase: h2 bog - shrubby

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Understory Tree				Ecological Status Score: 25-25
BLACK SPRUCE (<i>Picea mariana</i>)	1.0	0.0-2.0	50	Moisture Regime: Subhydric (moderately wet) (2)
Tall Shrub (2 to 5m)				Nutrient Regime: Submesotrophic (poor) (2)
BLACK SPRUCE (<i>Picea mariana</i>)	5.0	5.0-5.0	100	Elevation (range): 235 (230-240) M
Medium Shrub (0.5 to 2 m)				Slope (%): 0 - 0.49 (1), 2.5 - 5.99 (1)
WHITE BIRCH (<i>Betula papyrifera</i>)	1.5	0.0-3.0	50	Aspect: Level (1), Northerly (1)
SMALL BOG CRANBERRY (<i>Oxycoccus microcarpus</i>)	5.0	5.0-5.0	100	Topographic Position: Lower Slope (1), Depression (1)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	7.5	0.0-15.0	50	Soil Variables
NORTHERN LAUREL (<i>Kalmia polifolia</i>)	10.0	0.0-20.0	50	Soil Drainage: Very poorly drained (2)
BLACK SPRUCE (<i>Picea mariana</i>)	15.0	10.0-20.0	100	Soil Subgroup: FIBRISOL TYPIC (1), GLEYSOL REGO (1)
COMMON LABRADOR TEA (<i>Ledum groenlandicum</i>)	35.5	1.0-70.0	100	Surface Texture: Fibric (1)
LEATHERLEAF (<i>Chamaedaphne calyculata</i>)	40.0	10.0-70.0	100	Effective Texture: Sand (1)
Low Shrub (< 0.5m)				Depth to Mottles/Gley:
CLOUDBERRY (<i>Rubus chamaemorus</i>)	2.0	2.0-2.0	100	Organic Thickness: 26 - 39 cm (1)
Low Forb (< 30 cm)				Parent Material: Glaciofluvial (1), Undifferentiated Organic (2)
THREE-LEAVED SOLOMON'S-SEAL (<i>Smilacina trifolia</i>)	4.0	3.0-5.0	100	Humus Form
Graminoid				
WATER SEDGE (<i>Carex aquatilis</i>)	3.0	3.0-3.0	100	
SHEATHED COTTON GRASS (<i>Eriophorum vaginatum</i>)	4.0	0.0-8.0	50	
Moss				
JUNIPER HAIR-CAP (<i>Polytrichum juniperinum</i>)	1.0	0.0-2.0	50	
TUFTED MOSS (<i>Aulacomnium palustre</i>)	2.0	1.0-3.0	100	
N/A (<i>Sphagnum nemoreum</i>)	5.0	0.0-10.0	50	
MIDWAY PEAT MOSS (<i>Sphagnum magellanicum</i>)	17.5	0.0-35.0	50	
RUSTY PEAT MOSS (<i>Sphagnum fuscum</i>)	65.0	60.0-70.0	100	
Lichen				
REINDEER LICHEN (<i>Cladina mitis</i>)	12.5	0.0-25.0	50	

h3 bog -graminoid (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: h bog (subhydric/very poor)

General Description

This ecological site phase currently has no data and graminoid bogs have not been described in the Boreal Natural Region. This ecological site phase was created for photo interpretation of vegetation for Alberta Vegetation Inventory (AVI) and Primary Land Vegetation Inventory (PLVI) if the interpreters call for a mapcode of 9B (Hydric/very poor). If interpreted in the various vegetation land cover inventories it will provide a spatial location in which to sample.

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Characteristic Species

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

i poor fen (subhydric/medium) (n=3)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

The poor fen ecosite is intermediate in nutrient regime between the bog (h) and the rich fen (j) ecosites and as such has species characteristic of both. Drainage is poor to very poor, however, there is some movement of water through the substratum. This ecosite occupies level and depressional areas where impeded drainage or high water tables enhance the accumulation of organic matter. This organic matter consists of a combination of bog-type organic matter (peat moss) and rich fen-type organic matter (sedges, golden moss, tufted moss, and brown moss). Both the black spruce and/or tamarack that dominate a sparse canopy on the treed phase (i1) of the poor fen ecosite are stunted and generally considered unmerchantable. A number of saline influenced plant community types were described here (KUA4, KUA5). In the Northern Mixedwood subregion a saline lowland ecological site was described, but the saline ecological sites in the Kazan Upland are presently not adequately described that warrants a new ecological site.



Environmental Variables

Moisture Regime: Hygric (moist) (1), Subhygric (moderately moist) (1)
Nutrient Regime: Mesotrophic (medium) (1), Submesotrophic (poor) (1)
Elevation (range): 230 (205-250) M
Slope (%): level (3)
Aspect: Level (2)
Topographic Position: Lower Slope (1), Depression (2)

Soil Variables

Soil Drainage: Imperfectly drained (0), Poorly drained (2)
Soil Subgroup: FIBRISOL (1)
Surface Texture: Fibric (1)
Effective Texture: Fibric (1)
Depth to Mottles/Gley:
Organic Thickness: >= 80 cm (1)
Parent Material: Glaciolacustrine (0), Lacustrine (1), Undifferentiated Organic (1)
Humus Form

Successional Relationships

Succession within this ecosite occurs over periods of hundreds to thousands of years. Recovery from disturbance is extremely slow. Changing hydrologic regimes that can result from disturbance influence the direction and rate of succession. These systems depend on water flow through them, impeding this flow can result in reduction or elimination of tree cover and changes in the shrub, forb and grass layers (Beckingham and Archibald 1996).

Indicator Species

Tree

BLACK SPRUCE

Picea mariana

TAMARACK

Larix laricina

Shrub

COMMON LABRADOR TEA

Ledum groenlandicum

MYRTLE-LEAVED WILLOW

Salix myrtillofolia

RIVER ALDER

Alnus tenuifolia

Forb

SALINE SHOOTING STAR

Dodecatheon pulchellum

SEASIDE ARROW-GRASS

Triglochin maritima

Moss and Liverwort

BROWN MOSS

Drepanocladus exannulatus

GOLDEN MOSS

Tomenthypnum nitens

i1 poor fen - treed (n=1)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: i poor fen (subhydric/medium)

Characteristic Species

Tree

- [53.0] BLACK SPRUCE*
Picea mariana
- [18.0] TAMARACK*
Larix laricina

Shrub

- [30.0] COMMON LABRADOR TEA*
Ledum groenlandicum
- [6.0] RIVER ALDER*
Alnus tenuifolia
- [6.0] MYRTLE-LEAVED WILLOW*
Salix myrtillifolia
- [3.0] BEAKED WILLOW
Salix bebbiana
- [3.0] BOG CRANBERRY
Vaccinium vitis-idaea
- [2.0] BEAKED WILLOW
Salix bebbiana

Forb

- [2.0] SWAMP HORSETAIL
Equisetum fluviatile
- [1.0] COMMON HORSETAIL
Equisetum arvense

Moss and Liverwort

- [15.0] GOLDEN MOSS*
Tomenthypnum nitens

Graminoid

- [2.0] BLUEJOINT
Calamagrostis canadensis

Environmental Variables

Moisture Regime: Hygric (moist) (1)

Nutrient Regime: Submesotrophic (poor) (1)

Elevation (range): 205 (205-205) M

Slope (%): level (1)

Aspect: Level (1)

Topographic Position: Lower Slope (1)

Soil Variables

Soil Drainage: Poorly drained (1)

Soil Subgroup: FIBRISOL (1)

Surface Texture: Fibric (1)

Effective Texture: Fibric (1)

Depth to Mottles/Gley:

Organic Thickness: >= 80 cm (1)

Parent Material: Undifferentiated Organic (1), Lacustrine (1)

Humus Form

KUE9 Sb-Lt/Labrador tea/Golden moss (n=1)

(*Picea mariana-Larix laricina/Ledum groenlandicum/Tomenthypnum nitens*)

This community type is transitional between the rich fen and the bog ecological sites. Rich fens tend to be dominated by larch, willow and golden moss, whereas, bogs are dominated by black spruce, Labrador tea and peat moss.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: i poor fen (subhydric/medium)

Ecosite Phase: i1 poor fen - treed

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 25-25
TAMARACK (<i>Larix laricina</i>)	10.0	10.0-10.0	100	Moisture Regime: Hygric (moist) (1)
BLACK SPRUCE (<i>Picea mariana</i>)	30.0	30.0-30.0	100	Nutrient Regime: Submesotrophic (poor) (1)
Understory Tree				Elevation (range): 205 (205-205) M
TAMARACK (<i>Larix laricina</i>)	5.0	5.0-5.0	100	Slope (%): 0 - 0.49 (1)
BLACK SPRUCE (<i>Picea mariana</i>)	15.0	15.0-15.0	100	Aspect: Level (1)
Tall Shrub (2 to 5m)				Topographic Position: Lower Slope (1)
BEAKED WILLOW (<i>Salix bebbiana</i>)	3.0	3.0-3.0	100	Soil Variables
BLACK SPRUCE (<i>Picea mariana</i>)	5.0	5.0-5.0	100	Soil Drainage: Poorly drained (1)
RIVER ALDER (<i>Alnus tenuifolia</i>)	6.0	6.0-6.0	100	Soil Subgroup: FIBRISOL TERRIC (1)
Medium Shrub (0.5 to 2 m)				Surface Texture: Fibric (1)
BEAKED WILLOW (<i>Salix bebbiana</i>)	2.0	2.0-2.0	100	Effective Texture: Fibric (1)
TAMARACK (<i>Larix laricina</i>)	3.0	3.0-3.0	100	Depth to Mottles/Gley:
BLACK SPRUCE (<i>Picea mariana</i>)	3.0	3.0-3.0	100	Organic Thickness: >= 80 cm (1)
BOG CRANBERRY (<i>Vaccinium vitis-idaea</i>)	3.0	3.0-3.0	100	Parent Material: Lacustrine (1), Undifferentiated Organic (1)
RIVER ALDER (<i>Alnus tenuifolia</i>)	5.0	5.0-5.0	100	Humus Form
MYRTLE-LEAVED WILLOW (<i>Salix myrtillifolia</i>)	6.0	6.0-6.0	100	
COMMON LABRADOR TEA (<i>Ledum groenlandicum</i>)	30.0	30.0-30.0	100	
Tall Forb (>= 30 cm)				
COMMON HORSETAIL (<i>Equisetum arvense</i>)	1.0	1.0-1.0	100	
SWAMP HORSETAIL (<i>Equisetum fluviatile</i>)	2.0	2.0-2.0	100	
Graminoid				
BLUEJOINT (<i>Calamagrostis canadensis</i>)	2.0	2.0-2.0	100	
Moss				
GOLDEN MOSS (<i>Tomenthypnum nitens</i>)	15.0	15.0-15.0	100	

i2 poor fen - shrubby (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

This ecological site phase currently has no data, but is presented here because it has been described in adjacent subregions (Northern Mixedwood and Athabasca Plain).

Characteristic Species

Ecosite: i poor fen (subhydric/medium)

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

i3 poor fen - graminoid (n=2)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: i poor fen (subhydic/medium)

Characteristic Species

Forb

- [20.0] SALINE SHOOTING STAR*
Dodecatheon pulchellum
- [2.5] SEASIDE ARROW-GRASS*
Triglochin maritima
- [1.5] SEASIDE ARROW-GRASS
Triglochin maritima

Moss and Liverwort

- [27.5] BROWN MOSS*
Drepanocladus exannulatus

Graminoid

- [20.0] MAT MUHLY
Muhlenbergia richardsonis
- [4.0] AWNED SEDGE
Carex atherodes
- [1.0] WIRE RUSH
Juncus balticus

Environmental Variables

Moisture Regime: Hygric (moist) (0), Subhygric (moderately moist) (1)

Nutrient Regime: Mesotrophic (medium) (1)

Elevation (range): 242.5 (235-250) M

Slope (%): level (2)

Aspect: Level (1)

Topographic Position: Depression (2)

Soil Variables

Soil Drainage: Imperfectly drained (0), Poorly drained (1)

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material: Glaciolacustrine (0)

Humus Form

KUA4 Shooting star/Mat muhly (saline) (n=1)

(*Dodecatheon pulchellum*/*Muhlenbergia richardsonis*)

This community type is part of a saline meadow complex described within the La Butte Creek Wildland Provincial Park (Allen et al. 2002). Site 1 which is described was slightly drier and was dominated by mat muhly, shooting star and seaside arrow grass. The other site which was slightly wetter was dominated by brown moss, awned sedge and seaside arrowgrass on the drier margins. It was felt that the mat muhly dominated community was subjected to higher levels of salt through evaporation.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: i poor fen (subhydic/medium)

Ecosite Phase: i3 poor fen - graminoid

Plant Composition

Canopy Cover (%)

Environmental Variables

	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Medium Shrub (0.5 to 2 m)				Ecological Status Score: 0-0
HOARY WILLOW (<i>Salix candida</i>)	1.0	1.0-1.0	100	Moisture Regime: Subhydic (moderately moist) (1)
Tall Forb (>= 30 cm)				Nutrient Regime: Mesotrophic (medium) (1)
HIRSUTE FLEABANE (<i>Erigeron lonchophyllus</i>)	1.0	1.0-1.0	100	Elevation (range): 250 (250-250) M
SEASIDE ARROW-GRASS (<i>Triglochin maritima</i>)	5.0	5.0-5.0	100	Slope (%): 0 - 0.49 (1)
SALINE SHOOTING STAR (<i>Dodecatheon pulchellum</i>)	40.0	40.0-40.0	100	Aspect: Level (1)
Low Forb (< 30 cm)				Topographic Position: Depression (1)
SEA MILKWORT (<i>Glaux maritima</i>)	1.0	1.0-1.0	100	Soil Variables
Graminoid				Soil Drainage: Poorly drained (1)
SLENDER WHEAT GRASS (<i>Agropyron trachycaulum</i>)	1.0	1.0-1.0	100	Soil Subgroup:
MAT MUHLY (<i>Muhlenbergia richardsonis</i>)	40.0	40.0-40.0	100	Surface Texture:
				Effective Texture:
				Depth to Mottles/Gley:
				Organic Thickness:
				Parent Material: Glaciolacustrine (0)
				Humus Form

KUA5 Sedge/Brown moss (saline) (n=1)

(Carex atherodes/Drepanocladus exannulata)

This community represents the wetter portion of a saline meadow complex within the La Butte Creek Wildland Provincial Park (Allen et al. 2002). Site 1 which is slightly drier and was dominated by mat muhly, shooting star and seaside arrow grass. The other site described here was slightly wetter and was dominated by brown moss, awned sedge and seaside arrowgrass on the drier margins. Towards the areas of open water this community type integrated into wetland communities dominated by water and awned sedge.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: i poor fen (subhydric/medium)

Ecosite Phase: i3 poor fen - graminoid

Plant Composition

Canopy Cover (%)

Environmental Variables

	Canopy Cover (%)			Ecological Status Score: 0-0
	Mean	Range	Const.	
Low Forb (< 30 cm)				Moisture Regime: Hygric (moist) (0)
SEASIDE ARROW-GRASS <i>(Triglochin maritima)</i>	3.0	0.0-0.0	100	Nutrient Regime: Mesotrophic (medium) (0)
Graminoid				Elevation (range): 235 (235-235) M
WIRE RUSH <i>(Juncus balticus)</i>	2.0	0.0-0.0	100	Slope (%): 0 - 0.49 (1)
AWNED SEDGE <i>(Carex atherodes)</i>	8.0	0.0-0.0	100	Aspect: Level (0)
Moss				Topographic Position: Depression (1)
BROWN MOSS <i>(Drepanocladus exannulatus)</i>	55.0	0.0-0.0	0	Soil Variables

Soil Drainage: Imperfectly drained (0)

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material: Glaciolacustrine (0)

Humus Form

j rich fen (subhydric/rich) (n=3)

Natural Subregion: Kazan Upland

General Description

The rich fen ecosite is characterized by flowing water and alkaline nutrient-rich conditions. The soil is composed of organic matter derived from decomposing sedges, as well as golden, tufted, and brown mosses. This ecosite occupies level and depressional areas where the water table is at or near the surface for a portion of the growing season. Tamarack dominates the canopy on the treed phase while dwarf birch or willow form the canopy of the shrubby phase, and sedges dominate the graminoid phase of the rich fen ecosite. A number of sites within this ecological site were described with mineral soils. According to the Alberta Wetland Classification system (2015) mineral soils can only be swamps or marshes, however because of the species composition and moisture regime these willow and sedge dominated communities for this classification were placed within the rich fen ecological site.



Ecosection: KU Kazan Upland

Environmental Variables

Moisture Regime: Hydric (wet) (1), Subhydric (moderately wet) (1), Subhygric (moderately moist) (1)

Nutrient Regime: Mesotrophic (medium) (1), Permesotrophic (rich) (2)

Elevation (range): 264.5 (229-300) M

Slope (%): nearly level (1), level (2)

Aspect: Level (2)

Topographic Position: Depression (1), Level (2)

Soil Variables

Soil Drainage: Poorly drained (1), Very poorly drained (2)

Soil Subgroup: GLEYSOL (1), HUMIC GLEYSOL (1), MESISOL (1)

Surface Texture: Mesic (1), Sand (1), Silt (1)

Effective Texture: Loam (1), Sandy clay (1), Silt (1)

Depth to Mottles/Gley: 26 - 50 (1)

Organic Thickness: 26 - 39 cm (1), 0 - 5 cm (2)

Parent Material: Lacustrine (1), Rock (1), Undifferentiated Organic (1), Fluvial (2)

Humus Form

Successional Relationships

The rich fen is an early stage in hydrarch succession. Species composition, and direction and rate of succession changes with the changing hydrologic regime. As with other wetlands, rich fens have slow successional rates, so recovery from disturbance may also be slow.

Indicator Species

Shrub

FLAT-LEAVED WILLOW

Salix planifolia

BALSAM WILLOW

Salix pyrifolia

Moss and Liverwort

BROWN MOSS

Drepanocladus aduncus

Graminoid

BLUEJOINT

Calamagrostis canadensis

SMALL BOTTLE SEDGE

Carex utriculata

j1 rich fen - treed (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

General Description

This ecological site phase currently has no data, but is presented here because it has been described in adjacent subregions (Northern Mixedwood and Athabasca Plain).

Characteristic Species

Ecosite: j rich fen (subhydric/rich)

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

j2 rich fen - shrubby (n=3)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: j rich fen (subhydic/rich)

Characteristic Species

Tree

- [5.0] WHITE BIRCH
Betula papyrifera
- [3.3] ASPEN
Populus tremuloides

Shrub

- [30.0] BALSAM WILLOW*
Salix pyrifolia
- [31.6] FLAT-LEAVED WILLOW
Salix planifolia
- [3.2] SMOOTH WILLOW
Salix glauca

Moss and Liverwort

- [16.6] BROWN MOSS*
Drepanocladus aduncus

Graminoid

- [46.6] BLUEJOINT*
Calamagrostis canadensis
- [16.6] SMALL BOTTLE SEDGE*
Carex utriculata

Environmental Variables

Moisture Regime: Subhygric (moderately moist) (1), Hydric (wet) (1), Subhydic (moderately wet) (1)

Nutrient Regime: Mesotrophic (medium) (1), Permesotrophic (rich) (2)

Elevation (range): 264.5 (229-300) M

Slope (%): nearly level (1), level (2)

Aspect: Level (2)

Topographic Position: Depression (1), Level (2)

Soil Variables

Soil Drainage: Poorly drained (1), Very poorly drained (2)

Soil Subgroup: GLEYSOL (1), HUMIC GLEYSOL (1), MESISOL (1)

Surface Texture: Sand (1), Silt (1), Mesic (1)

Effective Texture: Silt (1), Sandy clay (1), Loam (1)

Depth to Mottles/Gley: 26 - 50 (1)

Organic Thickness: 26 - 39 cm (1), 0 - 5 cm (2)

Parent Material: Undifferentiated Organic (1), Rock (1), Lacustrine (1), Fluvial (2)

Humus Form

KUB3 Willow/Marsh reed grass (n=1)

(*Salix spp/Calamagrostis canadensis*)

This plant community is found along the edges of marsh reed grass and sedge fen meadows and in moist depressions. Willow will invade onto these graminoid fens to form the Willow/Marsh reed grass community type. Increased flooding and prolonged waterlogging may result in the disappearance of willow and a transition to a graminoid fen.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: j rich fen (subhydryc/rich)

Ecosite Phase: j2 rich fen - shrubby

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Understory Tree				Ecological Status Score: 40-40
SMOOTH WILLOW (<i>Salix glauca</i>)	5.0	5.0-5.0	100	Moisture Regime: Subhydryc (moderately moist) (1)
Medium Shrub (0.5 to 2 m)				Nutrient Regime: Permesotrophic (rich) (1)
SMOOTH WILLOW (<i>Salix glauca</i>)	5.0	5.0-5.0	100	Elevation (range): 0 (0-0) M
Tall Forb (>= 30 cm)				Slope (%): 0 - 0.49 (1)
COMMON FIREWEED (<i>Epilobium angustifolium</i>)	1.0	1.0-1.0	100	Aspect: Level (0)
Graminoid				Topographic Position: Level (1)
BLUEJOINT (<i>Calamagrostis canadensis</i>)	95.0	95.0-95.0	100	Soil Variables
				Soil Drainage: Poorly drained (1)
				Soil Subgroup: HUMIC GLEYSOL REGO (1)
				Surface Texture: Silt (1)
				Effective Texture: Silt (1)
				Depth to Mottles/Gley:
				Organic Thickness: 0 - 5 cm (1)
				Parent Material: Fluvial (1)
				Humus Form

KUB4 Willow/Sedge (n=1)

(*Salix planifolia*/*Carex utriculata*)

This community type is found along the edges of sedge fens (meadows) and in moist depressions. Willow becomes established at the edges of the sedge fens due to the shorter duration of standing water. Increased flooding and prolonged water-logging may result in the disappearance of willow and a transition to a sedge fen.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: j rich fen (subhydric/rich)

Ecosite Phase: j2 rich fen - shrubby

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Tall Shrub (2 to 5m)				Ecological Status Score: 40-40
FLAT-LEAVED WILLOW (<i>Salix planifolia</i>)	30.0	30.0-30.0	100	Moisture Regime: Hydric (wet) (1)
Medium Shrub (0.5 to 2 m)				Nutrient Regime: Permesotrophic (rich) (1)
BASKET WILLOW (<i>Salix petiolaris</i>)	2.0	2.0-2.0	100	Elevation (range): 229 (229-229) M
FLAT-LEAVED WILLOW (<i>Salix planifolia</i>)	65.0	65.0-65.0	100	Slope (%): 0 - 0.49 (1)
Graminoid				Aspect: Level (1)
BLUEJOINT (<i>Calamagrostis canadensis</i>)	40.0	40.0-40.0	100	Topographic Position: Level (1)
SMALL BOTTLE SEDGE (<i>Carex utriculata</i>)	50.0	50.0-50.0	100	Soil Variables
				Soil Drainage: Very poorly drained (1)
				Soil Subgroup: GLEYSOL REGO (1)
				Surface Texture: Sand (1)
				Effective Texture: Sandy clay (1)
				Depth to Mottles/Gley: 26 - 50 (1)
				Organic Thickness: 0 - 5 cm (1)
				Parent Material: Fluvial (1)
				Humus Form

KUC8 Willow/Brown moss/ Aw(Bw) (n=1)

(*Salix pyrifolia*/*Drepanocladus aduncus*)

This community type is found along the edges of sedge fens (meadows) and in moist depressions. Willow becomes established at the edges of the sedge fens due to the shorter duration of standing water. Increased flooding and prolonged water-logging may result in the disappearance of willow and a transition to a sedge fen. If the water table continues to decline the site will eventually become dominated by Aspen and birch.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: j rich fen (subhydric/rich)
Ecosite Phase: j2 rich fen - shrubby

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Overstory Tree				Ecological Status Score: 40-40
ASPEN (<i>Populus tremuloides</i>)	10.0	10.0-10.0	100	Moisture Regime: Subhydric (moderately wet) (1)
Understory Tree				Nutrient Regime: Mesotrophic (medium) (1)
BALSAM WILLOW (<i>Salix pyrifolia</i>)	1.0	1.0-1.0	100	Elevation (range): 300 (300-300) M
WHITE BIRCH (<i>Betula papyrifera</i>)	15.0	15.0-15.0	100	Slope (%): 0.5 - 2.49 (1)
Tall Shrub (2 to 5m)				Aspect: Level (1)
WHITE SPRUCE (<i>Picea glauca</i>)	1.0	1.0-1.0	100	Topographic Position: Depression (1)
BALSAM WILLOW (<i>Salix pyrifolia</i>)	90.0	90.0-90.0	100	
Medium Shrub (0.5 to 2 m)				Soil Variables
JACK PINE (<i>Pinus banksiana</i>)	1.0	1.0-1.0	100	Soil Drainage: Very poorly drained (1)
PRICKLY ROSE (<i>Rosa acicularis</i>)	1.0	1.0-1.0	100	Soil Subgroup: MESISOL TERRIC (1)
Low Shrub (< 0.5m)				Surface Texture: Mesic (1)
DEWBERRY (<i>Rubus pubescens</i>)	1.0	1.0-1.0	100	Effective Texture: Loam (1)
Graminoid				Depth to Mottles/Gley:
WATER SEDGE (<i>Carex aquatilis</i>)	1.0	1.0-1.0	100	Organic Thickness: 26 - 39 cm (1)
GRACEFUL MANNA GRASS (<i>Glyceria pulchella</i>)	2.0	2.0-2.0	100	Parent Material: Rock (1), Undifferentiated Organic (1), Lacustrine (1)
INLAND SEDGE (<i>Carex interior</i>)	3.0	3.0-3.0	100	Humus Form
BLUEJOINT (<i>Calamagrostis canadensis</i>)	5.0	5.0-5.0	100	
Moss				
TUFTED MOSS (<i>Aulacomnium palustre</i>)	2.0	2.0-2.0	100	
JUNIPER HAIR-CAP (<i>Polytrichum juniperinum</i>)	2.0	2.0-2.0	100	
SQUARROSE PEAT MOSS (<i>Sphagnum squarrosum</i>)	2.0	2.0-2.0	100	
BROWN MOSS (<i>Drepanocladus aduncus</i>)	50.0	50.0-50.0	100	

j3 rich fen - graminoid (n=0)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: j rich fen (subhydric/rich)

General Description

This ecological site phase currently has no data, but is presented here because it has been described in adjacent subregions (Northern Mixedwood and Athabasca Plain).

Characteristic Species

Environmental Variables

Moisture Regime:

Nutrient Regime:

Elevation (range):

Slope (%):

Aspect:

Topographic Position:

Soil Variables

Soil Drainage:

Soil Subgroup:

Surface Texture:

Effective Texture:

Depth to Mottles/Gley:

Organic Thickness:

Parent Material:

Humus Form

k marsh (hydric/rich) (n=3)

Natural Subregion: Kazan Upland

General Description

Marshes can be locally extensive in sheltered lake bays or along creek channels and are dominated by water and small bottle sedge, bulrushes and in deeper water pondweeds (Natural Regions Committee 2006). All the information for this ecological site is copied from the Central Mixedwood (marsh (I)) (Beckingham and Archibald 1996).

The marsh ecosite is found in level and depressional areas and around the shorelines of water bodies and riparian zones. The water is above the rooting zone for at least a portion of the growing season. These ecosites are dominated by a high diversity of emergent sedges and rushes.



Successional Relationships

The marsh ecosite is near the beginning stages of hydrarch succession. The marsh ecosite can be thought of as successional stable, but overtime will evolve into bogs and fens through the processes of terrestrialization (filling of shallow lakes) and palludification (blanketing of terrestrial ecosystems by overgrowth of wetland vegetation) (<http://www.aquatic.uoguelph.ca/wetlands/chapter2/bogpage2.htm>) both processes caused by accumulating organic matter.

Indicator Species

Forb

NORTHERN QUILLWORT

Isoetes echinospora

WATER ARUM

Calla palustris

WATER PARSNIP

Sium suave

Graminoid

NORTHERN MANNA GRASS

Glyceria borealis

Ecosection: KU Kazan Upland

Environmental Variables

Moisture Regime: Hydric (wet) (3)

Nutrient Regime: Eutrophic (very rich) (3)

Elevation (range): 231.33 (200-284) M

Slope (%): level (3)

Aspect: Level (2)

Topographic Position: Depression (3)

Soil Variables

Soil Drainage: Very poorly drained (3)

Soil Subgroup:

Surface Texture:

Effective Texture: Undifferentiated Organic (0)

Depth to Mottles/Gley:

Organic Thickness:

Parent Material: Water (0)

Humus Form

k1 marsh (n=3)

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: k marsh (hydric/rich)

Characteristic Species

Forb

- [31.6] WATER ARUM*
Calla palustris
- [6.6] NORTHERN QUILLWORT*
Isoetes echinospora
- [3.6] WATER PARSNIP*
Sium suave
- [3.3] NARROW-LEAVED BUR-REED
Sparganium angustifolium
- [1.6] ARUM-LEAVED ARROWHEAD
Sagittaria cuneata

Graminoid

- [30.0] NORTHERN MANNA GRASS*
Glyceria borealis
- [1.6] CYPERUS-LIKE SEDGE
Carex pseudo-cyperus

Environmental Variables

Moisture Regime: Hydric (wet) (3)
Nutrient Regime: Eutrophic (very rich) (3)
Elevation (range): 231.33 (200-284) M
Slope (%): level (3)
Aspect: Level (2)
Topographic Position: Depression (3)

Soil Variables

Soil Drainage: Very poorly drained (3)
Soil Subgroup:
Surface Texture:
Effective Texture: Undifferentiated Organic (0)
Depth to Mottles/Gley:
Organic Thickness:
Parent Material: Water (0)
Humus Form

KUA6 Water parsnip/Northern manna grass (n=1)

(*Sium suave*/*Glyceria borealis*)

This community was described in 2 m wide widths along stretches of the La Butte Creek in the La Butte Wildland Provincial Park (Allen et al. 2002). This community represents a floating-leaved aquatic community. The center of the creek was occupied by submerged vegetation usually dominated by clasping-leaf pondweed. Moving towards shore this community occupied the next zone of vegetation, followed by a band of emergent creeping spike-rush along the shoreline.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: k marsh (hydric/rich)
Ecosite Phase: k1 marsh

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Tall Forb (>= 30 cm)				Ecological Status Score: 0-0
ALPINE PONDWEED (<i>Potamogeton alpinus</i>)	1.0	1.0-1.0	100	Moisture Regime: Hydric (wet) (1)
VARIOUS-LEAVED PONDWEED (<i>Potamogeton gramineus</i>)	1.0	1.0-1.0	100	Nutrient Regime: Eutrophic (very rich) (1)
CLASPING-LEAF PONDWEED (<i>Potamogeton richardsonii</i>)	1.0	1.0-1.0	100	Elevation (range): 210 (210-210) M
ARUM-LEAVED ARROWHEAD (<i>Sagittaria cuneata</i>)	1.0	1.0-1.0	100	Slope (%): 0 - 0.49 (1)
WATER PARSNIP (<i>Sium suave</i>)	10.0	10.0-10.0	100	Aspect: Level (1)
NARROW-LEAVED BUR-REED (<i>Sparganium angustifolium</i>)	10.0	10.0-10.0	100	Topographic Position: Depression (1)
Graminoid				Soil Variables
NORTHERN MANNA GRASS (<i>Glyceria borealis</i>)	90.0	90.0-90.0	100	Soil Drainage: Very poorly drained (1)
Moss				Soil Subgroup:
N/A (<i>Fontinalis hypnoides</i>)	1.0	1.0-1.0	100	Surface Texture:
				Effective Texture: Undifferentiated Organic (0)
				Depth to Mottles/Gley:
				Organic Thickness:
				Parent Material: Water (0)
				Humus Form

KUA7 Northern quillwort (n=1)

(*Isoetes echinospora*)

This community type was described along La Butte (Allen et al. 2003). This community was found in a band close to the shoreline on a sandy substrate in water 0-30 cm deep. It was also noted that the consistency of the water depth for this community type was striking and suggests that this community type inhabits sands in shallow water around the entire lake. Other aquatic species noted in this community type include water milfoil and pondweeds.

Natural Subregion: Kazan Upland

Ecosection: KU Kazan Upland

Ecosite: k marsh (hydric/rich)

Ecosite Phase: k1 marsh

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Medium Shrub (0.5 to 2 m)				Ecological Status Score: 40-40
SWEET GALE (<i>Myrica gale</i>)	1.0	1.0-1.0	100	Moisture Regime: Hydric (wet) (1)
Tall Forb (>= 30 cm)				Nutrient Regime: Eutrophic (very rich) (1)
UNDIFFERENTIATED PONDWEED (<i>Potamogeton</i>)	1.0	1.0-1.0	100	Elevation (range): 284 (284-284) M
LARGE-LEAVED WHITE WATER CROWFOOT (<i>Ranunculus aquatilis</i>)	1.0	1.0-1.0	100	Slope (%): 0 - 0.49 (1)
ARUM-LEAVED ARROWHEAD (<i>Sagittaria cuneata</i>)	4.0	4.0-4.0	100	Aspect: Level (0)
NORTHERN QUILLWORT (<i>Isoetes echinospora</i>)	20.0	20.0-20.0	100	Topographic Position: Depression (1)
				Soil Variables
				Soil Drainage: Very poorly drained (1)
				Soil Subgroup:
				Surface Texture:
				Effective Texture:
				Depth to Mottles/Gley:
				Organic Thickness:
				Parent Material: Water (0)
				Humus Form

KUA8 Water arum/Sedge (n=1)

(*Calla palustris*/*Carex pseudocyperus*)

This community type was described around the open pools of water in an old meander channel along La Butte Creek near the Slave River in La Butte Creek Wildland Provincial Park (Allen et al. 2002). This community type was best expressed where the water table was about 1m deep. It was noted that open water covered about 50% of the plot.

Natural Subregion: Kazan Upland
Ecosection: KU Kazan Upland

Ecosite: k marsh (hydric/rich)
Ecosite Phase: k1 marsh

Plant Composition	Canopy Cover (%)			Environmental Variables
	Mean	Range	Const.	
Tall Forb (>= 30 cm)				Ecological Status Score: 0-0
NODDING BEGGARTICKS (<i>Bidens cernua</i>)	1.0	1.0-1.0	100	Moisture Regime: Hydric (wet) (1)
TUFTED LOOSESTRIFE (<i>Lysimachia thyrsoiflora</i>)	1.0	1.0-1.0	100	Nutrient Regime: Eutrophic (very rich) (1)
MARSH CINQUEFOIL (<i>Potentilla palustris</i>)	1.0	1.0-1.0	100	Elevation (range): 200 (200-200) M
MARSH SKULLCAP (<i>Scutellaria galericulata</i>)	1.0	1.0-1.0	100	Slope (%): 0 - 0.49 (1)
WATER PARSNIP (<i>Sium suave</i>)	1.0	1.0-1.0	100	Aspect: Level (1)
NARROW-LEAVED WATER-HEMLOCK (<i>Cicuta virosa</i>)	2.0	2.0-2.0	100	Topographic Position: Depression (1)
Low Forb (< 30 cm)				Soil Variables
SWEET-SCENTED BEDSTRAW (<i>Galium triflorum</i>)	1.0	1.0-1.0	100	Soil Drainage: Very poorly drained (1)
COMMON MARE'S-TAIL (<i>Hippuris vulgaris</i>)	1.0	1.0-1.0	100	Soil Subgroup:
COMMON DUCKWEED (<i>Lemna minor</i>)	1.0	1.0-1.0	100	Surface Texture:
WATER ARUM (<i>Calla palustris</i>)	95.0	95.0-95.0	100	Effective Texture:
Graminoid				Depth to Mottles/Gley:
SMALL BOTTLE SEDGE (<i>Carex utriculata</i>)	1.0	1.0-1.0	100	Organic Thickness:
CYPERUS-LIKE SEDGE (<i>Carex pseudo-cyperus</i>)	5.0	5.0-5.0	100	Parent Material: Water (0)
Moss				Humus Form
BROWN MOSS (<i>Drepanocladus aduncus</i>)	2.0	2.0-2.0	100	

Literature Cited

Adams, B.W., G. Ehlert, C. Stone, D. Lawrence, M. Alexander, M. Willoughby, C. Hincz, D. Moisey, and A. Bogen. 2003. Rangeland Health Assessment for Grassland, Forest and Tame Pasture. Alberta Sustainable Resource Development. Public Lands Division. Edmonton. AB. Pub. No. T/044. 104pp.

AGRASID. Version 4.0. [http://www1.agric.gov.ab.ca/\\$Department/deptdocs.nsf/All/sag14653](http://www1.agric.gov.ab.ca/$Department/deptdocs.nsf/All/sag14653)

Alberta Wetland Classification System. 2015. Alberta Environment and Sustainable Resource Development (ESRD), Water Policy Branch, Policy and Planning Division. Edmonton, AB. 66pp.

Allen, L., D. Johnson, K. Vujnovic. 2002. Small patch communities of La Butte Creek Wildland Provincial Park. A report prepared for Parks and Protected Areas, Alberta Community Development, Edmonton, Alberta.

Allen, L., J. D. Johnson, K. Vujnovic. 2004. Small Patch Communities of Colin-Cornwall Lakes Wildland Provincial Park. A report prepared for Parks and Protected Areas, Alberta Community Development, Edmonton, Alberta.

Bailey, A.W., M.G. Willoughby, R. Johansen and S. Smith. 1992. Management of Yukon Rangelands. Renewable Resources, Yukon Territorial Government, Whitehorse, Yukon. 55pp. ISBN-1-55018-138-6.

Beckingham, J. 1993. Ecosystem associations of Northern Alberta. Dept. of Environmental Protection, Alberta Forest Service, Edmonton.

Beckingham, J. and J.H. Archibald. 1996. Field guide to ecosites of Northern Alberta. Special report 5. Canadian Forest Service. Northwest Region. Edmonton, Alta.

Corns, I.G.W. and R.M. Annas. 1986. Field guide to forest ecosystems of West-Central Alberta. Northern Forestry Center, Canadian Forestry Service, Edmonton, Alta. 251pp.

Daubenmire, R. 1952. Forest vegetation of Northern Idaho and adjacent Washington and its bearing on concepts of vegetation classification. *Ecol. Mongr.* 22: 301-330.

Department of Environmental Protection. 1994. Natural Regions of Alberta. Alberta Environmental Protection. Edmonton, Alta. Pub. no.: I/531. 18pp.

Ecological Land Survey Site Description Manual (2nd). 2003. Resource Data Branch, Strategic Corporate Services Division. Alberta Sustainable Resource Development. Edmonton, AB. Pub. no. T/036. 112 pp.

Gauch, H.G. 1982. Multivariate analysis in community ecology. Cambridge University Press, Cambridge, 298pp.

<http://www.aquatic.uoguelph.ca/wetlands/chapter2/bogpage2.htm>. Chapter 2 - Canada's Aquatic Environments, University of Guelph, Ontario, Canada.

Marshall, I.B., Smith, C.A.S., Selby, C.J. 1996. A national framework for monitoring and reporting on environmental sustainability in Canada. *Environmental Monitoring and Assessment* 39: 25-38.

Meidinger, D. and J. Pojar (compilers and editors). 1991. Ecosystems of British Columbia. BC Min. For. Special Report Series 6. Victoria, BC. 330 pp.

Mueggler, W.F. 1988. Aspen community types of the Intermountain Region. U.S.D.A. Intermountain Research Station. INT-250. 133pp.

Natural Regions Committee 2006. Natural Regions and Subregions of Alberta. Compiled by D. J. Downing and W. W. Pettapiece. Government of Alberta. Pub. No. I/005.

Range Survey Manual. 1992. Range Management Section, Alberta Forest Service. Edmonton, Alta. 39pp.

Raup, H. 1935. Botanical Investigations in Wood Buffalo Park. Biological Series No. 20. Bulletin No. 74. Canada, Ottawa. 174pp.

Strong, W.L. and H.G. Anderson. 1980. Ecological Land Classification and Evaluation Reference Manual. Resource Inventory and Appraisal Resource Evaluation Branch, Alberta Energy and Natural Resources. Edmonton, AB. 160pp.

Strong, W.L. and J.M. Thompson. 1995. Ecodistricts of Alberta: Summary of Biophysical Attributes. Alberta Environmental Protection, Resource Data Division. Edmonton, Alta. Pub. no. T/319. 91pp.

Task Group on Unity and Concept. 1995. New concepts for assessment of rangeland condition. J. Range Manage. 38:220-225.

Thompson, W.H. and P.L. Hansen. 2002. Classification and management of riparian and wetland sites of the Alberta Grassland Natural Region and adjacent subregions. Bitterroot Restoration Inc. Prepared for the Alberta Riparian Habitat Management Program-Cows and Fish, Lethbridge, Alberta. 416pp.

Wallis, C. W. and C. Wershler. 1984. Kazan Upland resource assessment for ecological reserves planning in Alberta. Alberta Energy and Natural Resources, Edmonton, Alberta. 81 pp.

