

Limnanthes pumila* ssp. *grandiflora

Limnanthaceae
Woolly meadowfoam

Norman Jensen




Norman Jensen

Gerald D. Carr



Leaves sparsely
pubescent, leaflets 4-8
mm long



Flowers with white petals
and pubescent sepals

Stems sparsely pubescent,
5-15 cm long

Illustrated by Frank A. Lang
From: Threatened and Endangered
Vascular Plants of Oregon (1982)
R. J. Meinke

Low growing, annual herb. Stems sparsely pubescent, 5-15 cm long. Leaves 1-6 cm long, sparsely pubescent; leaflets linear to oblanceolate, 4-8 mm long. Flowers: sepals pubescent without at base, densely woolly pubescent within, 8-14 mm long; petals 7-10 mm long, white; filaments 4-5 mm long; anthers less than 1 mm. Fruit: nutlets, 3-5, 3-4.5 mm long, obovoid.


Lookalikes

Limnanthes floccosa ssp. *floccosa*
Limnanthes pumila ssp. *pumila*

differs from featured plant by

Ssp. *grandiflora* has sparsely pubescent stems

best survey times
J | F | M | A | M | J | J | A | S | O | N | D




***Limnanthes pumila* Howell ssp. *grandiflora* (Arroyo)**

S.C. Meyers & K.L. Chambers

Woolly meadowfoam

PLANTS symbol: LIFLG

August 2019 status Federal:LE; Oregon:LE; ORBIC: List 1



Distribution: The Agate Desert of central Jackson Co., Oregon.

Habitat: Wet edges of vernal pools.

Elevation: 370 - 400 m

Best survey time (in flower): March – April

Associated species:

Lupinus polycarpus (Smallflower lupine)

Trifolium depauperatum (Cowbag clover)

Lasthenia californica ssp. *californica* (Slender goldfields)

