

***Lachemilla mexiquense* (Rosaceae), a new species from Mexico**

Diego F. Morales-Briones¹

¹ Department of Biological Sciences and Stillinger Herbarium, University of Idaho, 875 Perimeter Drive MS 3051, Moscow, Idaho 83844-3051, U.S.A.

Corresponding author: Diego F. Morales-Briones (dfmoralesb@gmail.com)

Academic editor: Ali Dönmez | Received 27 January 2016 | Accepted 1 March 2016 | Published 25 March 2016

Citation: Morales-Briones DF (2016) *Lachemilla mexiquense* (Rosaceae), a new species from Mexico. PhytoKeys 62: 25–32. doi: 10.3897/phytokeys.62.7953

Abstract

A new species of *Lachemilla* (Rosaceae), *Lachemilla mexiquense* D.F. Morales-B., from Mexico is described and illustrated. This species is similar to *Lachemilla aphanoides* by its tripartite leaves and glomerulate inflorescence with entirely glabrous flowers, but it differs by its stoloniferous habit, persistent basal leaves and basal stipules, and smaller flowers with a campanulate-elongate hypanthium and single carpel. A key to the species of *Lachemilla* in Mexico is provided.

Resumen

Una nueva especie de *Lachemilla* (Rosaceae), *Lachemilla mexiquense* D.F. Morales-B., de México se describe e ilustra. Esta especie es similar a *Lachemilla aphanoides* por sus hojas tripartitas e inflorescencias glomeruladas con flores completamente glabras, pero difiere por su hábito estolonífero, hojas basales y estípulas basales persistentes y flores de menor tamaño con hipantio campanulado-elongado con un solo carpelo. Se provee una clave para las especies de *Lachemilla* en México.

Keywords

Lachemilla mexiquense, *Lachemilla*, Rosaceae, Mexico, new species

Introduction

Lachemilla Focke (Rydb.) is a morphologically highly variable group that includes perennial herbs, sub-shrubs, and shrubs. It comprises ca. 80 species and occurs between 2200 and 5000 m in elevation in the high mountains of the Neotropics, from northern Mexico to northern Argentina and Chile (Romoleroux 1996; Gaviria 1997), where it is one of the main elements of the páramo and superpáramo flora in South America. In Mexico the genus is represented by at least 10 species that can be found in sub-alpine and alpine habitats from the mountain pine forest to the high elevation zacatanales. *Lachemilla* has a nearly ubiquitous occurrence throughout the montane American tropics and remains a taxonomically complex group where species boundaries are often unclear and the infrageneric taxonomy is poorly defined. Since the only comprehensive revision of *Lachemilla* (Perry 1929), several works have tried to clarify its taxonomy (Rothmaler 1935, 1937; Notov and Kusnetzova 2004) and recently several regional treatments have been published (Romoleroux 2004; Barrie 2015), but a complete revision of the group is still needed. Recent taxonomic work aiming to produce a monographic treatment of *Lachemilla* has resulted in the description of several new species (Romoleroux 2009; Romoleroux and Morales-Briones 2012).

Here, a new species of *Lachemilla* is described and illustrated from Mexico. Material of the new species was collected in June 2015 during an expedition focusing solely on the genus *Lachemilla*. After detailed examination of the specimen, revision of species descriptions, and comparison with specimens at CAS, F, MEXU, MO, NY, TEX, and UC, it was established that the specimen collected in central Mexico represents a new species. The taxonomic treatment of this new species, including a key to species of *Lachemilla* in Mexico is provided.

Taxonomic treatment

Lachemilla mexiquense D.F. Morales-B., sp. nov.

urn:lsid:ipni.org:names:77153916-1

Figures 1, 2A

Diagnosis. *Lachemilla mexiquense* differs from *Lachemilla aphanoides* (Mutis ex L. f.) Rothm. by its caespitose and stoloniferous habit, creeping stems, basal leaves and basal stipule persistent, campanulate-elongate hypanthium and the presence of a single carpel.

Type. MEXICO. Estado de México, Municipio Ocuilan, 4 km NE of Santa Martha on road Santa Martha–Huitzilac, 19.07567°N, 99.36215°W, alt. 3,050 m, 30 June, 2015, Morales-Briones D. F. & Tenorio-Lezama P. 683. (holotype: ID!; isotype: MEXU!, QCA!).

Description. Caespitose herbs, stoloniferous; stems creeping, mat-forming, branches sometimes rooting, pilose. Basal leaves 3-parted, 6–20 × 5–15 mm, chartaceous, lateral segments bifid, segments obovate to cuneate, margin incised-dentate, lower surface




Figure 1. *Lachemilla mexiquense*. **A** Habit **B** Basal leaf and stipule **C** Flowering branch **D** Flower. Illustration by P. Lu-Irving.

pilose, upper surface sparsely pilose to glabrescent; petioles 12–35 mm long; stipules 5–15 mm long, adnate to the petiole at base, free, entire and acute at apex, membranaceous, greenish-white. Stem leaves 3-parted, 7–12 × 4–7 mm, chartaceous, lateral lobes entire or bifid, segments obovate to cuneate, margin deeply cleft, lower surface pilose, upper surface sparsely pilose to glabrescent; petioles 3–5 mm long; stipules 3–8 mm long, adnate to the petiole at base, free at apex, 6-lobed at apex, membranaceous and greenish-white at base, chartaceous and green at apex. Inflorescences axillary or terminal, glomerulate, 6–10 flowered cymes; floral bracts lobed, spreading; pedicels 1–1.5 mm long, pilose at apex. Flowers 1.2–1.5 mm long; hypanthium campanulate-elongate 1–1.2 × 0.6–0.8 mm, glabrous outside, glabrous inside, green when young, reddish at maturity; epipetalous 4, ovate, 0.6–0.7 × 0.5–0.7 mm, glabrous, apex acute; sepals 4,




Figure 2. *Lachemilla mexiquense*. **A** Habit **B** Type locality.

lanceolate, $0.5\text{--}0.6 \times 0.2\text{--}0.3$ mm, glabrous, apex acute; stamens 2, filaments $0.2\text{--}0.3$ mm long; carpels 1, stigma clavate. Achenes ovoid-globose, $0.9\text{--}1.1 \times 0.6\text{--}0.8$ mm, glabrous, one-seeded. Seeds ovate, $0.7\text{--}0.8 \times 0.4\text{--}0.6$ mm, pink, glabrous.




Figure 3. Geographic distribution of *Lachemilla mexiquense*.

Distribution and ecology. *Lachemilla mexiquense* is only known from the State of Mexico, municipality of Ocuilan, at ca. 3050 m altitude (Figures 2B, 3). *Lachemilla mexiquense* grows at the border of dense forest of various species of *Pinus*. This species lives in sympatry with *Lachemilla procumbens* (Rose) Rydb., *L. vulcanica* (Schltdl. & Cham.) Rydb., and *L. aphanoides* (Mutis ex L. f.) Rothm. It was collected in flower and fruit in late June.

Etymology. The specific epithet refers to the demonym for State of Mexico where the type specimen was collected.

Conservation status. *Lachemilla mexiquense* has a very limited geographic distribution, and is only known from the type locality (Figures 2B, 3). It occurs right outside the limits of the Cumbres del Ajusco National Park and Lagunas de Zempoala National Park. The type locality has been severely impacted by human activities, including conversion to agriculture (sheep and cow grazing). Following the IUCN (2014) guidelines, based on the reduced geographic distribution and altered land use at the type locality, this species should be categorized as endangered (EN), at least until other populations are discovered.

Notes. *Lachemilla mexiquense* resembles *L. aphanoides* by having tripartite leaves with bifid lateral segments and glomerulate inflorescence with entirely glabrous flowers. Nevertheless, *L. mexiquense* differs from *L. aphanoides* by its caespitose habit, creeping stems, and stolons that form dense mats. Also, the basal leaves and basal stipules are persistent, and flowers are smaller (1.2–1.5 mm long) with a campanulate-elongate hypanthium and single carpel. *Lachemilla rupestris* (Kunth) Rothm., a species from Andean South America with similar habit, differs from *L. mexiquense* by having

entire lateral segments of the leaves, yellow-brown membranaceous basal stipules, and larger flowers (2.5–3 mm long) with a turbinate-campanulate hypanthium, sericeous-hirsute pubescence, and 2–4 carpels.

In addition, phylogenetic analyses of chloroplast and nuclear DNA (Morales-Briones et al. unpubl. data) clearly separate *L. mexiquense* from *L. aphanoides* and *L. rupestris*. The chloroplast phylogeny place it as sister species of the ‘Orbiculate group,’ which encompasses species with stoloniferous habit, palmately lobed or cleft leaves, and flowers disposed in profuse terminal cymes, like *Lachemilla pectinata* (Kunth) Rothm. The nuclear phylogeny fails to confidently resolve the phylogenetic position of *L. mexiquense*, suggesting that it may be of hybrid origin, a common pattern seen throughout *Lachemilla*.

Key to the species of *Lachemilla* in Mexico

Notes. Adapted from Standley and Steyermark (1946), Romoleroux (2004), and Barrie (2015). Accepted taxa and synonymy follows the regional revisions of Romoleroux (2004) and Barrie (2015), with the exception of *L. sibbaldifolia* (Kunth) Rydb. and *L. pringlei* Rydb., which based on extensive field observations and the examination of herbarium material, are considered here as two distinct taxa.

- | | | |
|---|---|----------------------|
| 1 | Leaves pinnately divided | <i>L. pinnata</i> |
| — | Leaves simple or palmately divided or cleft | 2 |
| 2 | Basal leaves 5–11-lobed or 5–7-cleft | 3 |
| — | Basal leaves 3–5-cleft or 3–5-parted | 4 |
| 3 | Leaves shallowly 5–11-lobed, lobes triangular | <i>L. pectinata</i> |
| — | Leaves deeply 5–7-cleft, lobes elliptical to ovate | <i>L. venusta</i> |
| 4 | Inflorescence of loose cymes; hypanthium pubescent within | 5 |
| — | Inflorescence glomerulate, forming dense cymes; hypanthium glabrous within | 6 |
| 5 | Leaves 3-parted with bifid lateral segments, appearing 5-parted; stipules lobed or incised-dentate | <i>L. procumbens</i> |
| — | Leaves 3-parted with entire lateral segments, not appearing 5-parted; stipules bifid | <i>L. vulcanica</i> |
| 6 | Plants pilose to glabrate; flowers glabrous | 7 |
| — | Plants hirsute to sericeous; flowers pubescent, sometimes glabrate with age ... | 8 |
| 7 | Stems decumbent, ascending or erect; basal leaves and basal stipules often... caduceus; flowers 1.5–3.0 mm; 1–3 carpels | <i>L. aphanoides</i> |
| — | Stems creeping, stoloniferous; basal leaves and basal stipules persistent; flowers 1.2–1.5 mm; 1 carpel | <i>L. mexiquense</i> |
| 8 | Hypanthium densely pubescent with very short hairs; lower leaves short-petioled, the upper leaves sessile | <i>L. velutina</i> |

- Hypanthium sparingly pubescent with appressed hairs, lower and upper leaves petioled..... 9
- 9 Leaves appearing 5-lobed, the lateral lobes bifid; achenes subacute or subobtuse..... *L. sibbaldiifolia*
- Leaves appearing 3-lobed, the lateral lobes not bifid; achenes tapering to an acute apex..... *L. pringlei*

Acknowledgements

I am indebted to David S. Gernandt for facilitating collection permits and logistics, and to Pedro Tenorio-Lezama for assistance in the field. I am grateful to the curators of CAS, MEXU, TEX, and UC for access to their collections, Pat Lu-Irving for the illustrations, and Katya Romoleroux for access to additional specimens. I would like to thank Hannah E. Marx, David C. Tank and Katya Romoleroux for proving helpful comments on the manuscript. Funding for this work was provided by a Secretaría de Educación Superior, Ciencia, Tecnología e Innovación del Ecuador (SENESCYT) Doctoral scholarship, Graduate Student Research Grants from the Botanical Society of America (BSA), the American Society of Plant Taxonomists (ASPT) and the International Association of Plant Taxonomist (IAPT), the University of Idaho Stillinger Herbarium Expedition Funds, and a National Science Foundation Doctoral Dissertation Improvement Grant to David C. Tank and DFM-B (DEB-1502049).

References

- Barrie FR (2015) *Lachemilla* (Focke) Rydb. In: Davidse G, Sousa Sánchez M, Knapp S, Chiang Cabrera F (Eds) Flora Mesoamericana: Saururaceae a Zygophyllaceae 2(3): 1–347.
- Gaviria J (1997) Sinopsis del género *Lachemilla* (Focke) Rydberg (Rosaceae) para Venezuela. Plantula 1(3): 189–212.
- IUCN Standards and Petitions Subcommittee (2014) Guidelines for using the IUCN Red List Categories and Criteria. Version 11. Prepared by the Standards and Petitions Subcommittee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> [accessed: 15 Nov 2015]
- Notov AA, Kusnetzova TV (2004) Architectural units, axiality and their taxonomic implications in Alchemillinae. Wulfenia 11: 85–130.
- Perry LM (1929) A tentative revision of *Alchemilla* § *Lachemilla*. Contributions from the Gray Herbarium of Harvard University 84: 1–57.
- Romoleroux K (1996) 79. Rosaceae. In: Harling G, Andersson L (Eds) Flora of Ecuador 56: 1–152.
- Romoleroux K (2004) The genus *Lachemilla* (Rosaceae) in the northern Andes of South America. Lyonia 7(1): 21–32.
- Romoleroux K (2009) New Species of *Lachemilla* (Rosaceae) from South America. Novon: A Journal for Botanical Nomenclature 19(4): 502–506. doi: 10.3417/2006054

- Romoleroux K, Morales-Briones DF (2012) *Lachemilla jaramilloi* and *L. talamanquensis* spp. nov. (Rosaceae) from Ecuador and Costa Rica. Nordic Journal of Botany 30(6): 732–736. doi: 10.1111/j.1756-1051.2011.01413.x
- Rothmaler W (1935) Systematische Vorarbeiten zu einer Monographie der Gattung *Alchemilla*. (L.) Scop. emend. II. Die systematische Gliederung der Gattung. Repertorium novarum specierum regni vegetabilis 38: 409–412.
- Rothmaler W (1937) Systematische Vorarbeiten zu einer Monographie der Gattung *Alchemilla* (L.) Scop. VII. Aufteilung der Gattung und Nomenklatur. Repertorium novarum specierum regni vegetabilis 42: 164–173. doi: 10.1002/fedr.19370421106
- Standley P, Steyermark J (1946) Flora of Guatemala - Part IV. Fieldiana, Botany 24(4): 432–484.