

Effects of soil and drainage on the savanna vegetation in the northern Brazilian Amazonia

July 27 2017



The tree species *Xylopia aromatica* is a common representative of the Roraima's flora. Credit: Reinaldo Imbrozio Barbosa

It is a well-known fact that environmental factors such as soil texture and drainage determine to a very large degree the vegetation appearance, richness and composition at any site. However, there has been little research on how these variables influence the flora in the marvellous savannas—large open areas characterised by a complex and unique network of natural resources and life forms.

Consequently, a Brazilian research team, led by Dr. Maria Aparecida de Moura Araújo, Universidade Federal de Roraima, investigated the hydro-edaphic conditions in the [savanna](#) areas in the northern Brazilian Amazonia. Their study, complete with an openly available and ready for re-use dataset, is published in the open access *Biodiversity Data Journal*.

In the course of the Program for Biodiversity Research, managed by the Brazilian government, the scientists sampled 20 permanent plots in two savanna [areas](#) in the state of Roraima, located in the northern of the Brazilian Amazon. As a result, the team reports a total of 128 plant species classified into 34 families from three savanna habitats with different levels of hydro-edaphic restrictions.

Amongst the various factors playing a role in the soil characteristics of the area, are the tectonic events and past climatic fluctuations which have occurred in the most recent period of the Cenozoic era. Paleo, as well as modern fires are likely to be other culprits for the specific conditions.



The herb species *Merremia aturensis* is a common representative of the Roraima's flora. Credit: Reinaldo Imbrozio Barbosa

In conclusion, the authors suggest that the most restrictive savanna habitats - the wet grasslands, represent the home to less structurally complex plants, compared to the well-drained shrubby localities.

"The present study highlights the environmental heterogeneity and the biological importance of Roraima's savanna regarding the conservation of natural resources from the Amazon," say the scientists.

"In addition, it points out the need for greater investment in floristic

inventories associated with greater diversification of sites, since this entire ecosystem has been rapidly modified by agribusiness."



The herb species *Drosera roraimae* is a common representative of the Roraima's flora. Credit: Reinaldo Imbrozio Barbosa

More information: Maria Aparecida Araújo et al, Hydro-edaphic conditions defining richness and species composition in savanna areas of the northern Brazilian Amazonia, *Biodiversity Data Journal* (2017). [DOI:](#)

[10.3897/BDJ.5.e13829](https://doi.org/10.3897/BDJ.5.e13829)

Provided by Pensoft Publishers

Citation: Effects of soil and drainage on the savanna vegetation in the northern Brazilian Amazonia (2017, July 27) retrieved 2 May 2024 from <https://phys.org/news/2017-07-effects-soil-drainage-savanna-vegetation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.