



ORIGINAL ARTICLE

New circumscription of the genus *Gamochaeta* (Asteraceae, Gnaphalieae) inferred from nuclear and plastid DNA sequences

Estrella Urtubey¹ · Alicia López¹ · María A. Chemisquy² · Arne A. Anderberg³ ·
Carlos M. Baeza⁴ · Néstor D. Bayón⁵ · Leonardo P. Deble⁶ · Andrés Moreira-Muñoz⁷ ·
Guy L. Nesom⁸ · Mac H. Alford⁹ · Luciana Salomón¹ · Susana E. Freire¹

Received: 9 September 2015 / Accepted: 2 May 2016 / Published online: 1 July 2016
© Springer-Verlag Wien 2016

Abstract *Gamochaeta* (tribe Gnaphalieae, Asteraceae) is composed of ca. 60 species primarily distributed in tropical and subtropical America. Within the tribe Gnaphalieae, the genus is characterized by capitula arranged in spikes or head-like clusters, few hermaphroditic central florets, truncate style branches with apical sweeping trichomes, pappus bristles connate at the base into a ring falling as a unit, and achenes with globose twin trichomes. Previous molecular phylogenetic studies have suggested the paraphyly of the genus, but have not provided a basis for redefining generic limits due to incomplete taxon sampling. To address this problem, DNA sequences from the plastid (*trnL*-F) and nuclear (ETS and ITS) genomes were analyzed from a broad taxon sample representing the full range of morphological variation known in the genus. Our results affirm that *Gamochaeta* is paraphyletic as presently circumscribed. Two clades can be recognized: one clade that includes the majority of the species currently assigned to

Gamochaeta and a second clade that includes *Gamochaetopsis*, *Stuckertiella* and seven species of *Gamochaeta*. We present here a new circumscription of *Gamochaeta*, including two new combinations, *Gamochaeta alpina* and *Gamochaeta peregrina*, and the resurrection of *Gamochaeta capitata*. Our results also show *Omalotheca supina*, *O. norvegica* and *O. sylvatica*, which were placed by some authors in *Gamochaeta* or in *Gnaphalium*, form a monophyletic group distantly related to both genera.

Keywords *Gamochaetopsis* · *Omalotheca* · *Stuckertiella* · ETS · ITS · *trnL*-F

Introduction

Gamochaeta (tribe Gnaphalieae, Asteraceae) is represented by ca. 60 species characterized by their capitula arranged in spikes or head-like clusters, pistillate florets

Handling editor: Christoph Oberprieler.

✉ Estrella Urtubey
eurtubey@darwin.edu.ar

¹ Instituto de Botánica Darwiniún-CONICET, Labardén 200, CC 22, B1642HYD San Isidro, Buenos Aires, Argentina

² Centro Regional de Investigaciones Científicas y Transferencia Tecnológica de La Rioja (CRILAR)-Provincia de La Rioja, UNLaR, UNCa, SEGEMAR, CONICET, Entre Ríos y Mendoza s.n., 5301 Anillaco, La Rioja, Argentina

³ Department of Botany, Swedish Museum of Natural History, P. O. Box 50007, 104 05 Stockholm, Sweden

⁴ Departamento de Botánica, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Casilla 160-C, Concepción, Chile

⁵ Área de Botánica, Departamento de Ciencias Biológicas, Facultad de Ciencias Agrarias y Forestales, Universidad Nacional de La Plata, Avda. 60 entre 116 y 118, 1900 La Plata, Argentina

⁶ Curso de Licenciatura em Ciências da Natureza, Universidade Federal do Pampa, Dom Pedrito, RS CEP 96400-110, Brazil

⁷ Instituto de Geografía, Pontifícia Universidad Católica de Valparaíso, Brasil 2241, Valparaíso, Chile

⁸ 2925 Hartwood Drive, Fort Worth, TX 76109, USA

⁹ Department of Biological Sciences, University of Southern Mississippi, 118 College Drive #5018, Hattiesburg, Mississippi 39406-0001, USA

outnumbering the hermaphroditic disk florets, truncate style branches with apical sweeping trichomes, pappus bristles connate at the base into a ring that falls as a unit, and achenes with globose twin trichomes (e.g., Cabrera 1961; Anderberg 1991; Bayer et al. 2007; Ward et al. 2009). The greatest diversity is in South America, with ca. 45 species (Cabrera 1963, 1971, 1974, 1978; Aristeguieta 1964; Marticorena and Quezada 1985; Dillon and Sagástegui-Alva 1991a, b; Freire 1998; Deble and Marchiori 2007; Freire and Iharlegui 2008, 2014; Freire et al. 2011; Hind 2011). A few species are native to Central America, e.g., *G. irazuensis*, *G. standleyi* (Pruski 2011), and North America, e.g., *G. argyrinea* and *G. ustulata* (Nesom 1990a, 2004, 2006), and a few species are adventive or naturalized in Asia, Australia, Hawaii, New Zealand, Southern Africa, and Europe, e.g., *Gamochaeta calviceps* and *G. pensylvanica* (Drury 1971; Hilliard 1983; Chen and Bayer 2011; Alford 2012).

Gamochaeta was first proposed as a genus by Weddell (1855) to accommodate five species of *Gnaphalium* L. diagnosed by the presence of pappus bristles connate at the base into a ring, hence the name of the genus (from the Greek *gamos* “connate” and *xaite* “hairs”). In the following years, many authors considered *Gamochaeta* as a section of *Gnaphalium* (Bentham and Hooker 1873; Hoffmann 1890; Wagenitz 1965; Drury 1970, 1971; Merxmüller et al. 1977; Hilliard and Burtt 1981; Webb 1988), whereas others recognized it as a distinct genus (Cabrera 1961; Nesom 1990a, 2004, 2006; Anderberg, 1991; Dillon and Sagástegui-Alva 1991a, b; Freire and Iharlegui 1997, 2014; Bayer et al. 2007; Deble and Marchiori 2007; Chen and Bayer 2011; Hind 2011).

In the first morphological cladistic study of generic relationships in the Gnaphalieae, Anderberg (1991) included a cladogram in which *Gamochaeta* was sister to the “*Lucilia* group” (e.g., *Belloa* J.Rémy, *Chevreulia* Cass., *Facelis* Cass.) and made numerous combinations in *Gamochaeta*. Anderberg’s work demonstrated the heterogeneity of *Gnaphalium* as it was circumscribed in many floras (e.g., Rémy 1849; Aristeguieta 1964), as well as by Merxmüller et al. (1977). Anderberg (1991) also recognized many of the segregate genera from *Gnaphalium*, i.e., *Anaphaloïdes* (Benth.) Kirp., *Euchiton* Cass., *Gamochaeta*, *Gnaphaliothamnus* Kirp., *Homognaphalium* Kirp., and *Pseudognaphalium* Kirp., but still left a rather heterogeneous *Gnaphalium* including *Omalotheca* Cass. and *Synchaeta* Kirp. Different authors have variously recognized the genus *Omalotheca* (including *Synchaeta*) or placed *Omalotheca* species in *Gamochaeta* or *Gnaphalium*. Nesom (1990b) excluded *Omalotheca* from *Gnaphalium* and mentioned that *Omalotheca* was superficially similar to *Gamochaeta* in its spiciform capitulescences. In their

treatment for Flora of China, Chen and Bayer (2011) placed some species of *Omalotheca* in *Gamochaeta* (*O. nanchuanensis* (Y.Ling & Y.Q.Tseng) Holub, *O. sylvatica* (L.) Sch.Bip. & F.W.Schultz, *O. norvegica* (Gunnerus) Sch.Bip. & F.W.Schultz) but placed the type species of the genus (*O. supina* (L.) DC.) in *Gnaphalium*. However, Blöch et al. (2010) considered *Omalotheca hoppeana* (W.D.J.Koch) Sch.Bip. & F.W.Schultz, *O. norvegica*, *O. supina* and *O. sylvatica* as *Gnaphalium* species.

More recently, phylogenetic analyses utilizing molecular data have been used to address relationships within the tribe Gnaphalieae. Based on the plastid marker *trnL-F*, Blöch et al. (2010) presented a molecular phylogeny of the Asian-European *Leontopodium* R.Br. ex Cass. and found a relationship between *Gamochaeta* (*G. pensylvanica*), *Antennaria* Gaertn., *Ewartia* Beauverd, *Leucogenes* Beauverd and *Leontopodium*. Based on both nuclear (ETS, ITS) and plastid markers (*rpl32-trnL* intergeneric spacer + *trnL* intron + *trnL-F*), Galbany-Casals et al. (2010) recognized the “FLAG clade” including *Filago* Loefl., *Leontopodium*, *Antennaria* and *Gamochaeta*. In a study focused on ancient polyploidy in Gnaphalieae, Smissen et al. (2011) also recovered *Gamochaeta* [*G. coarctata*, *G. subfalcata*] in the “FLAG clade.” More recently, Freire et al. (2015) using DNA sequences from plastid (*rpl32-trnL*, *trnL-F*) and nuclear (ITS and ETS) markers, together with morphological characters, presented a cladogram of the South American “*Lucilia* group s.l.” in which *Gamochaeta* (three species) formed a well-supported clade together with *Gamochaetopsis* and *Stuckertiella*, hence providing initial evidence that *Gamochaeta*, as currently circumscribed, might not be monophyletic.

The aim of this study is to extend the sampling of *Gamochaeta* to provide a better supported and more reliable phylogenetic hypothesis for generic realignments using one plastid (*trnL-F*) and two nuclear DNA regions (ETS and ITS).

Materials and methods

Ingroup and outgroup

This study includes 33 species of *Gamochaeta* representing ca. 60 % of the species of the genus, all major morphological forms (concolorous and discolored leaves, spicate and head-like arrangement of capitula) and almost the entire distributional and elevational range of the genus. In order to test the monophyly of *Gamochaeta*, we included 22 species of 13 other genera of Gnaphalieae, ten of them belonging to the FLAG clade and two outside of the FLAG clade. The choice of these genera and species was based on

Table 1 Distinguishing characters of *Gamochaeta*, *Gamochaetopsis*, and *Stuckertiella* (exclusive character in boldface)

	<i>Gamochaeta</i>	<i>Gamochaetopsis</i>	<i>Stuckertiella</i>
Distribution	North, Central and South America	Southern Andes of Chile and Argentina	Central Andes of Peru, Bolivia and NW Argentina
Habit	Ascendent or erect multistemmed herbs (caespitose)	Caespitose multistemmed herbs	Ascendent multistemmed herbs
Capitula arrangement	Spikes (head-like clusters or solitary)	Head-like clusters	Head-like clusters
No. of ray florets	25–119 (12–20)	9–17	Many
Central florets	Hermaphrodite	Hermaphrodite	Male
No. corolla lobes of central florets	5(4)	5	4
Achenial twin hairs	Globose (clavate)	Clavate	Globose

results of previous investigations: (1) Based on Freire et al. (2015), representatives of the sister group of the *Gamochaeta-Stuckertiella* clade such as *Belloa chilensis* (Hook. & Arn.) J.Rémy, *Berroa gnaphalioides* (Less.) Beauverd, *Facelis plumosa* (Wedd.) Sch.Bip., *Jalcophila boliviensis* Anderb. & S.E.Freire, *Jalcophila ecuadorensis* M.O.Dillon & Sagást. and *Lucilia acutifolia* (Poir.) Cass. were included; (2) based on Galbany-Casals et al. (2010), Smissen et al. (2011), and Freire et al. (2015), other related Gnaphalieae species such as *Antennaria chilensis* J. Rémy, *Filago fuscescens* Pomel, *F. lutescens* Jord., *F. pyramidata* L., *Gnaphalium austroafricanum* Hilliard, *G. declinatum* L.f., *G. uliginosum* L., *Leontopodium alpinum* Cass., *L. microphyllum* Hayata, *Omalotheca norvegica*, *O. supina*, and *O. sylvatica* were included. Trees were rooted with *Achyrocline tomentosa* Rusby, selected from Freire et al. (2015). DNA samples were obtained from silica-preserved leaves and from herbarium specimens. Vouchers are deposited in CONC, LP, LPB, MNCS, MO, SI, SZU, USMS, and W. When no plant material was available, sequences were obtained from GenBank (“Appendix”).

Morphological characters and distribution of the species of *Gamochaeta* in Tables 1, 2 and 3 are from the literature and own observations.

DNA extraction, amplification, sequencing, sequence alignment and editing

DNA extraction used the modified CTAB protocol by Doyle and Dickson (1987), adapted for small amounts of plant material. When material preserved in silica gel was not available, DNA was extracted from herbarium specimens using the DNeasy Plant Mini Kit (QIAGEN Inc., Hilden, Germany).

The plastid intergenic spacer *trnL-F* (primers C and F; Taberlet et al. 1991) and nuclear regions ITS (primers ITS4 and ITS5; White et al. 1990) and ETS (primers ETS1 and 18S-ETS; Bayer et al. 2002 and Baldwin and Markos 1998, respectively) were selected for this study. Reactions were

performed in a final volume of 25 µl or rarely in 50 µl. Each reaction contained 50–100 ng of DNA, 1.5 units of Taq polymerase (Invitrogen Life Technologies, São Paulo, Brazil or TaKaRa ExTaq, Otsu, Shiga, Japan), 1 × PCR Buffer, 5 mM MgCl₂, 0.2 pmol of each primer and 0.025 mM of each dNTP. In species for which these protocols were unsuccessful, BSA 0.4 % and DMSO 1.6 % or a mixture of trehalose, BSA and polysorbate-20 (Samarakoon et al. 2013) were included to increase the yield of PCR. The annealing temperatures ranged between 48 and 52 °C for the plastid markers and 56–60 °C for the nuclear markers. Final extension at 72 °C for 6 min terminated the reactions. The quality of the PCR products was estimated by electrophoresis and visualized with ethidium bromide under UV light. A negative control with no template was included for each series of amplifications to test for contamination. PCR products were sequenced by Macrogen Inc. (Korea) or Eurofins MWG Operon (Louisville, KY).

Sequences were assembled and edited using the program ChromasPro version 1.34 (Technelysium Pty, Ltd, Tewantin, Australia). Matrices were edited using the program BioEdit (Hall 1999), and sequences were aligned using the application ClustalW, using multiple alignment with the option run ClustalW. Data matrices are deposited at TreeBase (TB2: S19294).

Phylogenetic analyses

Analyses of nuclear and plastid sequences were performed separately and combined. Parsimony analyses were conducted using the program TNT version 1.1 (Goloboff et al. 2008), with all characters equally weighted and considered unordered. Gaps were scored as missing data. In all analyses, parsimony-uninformative characters were omitted. Heuristic searches were performed using 1000 random addition replicates and tree bisection–reconnection (TBR) branch swapping, saving ten trees per replicate. Branch support was assessed with 10,000 parsimony jackknifing replicates (JK; Farris et al. 1996), using ten series of

Table 2 Key morphological characters of the species of *Gamochaeta*, *Gamochaetopsis* and *Stuckertiella* (species used in the analysis in boldface)

Genera <i>Gamochaeta</i>	Plant height (cm)	Leaf blade shape (cauline leaves)	Leaf surfaces	Capitula arrangement	Inflorescence spikes	Inner phyllary apex shape	No. of corolla lobes	No. of marginal florets	Achenial twin hairs
<i>G. affinis</i> Cabrera	5–12	Linear-ovovate	Concolorous	Head-like clusters	—	Acute-apiculate	5	ca. 30	Globose
<i>G. aliena</i> (Hook. & Arn.) Cabrera	16–17	Linear-ob lanceolate	Concolorous	Spikes	Continuous (interrupted basally)	Acute	5	ca. 18	Globose
<i>G. ambatensis</i> Ariza	ca. 25	Oblanceolate	Discolorous	Spikes	Continuous (interrupted basally)	Obtuse	5	ca. 50	Globose
<i>G. americana</i> (Mill.) Wedd.	15–35	Spathulate	Discolorous	Spikes	Continuous (interrupted basally)	Short acuminate	5	ca. 90	Globose
<i>G. andina</i> (Phil.) Cabrera	3–6	Narrowly linear	Concolorous	Head-like clusters	—	Acute	5	ca. 12	Globose
<i>G. antarctica</i> (Hook. f.) Cabrera	2–4	Obovate	Concolorous	Head-like clusters	—	Acute	5	Many	Globose
<i>G. antillana</i> (Urb.) Anderb.	15–25	Linear-lanceolate (basal)	Concolorous	Spikes	Continuous (interrupted basally)	Subobtuse	5	ca. 77	Globose
<i>G. argentina</i> Cabrera	10–30	Spathulate	Concolorous	Spikes	Interrupted	Acuminate	5	ca. 70	Globose
<i>G. argyrinea</i> G.L.Nesom	12–40	Oblanceolate	Discolorous	Spikes	Continuous (interrupted basally)	Truncate-rounded	5	ca. 118	Globose
<i>G. axillaris</i> (J.Rémy) Cabrera	ca. 15	Oblanceolate	Concolorous	Spikes	Interrupted	Obtuse	5	52–63	Globose
<i>G. badillana</i> (Ariteg.) Anderb.	ca. 4	Linear-lanceolate	Concolorous	Head-like clusters	—	Acute	5	Many	—
<i>G. beckii</i> Urubey & S.E.Freire	ca. 20	Oblanceolate	Slightly discolorous	Head-like clusters arranged in leafy spikes	—	Short acuminate	5	36–38	Globose
<i>G. berteroana</i> (DC.) Cabrera	ca. 15	Oblanceolate	Concolorous	Spikes	Continuous (interrupted basally)	Acute	5	Many	Globose
<i>G. cabrerae</i> Anderb.	ca. 6	Oblanceolate	Discolorous	Spikes	Continuous	Acute	5	42–45	Globose
<i>G. calviceps</i> (Fernald) Cabrera	20–40	Linear-ob lanceolate	Concolorous	Spikes	Continuous (interrupted basally)	Obtuse to rounded	5	ca. 50	Globose
<i>G. camaguensis</i> Deble	30–60	Spathulate	Concolorous	Spikes	Continuous (interrupted basally)	Acute	5	70–100	Globose

Table 2 continued

Genera <i>Gamochaeta</i>	Plant height (cm)	Leaf blade shape (caulin leaves)	Leaf surfaces	Capitula arrangement	Inflorescence spikes	Inner pharynx shape	No. of corolla lobes	No. of marginal florets	Achenial twin hairs
<i>G. chamissonis</i> (DC.) Cabreria	15–50	Linear- oblanceolate	Concolorous	Spikes	Interrupted	Obtuse	5	27–50	Globose
<i>G. chionesthes</i> G.L.Nesom	Up to 45	Oblanceolate	Discolorous	Spikes	Continuous (interrupted basally)	Acute to acute- acuminate	5	86–119	Globose
<i>G. coarctata</i> (Willd.) Kerguélen	20–50	Spathulate	Discolorous	Spikes	Continuous (interrupted basally)	Obtuse to rounded	5	80–85	Globose
<i>G. depilata</i> (Phil.) Cabreria	5–20	Obovate	Concolorous	Head-like clusters	–	Subacute	5	12–20	Globose
<i>G. deserticola</i> Cabreria	3–10	Oblanceolate	Concolorous	Spikes	Continuous (interrupted basally)	Subacute	5	ca. 25	Globose
<i>G. diffusa</i> Deble & Marchiori	15–30	Oblanceolate	Concolorous to slightly discolorous	Spikes	Interrupted	Obtuse- apiculate	5	40–60	Globose
<i>G. erecta</i> Deble	20–50	Oblanceolate to spathulate	Concolorous to slightly discolorous	Spikes	Interrupted	Acute	5	ca. 70	Globose
<i>G. erythractis</i> (Wedd.) Cabreria	1–6 (10)	Oblanceolate	Concolorous	Spikes	Continuous	Obtuse to subacute	5	ca. 30	Globose
<i>G. falcatia</i> (Lam.) Cabreria	10–50	Linear- oblanceolate	Concolorous	Spikes	Continuous	Acute	5	ca. 70	Globose
<i>G. flagaginea</i> (DC.) Cabreria	15–40	Oblanceolate	Concolorous	Spikes	Continuous (interrupted basally)	Acute	5	ca. 78	Globose
<i>G. foliosa</i> (Phil.) Anderb.	ca. 7	Broadly obovate	Slightly discolorous	Head-like clusters	–	Unknown	5	ca. 49	Oblong
<i>G. girardiana</i> Deble & A.S.Olivera	15–45	Oblanceolate	Concolorous	Head-like clusters arranged in leafy spikes	–	Acute	5	40–50	Globose
<i>G. graziaeae</i> (Rizzini) Deble	20–50	Oblanceolate	Concolorous	Spikes	Interrupted	Long- acuminate	5	ca. 101	Globose
<i>G. hiemalis</i> Cabreria	30–40	Linear- oblanceolate	Concolorous	Spikes	Continuous	Obtuse	5	30–50	Clavate
<i>G. humilis</i> Wedd. <i>G. irazuensis</i> G.L.Nesom	ca. 10 ca. 30	Oblanceolate Elliptic- oblanceolate	Discolorous	Spikes	Interrupted	Acute	5	35–40	Globose
			Discolorous	Spikes	Continuous	Acute	5	65–73	Globose

Table 2 continued

Genera <i>Gamochaeta</i>	Plant height (cm)	Leaf blade shape (caulin leaves)	Leaf surfaces	Capitula arrangement	Inflorescence spikes	Inner pharynx apex shape	No. of corolla lobes	No. of marginal florets	Achenial twin hairs
<i>G. longipedicellata</i> Cabrera	1–4	Oblong-spathulate	Concolorous	Solitary in the axils of the upper leaves	—	Acute	5	ca. 30	Globose
<i>G. lilioides</i> S.E.Freire & Tharr. V.M.Badillo	ca. 1	Oblong-ovate	Concolorous	Solitary	—	Acute	4	ca. 27	Globose
<i>G. meridensis</i> Reiche) Cabrera	1–4	Oblanceolate	Concolorous	Head-like clusters	—	Acute	5	Many	Globose
<i>G. monticola</i> (Phil. ex G. neuquensis Cabrera G. nigrevestis Deble & Marchiori	2–4	Oblanceolate	Concolorous	Head-like clusters	—	Acute	5	ca. 30	Globose
<i>G. linearis</i> Cabrera G. oligantha (Phil.) L.E.Nava	5–12 10–20	Oblanceolate	Concolorous	Head-like clusters	—	Acute	5	ca.12 50–70	Clavate Globose
<i>G. nivalis</i> Cabrera <i>G. paramoana</i> (S.F.Blake) Anderb.	4–7 2–5 7–15	Linear-ovate Oblanceolate	Concolorous	Head-like clusters	—	Subacute	5	13–15 More than 40	Globose Many
<i>G. pensylvanica</i> (Willd.) Cabrera	20–50	Spathulate	Concolorous	Spikes	Continuous (interrupted basally)	Obtuse or subacute	5	70–80	Globose
<i>G. procumbens</i> (Phil.) Cabrera	10–25	Linear-ovate	Concolorous	Head-like clusters	—	Acute or subacute	5	ca. 20	Globose
<i>G. purpurea</i> (L.) Cabrera	20–40	Oblanceolate	Slightly discolorous	Spikes	Continuous (interrupted basally)	Acute	5	70–80	Globose
<i>G. rizzini</i> Cabrera <i>G. serpyllifolia</i> Wedd. Cabrera	ca. 35 10–15	Oblanceolate Widely ovate or suborbicular	Discolorous Concolorous	Spikes Head-like clusters	Interrupted —	Acuminatae	5	Many ca. 20	Globose Globose
<i>G. simplicicaulis</i> (Willd. ex Spreng.) Cabrera	40–60	Oblanceolate	Discolorous	Spikes	Interrupted	Acuminatae	5	ca. 90	Globose
<i>G. sphacelata</i> (Kunth) Cabrera	15–20	Narrowly linear	Concolorous	Spikes	Continuous (interrupted basally)	Acute	5	ca. 30	Globose
<i>G. spiciformis</i> (Sch.Bip.) Cabrera	7–30	Widely obovate	Discolorous	Head-like clusters	—	Obtuse	5	ca. 50	Globose
<i>G. stachydfolia</i> (Lam.) Cabrera	10–40	Spathulate	Concolorous	Spikes	Continuous	Acute	5	ca. 50	Globose

Table 2 continued

Genera <i>Gamochaeta</i>	Plant height (cm)	Leaf blade shape (caulin leaves)	Leaf surfaces	Capitula arrangement	Inflorescence spikes	Inner phyllary apex shape	No. of corolla lobes	No. of marginal florets	Achenial twin hairs
<i>G. stagnalis</i> (L.M.Johnst.) Anderb.	2.5–30	ob lanceolate to oblong-ob lanceolate	Concolorous to slightly discolorous	Spikes	Continuous (interrupted basally)	Obtuse to rounded	5	ca. 50	?
<i>G. standleyi</i> (Steyermark)	1–4.5	Elliptic-ovate to ob lanceolate	Concolorous	Spike	Continuous	Subacute	5	ca. 65	Globose
<i>G. l.Nesom</i>									
<i>G. subfulvata</i> (Cabrera) Cabrera	15–35	ob lanceolate	Concolorous	Spikes	Continuous (interrupted basally)	Obtuse	5	ca. 100	Globose
<i>G. suffruticosa</i> (Phil.) Anderb.	ca. 10	Linear-ob lanceolate	Concolorous	Spikes	Interrupted	Obtuse	5	Unknown	Globose
<i>G. thomensis</i> (Spreng.) Anderb.	ca. 25	Spathulate	Slightly discolorous	Spikes	Continuous	Unknown	5	Unknown	Globose
<i>G. ustulata</i> (Nutt.) Holub	10–40	Broadly obovate to ob lanceolate	Discolorous	Spikes	Usually continuous	Rounded to obtuse	5	Many	?
<i>G. valparadisea</i> (Phil.) Anderb.	10–15	Spathulate	Concolorous	Spikes	Continuous	Acute	5	ca. 50	Globose
<i>G. villarroelii</i> (Phil.) Cabrera	15–20	Oblong-ob lanceolate	Concolorous	Spikes	Interrupted	Acute	5	38–49	Globose
<i>Gamochaetopsis</i> Anderb. & S.E.Freire									
<i>G. alpina</i> (Poep.) Anderb. & S.E.Freire	3–5	Widely obovate or suborbicular	Concolorous	Head-like clusters	—	Acute or subacute	5	9–17	Clavate
<i>Stuckertia</i> Beauverd									
<i>S. capitata</i> (Wedd.) Beauverd	8–30	Narrowly linear	Concolorous	Head-like clusters	—	Obtuse	4	50–75	Globose
<i>S. peregrina</i> Beauverd	4–15	Linear-ovate	Concolorous	Head-like clusters	—	Obtuse	4	Many	Globose

Table 3 Species of *Gamochaeta* (based on this study) with its distributions (species included in the analysis are in bold face)

Genera/species	Distribution (*adventive or naturalized)
<i>Gamochaeta</i>	
<i>G. affinis</i>	Falkland Islands (=Islas Malvinas)
<i>G. aliena</i>	C and S Chile. 250–2600 m a. s. l.
<i>G. alpina</i>	SW Argentina, S Chile. 1250–1900 m a. s. l.
<i>G. ambatensis</i>	NW Argentina. 2600–2700 m a. s. l.
<i>G. americana</i>	Central and South America. *New Zealand, *Hawaii. 0–3500 m a. s. l.
<i>G. andina</i>	C Chile. ca. 2300 m a. s. l.
<i>G. antarctica</i>	Falkland Islands (=Islas Malvinas)
<i>G. antillana</i>	S Brazil, Cuba, Uruguay, *USA, *Europe, *New Zealand
<i>G. argentina</i>	NE, C Argentina, S. Brazil, Uruguay
<i>G. argyrinea</i>	USA, West Indies, *Hawaii. 0–250 m a. s. l.
<i>G. axillaris</i>	C Chile
<i>G. badillana</i>	Venezuela. ca. 3000 m a. s. l.
<i>G. beckii</i>	Bolivia. ca. 3800 m a. s. l.
<i>G. berteroana</i>	C Chile
<i>G. cabrerae</i>	N and C Peru. 3100–4870 m a. s. l.
<i>G. calviceps</i>	Central and South America, *USA, *Australia, *New Zealand, *Europe, *Asia, *S Africa
<i>G. camaquaensis</i>	S Brazil
<i>G. capitata</i>	NW Argentina, Bolivia, Peru. 2000–3400 m a. s. l.
<i>G. chamissonis</i>	C and S Argentina, Chile. 1000–1800 m a. s. l.
<i>G. chionesthes</i>	USA
<i>G. coarctata</i>	Cosmopolitan. 0–3200 m a. s. l.
<i>G. depilata</i>	S Chile, SW Argentina. 500–2000 m a. s. l.
<i>G. deserticola</i>	NW Argentina and N Chile. 3000–4000 m a. s. l.
<i>G. diffusa</i>	S Brazil
<i>G. erecta</i>	S Brazil, N Uruguay
<i>G. erythractis</i>	NW Argentina, Bolivia, Peru. 3000–4000 m a. s. l.
<i>G. falcata</i>	NE Argentina, S Brazil, Paraguay, Uruguay
<i>G. filaginea</i>	NE Argentina, Brazil, Uruguay
<i>G. foliosa</i>	C Chile
<i>G. girardiana</i>	S Brazil
<i>G. grazielae</i>	SE Brazil
<i>G. hiemalis</i>	S and SE Brazil. 1400–1800 m a. s. l.
<i>G. humilis</i>	Bolivia, Chile, Peru. 3560–4000 m a. s. l.
<i>G. irazuensis</i>	Costa Rica. 3030–3430 m a. s. l.
<i>G. longipedicellata</i>	NW Argentina, Bolivia, Chile. 3500–4250 m a. s. l.
<i>G. lulioana</i>	Bolivia, Peru. 4500 m a. s. l.
<i>G. meridensis</i>	Venezuela
<i>G. monticola</i>	S Argentina, C Chile. 400–600 m a. s. l.
<i>G. neuquensis</i>	Argentina, S Chile. 780–1150 m a. s. l.
<i>G. nigrevestis</i>	S and SE Brazil, more than 1000 m a. s. l.
<i>G. nivalis</i>	SW Argentina. C and S Chile, 800–3000 m a. s. l.
<i>G. oligantha</i>	C Chile
<i>G. paramora</i>	Venezuela, probably Colombia. 2900–3500 m a. s. l.
<i>G. pensylvanica</i>	Cosmopolitan
<i>G. peregrina</i>	N and C Argentina
<i>G. procumbens</i>	C and S Chile. ca. 2000 m a. s. l.
<i>G. purpurea</i>	Cosmopolitan. 0–4300 m a. s. l.

Table 3 continued

Genera/species	Distribution (*adventive or naturalized)
<i>G. rizzini</i>	S and SE Brazil
<i>G. serpyllifolia</i>	Argentina, C and S Chile. 700–3200 m a. s. l. Chile
<i>G. simplicicaulis</i>	South America. *Australia, *New Zealand, *USA, *Hawaii. 0–3700 m a. s. l.
<i>G. sphacelata</i>	North and South America. ca. 2200 m a. s. l.
<i>G. spiciformis</i>	SW Argentina, S Chile. 250–1500 m a. s. l.
<i>G. stachydifolia</i>	Argentina, Brazil, Chile. Uruguay. 400–1800 m a. s. l.
<i>G. stagnalis</i>	Guatemala, Mexico, USA. 1000–2500 m a. s. l.
<i>G. standleyi</i>	Guatemala. 3500–4300 m a. s. l.
<i>G. subfalcata</i>	Argentina, Brazil, Chile, Uruguay. *New Zealand, *southern Africa. Up to 4100 m a. s. l.
<i>G. suffruticosa</i>	Chile
<i>G. thouarsii</i>	Tristan da Cunha
<i>G. ustulata</i>	USA, Canada, 0–1100 m a. s. l.
<i>G. valparadisea</i>	C Chile. 0–60 m a. s. l.
<i>G. villarroelii</i>	C Chile. 750–1200 m a. s. l.

random addition sequences, swapped using TBR and holding two optimal trees per series. Consistency index (CI; Kluge and Farris 1969) and retention index (RI; Farris 1989) were calculated as measures of homoplasy.

For each of the four markers, an appropriate model of evolution was selected with JModeltest v.2.1.1 (Darriba et al. 2012) based on the Akaike information criterion, AIC (Akaike 1973; Sugiura 1978; Hurvich and Tsai 1989). The best models were GTR + I+G for ITS; HKY + G for ETS and GTR + G for *trnL*-F. Bayesian inference was performed as implemented in Beast version 1.8.1 (Drummond et al. 2012). With BEAUTi v.1.6.2 (Drummond and Rambaut 2007) we created the input file with the nucleotide substitution models mentioned above, empirical base frequencies, four gamma categories, under an uncorrelated lognormal relaxed-clock model (Drummond et al. 2006), and a Yule process of speciation as prior. The MCMC analysis was performed for 10,000,000 generations and sampled every 1000th generation. Convergence of the chains was checked using Tracer v.1.5 (Drummond and Rambaut 2007). All trees obtained prior to convergence were discarded, and trees were summarized in a maximum clade credibility tree in TreeAnnotator v.1.6.2 (Drummond and Rambaut 2007). Trees were visualized and edited using FigTree version 1.4.2 (Rambaut 2014).

Analyses were performed for (1) plastid data only (*trnL*-F), (2) nuclear data only (ITS + ETS), and (3) combined nuclear + plastid data.

Results

Matrices

Sequences of the plastid *trnL*-F region were obtained for 47 taxa, including 27 *Gamochaeta* species, *Gamochaetopsis alpina*, *Stuckertiella capitata* (Wedd.) Beauverd, and ten other genera, *Achyrocline tomentosa* serving as the outgroup. The total length of the sequence ranged from 599 bp in *Gamochaeta grazielae* to 754 bp in *G. andina*, *G. serpyllifolia*, and *Gamochaetopsis alpina*. The aligned matrix consisted of 779 characters, of which 24 were parsimony informative.

Sequences of the nuclear ETS region were obtained for 55 taxa, including 33 *Gamochaeta* species, *Gamochaetopsis alpina* and *Stuckertiella capitata* and 20 other species, including the outgroup. The total length of the sequence ranged from 356 bp in *Leontopodium* to 474 bp in *Omalotheca norvegica* and *Antennaria chilensis*. The aligned matrix consisted of 482 characters, of which 112 were parsimony informative.

Sequences of the nuclear ITS region were obtained for 45 taxa, including 25 *Gamochaeta* species, *Gamochaetopsis alpina* and *Stuckertiella capitata* and 17 other species plus *Achyrocline tomentosa*. The total length of the sequence ranged from 516 bp in *Gamochaeta americana* to 527 bp in *Gnaphalium uliginosum*. The aligned matrix consisted of 532 characters, of which 92 were parsimony informative.

Parsimony and Bayesian analyses showed highly congruent results, with the latter providing more resolved nodes. Consensus trees obtained from the parsimony analyses of nuclear and nuclear + plastid data are illustrated, but some of the more resolved nodes from Bayesian analysis are also shown.

Relationships

The analysis of the plastid marker resulted in a highly polytomized consensus tree (CI 0.87, RI 0.95) due to low character state variability, showing little relevant information. Furthermore, the relationships do not correspond to any proposed hypotheses and even though they are poorly supported we decided to show the data obtained (Fig. 1). The combined analysis of the nuclear markers (hereafter referred to as nuclear analysis) resulted in 175 most parsimonious trees (CI 0.60, RI 0.79), and the combined analysis of the three markers (hereafter referred to as the combined analysis) resulted in 209 most parsimonious trees (CI 0.61, RI 0.79).

The hypothesis of relationships obtained from the nuclear and combined analyses shows that *Gamochaeta*, as currently circumscribed, is paraphyletic, since *Gamochaetopsis* and *Stuckertiella* also appear nested within *Gamochaeta* with 98/97 Jackknife support and a posterior probability of 1.00 (Figs. 2, 3).

The combined data, under Bayesian analysis, show the pair *Omalotheca norvegica*-*O. sylvatica* (maximum values) distant from *Gamochaeta*, forming a group with *Omalotheca supina* (Fig. 3) but with low support (54/0.91).

In the nuclear and combined analyses, *Gamochaeta* species are divided into two major groups: clade A (99/1.00) including *Gamochaetopsis alpina*, *Gamochaeta affinis*, *Gamochaeta affinis*, *Gamochaeta andina*, *Gamochaeta depilata*, *Gamochaeta nivalis*, *Gamochaeta procumbens*, *Gamochaeta serpyllifolia*, and *Gamochaeta spiciformis*; and clade B (*/0.86, 65/1.00) including all the remaining *Gamochaeta* species (Figs. 2, 3).

Clade A (Fig. 3) is principally characterized by having capitula arranged in head-like clusters (vs. clade B with capitula usually arranged in spikes). In the nuclear analysis *Stuckertiella* is placed as a basal member in the clade B (*/ 0.84, Fig. 2), and in the combined analysis *Stuckertiella* is placed as a basal member in the clade A (52/0.99, Fig. 3).

In both analyses, clade B showed some internal resolution, with the following clades present: *Gamochaeta ambatensis* + *G. falcata* (85/1.00 Fig. 2; 85/1.00 Fig. 3), principally characterized by having capitula with 50–70 marginal florets, and linear-oblanceolate or oblanceolate leaves (Table 2); *G. filaginea* + *G. pensylvanica* (61/1.00 Fig. 2, 61/1.00 Fig. 3), principally





Fig. 1 Strict consensus tree for plastid (*trnL-F*) data obtained under maximum parsimony. Numbers above branches are JK values, numbers below branches Bayesian PP.

characterized by having capitula with 70–80 marginal florets, acute or subacute to obtuse inner phyllaries, and oblanceolate or spatulate leaves (Table 2); *G. argyrinea* + *G. chionesthes* (61/0.98 Fig. 2; 62/0.99 Fig. 3b), principally characterized by having capitula with up 120 marginal florets, and discolorous, oblanceolate leaves (Table 2); *G. humilis* + *G. longipedicellata* + *G. lulioana* (63/0.93 Fig. 2, 63/0.94 Fig. 3c), principally characterized by having caespitose or subcaespitose habit, capitula with 27–40 marginal florets, acute inner phyllaries, and oblanceolate or oblong-ovate to oblong-spatulate leaves (Table 2); and *G. camaquaensis* + *G. erecta* + *G.*

Fig. 2 Strict consensus tree for nuclear (ETS + ITS) data obtained under maximum parsimony. Numbers above branches are JK values, numbers below branches Bayesian PP




grazielae + *G. stachydifolia* (75/1.00 Fig. 2, 71/1.00 Fig. 3), principally characterized by having capitula with 50–100 marginal florets, acute to acuminate inner phyllaries, and concolorous, oblanceolate to spathulate leaves (Table 2). Other small clades were also present within clade B in each analysis (Figs. 2, 3), e.g., *G. americana* clade (Fig. 3), which comprises *G. americana*, *G. coarctata* and *G. ustulata*, all with strongly discolored and broadly obovate to spathulate leaves.

The clade *Belloa chilensis*, *Berroa gnaphalioides*, *Facelis plumosa* and *Lucilia acutifolia* is placed as sister

group of clades A and B (53/0.88 Fig. 2) in both the nuclear and combined analyses (51/0.88 Fig. 3). They also conform to two morphologically well-defined groups. The genera of the clade of *Belloa chilensis*, *Berroa gnaphalioides*, *Facelis plumosa* and *Lucilia acutifolia* are defined by solitary or few together capitula and elongated (rarely clavate) twin hairs. Conversely, the clade A + B is defined by capitula arranged in head-like clusters or in spiciform inflorescences, and globose (sometimes clavate) twin hairs.

Finally, *Gamochaetopsis alpina* and *Stuckertiella capitata* appear nested within *Gamochaeta* in both analyses

Fig. 3 Strict consensus tree from combined molecular (ETS + ITS + trnL-F) data. Numbers above branches are JK values from the parsimony analysis, and numbers below branches Bayesian posterior probabilities (PP, asterisk indicates lack support). **a–c** Bayesian topologies. **Bold** lines indicate *Gamochaeta* and related genera



with Bayesian posterior probabilities and jackknife of 98 and 97, respectively (Figs. 2 and 3).

Discussion

The analyses affirm that *Gamochaeta* as currently circumscribed is paraphyletic, indicating the need of a revised circumscription at the generic level in the group.

Taxonomic position of *Stuckertiella* and *Gamochaetopsis* (Figs. 2, 3, 6)

Stuckertiella was described by Beauverd (1913) with two species: *Stuckertiella capitata* transferred from *Gamochaeta* and *S. peregrina*. The monotypic genus *Gamochaetopsis* was established by Anderberg and Freire (1991) to include *Gamochaetopsis alpina* from southern Chile and Argentina, which was principally diagnosed by

its achenes with short clavate twin hairs and capitula arranged in head-like clusters. The position of *Stuckertiella* and *Gamochaetopsis* in our analyses is congruent with morphological evidence. All the morphological characters that define *Stuckertiella* and *Gamochaetopsis* are also found in species of *Gamochaeta* (Table 1) with the exception of the autoapomorphy in *Stuckertiella* of the presence of functionally male central florets with four anthers (three with a small obtuse apical appendage and one with a long, acute apical appendage). The close similarity between *Gamochaetopsis* and *Gamochaeta*, and between *Stuckertiella* and *Gamochaeta* had previously been noted by Cabrera (1971) and Anderberg (1991), respectively. Furthermore, *Stuckertiella* shares with *Gamochaetopsis* clade its capitula arranged in head-like clusters, and with the remaining species of *Gamochaeta* its capitula with many florets (Table 1).

Unfortunately, we were unable to obtain living material of *Stuckertiella peregrina*, and herbarium specimens were not of sufficient quality for DNA extractions. However, we predict that this species will also group with *Gamochaeta*, given that it has character states like *Stuckertiella capitata* that unite this group with *Gamochaeta*.

Redefinition of *Gamochaeta* boundaries

(Figs. 2, 3, 4, 5, 6)

Gamochaeta is taxonomically difficult due to the fact that most species exhibit considerable morphological and ecological variability (Table 2). In this work, we included a broad geographical-taxonomic sampling, including species that grow at middle elevation (less than 3000 m), especially in dry hills, grasslands, sand hills, and disturbed habitats characterized by having ascending or erect stems, or more rarely, cespitose habit (e.g., *G. alpina*, *G. depilata*, *G. nivalis*, *G. procumbens*, *G. serpyllifolia*, and *G. spiciformis*). A small number of species (e.g., *G. deserticola*, *G. erythractis*, *G. meridensis*, *G. meridensis*, *G. paramora*) grow at high altitudes (3000–4000 m) in paramo, jalea and puna; and a few prostrate and acaulescent species (e.g., *G. cabrerae*, *G. humilis*, *G. longipedicellata*, *G. lulioana*) are adapted to life at even higher elevations between 3500–4500 m.

In order to have *Gamochaeta* monophyletic, we propose that the two species of *Stuckertiella* and the single species of *Gamochaetopsis* are transferred to *Gamochaeta* (Table 3). Synapomorphies for this large clade include the following: multistemmed perennial herbs, oblanceolate leaves, small capitula arranged in spikes or head-like clusters, marginal female florets outnumbering the hermaphroditic central florets, style branches truncate and penicillate, short-pilose achenes with globose twin hairs, and pappus bristles basally connate.

The low sequence divergence and lack of resolution within the *Gamochaeta* clade is probably due to a rapid and recent diversification in the Andes, as was postulated by Hughes and Eastwood (2006). In this way, two centers of diversity in the Andes could be suggested, one in the southern portion of South America and the second one mainly in the Andean-Brazilian region, with few species reaching North America.

Omalotheca (Figs. 2, 3)

Cassini published *Omalotheca* in Cuvier's Dictionnaire des Sciences Naturelles in 1828 to include the Eurasian *Gnaphalium supinum* L. and distinguished it from *Gnaphalium* based on a uniseriate pappus and obovoid, compressed achenes. Drury (1970) also distinguished *Omalotheca* from *Gnaphalium* and *Gamochaeta* based on terminal inflorescences in a leafy spikes and elongate twin hairs not emitting mucilage in water.

The molecular phylogeny we present here shows *Omalotheca supina*, *O. norvegica* and *O. sylvatica* forming a monophyletic group (Figs. 2, 3) distantly related to *Gamochaeta* and to *Gnaphalium uliginosum* (which is the generic type of *Gnaphalium*). This was pointed out by Galbany-Casals et al. (2010), who showed that *Gnaphalium supinum* was neither closely related to *Gnaphalium* s.str. nor to *Gamochaeta*. Similarly, Blöch et al. (2010) indicated that *Omalotheca* (sub *Gnaphalium*) was distantly related to *Gamochaeta*. The diagnostic characters for *Omalotheca* are the presence of connate pappus bristles (vs. free in *Gnaphalium*), lack of myxogenic twin hairs on the fruits (vs. usually myxogenic in *Gamochaeta*), spiciform capitulecence (vs. corymbs in *Gnaphalium*) and pistillate flowers with corollas filiform-tubular (vs. filiform in *Gamochaeta* and *Gnaphalium*). Further, as was pointed out by Nesom (1990b), *Gamochaeta* is strictly a New World genus.

Gamochaeta nanchuanensis (Y.Ling & Y.Q.Tseng) Y.S.Chen & R.J.Bayer was not included in our present analyses, so its phylogenetic position remains untested, although the original description of *Gnaphalium nanchuanense* Y.Ling & Y.Q.Tseng (Ling and Tseng 1978) states that it is very similar to *G. sylvaticum*, thus indicating that it, too, likely belongs in *Omalotheca*.

Taxonomic treatment

Gamochaeta Wedd., Chlor. And. 1(4–6): 151. 1856.
Gnaphalium L. subgen. *Gamochaeta* (Wedd.) Gren., Fl. Jurass. 2: 427. 1869. ≡ *Gnaphalium* L. sect. *Gamochaeta* (Wedd.) Benth. & Hook. f., Gen. Pl. 2(1): 306. 1873. ≡ *Gnaphalium* L. sect. *Gamochaeta* (Wedd.) O.Hoffm., Nat.




Fig. 4 Achenial hairs in *Gamochaeta*. **a** globose achenial hairs (*G. peregrina*, Cabrera 8009, SI), **b** clavate achenial hairs (*G. hiemalis*, Deble and Deble 4501, SI), **c** globose myxogenic achenial hairs (*G. americana*, Urtubey 187, SI), **d** globose achenial hairs (*G. girardiana*,

Deble and *Deble* 4501, SI), **e** style branches with pollen of disk floret (*G. americana*, Urtubey 187, SI), **f** clavate achenial hairs (*G. alpina*, Cabrera 5902, LP). Scale bars **a** 20 μ m, **b** 20 μ m, **c** 20 μ m, **d** 40 μ m, **e** 40 μ m, **f** 20 μ m


Pflanzenfam. 4(5): 188. 1894, **comb. illeg.** (Art. 53)—LECTOTYPE (designated by Cabrera 1961: 362): *G. americana* (Mill.) Wedd.

= *Stuckertiella* Beauverd, **syn. nov.** Bull. Soc. Bot. Genève sér. 2, 5: 205. 1913. —LECTOTYPE (designated by Cabrera, 1978: 128): *S. capitata* (Wedd.) Beauverd.

= *Gamochaetopsis* Anderb. & S.E.Freire, **syn. nov.** Bot. J. Linn. Soc. 106(2): 186. 1991—TYPE: *G. alpina* (Poepp.) Anderb. & S.E.Freire.

Sixty-one species distributed from the USA to Argentina: *G. affinis* Cabrera [*Gnaphalium affine* d'Urv., **nom. illeg.** (Art. 53); *G. malvinense* H.Koyama, **nom. illeg.** (Art. 52); *Gamochaeta malvinensis* (H.Koyama) T.R.Dudley, **comb.**

Fig. 5 Morphological diversity in *Gamochaeta* and allied genera: **a** *G. lulioana*, capitula solitary (photo S. Beck), **b** *G. longipedicellata*, capitula solitary (photo E. Urubey), **c** *G. stachydifolia*, capitula arranged in spikes (photo F.O. Zuloaga), **d** Spike basally discontinuous of *G. stachydifolia* (photo F.O. Zuloaga), **e** *G. coarctata*, capitula arranged in spikes (photo Pensiero), **f** *G. filaginea*, capitula arranged in spikes (photo F.O. Zuloaga)






Fig. 6 Capitula arranged in head-like clusters **a** *Gamochaeta spiciformis* (photo L. Zavala), **b** *Gamochaeta nivalis* (photo M.C. Baeza), **c** *Gamochaeta alpina* (photo N. Bayón), **d** *Gamochaeta capitata* (photo E. Urtubey)

illeg. (Art. 53)]; *G. aliena* (Hook. & Arn.) Cabrera (*Gnaphalium alienum* Hook. & Arn.); ***Gamochaeta alpina*** (Poepp.) S.E.Freire & Anderb., **comb. nov.** \equiv *Laennecia alpina* Poepp., Nov. Gen. Sp. Pl. 3: 56, tab. 262. 1845 [= *Lucilia alpina* (Poepp.) Cabrera, *Gamochaetopsis alpina* (Poepp.) Anderb. & S.E.Freire]; *G. ambatensis* Ariza; *G. americana* (Mill.) Wedd. [*Gnaphalium americanum* Mill.]; *G. purpureum* L. var. *americanum* (Mill.) Klatt; *G. consanguineum* Gaudich.; *G. guatemalense* Gand.; *Gamochaeta guatemalensis* (Gand.) Cabrera]; *G. andina* (Phil.) Cabrera (*Gnaphalium andinum* Phil.); *G. antarctica* (Hook.f.) Cabrera (*Gnaphalium antarcticum* Hook.f.); *G. antillana* (Urb.) Anderb. (*Gnaphalium antillanum* Urb.); *G. argentina* Cabrera; *G. argyrinea* G.L.Nesom; *G. axillaris* (J.Rémy) Cabrera (*Gnaphalium axillaris* J.Rémy); *G. badillana* (Aristeg.) Anderb. (*Gnaphalium badillanum* Aristeg.); *G. beckii* Urtubey & S.E.Freire; *G. berteroana* (DC.) Cabrera [*Gnaphalium berteroanum* DC.; *G. stachydifolium* Lam. var. *berteroanum* (DC.) DC.]; *G. cabrerae* Anderb.

[*Gamochaeta monticola* M.O.Dillon & Sagástegui, **nom. illeg.** (Art. 53); *G. oreophila* M.O.Dillon & Sagástegui, **nom. illeg.** (Art. 53)]; *G. calviceps* (Fernald) Cabrera (*Gnaphalium calviceps* Fernald); *G. camaquaensis* Deble; *Gamochaeta capitata* Wedd., Chlor. And. 1: 153. 1855 [= *Stuckertiella capitata* (Wedd.) Beauverd, *Gnaphalium capitatum* (Wedd.) Griseb., **comb. illeg.** (Art. 53), non Lamarck, 1786, nec Thunberg, 1799, *Gnaphalium weddelianum* Rusby, *Gamochaeta weddelliana* (Rusby) Anderb., **nom. illeg.** (Art. 52)]; *G. chamissonis* (DC.) Cabrera [*Gnaphalium chamissonis* DC.; *G. fernandezianum* Phil.; *G. julietii* Phil.; *G. polybotryum* Phil.; *G. purpureum* L. var. *julietii* (Phil.) Reiche; *G. serranoi* Phil.; *Gamochaeta fernandeziana* (Phil.) Anderb.; *G. julietii* (Phil.) Anderb.; *G. polybotrya* (Phil.) Cabrera; *G. serranoi* (Phil.) Cabrera]; *G. chionesthes* G.L.Nesom; *G. coarctata* (Willd.) Kerguélen [*Gnaphalium coarctatum* Willd.; *G. purpureum* L. var. *spicatum* Klatt; *G. spicatum* Lam., **nom. illeg.** (Art. 53); *Gamochaeta spicata* Cabrera]; *G. depilata* (Phil.) Cabrera

(*Gnaphalium depilatum* Phil.; *G. obscurum* Phil.); *G. deserticola* Cabrera; *G. diffusa* Deble & Marchiori; *G. erecta* Deble; *G. erythractis* (Wedd.) Cabrera [*Merope erythractis* Wedd.]; *Belloa erythractis* (Wedd.) Cabrera; *Gnaphalium erythractis* (Wedd.) Griseb.]; *G. falcata* (Lam.) Cabrera [*Gnaphalium falcatum* Lam., *G. purpureum* L. var. *falcatum* (Lam.) Torr. & A. Gray; *G. stachydifolium* Lam. var. *falcatum* (Lam.) Klatt]; *G. filaginea* (DC.) Cabrera (*Gnaphalium filagineum* DC.); *G. foliosa* (Phil.) Anderb. [*Gnaphalium foliosum* Phil.; *Gamochaeta chilensis* Deble, **nom. illeg.** (Art. 53)]; *G. girardiana* Deble & A.S.Oliveira; *G. grazielae* (Rizzini) Deble (*Gnaphalium grazielae* Rizzini); *G. hiemalis* Cabrera [*Gnaphalium hiemale* Rizzini, **nom. illeg.** (Art. 53); *Gamochaeta brasiliiana* Deble]; *G. humilis* Wedd.; *G. irazuensis* G.L.Nesom; *G. longipedicellata* Cabrera; *G. lulioana* S.E.Freire & Iharl.; *G. meridensis* V.M.Badillo; *G. monticola* (Phil. ex Reiche) Cabrera (*Gnaphalium monticola* Phil. ex Reiche); *G. neuquensis* Cabrera; *G. nigrevestis* Deble & Marchiori; *G. nivalis* Cabrera [*Gnaphalium affine* d'Urv. var. *pusillum* Speg.; *G. nivale* Phil., **nom. illeg.** (Art. 53); *Gamochaeta spiciformis* (Sch.Bip.) Cabrera var. *subaffinis* Cabrera]; *G. oligantha* (Phil.) L.E.Navas (*Gnaphalium oliganthum* Phil.); *G. paramora* (S.F.Blake) Anderb. [*Gnaphalium paramorum* S.F.Blake; *Lucilia paramora* (S.F.Blake) V.M.Badillo]; *G. pensylvanica* (Willd.) Cabrera [*Gnaphalium pensylvanicum* Willd.; *G. peregrinum* Fernald; *G. purpureum* L. subsp. *pensylvanica* (Willd.) O.Bolòs & Vigo; *G. purpureum* L. var. *spathulatum* Baker; *G. spathulatum* Lam., **nom. illeg.** (Art. 53); *G. platense* Cabrera; *Gamochaeta platensis* (Cabrera) Cabrera]; ***Gamochaeta peregrina*** (Beauverd) S.E.Freire & Anderb., **comb. nov.** = *Stuckertiella peregrina* Beauverd, Bull. Soc. Bot. Genève sér. 2, 5: 208. 1913. [= *Stuckertiella peregrina* Beauverd var. *fusca* Beauverd, *Stuckertiella peregrina* Beauverd var. *albida* Beauverd]; *G. procumbens* (Phil.) Cabrera (*Gnaphalium procumbens* Phil.); *G. purpurea* (L.) Cabrera [*Gnaphalium purpureum* L.; *G. rosaceum* I.M. Johnst.; *Gamochaeta rosacea* (I.M.Johnst.) Anderb.]; *G. rizzini* Cabrera; *G. serpyllifolia* Wedd. [*Gnaphalium serpyllifolium* J.Rémy, **nom. illeg.** (Art. 53); *Gamochaeta munnozii* Cabrera, **nom. illeg.** (Art. 52)]; *G. simplicicaulis* (Willd. ex Spreng.) Cabrera [*Gnaphalium simplicicaule* Willd. ex Spreng.; *G. purpureum* L. var. *simplicicaule* (Willd. ex Spreng.) Klatt]; *G. sphacelata* (Kunth) Cabrera [*Gnaphalium sphacelatum* Kunth; *G. purpureum* L. var. *sphacelatum* (Kunth) Speg.]; *G. spiciformis* (Sch.Bip.) Cabrera [*Gnaphalium spiciforme* Sch.Bip.]; *G. affine* d'Urv. var. *medium* Speg.; *G. mucronatum* Phil.; *G. ptereoanum* Phil.; *Gamochaeta ptereoana* (Phil) Anderb.]; *G. stachydifolia* (Lam.) Cabrera [*Gnaphalium stachydifolium* Lam.]; *G. purpureum* L. var. *stachydifolium* (Lam.) Baker]; *G. stagnalis* (I.M.Johnst.) Anderb. (*G. stagnale* I.M.Johnst.); *G. standleyi* (Steyermark) G.L.Nesom

(*Gnaphalium standleyi* Steyermark); *G. subfalcata* (Cabrera) Cabrera (*Gnaphalium subfalcatum* Cabrera); *G. suffruticosa* (Phil.) Anderb. (*Gnaphalium suffruticosum* Phil.); *G. thouarsii* (Spreng.) Anderb. (*Gnaphalium thouarsii* Spreng.); *G. ustulata* (Nutt.) Holub [*Gnaphalium ustulatum* Nutt.; *G. pannosum* Gand.; *G. purpureum* L. var. *ustulatum* (Nutt.) B.Boivin; *Gamochaeta ustulata* (Nutt.) G.L.Nesom]; *G. valparadisea* (Phil.) Anderb. (*Gnaphalium valparadiseum* Phil.); *G. villarroelii* (Phil.) Cabrera (*Gnaphalium villarroelii* Phil.).

Concluding remarks

Our study, using a plastid marker (*trnL-F*) and two nuclear markers (ETS and ITS), suggested that the genus *Gamochaeta* is paraphyletic and that *Stuckertiella* and the monotypic genus *Gamochaetopsis* should be included for the establishment of the monophyly in the genus. This American genus *Gamochaeta* is diagnosed by having slender pappus bristles that are basally connate, central florets hermaphroditic (rarely functionally male), capitula arranged in terminal head-like clusters or more usually crowded into spiciform inflorescences, and achenes pilose (occasionally glabrous), with short globose or sometimes clavate, duplex hairs. Two new combinations are proposed under *Gamochaeta*: *Gamochaeta alpina* (Poepp.) S.E.Freire & Anderb. and *Gamochaeta peregrina* (Beauverd) S.E.Freire & Anderb., and *Gamochaeta capitata* Wedd. is resurrected from synonymy of *Stuckertiella capitata* (Wedd.) Beauverd.

Acknowledgments We express our thanks to the reviewers for critical reading of the manuscript; to S. Beck (LPB, Bolivia) for his assistance during field work in Bolivia; to F. Zuloaga (SI, Argentina) and A. Tribsch (SZU, Austria) who provided material from their own collections; to L. Aagesen for comments on early drafts of the manuscript; to the curators of CONC, LP, LPB, MNCS, MO, SI, SZU, USMS and W for providing materials; to F. Zuloaga and S. Beck for permission to use their photographs included in Fig. 6; J. Hurst for technical assistance. This work was supported by Consejo Nacional de Investigaciones Científicas y Técnicas for financial support (PIP 112-200801-02196) to S. E. Freire and FONDECYT No. 1150425 to A. Moreira Muñoz.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Appendix

Species used in the molecular studies with voucher information and GenBank accession numbers (ETS, ITS and *trnL-F*). Accession number in bold indicates data obtained

from GenBank. * indicates sequence not obtained for that marker.

Achyrocline tomentosa. KM091358, KM091389, KM091420. *Antennaria chilensis* var. *magellanica*. KM091373, KM091395, *. *Antennaria dioica*. FN645610, FJ639964, FN645790. *Belloa chilensis* KM091349, KM091388, KM091430. *Berroa gnaphalioides*. KM091355, KM091386, KM091418. *Facelis plumosa*. KM091372, KM091394, KM091428. *Filago fuscescens*. FN645579, FN645846, FN645764. *Filago lutescens*. FN645596, FN645883, FN645779. *Filago pyramidalis*. FN645589, FN645872, HM364535. *Gamochaeta affinis*. Argentina. Tierra del Fuego: Ushuaia, Bahía Valentín, 16 Oct 1971, Dudley et al. 303 (MO). KX078016, *, *. *Gamochaeta alpina*. (under *Gamochaetopsis alpina*). KM091356, KM091390, KM091417. *Gamochaeta ambatensis*. Argentina. Jujuy: Valle Grande, de San Francisco a Alto de Calilegua, 21 Feb 2008, Zuloaga et al. 10324 (SI). KX078017, *, KX078049. *Gamochaeta americana*. KM091382, KM091411, KM091437. *Gamochaeta andina*. Chile. Maule: Maule, Lueberg and Teiller 2290 (CONC). KX078018, KX077990, KX078050. *Gamochaeta antillana*. USA. Mississippi: Lamar County, Alford 3892 (USMS). KX078019, KX077991, KX078051. *Gamochaeta argyrinea*. USA. Nesom WMGT8 (LP). KX078020, *, *. *Gamochaeta beckii*. Bolivia. La Paz: Nor Yungas, camino a Coroico, entrada del camino viejo hacia la Mina Lourdes, 30 Mar 2010, Urtubey et al. 508 (SI). KX078021, KX077992, KX078052. *Gamochaeta calviceps*. Argentina. Entre Ríos: Colón, camino al arroyo El Palmar, 51 m a. s. l., 7 Dec 2008, Urtubey and Bazzarri 384 (SI). KX078022, *, *. *Gamochaeta camaquensis*. Brasil. Rio Grande do Sul: Cacapava do Sul, 30 Sep 2009, Deble and Deble 11879 (SI). KX078023, KX077993, KX078053. *Gamochaeta capitata* (under *Stuckertiella capitata*) KM091369, KM091398, KM091412. *Gamochaeta chamissonis*. Chile. Metropolitana: El Yeso, 1283 m a. s. l., 26 Jan 2012, Urtubey and Freire 713 (SI). KX078024, KX077994, KX078054. *Gamochaeta chionesthes*. USA. Georgia: 15 Apr 2001, Nesom GASCO4-26 (LP). KX078025, KX077995, KX078055. *Gamochaeta coarctata*. Argentina. Neuquén: Minas, de Chos Malal a Andacollo, ca. 40 km junto al arroyo El Manzanito, 1443 m a. s. l., 12 Jan 2009, Urtubey et al. 409 (SI). KX078026, KX077996, KX078056. *Gamochaeta depilata*. Chile. Biobío: Mihoc et al. 4092 (CONC). KX078027, KX077997, KX078057. *Gamochaeta deserticola*. Bolivia. La Paz: Murillo, 4100 m a. s. l., 4.8 km al NE del autopista por el camino subiendo el Valle del Río Kaluyo 28 Feb 1987, Solomon 16209 (LPB). KX078028, *, KX0780580. *Gamochaeta erecta*. Brasil. Rio Grande do Sul: Cacapava do Sul, 28 Oct 2009, Deble and Deble 11874 (SI). KX078029, KX077998, KX078060. *Gamochaeta erythractis*. Argentina. Jujuy: Humahuaca, camino a la Mina Aguilar, 3940 m a. s. l., 23 Mar 2009, Urtubey and Freire 442 (SI). KX078030, KX077999, KX078061. *Gamochaeta falcata*. Argentina. Córdoba: Calamuchita, desde Yacanto de Calamuchita al Champaquí, ca. 2100 m a. s. l., 18 Jan 2006, Urtubey and Bazzarri 197 (SI). *, KX078000, *. Uruguay. Colonia: Conchillares, playa, 18 Nov 2010, Urtubey 520 (SI). KX0780310, *, *. *Gamochaeta filaginea*. Argentina. Buenos Aires: Isla Martín García, 24 Sep 2000, Hurrell 4363 (LP). KX078032, *, *. *Gamochaeta grazielae*. Brasil. Río de Janeiro: Nov 2006, Deble and Deble 6018 (SI). KX078033, KX078001, KX078062. *Gamochaeta humilis*. Bolivia. La Paz: Nor Yungas, camino a Coroico, bajando de la ruta en la entrada del camino viejo hacia la Mina Lourdes, 3919 m a. s. l., 30 Mar 2010, Urtubey et al. 506 (SI). KX078034, KX078002, KX078059. *Gamochaeta longipedicellata*. Bolivia. La Paz: Murillo, 24 Mar 2010, Urtubey et al. 473 (SI). KM091381, KM091410, KX078063. *Gamochaeta lulioana*. Bolivia. La Paz: Murillo, en el borde de bofedal, 4661 m a. s. l., 24 Mar 2010, Urtubey et al. 480 (SI). KX078035, KX078003, KX078064. *Gamochaeta nivalis*. Argentina. Río Negro: Bariloche, Cerro Catedral, 16 Jan 2009, Urtubey et al. 415 (SI). KX078036, *, KX078065. *Gamochaeta pensylvanica*. Argentina. Buenos Aires: La Plata, ciudad Vieja, 6 Oct 2012, Urtubey 755 (SI). *, *, KX078066. USA. Mississippi: Lamar County, Alford 4156 (USMS). KX078037, KX078004, *. *Gamochaeta procumbens*. Argentina. Tierra del Fuego: Rio Grande, Estancia Cullén, 6 Jan 1971, Goodall 3118 (Herb. Goodall). KX078038, KX078005, *. *Gamochaeta purpurea*. USA, Nesom WMGT13 (LP) KX078039, KX078006, KX078067. *Gamochaeta serpyllifolia*. Chile. Los Ríos: Valdivia, Comuna de Panguipulli, faldeos Volcán Chosuenco, 39°55'71°59', 1558 m a. s. l., 5 Feb 2012, Baeza 4357 (CONC). KM091380, KX078010, KX078068. *Gamochaeta simplicicaulis*. Argentina. Jujuy: Tilcara, camino de Molulo a Huairahuasi, 3100 m a. s. l., 11 Feb 2010, Zuloaga et al. 11702 (SI). KX078040, KX078007, KX078069. *Gamochaeta sphacelata*. Argentina. Salta: Santa Victoria, Nazareno, 3308 m a. s. l., Adler 4 (MNCS 1665). KX078041, *, KX078070. *Gamochaeta spiciformis*. Chile. Magallanes: Domínguez 984 (CONC). *, KX078011, KX078071. Argentina. Tierra del Fuego: Ushuaia, Brown Sawmill, Goodall 286 (LP). KX078042, *, *. *Gamochaeta stachydifolia*. Argentina. Buenos Aires: Tandil, hacia sierras de Las Animas desde el Hotel Elegance, 12 Nov 2005, Urtubey and Bazzarri 179 (SI) KX078043, *, *. *Ibid*. Frente al hotel Elegance, 1 Oct 2010, Urtubey 513 (SI). *, KX078008, KX078072. *Gamochaeta subfalcata*. Uruguay. Canelones: La Floresta, 19 Nov 2010, Urtubey 524 (SI). KX078044, KX078009, KX078073. *Gamochaeta ustulata*. USA. Oregon: Lincoln

County, *Shannon Straub* 201 (USMS). KX078045, KX078012, KX078074. ***Gnaphalium austroafricanum*. FN645630, FN645830, FN645756.** *Gnaphalium declinatum* FR821617, *, FR822634. *Gnaphalium uliginosum* Austria. Upper Austria: Innviertel, Reichersberg am Inn, Hübing, 22 Aug 2010, *Pflugbeil* 936 (SZU). KX078048, KX078015, KX07807077. *Jalcophila boliviensis*. KM091370, KM091402, KM091429. *Jalcophila ecuadorensis*. KM091383*,*. *Leontopodium alpinum* FM173135, GU943413, AF141821 + AF141733. *Leontopodium microphyllum*. FJ640015, FJ639947, FJ640047 + FJ639977. *Lucilia acutifolia*. KM091374, KM091396, KM091432. *Omalotheca norvegica* (Gunnerus) Sch.-Bip. & F.W.Schultz. Russia. Republic Altai: surroundings of Zeminsky (Zeminskij) pass to area NE of the pass, ca. 105 km S Gorno Altaysk (Altajsk), subalpine Siberian pine forest, 1700-1730 m a. s. l., 11 Aug 2003, *Tribsch* and *Essl* 10383 (W). KX078047, KX078013, KX078075. *Omalotheca supina* FN645558, AY445230, FN645789. *Omalotheca sylvatica*. Austria. Carinthia: Lavanttal, Koralpe, Hartelsberg SE Wolfsberg, 1345 m a. s. l., *Riegler-Hager* 921 (W). KX078046, KX078014, KX078076.

References

- Akaike H (1973) Information theory and an extension of the maximum likelihood principle. In: Petrov BN, Csaki F (eds) Second international symposium on information theory. Akadémiai Kiado, Budapest, pp 267–281
- Alford MH (2012) New records of *Gamochaeta* (Asteraceae) in the Hawaiian Archipelago. In: Evenhuis NL, Eldredge LG (eds), Records of the Hawaii Biological Survey for 2011. Bishop Mus Occas Pap 113:1–6
- Anderberg AA (1991) Taxonomy and phylogeny of the tribe Gnaphalieae (Asteraceae). *Opera Bot* 104:1–195
- Anderberg AA, Freire SF (1991) A cladistics and biogeographic analysis of the *Lucilia* group (Asteraceae, Gnaphalieae). *Bot J Linn Soc* 106:173–198
- Aristeguieta L (1964) Compositae. In: Lasser T (ed) Flora de Venezuela 10(1). Dirección de Recursos Naturales Renovables MAC, Caracas, pp 1–483
- Baldwin BG, Markos S (1998) Phylogenetic utility of the external transcribed spacer (ETS) of 18S–26S rDNA: congruence of ETS and ITS trees of *Calycadenia* (Compositae). *Molec Phylogen Evol* 10:449–463
- Bayer RJ, Greber DG, Bagnall NH (2002) Phylogeny of Australian Gnaphalieae (Asteraceae) based on four sequences, the *trnL* intron, *trnL/trnF* intergenic spacer, matK, and ETS. *Syst Bot* 27:801–814
- Bayer RJ, Breitwieser I, Ward J, Puttock C (2007) Tribe Gnaphalieae. In: Kadereit JW, Jeffrey C (eds) The families and genera of vascular plants, flowering plants—Eudicots: Asterales, vol 8. Springer, Berlin, pp 246–283
- Beauverd G (1913) Contribution à l'étude des Composées 8(1). Le genre *Stuckertia* Beauverd. *Bull Soc Bot Genève* 2:205–209
- Bentham G, Hooker JD (1873) Compositae. In: Bentham G, Hooker JD (eds) Genera plantarum 2. Lovell Reeve & Co, London, pp 163–533
- Blöch C, Dickoré WB, Samuel R, Stuessy TF (2010) Molecular phylogeny of the edelweiss (*Leontopodium*, Asteraceae-Gnaphalieae). *Edinburgh J Bot* 67:235–264
- Cabrera AL (1961) Observaciones sobre las Inuleae-Gnaphalineae (Compositae) de América del Sur. *Bol Soc Argent Bot* 9:359–386
- Cabrera AL (1963) Compuestas: *Gamochaeta*. In: Cabrera AL (ed), Flora de la Provincia de Buenos Aires, Colecc Ci Inst Nac Tecnol Agropecu 4(6), Buenos Aires, pp 166–178
- Cabrera AL (1971) Compuestas: *Gamochaeta*. In: Correa MN (ed), Flora Patagónica, Colecc Ci Inst Nac Tecnol Agropecu 8(7), Buenos Aires, pp 117–127
- Cabrera AL (1974) Compuestas: *Gamochaeta*. In: Burkart A (ed), Flora Ilustrada de Entre Ríos, Colecc Ci Inst Nac Tecnol Agropecu 6(6), Buenos Aires, pp 319–327
- Cabrera AL (1978) Compuestas: *Gamochaeta*. In: Cabrera AL (ed), Flora de la Provincia de Jujuy, Colecc Ci Inst Nac Tecnol Agropecu 13(10), Buenos Aires, pp 302–311
- Chen YS, Bayer RA (2011) *Gamochaeta*. In: Wu ZY, Raven PH, Hong DY (eds), Flora of China, Fam 276 Asteraceae, vol. 20–21, Science Press and Missouri Bot. Gard. Press, Beijing, & St. Louis, pp 776–778
- Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nature Meth* 9:772. doi:[10.1038/nmeth.2109](https://doi.org/10.1038/nmeth.2109)
- Deble LP, Marchiori JNC (2007) Sinopse do género *Gamochaeta* Weddell (Asteraceae-Gnaphalieae) no Brasil. *Balduinia* 10:21–31
- Dillon MO, Sagástegui-Alva A (1991a) Sinopsis de los géneros de Gnaphaliinae (Asteraceae-Inuleae) de Sudamérica. *Arnaldoa* 1:5–91
- Dillon MO, Sagástegui-Alva A (1991b) *Gamochaeta*. In: Macbride JF & collab (eds), Flora of Peru, Asteraceae: Part V. Fieldiana, Bot, n.s. 26:27–32
- Doyle JJ, Dickson EE (1987) Preservation of plant samples for DNA restriction endonuclease analysis. *Taxon* 36:715–722
- Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol Biol* 7:214. doi:[10.1186/1471-2148-7-214](https://doi.org/10.1186/1471-2148-7-214)
- Drummond AJ, Ho SYW, Phillips MJ, Rambaut A (2006) Relaxed phylogenetics and dating with confidence. *PLoS Biol* 4:e88. doi:[10.1371/journal.pbio.0040088](https://doi.org/10.1371/journal.pbio.0040088)
- Drummond AJ, Suchard MA, Xie D, Rambaut A (2012) Bayesian phylogenetics with BEAUTi and the BEAST 1.7. *Molec Biol Evol* 29:1969–1973
- Drury DG (1970) A fresh approach to the classification of the genus *Gnaphalium* with particular reference to the species present in New Zealand (Inuleae-Compositae). *New Zealand J Bot* 8:222–248
- Drury DG (1971) The American spicate cudweeds adventive to New Zealand (*Gnaphalium* Section *Gamochaeta*-Compositae). *New Zealand J Bot* 9:157–185
- Farris JS (1989) The retention index and the rescaled consistency index. *Cladistics* 5:417–419. doi:[10.1111/j.1096-0031.1989.tb00573.x](https://doi.org/10.1111/j.1096-0031.1989.tb00573.x)
- Farris JS, Albert V, Källersjö M, Lipscomb D, Kluge A (1996) Parsimony jackknifing outperforms neighbor-joining. *Cladistics* 12:99–124
- Freire SE (1998) Inuleae. Compositae. In: Bocquet GF, Crosby MR (eds) Flora del paraguay, vol 27. Conservatoire et Jardin botaniques de la Ville de Genève-Missouri Botanical Garden, Geneve & St Louis, pp 1–100

- Freire SE, Iharlegui L (1997) Sinopsis preliminar del género *Gamochaeta* (Asteraceae, Gnaphalieae). *Bol Soc Argent Bot* 33:23–35
- Freire SE, Iharlegui L (2008) Asteraceae: *Gamochaeta*. In: Zuloaga FO, Morrone, O, Belgrano, MJ (eds), *Catálogo de las Plantas Vasculares del Cono Sur*, Monogr Syst Bot Miss Bot Gard, Missouri Botanical Garden, St Louis 107:1307–1312
- Freire SE, Iharlegui L (2014) *Gamochaeta*. In: Anton, AM, Zuloaga, FO, Belgrano, MJ (eds), Freire, SE (coord), *Flora Vascular Argentina, Asteraceae: Anthemideae-Gnaphalieae*, 7(1). Estudio Sigma, Buenos Aires, pp 463–482
- Freire SE, Paz Deble L, Iharlegui L (2011) Tribe Inuleae. In: Reis A (ed) *Flora Ilustrada Catarinense. Herbário Barbosa Rodrigues, Itajai*, pp 1067–1197
- Freire SE, Chemisquy MA, Anderberg AA, Beck SG, Meneses RI, Loeuille B, Urtubey E (2015) The *Lucilia* group (Asteraceae, Gnaphalieae): phylogenetic and taxonomic considerations based on molecular and morphological evidence. *Pl Syst Evol* 301:1227–1248. doi:[10.1007/s00606-014-1147-0](https://doi.org/10.1007/s00606-014-1147-0)
- Galbany-Casals M, Andres-Sanchez S, Garcia-Jacas N, Susanna A, Rico E, Martinez-Ortega MM (2010) How many of Cassini anagrams should there be? Molecular systematics and phylogenetic relationships in the *Filago* group (Asteraceae, Gnaphalieae), with special focus on the genus *Filago*. *Taxon* 59:1671–1689
- Goloboff PA, Farris JS, Nixon KC (2008) TNT, a free program for phylogenetic analysis. *Cladistics* 24:774–786
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Series* 41:95–98
- Hilliard OM (1983) *Gamochaeta* (sub *Gnaphalium*). In: Leistner OA (ed) *Flora of southern Africa* 33(7) Asteraceae-Inuleae 2-Gnaphaliinae 1. Botanical Research Institute, Pretoria, pp 27–29
- Hilliard OM, Burtt BL (1981) Some generic concepts in Compositae—Gnaphaliinae. *Bot J Linn Soc* 82:181–232
- Hind DJN (2011) An annotated preliminary checklist of the Compositae of Bolivia. version 2. Available at: <http://www.kew.org/science/tropamerica/boliviacompositae>. Accessed 11 May 2012
- Hoffmann O (1890) Compositae-Inuleae. In: Engler A, Prantl K (eds) *Die Naturlichen Pflanzenfamilien* 4(5). Wilhelm Engelmann, Leipzig, pp 172–210
- Hughes CE, Eastwood RJ (2006) Island radiation on a continental scale: exceptional rates of plant diversification after uplift of the Andes. *Proc Natl Acad Sci USA* 103:10334–10339. doi:[10.1073/pnas.0601928103](https://doi.org/10.1073/pnas.0601928103)
- Hurvich CM, Tsai CL (1989) Regression and time series model selection in small samples. *Biometrika* 76:297–307. doi:[10.1093/biomet/76.2.297](https://doi.org/10.1093/biomet/76.2.297)
- Kluge AG, Farris SJ (1969) Quantitative phyletics and the evolution of anurans. *Syst Zool* 18:1–32. doi:[10.2307/2412407](https://doi.org/10.2307/2412407)
- Ling Y, Tseng YQ (1978) Miscellaneous notes on Chinese compositae. *Acta Phytotax Sin* 16:82–86
- Marticorena C, Quezada M (1985) Catálogo de la flora vascular de Chile. *Gayana, Bot* 42:5–157
- Merxmüller H, Leins P, Roessler H (1977) Inuleae-systematic review. In: Heywood VH, Harborne JB, Turner BL (eds) *The Biology and Chemistry of the Compositae*. Academic Press, London, pp 577–602
- Nesom GL (1990a) Taxonomic status of *Gamochaeta* (Asteraceae: Inuleae) and the species of the United States. *Phytologia* 68:186–198
- Nesom GL (1990b) Taxonomic summary of *Omalotheca* (Asteraceae: Inuleae). *Phytologia* 68:241–246
- Nesom GL (2004) New species of *Gamochaeta* (Asteraceae: Gnaphalieae) from the eastern United States and comments on similar species. *Sida* 21:717–742
- Nesom GL (2006) *Anaphalis*, *euchiton*, *facelis*, *gamochaeta*, *gnaphalium*, *omalotheca*, *pseudognaphalium*, *xerochrysum* (Gnaphalieae). In: *Flora of North America Editorial Committee* (ed) 1993+. *Flora of North America North of Mexico*, vol 19. Oxford University Press, New York, pp 415–442
- Pruski JF (2011) Asteraceae: *Gamochaeta*. In: Davids G, Sousa Sánchez M, Knapp S, Chiang Cabrera F (eds), *Flora Mesoamericana* 5(2). Missouri Botanical Garden Press, México DF, pp 598–610
- Rambaut A (2014) Tree Figure Drawing Tool, version 1.4.2, Institute of Evolutionary Biology, University of Edinburgh
- Rémy EJ (1849) Compuestas. In: Gay C (ed), *Historia física y política de Chile. Botánica* 4, en casa del autor, Paris, pp 5–317
- Samarakoon T, Wang SY, Alford MH (2013) Enhancing PCR amplification of DNA from recalcitrant plant specimens using a trehalose-based additive. *Appl Plant Sci* 1:1200236. doi:[10.3732/apps.1200236](https://doi.org/10.3732/apps.1200236)
- Smissen RD, Galbany-Casals M, Breitwieser I (2011) Ancient allopolyploidy in the everlasting daisies (Asteraceae: Gnaphalieae): complex relationships among extant clades. *Taxon* 60:649–662
- Sugiura N (1978) Further analysis of the data by Akaike's information criterion and the finite corrections. *Commun Stat- Theory Methods* A7:13–26
- Taberlet P, Gielly L, Pautou G, Bouvet J (1991) Universal primers for amplification of three non-coding regions of chloroplast DNA. *Pl Molec Biol* 17:1105–1109. doi:[10.1007/BF00037152](https://doi.org/10.1007/BF00037152)
- Wagenitz G (1965) Compositae (Korbblütler) In: Hegi G (ed), *Illustrierte Flora von Mitteleuropa* 2. Augflage, 6(3) Lieferung 2, Carl Hanser Verlag, München
- Ward J, Bayer RJ, Breitwieser I, Smissen R, Galbany-Casals M, Unwin M (2009) Gnaphalieae. In: Funk VA, Susanna A, Stuessy TF, Bayer RJ (eds) *Systematics, evolution, and biogeography of compositae*. International Association for Plant Taxonomy, Vienna, pp 539–588
- Webb CJ (1988) Asteraceae: *Gnaphalium*. In: Webb CJ, Sykes WR, Garnock-Jones PJ (eds) *Flora of New Zealand*, vol. 4, Botany Division D.S.I.R., Christchurch, pp 234–249
- Weddell HA (1855) *Gamochaeta* pars 4–6, pp 151–154 in Chloris Andina, P. Bertrand, Paris, pp 473–484
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Shinsky JJ, White TJ (eds) *PCR protocols: a guide to methods and applications*. Academic Press, San Diego, pp 315–322