

A new species of *Bufonia* L. (Caryophyllaceae)
from the Kordestan province (Iran)

Farrokh GHAHREMANINEJAD, Zahra ROSTAMI GHADI,
Mostafa ASSADI & Atiye NEJAD FALATOURY

DIRECTEUR DE LA PUBLICATION : Bruno David
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF*: Thierry Deroin

RÉDACTEURS / *EDITORS*: Porter P. Lowry II; Zachary S. Rogers

ASSISTANTS DE RÉDACTION / *ASSISTANT EDITORS*: Emmanuel Côté (adanson@mnhn.fr) ; Anne Mabilie

MISE EN PAGE / *PAGE LAYOUT*: Emmanuel Côté

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD*:

P. Baas (Nationaal Herbarium Nederland, Wageningen)
F. Blasco (CNRS, Toulouse)
M. W. Callmander (Conservatoire et Jardin botaniques de la Ville de Genève)
J. A. Doyle (University of California, Davis)
P. K. Endress (Institute of Systematic Botany, Zürich)
P. Feldmann (Cirad, Montpellier)
L. Gautier (Conservatoire et Jardins botaniques de la Ville de Genève)
F. Ghahremaninejad (Kharazmi University, Téhéran)
K. Iwatsuki (Museum of Nature and Human Activities, Hyogo)
K. Kubitzki (Institut für Allgemeine Botanik, Hamburg)
J.-Y. Lesouef (Conservatoire botanique de Brest)
P. Morat (Muséum national d'Histoire naturelle, Paris)
J. Munzinger (Institut de Recherche pour le Développement, Montpellier)
S. E. Rakotoarisoa (Millenium Seed Bank, Royal Botanic Gardens Kew, Madagascar Conservation Centre, Antananarivo)
É. A. Rakotobe (Centre d'Applications des Recherches pharmaceutiques, Antananarivo)
P. H. Raven (Missouri Botanical Garden, St. Louis)
G. Tohmé (Conseil national de la Recherche scientifique Liban, Beyrouth)
J. G. West (Australian National Herbarium, Canberra)
J. R. Wood (Oxford)

COUVERTURE / *COVER*:

Bufonia iranica Z.Rostami, Assadi & F.Ghahrem., sp. nov. From holotype: *Mozaffarian 70066*.

Adansonia est indexé dans / *Adansonia is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Agriculture, Biology, and Environmental Sciences®
- Scopus®

Adansonia est distribué en version électronique par / *Adansonia is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Adansonia est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Adansonia is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*:

Geodiversitas, Zoosystema, Anthropozoologica, European Journal of Taxonomy, Naturae, Cryptogamie sous-sections *Algologie, Bryologie, Mycologie*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2019
ISSN (imprimé / *print*): 1280-8571/ ISSN (électronique / *electronic*): 1639-4798

A new species of *Bufonia* L. (Caryophyllaceae) from the Kordestan province (Iran)

Farrokh GHahremaninejad

Department of Plant Sciences, Faculty of Biological Sciences, Kharazmi University, Postal
Code 15719-14911, Tehran (Iran)
ghahremaninejad@khu.ac.ir (corresponding author)
<https://orcid.org/0000-0001-5860-9976>

Zahra ROSTAMI GHADI

Department of Plant Sciences, Faculty of Biological Sciences, Kharazmi University, Postal
Code 15719-14911, Tehran (Iran)
rostamighadii@gmail.com

Mostafa ASSADI

Research Institute of Forests and Rangelands,
Agricultural Research Education and Extension Organization (AREEO),
P.O. Box 13185-1166, Tehran (Iran)
assadi@rifr-ac.ir
<https://orcid.org/0000-0003-3522-3717>

Atiye Nejad FALATOURY

Iranian Research Institute of Plant Protection, Agricultural Research,
Education and Extension Organization (AREEO), postal code: 19858-13111 Tehran (Iran)
a.falatoury@areeo.ac.ir
<https://orcid.org/0000-0001-9014-585X>

Submitted on 8 March 2019 | accepted on 23 May 2019 | published on 21 October 2019

Ghahremaninejad F., Rostami Ghadi Z., Assadi M. & Falatoury A. N. 2019. — A new species of *Bufonia* L. (Caryophyllaceae) from the Kordestan province (Iran). *Adansonia*, sér. 3, 41 (13): 177-182. <https://doi.org/10.5252/adansonia-a2019v41a13>. <http://adansonia.com/41/13>

ABSTRACT

During investigation of the genus *Bufonia* L., in various herbaria of Iran, a new species, *B. iranica* Z.Rostami, Assadi & F.Ghahrem., sp. nov., was recognized and thus is described here. This new species is closely related to *B. enervis*, with some morphological differences such as size and shape of the perianth, and the seed shape. A key for the species occurring in Iran, illustrations and a distribution map of the new and related species are provided. The seed morphology was studied using a scanning electron microscope.

KEY WORDS

Alsinoideae,
Flora of Iran,
Sagineae,
seed,
new species.

RÉSUMÉ

Une espèce nouvelle de Bufonia L. (Caryophyllaceae) de la province iranienne du Kurdistan.

Au cours de l'examen du genre *Bufonia* L. dans différents herbiers d'Iran, une nouvelle espèce (*B. iranica* Z.Rostami, Assadi & F.Ghahrem., sp. nov.) a été reconnue et est ici décrite. Elle est étroitement apparentée à *B. enervis* Boiss., avec quelques différences dans la taille et la forme du périanthe, ainsi que dans la forme de la graine. Une clé des espèces du genre en Iran, des illustrations et une carte de répartition sont fournies. La morphologie de la graine a été observée au microscope à balayage.

MOTS CLÉS

Alsinoideae,
Flore d'Iran,
Sagineae,
graine,
espèce nouvelle.

INTRODUCTION

Since the publishing of *Flora Iranica's* treatment for the genus *Bufonia* L., the present work is the first taxonomic study on the Iranian species of *Bufonia*. The genus belongs to the family Caryophyllaceae, subfamily Alsinoideae (Rechinger 1988). The genus *Bufonia* is taxonomically close to the genera *Cerastium* L. and the genus *Sagina* L. In more recent phylogenetic studies *Bufonia*, *Colobanthus* Bartl., *Sagina*, and *Drypis spinosa* L. are placed in a clade that represents tribe Sagineae sensu Harbaugh (Harbaugh *et al.* 2010; Greenberg & Donoghue 2011).

Diagnostic morphological characteristics of this genus are having flowers with four sepals, four tepals and a bivalve compressed lenticular capsule. The genus is morphologically such a complex taxon that finding the proper characters for accurately separating species is difficult. During our work on this genus in several Iranian herbaria, it appears that many specimens, especially from W Iran, were mistakenly determined at the species level. *Bufonia* includes nearly 35 species in the world (Içim & Behçet 2013), it is widely distributed from Macaronesia through Turkey, and Iran to Afghanistan; the northern boundaries of the distribution range are in the western Iberian Peninsula, central France and southern Switzerland and east Mediterranean region.

It continues from Serbia to Macedonia, Romania to the southern Ukraine. In Africa, it occupies lands of Maghreb, in Libya but it is not present in the African part of Egypt. Southern limits of the Asian range are Sinai Peninsula, Jordan, Syria, mountainous parts of Iraq to Iran. At the eastern edge, it grows in southern Pakistan, reaching Turkmenistan and Transcaucasian republics of Armenia and Azerbaijan in the north. The highest species richness is found in Iran and NW Africa. In Europe, the genus is represented by a few species only, often of rare occurrence (Italy, Greece). Its complete absence in some Mediterranean islands is striking (e.g. Sardinia, Sicily, Cyprus). Most species grow on dry stony slopes up to alpine area (e.g. Morocco and Iran above 3000 m) (Chrtek & Křisa 1999a). The number of species decreases to the east.

According to *Flora Iranica*, *Bufonia* includes 14 species in Iranian plateau of which 13 grow in Iran, 10 of them are endemics for Iran (Rechinger 1988). Based on newer studies, *B. koelzii* Rech.f., an endemic species for Iran, was transferred to genus *Arenaria* as *A. polycnemifolia* Boiss. because of having pentamerous perianth and six-part splitting capsule (Chrtek & Křisa 1999b). According to molecular and anatomical evidences, *B. koelzii* is a synonym of *Eremogone polycnemifolia* Boiss. (Sadeghian *et al.* 2015; Mousavi *et al.* 2019). Additionally, *B. kotschyana* Boiss. was

A NEW IDENTIFICATION KEY FOR *BUFONIA* L. SPECIES IN IRAN

- 1. Annual 2
— Perennial, woody at base 3
- 2. Inflorescences are often limited to the upper part of the stem, stamens 3-4 *B. parviflora* Grisb.
— The flowers are all along the stem, stamens 8 *B. oliveriana* Ser.
- 3. Inflorescence not congested, pedicels usually longer than sepals, up to 30 mm long 4
— Inflorescence congested, pedicels shorter up to 10 mm long 6
- 4. Pedicel filiform 5
— Pedicel not filiform *B. capsularis* Boiss. & Hausskn.
- 5. Sepals 2-3 mm long, ovate-orbulate, apiculate, ovule 4 *B. macrocarpa* Ser.
— Sepals 2.5-4 mm long, ovate-lanceolate, acute, ovule 4-8 *B. sintenisii* Freyn.
- 6. Sepals puberulent or velutinous 7
— Sepals glabrous 9
- 7. Pedicel glabrous, sepals up to 6 mm long *B. iranica* Z.Rostami, Assadi & F.Ghahrem., sp. nov.
— Pedicel pubescent, sepals up to 4 mm long 8
- 8. Stems branched, sepals velutinous *B. hebecalyx* Boiss.
— Stems usually not branched, sepals glabrous or puberulent *B. enervis* Boiss.
- 9. Inflorescences with 5-12 flowers *B. capitata* Bornm.
— Inflorescences with up to 5 flowers 10
- 10. Sepals 2-3 mm long *B. micrantha* Boiss. & Hausskn.
— Sepals longer 11
- 11. Sepals ± 6 mm long *B. calycina* Boiss. & Hausskn.
— Sepals 3- 5.5 mm long 12
- 12. Ovules 2 *B. stapfii* Bornm.
— Ovules 4 *B. kotschyana* Boiss.



FIG. 1. — *Bufonia iranica* Z.Rostami, Assadi & F.Ghahrem., sp. nov. From holotype: Mozaffarian 70066 (TARI).




FIG. 2. — **A-E**, Details of morphological characters of *Bufonia iranica* Z.Rostami, Assadi & F.Ghahrem., sp. nov. (*Mozaffarian 70066*): **A**, calyx; **B**, upper part of sepal; **C**, mature ovary; **D**, mature open ovary; **E**, petal; **F, G**, *Bufonia enervis* Boiss. (*Iranshahr & Termeh 12159-E*): **F**, calyx; **G**, petal; **H-M**, *B. iranica* Z.Rostami, Assadi & F.Ghahrem., sp. nov.: **H**, seed; **I**, close view of the seed surface from median part of the seed at the position between hilum and abfunicular side; **J, K**, close views of the seed surface at the abfunicular side; **L**, close view of the seed surface from median part of the seed at the hilum; **M**, close view of anticlinal border between testa cells. Scale bars: A, D, E, 1.2 mm; B, 0.38 mm; C, 0.97 mm; F, 0.87 mm; G, 0.82 mm; H, 1 mm; I-L, 100 μ m; M, 10 μ m.

recorded from Syria (Post 1932), but was not yet mentioned in *Flora Iranica*. Most of the nine endemic species for Iran are narrow endemic taxa. The most species of *Bufonia* are distributed in the west and north west of Iran. However, some species are also distributed in the mountainous areas of the northeast and center of Iran. The distribution of this genus in Iran is often limited to over 1500 meters in the Irano-Touranian region.

LIST OF IRANIAN SPECIES OF *BUFONIA*

- B. macrocarpa* Ser.; distribution: Iran. Endemic.
- B. sintenisii* Freyn.; dist.: Iran, Afghanistan, Turkmenistan.
- B. capsularis* Boiss. & Hausskn; dist.: Iran. Endemic.
- B. capitata* Bornm.; dist.: Iran. Endemic.
- B. hebecalyx* Boiss.; dist.: Iran. Endemic.
- B. iranica* Z.Rostami, Assadi & F.Ghahrem., sp. nov.; dist.: Iran. Endemic.




FIG. 3. — Distribution of *Bufo iranica* Z.Rostami, Assadi & F.Ghahrem., sp. nov. (■) and *Bufo enervis* Boiss. (●).

- B. enervis* Boiss.; dist.: Iran. Endemic.
B. stapfi Bornm.; dist.: Iran. Endemic.
B. kotschyana Boiss.; dist.: Iran, Syria.
B. calycina Boiss. & Hausskn.; dist.: Iran. Endemic.
B. micrantha Boiss. & Hausskn.; dist.: Iran. Endemic.
B. parviflora Griseb.; dist.: Anatolia, Syria, Lebanon, Jordan, Iran, Armenia, Azerbaijan, Europe Macedonia, Greece, Bulgaria, Ukraine, Serbia.
B. oliveriana Ser.; dist.: Iraq, Iran, Turkmenistan, Afghanistan.

MATERIAL AND METHODS

For this study, we used several important references for determination of the genus, like *Flora Orientalis* (Boissier 1867), *Flora of Turkey* (Cullen 1967), *Flora Europaea* (Halliday 1966), *Flora of the U.S.S.R.* (Schischkin 1936), *Flora Iranica* (Rechinger 1988), etc. Also several type specimens photos have been studied from BM, G, JE, P, S, W and WU herbaria. All the specimens of *Bufo*, nearly 230, preserved in five Iranian herbaria, i.e. FAR, FUMH, IRAN, T and TARI, have been determined.

The authors did a comparison between diagnostic characters of the new species and its most closely related species (Table 1). Most differences were related to reproductive structures (length and shape of outer and inner sepals, their margin width, length and shape of petal, its apex shape and seed size). Most similarities were related to plant inflorescence and growth form. Seed samples were scanned with a KYKY SEM, EM3200 model. Vouchers were scanned with Olympus SZH stereomicroscope equipped with Canon EoS 700 D camera.

SYSTEMATICS

Bufo iranica

Z.Rostami, Assadi & F.Ghahrem., sp. nov.
 (Figs 1; 2)

Species nova affinis *Bufo* *enervis* Boiss. sed differt a *pedicellis glabris* (nec *puberulis*), *sepalis late ovatis ad margines 0.5-1 mm late membranaceis* (nec *ovato-lanceolatis et minus quam 0.5 mm late membranaceis*), *4-6 mm longis* (nec *3-4 mm longis*), *petalis late ovatis, acutis, 5-5.5 mm longis* (nec *ovato-lanceolatis, obtusis, 2.5-3.5 mm longis*), *seminibus orbicularibus, 2 × 2 mm* (nec *late ellipticis, 2.5 × 1.5 mm*).

TYPUS. — Iran. Kordestan province: Saghghez, between Sonate and Divandarre, Kuh-e Ghaleh, between Bashmagh and Jafarabad, 2300 m, 14.VII.1991, V. Mozaffarian 70066 (holo-, TARI!).

SEED MORPHOLOGY. — The seeds of *B. iranica* Z.Rostami, Assadi & F.Ghahrem., sp. nov. are brown, orbicular with lateral hilum, and about 2 mm diameter. The surface has elongated testa cells on the wing and hilum of the seed (Fig. 2H-M). The periclinal wall of the testa cells is slightly convex. The anticlinal walls are Zip-shaped.

DISTRIBUTION. — This endemic species was collected from west of Iran, Kordestan province (Fig. 3). The locality of type specimen in label was erroneously mentioned as Azarbaiejan province, the other notes are accurate. It grows on dry rocky slopes of a mountainous area.

ETYMOLOGY. — The specific epithet refers to the country of the type specimen, Iran.

DESCRIPTION

Perennial, woody at base. Stems numerous, 12-25 cm high, erect, rigid, glabrous puberulent, simple or with short branches parallel to the stem; internodes 35-40 mm long. Basal leaves connate and sheathing, lower leaves 8-10 mm

TABLE 1. — Comparison of the new species with its close species.

Characters	<i>B. iranica</i> Z.Rostami, Assadi & F.Ghahrem, sp. nov.	<i>B. enervis</i>
Pediceal	Glabrous	Puberulent
Length sepals	4-6 mm	3-4 mm
Shape sepals	Broadly ovate	Ovate-lanceolate
Length petals	5-5.5 mm	2.5-3.5 mm
Shape petals	Ovate	Obovate
Seed shape	Orbicular	Elliptical

long, upper leaves shorter, subulate, rigid, adhesive to stem, ciliolate on the margin. Bracts 3 mm long, ovate-oblong, acute, ciliolate on the margin. Inflorescence compact. Flowers crowded in few flowered (2-5 flowers) terminal or lateral cymes, subsessile or very shortly pedicellate. Pedicels very short, rigid, glabrous. The outer pair sepals 4-5 mm long, broadly ovate, acute, 5-nerved, with white 0.5-0.7 mm wide scarious margin, straw yellow or black red. The inner pair sepals 5-6 mm long, broadly ovate, acute, 5-nerved, middle nerve excurrent, with 1 mm broad white scarious margin, straw color, pale purple or pallid purpurine or reddish, glabrous. Petals 5-5.5 mm long, broadly ovate, acute. Stamens 8, 3.5 mm long. Ovary with 4 ovules. Capsule compressed, 3.5 mm long, semi-elliptic. Seeds brown, 2 × 2 mm, orbicular, laterally compressed, with few tubercles. In terms of exomorphology, the surface of the seed is constructed of regularly arranged elongate cells with zip-shaped margins.

REMARKS

The new species is close to *Bufonia enervis* but it differs with having glabrous pedicels; longer sepals (4-6 not 3-4 mm long), and wider sepals (>0.5 not 0.5 mm wide) broadly ovate in shape (not ovate-lanceolate); longer petals (5-5.5 not 2.5-3.5 mm long), broadly ovate in shape (not obovate), and acute at the top (not obtuse); and different seeds (2 × 2 not 2.5 × 1.5 mm; orbicular not broadly elliptical). The type specimen was determined as *Arenaria* cf. *szowitzii* Boiss., but with having four sepals, four petals and bivalve compressed lenticular capsule, it definitely belongs to *Bufonia*.

Acknowledgements

The authors appreciate all the following herbaria for helping to access to the plant specimens or sending photo of the type material or critical specimens: BM, FAR, FUMH, G, IRAN,

JE, P, S, T, TARI, W, and WU. Special thanks to Dr John Edmondson (Kew, UK) for English editing of the manuscript. The two referees helped us in revising the manuscript through their useful comments, which are highly appreciated.

REFERENCES

BOISSIER E. 1867. — *Flora Orientalis*. Vol. 1. Basileae, Genevaae: 664-668. <https://doi.org/10.5962/bhl.title.20323>

CHRTEK J. & KRISA B. 1999a. — A revision of Asian species of the genus *Bufonia* L. *Acta Universitatis Carolinae-Biologica* 43 (2): 77-118.

CHRTEK J. & KRISA B. 1999b. — Taxonomic notes to the species of *Bufonia koelzii*. *Novitates Botanicae Universitatis Carolinae* 13: 35-38.

CULLEN J. 1967. — *Bufonia* L., in DAVIS P. H. (ed.), *Flora of Turkey and the East Aegean Islands*. Vol. 2. Edinburgh University Press, Edinburgh: 89, 90.

GREENBERG A. K. & DONOGHUE M. J. 2011. — Molecular systematics and character evolution in Caryophyllaceae. *Taxon* 60 (6): 1637-1652. <https://doi.org/10.1002/tax.606009>

HALLIDAY G. 1966. — *Bufonia* L., in TUTIN T. G. *et al.* (eds), *Flora Europaea*. Ed. 2, Vol. 1. Cambridge University Press, Cambridge: 160-161.

HARBAUGH D. T., NEPOKROEFF M., RABELER R. K., MCNEILL J., ZIMMER E. A. & WAGNER W. L. 2010. — A new lineage-based tribal classification of the family Caryophyllaceae. *International Journal of Plant Sciences* 171 (2): 185-198. <https://doi.org/10.1086/648993>

ILÇİM A. & BEHÇET L. 2013. — *Bufonia yildirimhanii* sp. nov. (Caryophyllaceae) from Turkey. *Nordic Journal of Botany* 31 (1): 84-86. <https://doi.org/10.1111/j.1756-1051.2012.01410.x>

POST G. E. 1932. — *Flora of Syria, Palestine and Sinai*. Vol. 1. Syrian Protestant College, Beirut: 192, 193. <https://doi.org/10.5962/bhl.title.20319>

RECHINGER K. H. 1988. — *Bufonia* L., in RECHINGER K. H. (ed.), *Flora Iranica*. Vol. 163. Akademische Druck- u. Verlagsanstalt, Graz: 114-119.

MOUSAVI S., PIRANI A. & ZARRE S. 2019. — Stem anatomy and its systematic implication in *Bufonia* (Caryophyllaceae, Saginaceae) and related genera. *Phytotaxa* 394 (2): 148-160. <https://doi.org/10.11646/phytotaxa.394.2.3>

SADEGHIAN S., ZARRE S., RABELER R. K. & HEUBL G. 2015. — Molecular phylogenetic analysis of *Arenaria* (Caryophyllaceae: tribe Arenarieae) and its allies inferred from nuclear DNA internal transcribed spacer and plastid DNA rps16 sequences. *Botanical Journal of the Linnean Society* 178 (4): 648-669. <https://doi.org/10.1111/boj.12293>

SCHISCHKIN B. K. 1936. — *Bufonia* L., in KOMAROV V. L. (ed.), *Flora of the U.S.S.R.* Vol. 6. Izdatel'stvo Akademii Nauk SSSR, Moscow, Leningrad: 475-479. <https://biodiversitylibrary.org/page/30223812>

Submitted on 8 March 2019;
 accepted on 23 May 2019;
 published on 21 October 2019.