

Generic and infrageneric classification in *Acacia* (Leguminosae: Mimosoideae): a list of critical species on which to build a comparative data set

Bruce R. Maslin^a and Charles H. Stirton^b

^a Western Australian Herbarium, Department of Conservation and Land Management, Locked Bag 104, Bentley Delivery Centre, W.A 6983, Australia (Corresponding author.)

^b The National Botanic Garden of Wales, Middleton Hall, Llanarthne, Carmarthenshire, Wales, UK, SA32 8HW

SUMMARY

Some recent proposals argue for the subdivision of the large pantropical genus *Acacia*. As a basis for making informed decisions on this important matter it is considered necessary that further broad-based, comparative studies be conducted. In order to assist researchers wishing to undertake such studies a list of "critical species", representative of many of the currently recognized major subgroups within the genus, is presented. Also, synoptic taxonomic notes are given for many of the currently recognized formal and informal infrageneric groups within *Acacia*.

INTRODUCTION

A number of recent papers have discussed or proposed the possible subdivision of *Acacia* (see Maslin 1988, 1989; Chappill and Maslin (1995); Pedley 1986, 1987a, 1987b). As currently conceived the genus comprises three large subgenera, *Acacia*, *Aculeiferum* and *Phyllodineae*, the first two pantropical in distribution and the third largely confined to the Australian region. A consensus now appears to be emerging that *Acacia* should be divided, but no agreement has been reached on where the dividing lines should be drawn. As discussed by Chappill and Maslin (1995) much of the problem appears to lie with the status of subgenus *Aculeiferum* and its relationship to subgenus *Phyllodineae*. What has become clear is that in many instances taxonomically relevant data are available for only a few species, for species in a restricted geographical area, or not at all. To fill these gaps in our knowledge, corroborative data should be obtained from as many representative species-groups and via as many disciplines as possible.

The genus *Acacia* comprises over 1200 species. Therefore, in undertaking comparative studies within a reasonable time it is not possible to examine all species. A practical solution is to choose representative species from each of the currently recognized major subgroups, bearing in mind that even the delimitation of many of these subgroups is still uncertain. Multidisciplinary studies of these "critical species" will assist in devising a more meaningful classification of *Acacia sensu lato* and/or of the three subgenera which comprise it. Additionally, the data will be valuable in assessing the relationship of these taxa to other genera within the Mimosoideae (especially tribes Mimoseae and Ingeae) which, hopefully, will lead to a resolution of the taxonomic status of the tribe Acacieae.

We therefore propose a list of "critical species" (Appendix 1 and 2), based on an evaluation by various experts on *Acacia* (see Acknowledgements) and urge those working on the genus to include as many of these taxa as possible in their studies. The "classification" used in Appendix

1 is in many ways a very pragmatic one (an over-view of the more commonly used formal classifications of *Acacia* is given in Table 1 of Maslin 1988 and Table 1 of Chappill and Maslin 1995). At the higher levels we adopt a combination of the schemes proposed by Vassal (1972) and Pedley (1978). Because of their high species numbers the Australian sections *Plurinerves* plus *Juliflorae*, and *Phyllodineae*, have been divided into informal sub-groups, namely, "microneurous" vs. "oligoneurous" (Cowan and Maslin 1990) and racemose vs. non-racemose respectively. Within each higher-order taxon the species are grouped according to their presumed taxonomic affinities.

The selection of species for inclusion on the "critical list" was based on the following criteria.

- The species in total should represent, as far as possible, the taxonomic variation within the genus.
- It should be possible to acquire experimental material of the nominated species with minimal cost and effort.
- Other things being equal, preference is given to taxa that have already been studied intensively using chemistry, pollen, ontogeny, etc.

It has not been possible, however, to include only species that meet all these criteria. Some taxa have been included on their scientific merit or because of their anomalous taxonomic placement. Other species have been included simply because collections have already been made and distributed to specialists for detailed study. Species not on the list are not precluded from study; if authors contact us, we will attempt to assign the taxa to their appropriate taxonomic subgroup. This is important because some researchers may already have material which could substitute for taxa we have listed. We emphasise that this is a preliminary list of species and undoubtedly other taxa will need to be added in due course. We hope that the proposal presented in this article will both stimulate further research and facilitate a more unified approach to *Acacia* systematics. We would therefore urge workers to consider the species included on our preliminary list of "critical species" and either contact us to effect changes or to publish modifications themselves in subsequent issues of this Bulletin.

Within *Acacia* a number of broad-based comparative research programs are currently being carried out on chemistry (proteinase inhibitors, DNA-sequencing, gums, and immunology), morphology and anatomy (spinescence, foliar anatomy, seed morphology, seedling anatomy, pollen) and systematics. However, there is a general consensus among workers that new comparative studies are needed (Maslin 1987) and the following have been suggested as desirable topics for research:

- molecular systematics (DNA markers and allozyme markers)
- inflorescence structure
- floral organogenesis
- lipid chemistry
- phyllode anatomy/morphogenesis
- legume anatomy
- legume morphology
- wood anatomy

TAXONOMIC NOTES ON ACACIA AND FAIDHERBIA

The following taxonomic notes are presented as an aid to researchers intending to work on *Acacia* and in order to provide a contextual framework for the list of "critical species" (Appendix 1). This is not an exhaustive discussion of the classification of *Acacia* but rather synoptic notes designed to introduce researchers to some of the more important recent taxonomic literature on *Acacia* and *Faidherbia* and to provide some insights into current taxonomic thinking, especially with respect to the main species-groups that are currently recognized within *Acacia*. Additional relevant information will be found in the references that we cite.

1. Genus *Faidherbia*

Until recently this monotypic African genus was treated as a species of *Acacia*, namely, *A. albida*. However, as summarized by Ross (1979) this taxon is not closely related to the other African species of *Acacia*. Also, the results of recent immunological studies by Brain (1987 and 1990) and cladistic analyses of Chappill and Maslin (1995) clearly indicate that this taxon should be removed from *Acacia*. Although *Faidherbia* was placed in tribe Acacieae by Vassal (1981) it has shortly united filaments and pollen characters similar to the *Enterolobium* - *Albizia* complex (*Samanea* group) within Ingeae (Guinet and Lugardon 1976, Guinet 1981 and 1990).

2. Genus *Acacia*

In recent years there have been many studies of the phylogeny and classification of *Acacia* (e.g. Brain 1987 and 1990, Conn *et al.* 1989, Evans *et al.* 1977, Guinet 1969, 1979 and 1990, Guinet and Vassal 1978, Maslin *et al.* 1988, Chappill and Maslin (1995), Pedley 1978, 1986 and 1987, Pettigrew and Watson 1975, Ross 1973, 1979 and 1981, Tindale and Roux 1969 and 1974, Vassal 1972, Vassal and Rouane 1987 and 1991). As currently circumscribed the genus includes more than 1200 species contained in three subgenera, namely, subgenus *Acacia*, subgenus *Aculeiferum* and subgenus *Phyllodineae* (syn. subgenus *Heterophyllum*). Although Pedley (1986) attributed generic rank to these taxa, namely, *Acacia*, *Senegalia* and *Racosperma* respectively, these have not been widely adopted. Pedley's proposal has widespread nomenclatural implications which is one of the reasons why it is not generally accepted. Another reason is that his assessment of *Acacia* did not include a comparative study of allied genera in tribes Ingeae and Mimoseae. This latter deficiency was addressed in a preliminary broad-based cladistic analysis of tribe Acacieae by Chappill and Maslin (1995) where it was shown that *Faidherbia* and subgenus *Acacia* were very different from subgenus *Aculeiferum* and subgenus *Phyllodineae*.

Global distributions for the three subgenera of *Acacia* are given in Ross (1981).

A. *Acacia* subgenus *Acacia*

This pan-tropical group includes approximately 120-130 species (Madsen 1990). Recent studies have shown that subgenus *Acacia* is distinct from the other two subgenera of *Acacia* and that it probably evolved independently of them, fide Guinet and Vassal (1978), Pedley (1986), Vassal (1988), Brain (1987 and 1990) and Chappill and Maslin (1995). However, for reasons

discussed in Maslin (1988) and Chappill and Maslin (1995) separate generic status for this taxon is not advocated at present.

Bentham (1875) divided subgenus *Acacia* into three groups based on the position of the involucl on the peduncle; however, this has proven an extremely variable character and is no longer considered suitable for discriminating infra-generic groups (Ross 1979). Although Britton and Rose (1928) divided the subgenus into 12 genera, primarily on the basis of fruit characteristics, most modern classifications do not recognize formal subordinate categories within subgenus *Acacia*. Vassal (1972), however, did divide subgenus *Acacia* section *Acacia* into two subsections, namely, *Pluriseriae* and *Uniseriae*, but his classification has not been generally adopted. Most African species of subgenus *Acacia* occur in the *Uniseriae* and, as discussed by Ross (1979), no ways have yet been found to meaningfully subdivide this group. In the New World, however, where the subgenus *Acacia* is represented by about 50 species (Lee *et al.*, 1989), seven reasonably distinct, albeit small, species-groups can be recognized, while in Asia another two groups occur. (NB. These nine groups do not account for all species of subgenus *Acacia* which occur in these two regions.)

New World Groups

(a) ***Acacia rigidula* group** (comprising *A. bilimekii*, *A. brandegeana*, *A. californica*, *A. pringlei* and *A. rigidula*, fide Lee *et al.* 1989 and Clarke 1995). These species are very distinct on account of their spicate inflorescences with 4-merous, white flowers; they also have a distinctive flavonoid chemistry (Seigler, pers. comm.) This group, together with the *A. constricta* group (see below) comprised *Acaciopsis* Britton and Rose, a "genus" recognized by a strip of papery endocarp on the inner surface of each legume valve. There is a superficial resemblance of several members of the *A. rigidula* group to species of *Pithecelobium*, especially *P. dulce*.

(b) ***Acacia constricta* group** (comprising *A. biaciculata*, *A. constricta*, *A. neovernicosa* and *A. schottii*, fide Clarke *et al.* 1990); *A. glandulifera*, a species included by Britton and Rose in the genus *Poponax* Raf. is included in the *Acacia constricta* group by Clarke (1995). *Acacia pacensis* appears to be intermediate between the *A. constricta* and *A. farnesiana* groups (Seigler, pers. comm.) although it was placed in the *A. farnesiana* group by Clarke (1995). Unlike other species of subgenus *Acacia* with globular heads, members of the *A. constricta* group sometimes have a single or scattered bracts on their peduncle instead of the usual multibracteate, whorled involucl; they also show distinctive flavonoid patterns (Seigler, pers. comm.).

(c) ***Acacia farnesiana* group** (comprising the New World species *A. albicorticata*, *A. atramentaria*, *A. caven*, *A. cupeyensis*, *A. farnesiana*, *A. pinetorum*, *A. polypyrgigenes*, *A. schaffneri*, *A. seifriziana*, *A. smallii* and *A. tortuosa*, fide Clarke *et al.* 1989 and Clarke 1995; also the African species *A. erioloba* and *A. haematoxylon*, fide Guinet 1990). In the past a number of these species were sometimes referred to the genus *Vachellia* Wight and Arn. or *Poponax* Raf. Guinet (loc. cit.) alluded to unusual pollen characters found in members of the *A. farnesiana* group; he also noted that most of these species have stamen filaments irregularly connate at the base (no stamen tube though), a floral involucl and +/- indehiscent pods.

(d) **Acacia acuífera group** (comprising *A. acuífera*, *A. anegadensis*, *A. barahonensis*, *A. belairioides*, *A. bucheri*, *A. caurina*, *A. cucuyo*, *A. daemon*, *A. roigii* and *A. zapatensis*, fide Clarke and Seigler 1991 and Clarke 1995). These species are restricted to the Caribbean and are characterized by low stamen number, fruit structure and a ramified stipular spine systems.

(e) **Acacia choriophylla group** (comprising only *A. choriophylla*, fide Clarke 1995). This species is restricted to the Caribbean. It was placed in the group containing *A. acuífera* by Clarke and Seigler (1991) but is now known not to be related to that group; perhaps related to the Ant *Acacia* group (Seigler, pers. comm.)

(f) **Ant Acacia group** (comprising *A. allenii*, *A. chiapensis*, *A. collinsii*, *A. cookii*, *A. cornigera*, *A. gentlei*, *A. gladiata*, *A. globulifera*, *A. hindsii*, *A. hirtipes*, *A. janzenii*, *A. mayana*, *A. melanoceras*, *A. oaxacanum*, *A. sphaerocephala*, *A. standleyi* and *A. ruddiae*, fide Janzen 1974, Clarke 1995 and Seigler pers. comm.).

(g) **Acacia macracantha group**. The three Central American non-Ant Acacias, *A. macracantha*, *A. pennatula* and *A. cochliacantha* are closely related to the Ant *Acacia* group, fide Ebinger and Seigler (1990) and Clarke (1995) and were among the 12 species included by Britton and Rose (1928) in the genus *Poponax*. The southern South American species, *A. aroma*, *A. huarango*, *A. rorudiana* and an undescribed species are also included in the *A. macracantha* group by Clarke (1995).

Asian Groups

Members of the following two groups differ most obviously from other Asian species of subgenus *Acacia* by their paniculate inflorescences, and, in the case of the *A. harmandeana* group at least, their low number of stamens per flower (10-15).

(a) **A. harmandeana group** (comprising *A. harmandeana*, *A. leucophloea* and *A. myaingii*). Inflorescences paniculate; involucre present; stipules spiny; stamens 10-15 per flower. The genus *Delaporteia* Gagnepain, which was segregated from *Acacia* on the basis of a small gland present at the top of the anther, is treated by Nielsen (1980) as a synonym of *A. harmandeana* (see also Guinet 1981). Nielsen noted that gland-bearing anthers occur in most other Asian acacias.

(b) **A. inopinata group** (comprising *A. craibii* and *A. inopinata*). Inflorescences paniculate; involucre present; stipules spiny. *Acacia inopinata* was originally described as *Nimiria inopinata*.

B. Acacia subgenus Aculeiferum

Subgenus *Aculeiferum* was established by Vassal (1972) and currently includes about 180-190 species (Madsen 1990) contained in three sections, namely, *Aculeiferum*, *Monacantha* and *Filicinae*. As shown in Table 1 of Maslin (1988) and Chappill and Maslin (1995) the subgenus corresponds to the combined series *Vulgares* and *Filicinae* of Bentham (1875). Species of this pan-tropical subgenus are widespread in both North and South America and in Africa; a small number of species are found in Asia and only one in Australia. As already noted, Pedley (1986) treated subgenus *Aculeiferum* as a distinct genus, *Senegalia*, but this change has not been widely adopted.

A number of recent studies have shown that subgenus *Aculeiferum* is more closely related to subgenus *Phyllodineae* than it is to subgenus *Acacia* with which it shares a common geographical area. Data from studies of pollen (Guinet 1990) and free amino acids (Evans *et al.* 1977) suggest that the differences between the two subgenera are relatively minor and this is supported by the results of a preliminary cladistic analysis by Chappill and Maslin (1995). This latter study, however, did not support the monophyly of subgenus *Aculeiferum*.

(a) Section *Aculeiferum*

Species of section *Aculeiferum*, which are confined to Africa and Asia, form a natural group, distinct from section *Monacantha* (Evans *et al.* 1977, Brain 1990, Chappill and Maslin 1995). These studies also showed *A. ataxacantha* (an African species which Vassal placed in section *Monacantha*) as having close affinities with species of this section; the sectional placement of this species should therefore be re-assessed. According to Ross (1979) all the African species of section *Aculeiferum* have spicate inflorescences and it is desirable to divide them into two groups, namely, those with prickles in pairs near the nodes and those with prickles in threes or solitary near the nodes.

(i) **Prickles in pairs.** *Acacia andongensis*, *A. burkei*, *A. caffra*, *A. erubescens*, *A. fleckii*, *A. galpinii*, *A. goetzei*, *A. gourmaensis*, *A. hecatophylla*, *A. hereroensis*, *A. laeta*, *A. manubensis*, *A. mellifera*, *A. montis-usti*, *A. nigrescens*, *A. persiciflora*, *A. polyacantha*, *A. pseudonigrescens*, *A. robyniana*, *A. royumae*, *A. schlechteri*, *A. tanganyikensis*, *A. venosa* and *A. welwitschii*.

(ii) **Prickles solitary or in threes.** *Acacia ankokib*, *A. asak*, *A. caraniana*, *A. cheilanthifolia*, *A. condyloclada*, *A. dudgeoni*, *A. hamulosa*, *A. ogadensis*, *A. oliveri*, *A. senegal*, *A. somalensis*, *A. thomasii* and *A. zizyphisina*.

(b) Section *Monacantha*

Species of section *Monacantha* occur in the Americas and from Africa through Asia to Australia. No species is common to both Africa and the Americas, however, some show close similarities (Ross 1981). All non-American species of section *Monacantha* which have globular heads of flowers belong to the "*Acacia pennata* group" (Brenan and Exell 1957); this group seemingly also occurs in the New World (see the *A. riparia* group below).

New World Groups

According to Seigler (*pers. comm.*) subgenus *Aculeiferum* in the New World has three main centres of diversity: (1) Mexico and adjacent southwestern U.S. and Central America; (2) southern Brazil, northern Argentina, Paraguay and Bolivia; (3) northeastern Brazil. Preliminary studies by Seigler suggest that the 50 or so taxa from area (1) can be divided into five major species-groups:

(i) ***Acacia acatlensis* group** (comprising *Acacia acatlensis*, *A. centralis*, *A. compacta*, *A. coulteri*, *A. dolichostachya*, *A. millefolia*, *A. sericea*, *A. usmacintensis*, *A. willardiana* and *Senegalia durangensis*). Members this group lack prickles but the stipules (when present) may occur as small subulate spines (e.g. *A. millifolia*, *A. centralis*) or as spinose horn-like projections (e.g. *A. coulteri*). Spinose stipules are a constant feature of species in subgenus *Acacia* but are otherwise rare (? or absent) elsewhere in subgenus *Aculeiferum*. *Acacia*

willardiana differs from all other members of subgenus *Aculeiferum* in having leaves reduced to phyllode-like structures (see Vassal and Guinet 1972). Evans *et al.* (1977) showed that free amino acids of *A. coulteri* are the same as in species of section *Aculeiferum* (this section is seemingly not represented in the New World).

(ii) **Acacia berlandieri group** (comprising *Acacia anisophylla*, *A. berlandieri*, *A. crassifolia*, *A. emoryana*, *A. greggii*, *A. occidentalis*, *A. parviflora* (syn. *A. micrantha*), *peninsularis*, *A. reniformis*, *A. roemeriana*, *A. sororia*, *A. subangulata*, *A. wrightii* and *Senegalia painteri*). This is a somewhat diverse group of species; *A. berlandieri* hybridizes with a number of the taxa listed here (Seigler, pers. comm.). *Acacia greggii* was placed in the monotypic subsection *Cryptocotylae* by Vassal (although *A. roemeriana* was also referred to this subsection by Conn *et al.* 1989, its seedling ontogeny is unknown). The only other species of this group studied by Vassal was *A. berlandieri* which he placed in subsection *Phanerocotylae* (along with one or two species from the *A. acatensis* and *A. glomerosa* groups).

(iii) **Acacia riparia group** (comprising *Acacia hayesii*, *A. macilenta*, *A. riparia*, *A. tenuifolia* and *Senegalia acapulcensis*). These species have either globular heads (e.g. *A. riparia*, *A. tenuifolia*) or cylindrical spikes (e.g. *A. macilenta*, *A. hayesii*; *A. bonariensis* from Argentina seems close to these species). Members of the *A. riparia* group have prickles arranged in rows like members of the "*A. pennata* group" from Africa and Asia. Evans *et al.* (1977) showed that *A. riparia* accumulated the same unknown free amino acid as members of the "*A. pennata* group".

(iv) **Acacia glomerosa group** (comprising *Acacia gaumeri*, *A. glomerosa*, *A. interior*, *A. mammifera*, *A. mirandae*, *A. picachensis* and *A. purpursii*). Closely related to the *A. riparia* group.

(v) **Acacia amazonica group** (comprising *Acacia amazonica*, *A. kuhlmannii*, *A. lacerans*, *A. velutina*, and probably *A. serra*).

African Groups

According to Ross (1979: 9) the African species of section *Monacantha* share the following characters: pale yellowish-white flowers, stipitate ovaries, most are climbers or scandent shrubs, all lack secondary leaves, and all have pods which are +/- umbonate over the seeds and have a fairly conspicuous transverse venation. Ross considered it desirable to divide the African *Monacantha* into those species with capitate inflorescences and those with spicate inflorescences (Vassal 1972 did not make this distinction).

(i) **Spicate inflorescences** (*Acacia ataxacantha*, *A. chariessa*, *A. eriocarpa* and *A. macrostachya*). Ross (1979: 10) noted that these four species differed from those of group (ii) below in pollen, seedling and seed characters characters. Serological studies by Brain (1990) showed *A. ataxacantha* as being close to (spicate) members of section *Aculeiferum* and distinct from the capitate species of section *Monacantha*. Furthermore, Evans *et al.* (1977) included *A. ataxacantha* in their free amino acid Group 3 (which otherwise included species of section *Aculeiferum*). This species is therefore anomalous within section *Monacantha*.

(ii) **Capitate inflorescences** (*Acacia adenocalyx*, *A. brevispica*, *A. ciliolata*, *A. erythrocalyx*, *A. kamerunensis*, *A. kraussiana*, *A. latistipulata*, *A. lujae*, *A. montigena*, *A. pentagona*, *A.*

schweinfurthii, *A. taylorii* and *A. tephrodermis*). Brain (1990) showed that *A. brevispica* and *A. schweinfurthii* were quite distinct serologically from the African spicate members of both section Monacantha and section Aculeiferum. *Acacia kraussiana* on the other hand was, serologically, like a spicate species.

(c) Section Filicinae

According to Pedley (1987a) the morphological and chemical attributes of this endemic New World group suggest that it could well be treated as a distinct genus; a similar view was expressed by Guinet (in Maslin 1987) and by Seigler (pers. comm.). Guinet and Vassal (1978) were of the opinion that section Filicinae preserves the morphological characters closest to those postulated as being ancestral in the genus. This idea was supported by the cladistic analysis of Chappill and Maslin (1995). According to Guinet (1990) section Filicinae is closely related to the *Piptadenia* group in tribe Mimoseae; however, Chappill and Maslin suggest that these similarities may be interpreted as plesiomorphies.

C. *Acacia* subgenus Phyllodineae

This subgenus with in excess of 900 species is largely confined to Australia. In a recent cladistic analysis of tribe Acacieae by Chappill and Maslin (1995) subgenus Phyllodineae was supported as monophyletic with subgenus Aculeiferum section Aculeiferum the sister group. As already noted Pedley's (1986) proposal that subgenus Phyllodineae be treated as a distinct genus, *Racosperma*, has not been widely adopted.

The geographic patterns of species-richness for subgenus Phyllodineae and its seven sections are mapped and discussed by Hnatiuk and Maslin (1988) and Maslin and Pedley (1988). Individual species distributions are given in Maslin and Pedley (1982) but this work is now considerably dated, especially because of the large numbers of new species which have been described (or are in preparation) since 1982. The species numbers given below follow those of Maslin (1995) and update those given by Maslin and Pedley (1988). Most of the 20 extra-Australian species of subgenus Phyllodineae are discussed by (Pedley 1975). Coulaud *et al.* (1995) recently published the results of a cytogenetic investigation on *A. heterophylla* (La Reunion Island) and its close relative, *A. melanoxyton* (Australia)

As will become evident from the brief discussions which follow there is a critical need to reassess the classification of subgenus Phyllodineae in order to devise a system of meaningful categories which reflect the evolutionary history of this very large and important group.

Currently the most generally used classification is that of Pedley (1978) which represents a good, pragmatic attempt at rationalizing earlier schemes by Bentham (1842, 1864, 1875) and Vassal (1972). The seven sections recognized by Pedley correspond fairly well to groupings of series or subseries defined earlier by Bentham (see Table 1 of Maslin 1988 and Table 1 of Chappill and Maslin 1995). Although we have adopted Pedley's scheme here it does not follow that we consider all these groups to be "good" taxonomic entities. Indeed, as has been demonstrated in the recent cladistic and immunology studies by Chappill and Maslin (1995) and Brain and Maslin (in press) respectively, considerable work is still needed to elucidate the classification of subgenus Phyllodineae. Brain and Maslin in particular challenge many of the conventional beliefs concerning the classification of subgenus Phyllodineae (see discussions below).

Additional to the species-groups that are noted below there are a number from the Arid Zone which are listed in Maslin and Hopper (1982).

(a) Section Botrycephalae (42 species). Species of this section are confined to eastern Australia and recent studies have suggested that it is most closely related to certain racemose species of section Phyllodineae from eastern Australia (Tindale and Roux 1969 and 1974, Vassal 1972 and Pettigrew and Watson 1975, Brain and Maslin in press).

Based on chemical data a number of subgroups have been recognized within section Botrycephalae. For example, studies by Tindale and Roux (1969) of flavonoid and condensed-tannin contents of the heartwood and bark from 18 species recognized the following species-groups: (1) *A. constablei*, *A. decurrens*, *A. irrorata* subsp. *velutinella*, *A. mearsii*, *A. parramattensis* and *A. trachyphloia*; (2) *A. filicifolia*, *A. irrorata* and *A. silvestris*; (3) *A. botrycephala*, *A. cardiophylla*, *A. chrysotricha*, *A. leuoclada* subsp. *argentifolia* and *A. oshanesii*; (4) *A. baileyana* and *A. dealbata*. Also, from detailed analyses of gum from 10 species Anderson (1978) and Anderson *et al.* (1971 and 1984) proposed that section Botrycephalae be divided into two distinct groups. Group A included six taxa (*A. deanei* and *A. deanei* subsp. *paucijuga*, *A. irrorata* subsp. *irrorata*, *A. parramattensis*, *A. parvipinnula* and *A. trachyphloia*) which differed greatly in their gum composition from the phyllodinous species of Bentham's (1864) "Division" Phyllodineae; Group B also contained six taxa (*A. dealbata* subsp. *dealbata* and subsp. *subalpina*, *A. filicifolia*, *A. leuoclada*, *A. silvestris* and *A. terminalis*) but these showed a strong chemical resemblance to the phyllodinous taxa examined.

Pedley (1986) suggested that perhaps *A. leptoclada* and *A. mitchellii* might constitute a third group. *Acacia mitchellii*, which differs from all other species of Botrycephalae in having non-racemose inflorescences, was excluded from section Pulchellae by Guinet *et al.* (1980).

(b) Section Pulchellae (27 species). This group is endemic in south-west Western Australia (Maslin 1975 and 1979) and as discussed by Guinet *et al.* (1980) comprises a "core" of interrelated species centered around *A. pulchella*, *A. browniana* and *A. drummondii*. The affinities of the few additional species included in section Pulchellae are unclear although some seem related to species of section Phyllodineae, e.g. *A. gilbertii* (with *A. myrtifolia*) and *A. insolita* (with *A. squamata*). The relationship of the "core" species to the rest of the genus is unknown. Vassal (1972) included some in his section Pulchelloidea, however, this section was not supported by Pettigrew and Watson (1975). *Acacia mitchellii*, the only eastern Australian species referred by Bentham to section Pulchellae, is better placed in section Botrycephalae.

(c) Section Alatae (21 species). Although this section was recognized in Pedley's (1978) classification he correctly relegated it to synonymy under section Phyllodineae (as *Racosperma* section *Racosperma*) in his 1986 scheme. For convenience this section is retained in the "critical list" below. Some of the species of this section were included in Vassal's section Pulchelloidea (see above).

(d) Section Lycopodiifoliae (17 species). The tropical/arid zone species of section Lycopodiifoliae probably constitute a natural group. It is not clear how this section relates to the rest of subgenus Phyllodineae, however, as noted by Pedley (1987c) and Cowan and Maslin (1990a) there are certain floral details which suggest some affinities to the *A. deltoidea* group (see under section Juliflorae below). Although *A. baueri* (central east coast of

Australia) was included in the section by Pedley (1978) this species is probably better placed in section Phyllodineae somewhere near *A. brunioides*. Relationships between species of section Lycopodiifoliae are shown in Pedley (1972). A detailed, elegant study by Rutishauser and Sattler (1986) examined phyllode architecture and development in one member of section Lycopodiifoliae, namely, *A. longipedunculata*.

(e) Section Phyllodineae (387 species). This is the largest group within subgenus Phyllodineae. It is widespread in Australia with the most species-rich areas occurring south of the Tropic of Capricorn in the temperate and adjacent semiarid areas of eastern, south-eastern and south-western Australia; the section is poorly represented in the tropical/sub-tropical north of the continent. There are no indigenous species of section Phyllodineae occurring outside Australia.

Although Bentham (1864) recognized six subseries within his series Uninerves (= section Phyllodineae) these are largely artificial groups and are generally not adopted by modern workers. Furthermore, certain species traditionally referred to this section are clearly related to species currently placed in section Plurinerves. Excluding these taxa (some of which are discussed below) most published evidence suggests that there is a major difference within subgenus Phyllodineae between species with so-called "1-nerved" phyllodes (i.e. section Phyllodineae) and those with plurinerved phyllodes (i.e. the combined sections Plurinerves and Juliflorae). Pedley (1986) recognized this division but refrained from attributing subgeneric rank to the two groups. Unpublished data on hybridity in *Acacia* (Kenrick, Knox, Guinet and Maslin) show no natural or artificial hybrids recorded between species of these two groups. Although this division was evident in the cladistic analysis by Chappill and Maslin (1995) it was not able to be substantiated in the immunology study by Brain and Maslin (in press).

While inflorescence structure seems fundamentally important in defining natural groups within section Phyllodineae the traditional method of dividing species into racemose vs. non-racemose subgroups is a somewhat over-simplified approach. Indeed, the immunology study by Brain and Maslin (in press) suggests that these two subgroups are far more distinct from one another than hitherto recognized. Bentham (1864) included species with elongated racemes in subseries Racemosae, while species with simple inflorescences or extremely reduced racemes were distributed among five other subseries. However, recent studies show examples where members of closely related species-pairs may have different inflorescence types, or where a single species may possess two different inflorescence types, e.g. *A. victoriae* (elongated racemes) - *A. synchronicia* (simple inflorescences), fide Maslin (1992), *A. calamifolia* (very reduced racemes) - *A. nematophylla* (simple inflorescences), fide Maslin and Whibley (1987), *A. bivenosa* (both racemes and simple inflorescences), fide Chapman and Maslin (1992).

Based on a study of flavonoid patterns in *Acacia* Tindale and Roux (1974) showed that subseries Racemosae could be divided into at least two basic groups. Although not all species referable to this subseries were included in their study they demonstrated that arid/ semi-arid species such as *A. blakelyi*, *A. murrayana* (syn. *A. frumentacea*), *A. pruinocarpa*, *A. sclerosperma*, *A. victoriae*, etc. contained pyrogallol-type compounds (which also occurred extensively in species of sections Juliflorae and Plurinerves); on the other hand species predominantly from eastern Australia, e.g. *A. falcata*, *A. falciformis*, *A. fimbriata*, *A. kettlewelliae*, *A. pycnantha*, *A. retinodes*, etc. contained resorcinol-type compounds (which also occurred in species of section Botrycephalae). Based on seed characters it is possible to divide this latter group into species with long funicles which encircle the seeds, e.g. *A. falcata*, *A.*

retinodes, *A. rubida* (there are over 40 species Australia-wide with this sort of funicle, most of these are referable to the "*A. microbotrya* group", fide Maslin 1995a) and those with short, non-encircling funicles, e.g. *A. adunca*, *A. pycnantha*, *A. williamsonii*, etc. (Maslin, unpublished data). The Brain and Maslin immunology study showed the racemose group as comprising three subgroups (centered on *A. ligulata*, *A. pyrifolia* and *A. podalyriifolia*).

The non-racemose members of section Phyllodineae are generally not well-studied. However, the work by Brain and Maslin (in press) has provided some surprising insights on possible affinities for this group. In this study Brain and Maslin were unable to confirm the traditional view that the racemose and non-racemose members of section Phyllodineae are closely related; indeed, based on their serological evidence at least some non-racemose taxa were shown to be related to species currently placed in sections Plurinerves and Juliflorae. Clearly, much work is still needed to be done on the "non-racemosae" in order to clarify the classification and phylogeny of the complex. To date a few discrete groups have been recognized within this assemblage, e.g., the Triangulares group which Maslin (1978a) showed as comprising two distinct sub-groups centered around *A. biflora* and *A. horridula*, and the *A. ulicifolia* group which Pedley (1969) recognized (we would add the following three non-Queensland species to those listed by Pedley: *A. asparagoides*, *A. brownii* and *A. gunnii*).

As noted above there are some instances where the traditional division of species of subgenus Phyllodineae into uninerved vs. plurinerved groups does not result in a natural clustering of taxa. Examples of this can be found in taxonomically distant groups, e.g. the "*A. wilhelmiana* group" (Maslin 1990), *A. murrayana* (1-nerved phyllodes) - *A. praelongata* (plurinerved phyllodes) and the "*A. paradoxa* group" (where *A. aspera*, *A. ausfeldii*, *A. dodonaeifolia*, *A. leprosa*, *A. paradoxa* and *A. rhotinocarpa* all have 1-nerved phyllodes whereas *A. cognata*, *A. glandulicarpa*, *A. howittii*, *A. montana*, *A. subporosa* and *A. verniciflua* all have more than 1-nerved phyllodes). Also, diaphyllodinous species such as *A. basedowii*, *A. diaphyllodinea*, *A. leptospermoides*, etc. (fide Maslin 1978 and Vassal and Maslin 1979) are anomalous in having horizontally flattened phyllodes that are 1-nerved adaxially and 3-nerved abaxially.

(f) **Section Plurinerves** (212 species) is discussed under section Juliflorae below.

(g) **Section Juliflorae** (235 species). A number of recent studies have suggested that this section is closely related to section Plurinerves, fide Tindale and Roux 1969 and 1974, Vassal 1972, Pettigrew and Watson 1975 and Tindale 1980. Indeed, in his most recent classification Pedley (1986) combined the two groups under *Racosperma* section Plurinerves (see Table 1 of Maslin 1988 and Chappill and Maslin 1995). However, the immunology study by Brain and Maslin (in press) in part at least challenges this view. Their results suggest that while section Plurinerves seems to be very coherent, doubt is cast on section Juliflorae as a natural group. One subgroup of species within section Juliflorae showed strong affinities with species of section Plurinerves the other subgroup (surprisingly) showed affinities with certain non-racemose species of section Phyllodineae.

These two sections are widespread in Australia and show similar geographic patterns; however, there are differences, the most important being that section Plurinerves has fewer species north of the Tropic of Capricorn and more species in inland areas of south-eastern Australia. About 20 species from these two sections occur outside Australian - see Pedley 1975 for review. *Note.* Subsequent to Pedley's work another two extra-Australian species referable to section Juliflorae have come to light, namely, *A. pubirachis* (occurs in northern Australia and in

Papua New Guinea) and a seemingly new species (allied to *A. cowleana*) endemic to the Indonesian Island of Wetar and which was noted by Nielsen (1992) under *A. leptocarpa*.

Future classifications will need to determine appropriate infrasectional categories for species encompassed by the combined sections Juliflorae and Plurinerves, and it appears that phyllode nervature will prove important in this work. In general terms these plurinerved species can be conveniently grouped according to the number and relative spacing of nerves on their phyllodes. This method was adopted by Pedley (1987d), Maslin and Pedley (1988) and Cowan and Maslin (1990) where the following main trends were recognized:

1. *Microneurous phyllodes*. Phyllodes with numerous, very fine, closely approximating longitudinal nerves without anastomoses between them, e.g. *A. aneura*, *A. lasiocalyx*, *A. tumida*, etc. (section Juliflorae), *A. calcicola*, *A. coriacea*, *A. pendula*, etc. (section Plurinerves).
2. *Oligoneurous phyllodes*. Phyllodes with only a few, distant longitudinal nerves. Within this pattern of nervature two subgroups can be recognized:
 - (i) anastomoses absent or few between the main longitudinal nerves, e.g. *A. oncinophylla*, *A. jibberdingensis* (section Juliflorae, there are relatively few species of this section with this pattern), *A. cochlearis*, *A. confusa*, *A. complanata*, *A. elongata*, etc. (section Plurinerves).
 - (ii) anastomoses numerous between the main longitudinal nerves, e.g. *A. crassa*, *A. dimidiata*, *A. holosericea* (section Juliflorae), *A. ixiophylla*, *A. koa* (from Hawaii), *A. melanoxylon*, *A. platycarpa*, etc. (section Plurinerves).

This system is primarily a mnemonic device for informally arranging species within this large group, nevertheless recent studies have recognized a number of species-groups within these broad categorizations, e.g. the *A. holosericea* group (Maslin and Thomson 1992: Group 2ii), the "*A. cunninghamii*" group (Pedley 1974: Group 2ii), the *A. verricula* and *A. flavipila* groups (Cowan and Maslin 1990: Group 2ii), the *A. aneura* group (Randell 1992: Group 1) and the *A. multilineata* group (Cowan and Maslin 1990b: Group 1). However, there are some taxa that cannot be readily ascribed to any of the above groups, e.g. the *A. deltoidea* group (fide Pedley 1987c and Cowan and Maslin 1990a).

Apart from phyllode nervature some groups are defined primarily by their carpological and/or inflorescence characters. For example species of the *A. stigmatophylla* group (Tindale 1980) have very distinctive pods (not unlike those of *Calliandra* spp.) while those of the *A. longifolia* group (i.e. Bentham's series Juliflorae subseries Tetramerae and series Pungentes subseries Spicatae pro parte) are united by their 4-merous flowers (see Pedley 1964). Species from the latter group possess phyllodes that fall into a number of the categories outlined above (or they may even have 1-nerved phyllodes as in species of section Phyllodineae, e.g. *A. verticillata*).

CONCLUSION

This synoptic survey has indicated a number of critical problem areas in *Acacia* taxonomy. By a careful resolution across a wide range of critical species it will be possible to come to a more meaningful natural classification of this large important genus.

ACKNOWLEDGEMENTS

The compilation of the "critical list" has evolved from careful analyses by many people and we wish to thank in particular I. Nielsen, L. Rico Arce, J. Ross, J. Vassal and especially D. Seigler. We also thank P. Brain for encouraging us to prepare the list and R.S. Cowan, G.P. Lewis, J.M. Lock, L. Rico Arce and D. Seigler for critically reading the manuscript.

REFERENCES

- Anderson, D.M.W. (1978). Chemotaxonomic aspects of the chemistry of *Acacia* gum exudates. *Kew Bull.* 32: 529-536.
- Anderson, D.M.W., Bell, P.C. and McNab, C.G.A. (1971). An analytical study of some *Acacia* gum exudates of the series Botryocephalae. *Carbohyd. Res.* 20: 269-274.
- Anderson, D.M.W., Farquhar, J.G.K. and McNab, C.G.A. (1984). The gum exudates from some *Acacia* subspecies of the series Botryocephalae. *Phytochem.* 23: 579-580.
- Bentham, G. (1842). Notes on Mimoseae, with a Synopsis of Species. *London J. Bot.* 1: 318-528.
- Bentham, G. (1864). *Flora Australiensis*. vol. 2. (Lovell Reeve: London.)
- Bentham, G. (1875). Revision of Suborder Mimoseae. *Trans. Linn. Soc. London* 30: 335-664.
- Brain, P. (1987). Immunology and phylogeny: a preliminary study of *Acacia*. *S. Afr. J. Sci.* 83: 422-427.
- Brain, P. (1990). Immunology and phylogeny II: further studies on *Acacia*. *Suid-Afrikaanse Tydskrifvir Wetenskap* 86: 195-199.
- Brenan, J.P.M. and Exell, A.W. (1957). *Acacia pennata* (L.) Willd. and its relatives in tropical Africa. *Bol. Soc. Brot. ser. 2*, 31: 99-140.
- Britton, N.L. and Rose, J.N. (1928). *North American Flora*. Vol.23, part 3. (The New York Botanical Garden: New York.)
- Chapman, A.R. and Maslin, B.R. (1992). *Acacia* Miscellany 5. A review of the *A. bivenosa* group (Leguminosae: Mimosoideae: Section Phyllodineae). *Nuytsia* 8: 249-283.
- Chappill, J.A. and Maslin, B.R. (1995). A phylogenetic assessment of tribe Acacieae. In Crisp, M. and Doyle J.J. (eds) *Advances in Legume Systematics 7: Phylogeny*, pp. 77-99. (Royal Botanic Gardens, Kew: London.)
- Clarke, H.D. (1995). Systematic and evolutionary studies of *Acacia* series Gummiferae (Fabaceae: Mimosoideae). PhD Thesis, University of Illinois at Urbana-Champaign.
- Clarke, H.D. and Seigler, D.S. (1991). The systematics of *Acacia* series Gummiferae in the Caribbean. *Amer. J. Bot.* 78(6): 173-174.

- Clarke, H.D., Seigler, D.S. and Ebinger, J.E. (1989). *Acacia farnesiana* (Fabaceae: Mimosoideae) and related species from Mexico, the southwestern U.S. and the Caribbean. *Syst. Bot.* 14(4): 549-564.
- Clarke, H.D., Seigler, D.S. and Ebinger, J.E. (1990). *Acacia constricta* (Fabaceae: Mimosoideae) and related species from the southwestern U.S. and Mexico. *Amer. J. Bot.* 77(3): 305-315.
- Conn, E.E., Seigler, D.S., Maslin, B.R. and Dunn, J. (1989). Cyanogenesis in *Acacia* subgenus *Aculeiferum*. *Phytochemistry* 28(3): 817-820.
- Coulaud, J. Brown, S.C. and Siljak-Yakovlev, S. (1995). First cytogenetic investigation in populations of *Acacia heterophylla*, endemic from La Reunion Island, with reference to *A. melanoxylon*. *Ann. Bot.* 75: 95-100.
- Cowan, R.S. and Maslin, B.R. (1990). *Acacia* Miscellany 1. Some oligoneurous species of *Acacia* (Leguminosae: Mimosoideae: Section Plurinerves) from Western Australia. *Nuytsia* 7: 183-199.
- Cowan, R.S. and Maslin, B.R. (1990a). *Acacia* Miscellany 2. Species related to *A. deltoidea* (Leguminosae: Mimosoideae: section Plurinerves). *Nuytsia* 7(2): 201-208.
- Cowan, R.S. and Maslin, B.R. (1990b). *Acacia* Miscellany 3. Some new microneurous taxa of Western Australia related to *A. multilineata* (Leguminosae: Mimosoideae: Section Plurinerves). *Nuytsia* 7: 209-219.
- Ebinger, J.E. and Seigler, D.S. (1990, unpublished ms). Ant-*Acacia* hybrids of Central America.
- Evans, C.S., Qureshi, M.Y. and Bell, E.A. (1977). Free amino acids in the seeds of *Acacia* species. *Phytochemistry* 16: 565-570.
- Guinet, P. (1969). Les Mimosaceae, etude de palynologie fondamentale, correlations, evolution. *Trav. Sec. Sci. Techn. Pondichery* 9: 1-293.
- Guinet, P. (1979). Biologie evolutive dans le genre *Acacia*: arguments palynologiques. *Mem. Trav. Inst. Montpellier* 4: 41-51.
- Guinet, P. (1981). Mimosoideae: the Characters of their Pollen Grains, pp. 835-855. In: R.M. Polhill and P.H. Raven (eds.), *Advances in Legume Systematics Part 2* (Royal Botanic Gardens, Kew: England.)
- Guinet, P. (1990). The genus *Acacia* (Leguminosae, Mimosoideae): its affinities as borne out by its pollen characters. *Pl. Syst. Evol. [Suppl. 5]*: 81-90.
- Guinet, P. and Lugardon, (1976). Diversite des structures de l'exine dans le genre *Acacia* (Mimosaceae). *Pollen et Spores* 18: 483-511.
- Guinet, P. and Vassal, J. (1978). Hypotheses on the differentiation of the major groups in the genus *Acacia* (Leguminosae). *Kew Bull.* 32: 509-527.
- Guinet, P., Vassal, J., Evans, C.S. and Maslin, B.R. (1980). *Acacia* (Mimosoideae): composition and affinities of the series Pulchellae Benth. *Bot. J. Linn. Soc.* 80: 53-68.
- Guinet, P. (1979). Biologie evolutive dans le genre *Acacia*: arguments palynologiques. *Mem. Trav. Inst. Montpellier* 4: 41-51.
- Hnatiuk, R.J. and Maslin, B.R. (1988). Phytogeography of *Acacia* in Australia in relation to climate and species-richness. *Austral. J. Bot.* 36: 361-383.

- Janzen, D.H. (1974). Swollen-Thorn Acacias of Central America. *Smithsonian Contr. Bot.* 15: 1-131.
- Lee, Y.S., Seigler, D.S. and Ebinger, J.E. (1989). *Acacia rigidula* (Fabaceae) and related species in Mexico and Texas. *Syst. Bot.* 14(1): 91-100.
- Madsen, E.B. (1990). Flora of Ecuador. *Acacia* Mill. (Leguminosae: Mimosoideae). MSc Thesis, Aarhus.
- Maslin, B.R. (1975). Studies in the genus *Acacia* (Mimosaceae) 4. A revision of the series Pulchellae. *Nuytsia* 1: 388-492.
- Maslin, B.R. (1978). Studies in the genus *Acacia* (Mimosaceae) - 7. The taxonomy of some diaphyllodinous species. *Nuytsia* 2(4): 200-219.
- Maslin, B.R. (1978a). Studies in the genus *Acacia* (Mimosaceae) - 8. A revision of the Uninerves-Triangulares, in part (the tetramerous species). *Nuytsia* 2: 266-333.
- Maslin, B.R. (1979). Studies in the genus *Acacia* (Mimosoideae) - 9. Additional notes on the Series Pulchellae Benth. *Nuytsia* 2: 354-367.
- Maslin, B.R. (1987). Report of discussions at I.G.S.M. meeting no. 6 on the proposal to split *Acacia*. *Bulletin of the International Group for the Study of Mimosoideae* 15: 108-118.
- Maslin, B.R. (1988). Should *Acacia* be divided? *Bulletin of the International Group for the Study of Mimosoideae* 16: 54-76.
- Maslin, B.R. (1989). Wattle become of *Acacia*? *Australian Systematic Botany Society Newsletter* 58: 1-13.
- Maslin, B.R. (1990). *Acacia* Miscellany 4. Three new Western Australian species with affinities to *A. wilhelmiana* (Leguminosae: Mimosoideae: section Plurinerves) from Western Australia. *Nuytsia* 7(2): 221-228.
- Maslin, B.R. (1992). *Acacia* Miscellany 6. A review of *Acacia victoriae* and related species (Leguminosae: Mimosoideae: Section Phyllodineae). *Nuytsia* 8: 285-309.
- Maslin, B.R. (1995). Systematics and phylogeography of Australian species of *Acacia*: an overview. *IFA Newsletter* 36(2): 2-5.
- Maslin, B.R. (1995a). *Acacia* miscellany 14. Taxonomy of some Western Australian "Uninerves-Racemosae" species (Leguminosae: Mimosoideae: section Phyllodineae) *Nuytsia* 10(2): 181-203.
- Maslin, B.R., Dunn, J.E. and Conn, E.E. (1988). Cyanogenesis in Australian species of *Acacia*. *Phytochemistry* 27(2): 421-428.
- Maslin, B.R. and Hopper, S.D. (1982). Phylogeography of *Acacia* (Leguminosae: Mimosoideae) in Central Australia. pp. 301-316. In: Barker, W.R. and Greenslade, P.J.M. (eds.) *Evolution of the Flora and Fauna of Arid Australia* (Peacock Publications: Adelaide.)
- Maslin, B.R. and Pedley, L. (1982). The distribution of *Acacia* (Leguminosae: Mimosoideae) in Australia. Part 1. Species distribution maps. *W. Austral. Herb. Research Notes* 6: 1-128.
- Maslin, B.R. and Pedley, L. (1988). Patterns of distribution of *Acacia* in Australia. *Austral. J. Bot.* 36: 385-393.

- Maslin, B.R. and Thomson, L.A.J. (1992). Re-appraisal of the taxonomy of *Acacia holosericea* A. Cunn. ex Don, including the description of a new species, *A. colei*, and the reinstatement of *A. neurocarpa* A. Cunn. ex Hook. Austral. Syst. Bot. 5: 729-743.
- Maslin, B.R. and Whibley, D.J.E. (1987). The taxonomy of some South Australian *Acacia* section Phyllodineae species (Leguminosae: Mimosoideae). Nuytsia 6(1): 19-32.
- Nielsen, I. (1980). Notes on Indo-chinese Mimosaceae. Adansonia ser. 2, 19: : 339-363.
- Nielsen, I. (1992). Mimosaceae (Leguminosae - Mimosoideae). Fl. Malesiana ser. 1, 11(1): 1-226.
- Pedley, L. (1964). Notes on *Acacia*, chiefly from Queensland, II. Proc. Roy. Soc. Qd 75: 29-35.
- Pedley, L. (1969). Notes on *Acacia*, chiefly from Queensland, III. Contr. Qd Herb. 4: 1-7.
- Pedley, L. (1972). Revision of *Acacia lycopodiifolia* A.Cunn. ex Hook. and its allies. Contr. Qd Herb. No. 11: 1-23.
- Pedley, L. (1974). Notes on *Acacia*, chiefly from Queensland, IV. Contr. Qd Herb. 15: 1-27.
- Pedley, L. (1975). Revision of the extra-Australian species of *Acacia* subg. *Heterophyllum*. Contr. Qd Herb. No. 18: 1-24.
- Pedley, L. (1978). A revision of *Acacia* Mill. in Queensland. Austrobaileya 1: 75-234.
- Pedley, L. (1986). Derivation and dispersal of *Acacia* (Leguminosae), with particular reference to Australia, and the recognition of *Senegalia* and *Racosperma*. Bot. J. Linn. Soc., 92: 219-254.
- Pedley, L. (1987). Australian acacias - taxonomy and phytogeography. In J. Turnbull (ed.) *Australian acacias in developing countries: proceedings of an international Workshop held at the Forestry Training Centre, Gympie, Qld, Australia, 4-7 Aug. 1986*. Publ. Australian Centre for International Agricultural Research. Proceedings No. 16.
- Pedley, L. (1987a). In defence of *Racosperma*. Bulletin of the International Group for the Study of Mimosoideae 15: 123-129.
- Pedley, L. (1987b). Generic status of *Acacia sensu lato*. Australian Systematic Botany Society Newsletter 53: 87-91.
- Pedley, L. (1987c). *Racosperma deltoideum* (Cunn. ex G. Don) Pedley (Leguminosae: Mimosoideae) and related species on northern Australia. Austrobaileya 2(4): 314-320.
- Pedley, L. (1987d). Notes on *Racosperma* Martius (Leguminosae: Mimosoideae), 1. Austrobaileya 2(4): 321-327.
- Pettigrew, C.J. and Watson, L. (1975). On the classification of Australian Acacias. Austral. J. Bot. 23: 833-347.
- Randell, B.R. (1992). Mulga, a revision of the major species. J. Adelaide Bot. Gard. 14: 105-132.
- Ross, J.H. (1973). Towards a classification of African Acacias. Bothalia 11: 107-113.
- Ross, J. (1979). A conspectus of the African *Acacia* species. Mem. Bot. Survey S. Africa No. 44: 1-155.
- Ross, J.H. (1981). An analysis of the African *Acacia* species: their distribution, possible origins and relationships. Bothalia 13: 389-413.

- Rutishauser, R. and Sattler, R. (1986). Architecture and development of the phyllode - stipule whorls of *Acacia longipedunculata*: controversial interpretations and continuum approach. *Canadian J. Bot.* 64: 1987-2019.
- Tindale, M.D. (1980). Notes on Australian taxa of *Acacia* No. 7. *Telopea* 2: 113-125.
- Tindale, M.D. and Roux, D.G. (1969). A phytochemical survey of the Australian species of *Acacia*. *Phytochem.* 8: 1713-1727.
- Tindale, M.D. and Roux, D.G. (1974). An extended phytochemical survey of Australian species of *Acacia*: chemotaxonomic and phylogenetic aspects. *Phytochem.* 13: 829-839.
- Vassal, J. (1972). Apport des recherches ontogeniques et seminologiques a l'etude morphologique, taxonomique et phylogénique du genre *Acacia*. *Bull. Soc. Hist. Nat. Toulouse* 108: 105-247.
- Vassal, J. (1981). Acacieae. In: R.M. Polhill and P.H. Raven (eds.), "Advances in Legume Systematics Part I" pp. 169-171. (Royal Botanic Gardens, Kew: England.)
- Vassal, J. (1988). Some remarks about the taxonomy of *Acacia*. *International Group for the Study of Mimosoideae, Bulletin No. 16*: 77-82.
- Vassal, J. and Guinet, P. (1972). Un *Acacia* Americain a petiole diaphyllocline *A. willardiana* Rose. *Adansonia ser. 2*, 12: 421-428.
- Vassal, J. and Maslin, B.R. (1979). Phylloides et phyllocladisation des le genre *Acacia*. *Bull. Soc. Hist. Nat. Toulouse t.115, fasc. 3-4*, pp. 393-401.
- Vassal, J. and Rouane, P. (1987). Testing Bentham's series of Australian Acacias s.l. (Pulchellae, Botrycephalae, Phyllodineae) on the basis of juvenile characters. *International Group for the Study of Mimosoideae, Bulletin No. 15*: 104-105.
- Vassal, J. and Rouane, P. (1991). Apport des caracteres juveniles a la classification des Acacias australiens. *Bull. Soc. Hist. Nat. Toulouse* 127: 101-109.

Appendix 1 List of "critical species" recommended for use in comparative studies on the generic status of *Acacia*. See Appendix 2 for alphabetic listing of the species given here.

FAIDHERBIA

F. albida (Africa)

ACACIA

1. SUBGENUS ACACIA

(GROUP 1)

- a. *A. anegadensis* or *A. daemon* (New World)
- b. *A. bilimekii* or *A. pringlei* or *A. rigidula* (New World)
- c. *A. bidwillii* (Australia)
- d. *A. caven* or *A. farnesiana* or *A. schaffneri* (New World) and/or *A. erioloba* or *A. haematoxylon* (Africa).
- e. *A. choriophylla*

- f. *A. cochliacantha* or *A. macracantha* or *A. pennatula* and *A. collinsii* or *A. cookii* or *A. cornigera* (New World)
- g. *A. constricta* or *A. glandulifera* or *A. neovernicosa* or *A. schottii* (New World)
- h. *A. craibii* or *A. inopinnata* (Asia)
- i. *A. dolichocephala* (Africa)
- j. *A. drepanolobium* (Africa)
- k. *A. harmandiana* or *A. leucopholea* (Asia)
- l. *A. karroo* or *A. seyal* (Africa)
- m. *A. nilotica* (Africa, Asia)
- n. *A. pacensis* (New World)
- o. *A. sphaerocephala* (New World)
- p. *A. tortilis* (Africa)

2. SUBGENUS ACULEIFERUM

A. Section Aculeiferum (GROUP 2)

- a. *A. caffra* or *A. goetzei* or *A. nigrescens* (Africa)
- b. *A. catechu* (India)
- c. *A. mellifera* (Africa)
- d. *A. modesta* (India)
- e. *A. senegal* (Africa, India)

B. Section Monacantha (GROUP 3)

- a. *A. acapulcensis* or *A. bonariensis* or *A. hayesii* or *A. riparia* and/or *A. tenuifolia* (New World)
- b. *A. acatlensis* or *A. centralis* or *A. coulteri* or *A. mammifera* or *A. usumacintensis* (New World)
- c. *A. amazonica* or *A. kuhlmannii* or *A. lacerans* or *A. velutina* (New World)
- d. *A. anisophylla* or *A. roemeriana* (New World)
- e. *A. articulata* or *A. bahiensis* (New World)
- f. *A. berlandieri* and/or *A. subangulata* (New World)
- g. *A. brevispica* (Africa) or *A. megaladena* (Asia) or *A. pennata* (Asia) or *A. schweinfurthii* (Africa)
- h. *A. donnaiensis* (New World)
- i. *A. eriocarpa* (Africa)
- j. *A. glomerata* or *A. purpusii* (New World)
- k. *A. greggii* and/or *A. wrightii* (New World)
- l. *A. thailandica* (Asia)
- m. *A. willardiana* (New World)

C. Section Filicinae (GROUP 4)

- a. *A. angustissima* or *A. boliviana* (New World)
- b. *A. sousae* or *A. tequilana* (New World)
- c. *A. villosa* (New World)
- d. *A. velvae* (New World)

D. Section ? (GROUP 5)

- a. *A. ataxacantha* (Africa)
- b. *A. furcata* (New World)

- c. *A. muricata* (New World)
- d. *A. visco* (New World)

3. SUBGENUS PHYLLODINEAE (Australian species unless otherwise indicated)

A. Section Botrycephalae (GROUP 6)

- a. *A. baileyana* or *A. dealbata*
- b. *A. chrysotricha*
- c. *A. cardiophylla* or *A. leucoclada*
- d. *A. decurrens* or *A. irrorata* or *A. mearnsii* or *A. parramattensis*
- e. *A. leptoclada* and/or *A. mitchellii*
- f. *A. silvestris* or *A. filicifolia*
- g. *A. schinoides* or *A. pruinosa* or *A. spectabilis*
- h. *A. elata*

B. Section Pulchellae (GROUP 7)

- a. *A. browniana* or *A. lateriticola* or *A. pentadenia*
- b. *A. drummondii*
- c. *A. gilbertii*
- d. *A. lasiocarpa* or *A. pulchella*

C. Section Alatae (an un-natural assemblage) (GROUP 8)

- a. *A. alata* or *A. willdenowiana*
- b. *A. aphylla*

D. Section Lycopodiifoliae (GROUP 9)

- a. *A. adoxa* or *A. hippuroides* or *A. spondylophylla*

E. Section Phyllodineae

(a) "Racemose species" (GROUP 10)

- a. *A. ampliceps* or *A. bivenosa* or *A. ligulata* or *A. rostellifera* or *A. sclerosperma*
- b. *A. calamifolia* or *A. falcata* or *A. microbotrya* or *A. notabilis* or *A. retinodes* or *A. rubida* or *A. steedmanii*
- c. *A. bancroftii*
- d. *A. binervata*
- e. *A. blakelyi* or *A. scirpifolia*
- f. *A. buxifolia* or *A. cultriformis*
- g. *A. celastrifolia* or *A. myrtifolia*
- h. *A. extensa* and/or *A. urophylla*
- i. *A. fimbriata* or *A. linifolia* or *A. podalyriifolia* or *A. prominens* or *A. pycnantha* or *A. vestita*
- j. *A. fasciculifera*
- k. *A. murrayana* or *A. praelongata* (phyllodes plurinerved)
- l. *A. inaequilatera* or *A. pyrifolia*
- m. *A. synchronicia* or *A. victoriae*
- n. *A. penninervis* or *A. falciformis*
- o. *A. pruinocarpa* or *A. ensifolia*
- p. *A. suaveolens* and/or *A. hemiteles* or *A. iteaphylla* or *A. prainii*

(b) "Non-racemose species" (or racemes very short)**(GROUP 11)**

- a. *A. acanthoclada* or *A. bidentata*
- b. *A. acinacea* or *A. flexifolia* or *A. lineata*
- c. *A. alleniana* or *A. juncifolia*
- d. *A. amblygona* or *A. pravifolia*
- e. *A. andrewsii* or *A. barbinervis*
- f. *A. baueri*
- g. *A. biflora* or *A. divergens*
- h. *A. brachyphylla* or *A. nitidula* or *A. sulcata*
- i. *A. brownii* or *A. gunnii* or *A. ulicifolia*
- j. *A. brunioides* or *A. conferta* or *A. tindaleae*
- k. *A. burbridgeae* or *A. gnidium* or *A. islana* or *A. johnsonii*
- l. *A. dentifera*
- m. *A. dictyophleba*
- n. *A. genistifolia* and *A. verticillata* (sect. *Juliflorae*)
- o. *A. handonis* and/or *A. rossei* or *A. sedifolia*
- p. *A. horridula* or *A. pycnocephala*
- q. *A. insolita*
- r. *A. leptospermoides*
- s. *A. littorea* or *A. truncata*
- t. *A. maitlandii* or *A. siculiformis*
- u. *A. merrallii* or *A. sericocarpa*
- v. *A. paradoxa* (syn. *A. armata*) and/or *A. stricta*
- w. *A. leprosa* or *A. verniciflua*
- x. *A. peuce*
- y. *A. tetragonophylla*
- z. *A. viscifolia* or *A. wilhelmiana*

F. Section Plurinerves**(a) "Microneurous species"****(GROUP 12)**

- a. *A. assimilis* or *A. lineolata* or *A. rigens*
- b. *A. calcicola* or *A. cambagei* or *A. melvillei* or *A. omalophylla* or *A. papyrocarpa* or *A. pendula*
- c. *A. colletioides* or *A. nyssophylla*
- d. *A. coriacea*
- e. *A. densiflora* or *A. eremophila* or *A. mackeyana*
- f. *A. oswaldii*
- g. *A. stenophylla*
- h. *A. translucens* or *A. nuperrima*

(b) "Oligoneurous species"**(GROUP 13)**

- a. *A. anaticeps*
- b. *A. adenogonia* or *A. deltoidea*
- c. *A. cochlearis* or *A. comans* or *A. latipes*
- d. *A. cognata* or *A. subporosa* or *A. veronica*
- e. *A. complanata* or *A. estropholata* or *A. excelsa*
- f. *A. confusa* (Taiwan, Philippines)
- g. *A. consobrina* or *A. flavipila* or *A. ixiophylla*

- h. *A. cyclops*
- i. *A. dunnii* or *A. latescens* or *A. platycarpa*
- j. *A. elongata*
- k. *A. melanoxydon* (Australia) and *A. heterophylla* (Mascarene Islands) and/or *A. koa* (Hawaii): see Coulaud *et al.* (1995).
- l. *A. multisiliqua* or *A. simsii*
- m. *A. redolens* or *A. trineura*
- n. *A. retivenea*

G. Section Juliflorae

(a) "Microneurous species" (GROUP 14)

- a. *A. acuminata* or *A. burkittii* or *A. multispicata*
- b. *A. ammobia* or *A. doratoxydon* or *A. lasiocalyx*
- c. *A. aneura* or *A. brachystachya* or *A. ramulosa* or *A. coolgardiensis*
- d. *A. blakei* or *A. burrowii* or *A. cheelii*
- e. *A. citrinoviridis* or *A. olgana* or *A. yorkkrakinensis*
- f. *A. cyperophylla* or *A. rhodophloia*
- g. *A. ephedroides* or *A. fauntleroyi* or *A. inophloia*
- h. *A. eriopoda* or *A. tumida*
- i. *A. guymeri* or *A. hyaloneura* or *A. jakesiana*
- j. *A. plectocarpa*
- k. *A. jibberdingensis*
- l. *A. kempeana* or *A. stowardii*
- m. *A. xiphophylla*

(b) "Oligoneurous species" (GROUP 15)

- a. *A. ancistrocarpa* or *A. stigmatophylla*
- b. *A. chisholmii* or *A. lysiphloia* or *A. monticola* (sect. Plurinerves) or *A. trachycarpa*
- c. *A. coleii* or *A. cowleana* or *A. holosericea* or *A. mangium*
- d. *A. concurrens* or *A. crassa* or *A. leiocalyx* or *A. longispicata*
- e. *A. denticulosa*
- f. *A. longifolia* or *A. sophorae*
- g. *A. neurophylla*
- h. *A. pachycarpa*

Appendix 2 An alphabetic list of taxa included in Appendix 1. The alpha-numeric designations in parentheses following species names refer to the *Acacia* subgeneric GROUP that is given in Appendix 1.

<i>A. acanthoclada</i> (11a)	<i>A. amazonica</i> (3c)	<i>A. angustissima</i> (4a)
<i>A. acapulcensis</i> (3a)	<i>A. amblygona</i> (11d)	<i>A. anisophylla</i> (3d)
<i>A. acatensis</i> (3b)	<i>A. ammobia</i> (14b)	<i>A. aphylla</i> (8b)
<i>A. acinacea</i> (11b)	<i>A. ampliceps</i> (10a)	<i>A. articulata</i> (3e)
<i>A. acuminata</i> (14a)	<i>A. anatriceps</i> (13a)	<i>A. assimilis</i> (12a)
<i>A. adenogonia</i> (13b)	<i>A. ancistrocarpa</i> (15a)	<i>A. ataxacantha</i> (5a)
<i>A. adoxa</i> (9a)	<i>A. andrewsii</i> (11e)	<i>A. bahiensis</i> (3e)
<i>A. alata</i> (8a)	<i>A. anegadensis</i> (1a)	<i>A. baileyana</i> (6a)
<i>A. allentiana</i> (11c)	<i>A. aneura</i> (14c)	<i>A. bancroftii</i> (10c)

- A. barbinervis* (11e)
A. baueri (11f)
A. berlandieri (3f)
A. bidenata (11a)
A. bidwillii (1c)
A. biflora (11g)
A. bilimekii (1b)
A. binervata (10d)
A. bivenosa (10a)
A. blakei (14d)
A. blakelyi (10e)
A. boliviana (4a)
A. bonariensis (3a)
A. brachyphylla (11h)
A. brachystachya (14c)
A. brevispica (3g)
A. browniana (7a)
A. brownii (11i)
A. brunioides (11j)
A. burbidgeae (11k)
A. burkittii (14a)
A. buxifolia (10f)
A. burrowii (14d)
A. caffra (2a)
A. calamifolia (10b)
A. calcicola (12b)
A. cambagei (12b)
A. cardiophylla (6c)
A. catechu (2b)
A. caven (1d)
A. celastrifolia (10g)
A. centralis (3b)
A. cheelii (14d)
A. chisholmii (15b)
A. choriophylla (1e)
A. chrysotricha (6b)
A. citrinoviridis (14e)
A. cochlearis (13c)
A. cochliacantha (1f)
A. cognata (13d)
A. coleii (15c)
A. colletioides (12c)
A. collinsii (1f)
A. comans (13c)
A. complanata (13e)
A. concurrens (15d)
A. conferta (11j)
A. confusa (13f)
- A. consobrina* (13g)
A. constricta (1g)
A. cookii (1f)
A. coolgardiensis (14c)
A. coriacea (12d)
A. cornigera (1f)
A. coulteri (3b)
A. cowleana (15c)
A. craibii (1h)
A. crassa (15d)
A. cultriformis (10f)
A. cyclops (13h)
A. cyperophylla (14f)
A. daemon (1a)
A. dealbata (6a)
A. decurrens (6d)
A. deltoidea (13b)
A. densiflora (12e)
A. denticulosa (15e)
A. dentifera (11l)
A. dictyophleba (11m)
A. divergens (11g)
A. dolichocephala (1i)
A. donnaiensis (3h)
A. doratoxylon (14b)
A. drepanolobium (1j)
A. drummondii (7b)
A. dunnii (13i)
A. elata (6h)
A. elongata (13j)
A. ensifolia (10o)
A. ephedroides (14g)
A. eremophila (12e)
A. eriocarpa (3i)
A. erioloba (1d)
A. eriopoda (14h)
A. estrophiolata (13e)
A. excelsa (13e)
A. extensa (10h)
A. falcata (10b)
A. falciformis (10n)
A. farnesiana (1d)
A. fasciculifera (10j)
A. fauntleroyi (14g)
A. filicifolia (6f)
A. fimbriata (10i)
A. flavipila (13g)
A. flexifolia (11b)
- A. furcata* (5b)
A. genistifolia (11n)
A. gilbertii (7c)
A. glandulifera (1g)
A. glomerosa (3j)
A. gnidium (11k)
A. goetzei (2a)
A. greggii (3k)
A. gunnii (11i)
A. guymeri (14i)
A. haematoxylon (1d)
A. handonis (11o)
A. harmandiana (1k)
A. hayesii (3a)
A. hemiteles (10p)
A. heterophylla (13k)
A. hippuroides (9a)
A. holosericea (15c)
A. horridula (11p)
A. hyaloneura (14i)
A. inaequilatera (10l)
A. inophloia (14g)
A. inopinata (1h)
A. insolita (11q)
A. irrorata (6d)
A. islana (11k)
A. iteaphylla (10p)
A. ixioophylla (13g)
A. jakesiana (14i)
A. jibberdingensis (14k)
A. johnsonii (11k)
A. juncifolia (11c)
A. karroo (1l)
A. kempeana (14l)
A. koa (13k)
A. kuhlmannii (3c)
A. lacerans (3c)
A. lasiocalyx (14b)
A. lasiocarpa (7d)
A. lateriticola (7a)
A. latescens (13i)
A. latipes (13c)
A. leiocalyx (15d)
A. leprosa (11w)
A. leptoclada (6e)
A. leptospermoides (11r)
A. leucoclada (6c)
A. leucopholea (1k)

- A. ligulata* (10a)
A. lineata (11b)
A. lineolata (12a)
A. linifolia (10i)
A. littorea (11s)
A. longifolia (15f)
A. longispicata (15d)
A. lysiphloia (15b)
A. mackeyana (12e)
A. macracantha (1f)
A. maitlandii (11t)
A. mammifera (3b)
A. mangium (15c)
A. mearnsii (6d)
A. megaladena (3g)
A. melanoxydon (13k)
A. mellifera (2c)
A. melvillei (12b)
A. merrallii (11u)
A. microbotrya (10b)
A. mitchellii (6e)
A. modesta (2d)
A. monticola (15b)
A. multisiliqua (13l)
A. multispicata (14a)
A. murrayana (10k)
A. muricata (5c)
A. myrtifolia (10g)
A. neovernicosa (1g)
A. neurophylla (15g)
A. nigrescens (2a)
A. nilotica (1m)
A. nitidula (11h)
A. notabilis (10b)
A. nuperrima (12h)
A. nyssophylla (12c)
A. olgana (14e)
A. omalophylla (12b)
A. oswaldii (12f)
A. pacensis (1n)
A. pachycarpa (15h)
A. papyrocarpa (12b)
A. paradoxa (11v)
A. parramattensis (6d)
A. pendula (12b)
A. pennata (3g)
A. pennatula (1f)
A. penninervis (10n)
- A. pentadenia* (7a)
A. peuce (11x)
A. platycarpa (13i)
A. plectocarpa (14j)
A. podalyriifolia (10i)
A. praelongata (10k)
A. prainii (10p)
A. pravifolia (11d)
A. pringlei (1b)
A. prominens (10i)
A. pruinocarpa (10o)
A. pruinosa (6g)
A. pulchella (7d)
A. purpusii (3j)
A. pycnantha (10i)
A. pycnocephala (11p)
A. pyrifolia (10l)
A. ramulosa (14c)
A. redolens (13m)
A. retinodes (10b)
A. retivenea (13m)
A. rhodophloia (14f)
A. rigens (12a)
A. rigidula (1b)
A. riparia (3a)
A. roemeriana (3d)
A. rossei (11o)
A. rostellifera (10a)
A. rubida (10b)
A. schaffneri (1d)
A. schinoides (6g)
A. schottii (1g)
A. schweinfurthii (3g)
A. scirpifolia (10e)
A. sclerosperma (10a)
A. sedifolia (11o)
A. senegal (2e)
A. sericocarpa (11u)
A. seyal (11)
A. siculiformis (11t)
A. silvestris (6f)
A. simsii (13l)
A. sophorae (15f)
A. sousae (4b)
A. sphaerocephala (1o)
A. spectabilis (6g)
A. spondylophylla (9a)
A. steedmanii (10b)
- A. stenophylla* (12g)
A. stigmatophylla (15a)
A. stowardii (14b)
A. stricta (11v)
A. suaveolens (10p)
A. subangulata (3f)
A. subporosa (13d)
A. sulcata (11h)
A. synchronicia (10m)
A. tenuifolia (3a)
A. tequilana (4b)
A. tetragonophylla (11y)
A. thailandica (3l)
A. tindaleae (11j)
A. tortilis (1p)
A. trachycarpa (15b)
A. translucens (12h)
A. trineura (13m)
A. truncata (11s)
A. tumida (14h)
A. ulicifolia (11i)
A. urophylla (10h)
A. usumacintensis (3b)
A. velutina (3c)
A. velvae (4d)
A. verniciflua (11w)
A. veronica (13d)
A. verticillata (11n)
A. vestita (10i)
A. victoriae (10m)
A. villosa (4c)
A. viscifolia (11z)
A. visco (5d)
A. wilhelmiana (11z)
A. willardiana (3m)
A. willdenowiana (8a)
A. wrightii (3k)
A. xiphophylla (14m)
A. yorkkrakinensis (14e)
Faidherbia albida

