

NOTULAE HEPATICARUM POLYNESIAE

Harvey A. Miller

Department of Biological Sciences
University of Central Florida
Orlando, Florida 32816

Preparation of the catalogue of species for the *Prodromus Florae Hepaticarum Polynesiae* resulted in discovery of nomenclaturally unacceptable names. Many were presented as *nomina nuda* and could be listed without further attention. Others, however, were assigned generic names now considered invalid and revisions were necessary in order that correct names could be provided for each taxon, as well as synonyms. Several later homonyms required new names.

The area included in the *Prodromus Florae Hepaticarum Polynesiae* extends from Hawaii to the Bonin Islands, southward east of the Philippines and New Guinea to Lord Howe, eastward north of New Zealand to Easter. Thus, all tropical Pacific islands outside the indomalayan archipelagoes have been treated with synonyms cross-referenced to the page number of the original report as well as to the source document itself. The following list provides acceptable names for liverwort taxa lacking them in the Pacific area.

Anthoceros javanicoides nom. nov.

Anthoceros javanicus Steph. 1916. Spec. Hep. 5: 988.
nec Anthoceros javanicus Nees 1830. Enum. Pl. Crypt. Javae: 1.
Type specimen: Java, Dien Gbze, Ernst 18 in hb G.

Asterella boryana (Mont.) comb. nov.

Fimbriaria boryana Mont. 1838. Ann. Sci. Nat. 9: 41.

Asterella coronata (Steph.) comb. nov.

Fimbriaria coronata Steph. 1917. Spec. Hep. 6: 12.

Asterella dioica (Steph.) comb. nov.

Fimbriaria dioica Steph. 1914. Jour. Proc. Roy. Acad. New South Wales 48: 104.

Asterella dognyensis nom. nov.

Fimbriaria umbonata Steph. 1917. Spec. Hep. 6: 17.
nec Fimbriaria umbonata (Wallr.) Wallr. in Gott. et al. 1846.
Syn. Hep.: 559.

Type specimen: Nova Caledonia, in jugo Dogny, 1060 m., IX 1909,
Le Rat 219 ex hb Gen. Paris, in hb G.

Asterella heteroflora (Steph.) comb. nov.

Fimbriaria heteroflora Steph. 1917. Spec. Hep. 6: 14.

Asterella latifrons (Steph.) comb. nov.

Fimbriaria latifrons Steph. 1917. Spec. Hep. 6: 15.

Asterella levispora (Steph.) comb. nov.

Fimbriaria levispora Steph. 1917. Spec. Hep. 6: 15.

Asterella longebarbata (Steph.) comb. nov.

Fimbriaria longebarbata Steph. 1889. Hedwigia 28: 156.

Asterella tenerima (Steph.) comb. nov.

Fimbriaria tenerima Steph. 1917. Spec. Hep. 6: 17.

Bazzania confertissima (Steph.) comb. nov.

Mastigobryum confertissimum Steph. 1911. Denkschr. K. Akad. Wiss., Math.-Nat. Kl. Wien 88: 32.

Bazzania conistipula (Steph.) comb. nov.

Mastigobryum conistipulum Steph. 1924. Spec. Hep. 6: 458.

Bazzania consociata (Steph.) comb. nov.

Mastigobryum consociatum Steph. 1924. Spec. Hep. 6: 458.

Bazzania falcifolia (Steph.) comb. nov.

Mastigobryum falcifolium Steph. 1911. Denkschr. K. Akad. Wiss., Math.-Nat. Kl. Wien 88: 33.

Bazzania gunniana (Steph.) comb. nov.

Mastigobryum gunnianum Steph. 1914. Journ. Proc. Roy. Soc. New South Wales 48: 123.

Bazzania hamatifolia (Steph.) comb. nov.

Mastigobryum hamatifolium Steph. 1908. Spec. Hep. 3: 486.

Bazzania hebridensis (Steph.) comb. nov.

Mastigobryum hebridense Steph. 1914. Journ. Proc. Roy. Soc. New South Wales 48: 124.

Bazzania integrifolia (Steph.) comb. nov.

Mastigobryum integrifolium Steph. 1911. Denkschr. K. Akad. Wiss., Math.-Nat. Kl. Wien 88: 33.

Bazzania leratii (Beauverd) comb. nov.

Mastigobryum leratii Beauverd in Steph. 1924. Spec. Hep. 6: 477.

Bazzania lessonii (Steph.) comb. nov.

Mastigobryum lessonii Steph. 1909. Spec. Hep. 3: 531.

Bazzania obcuneata (Steph.) comb. nov.

Mastigobryum obcuneatum Steph. 1908. Spec. Hep. 3: 487.

Bazzania paucidens (Steph.) comb. nov.

Mastigobryum paucidens Steph. 1908. Spec. Hep. 3: 484.

Bazzania pulchella (Steph.) comb. nov.

Mastigobryum pulchellum Steph. 1908. Spec. Hep. 3: 442.

Bazzania quadratistipula nom. nov.

Mastigobryum quadratum Steph. 1924. Spec. Hep. 6: 477.

nec Mastigobryum quadratum Colenso 1886. Trans. New Zealand Inst. 18: 246.

nec Bazzania quadrata (Colenso) Mart. et Hodgs. 1950. Trans. Proc. Roy. Soc. New Zealand 78: 499.

Type specimen: Nova Caledonia, Le Rat legit.

Bazzania squarrosa (Steph.) comb. nov.

Mastigobryum squarrosum Steph. 1908. Spec. Hep. 3: 460.

Bazzania subserrifolia (Beauverd) comb. nov.

Mastigobryum subserrifolium Beauverd in Steph. 1908. Spec. Hep. 480.

Bazzania temariana (Steph.) comb. nov.

Mastigobryum temarianum Steph. 1908. Spec. Hep. 3: 532.

Bazzania upoluensis (Steph.) comb. nov.

Mastigobryum upoluense Steph. 1911. Denkschr. K. Akad. Wiss., Math.-Nat. Kl. Wien 88: 33.

Chiloscyphus rotundiphyllus nom. nov.

Chiloscyphus rotundifolius Steph. 1895. Hedwigia 34: 236.

nec Chiloscyphus rotundifolius Mitt. 1884. Challenger Exped. 1 (3): 85.

Type specimen: New Caledonia, Dent de St. Vincent, VII 1909, Le Rat 189 ex hb Theriot in hb G.

Cololejeunea comptonii (Pears.) comb. nov.

Leptocolea comptonii Pears. 1922. Journ. Linn. Soc. Bot. 46: 40.

Cololejeunea crenulata (Pears.) comb. nov.

Leptocolea crenulata Pears. 1922. Journ. Linn. Soc. Bot. 46: 41.

nec Cololejeunea crenulata (Herz.) Benedix 1953.

Cololejeunea mouensis (Tixier) comb. nov.

Campylolejeunea mouensis Tixier 1979. *Nova Hedwigia* 31: 727.

Cololejeunea polyantha (Mitt.) comb. nov.

Lejeunea polyantha Mitt. in Seem. 1871. *Flora Vitiensis*: 416.
Physocolea polyantha (Mitt.) Steph. 1916. *Spec. Hep.* 5: 901.

Cololejeunea retusula (Mitt.) comb. nov.

Lejeunea retusula Mitt. 1871. *Flora Vitiensis*: 416.
Physocolea retusula (Mitt.) Steph. 1916. *Spec. Hep.* 5: 904.

Cololejeunea spathulifolia (Steph.) comb. nov.

Leptocolea spathulifolia Steph. 1916. *Spec. Hep.* 5: 855.

Cololejeunea vitiensis (E. O. Campbell) comb. nov.

Leptocolea vitiensis E. O. Campbell. 1971. *Jour. Roy. Soc. New Zealand* 1: 15.

Colura tutuilana (Pears.) comb. nov.

Colurolejeunea tutuilana Pears. 1924. *Carnegie Inst. Washington Dept. Marine Biol. Publ.* 20: 151.

Frullania matafaoica nom. nov.

Frullania minutissima Pears. 1924. *Carnegie Inst. Washington Dept. Marine Biol. Publ.* 20: 140.

nec Frullania minutissima Colenso 1887.

nec Frullania minutissima Schiffner 1900.

Type specimen: "Trail to Mount Matafao, M. C. Collarino, June-July 1920 (382a)."

Frullania novopommeriensis nom. nov.

Frullania fauriana Steph. in Rechinger 1909. *Denkschr. K. Akad. Wiss., Math.-Nat. Kl. Wien* 85: 196.

nec Frullania fauriana Steph. 1894. *Hedwigia* 33: 144.

Type specimen: "Insel Neu-Pommern... am Wege von Paparatawa nach Toma, nr. 4598." (Leg. Rechinger)

Jungermannia comptonii (Pears.) comb. nov.

Haplozia comptonii Pears. 1922. *Journ. Linn. Soc. Bot.* 46: 20.

Jungermannia dubioides nom. nov.

Jungermannia dubia Nees in Endlicher 1833. *Prodr. Fl. Norfolk*: 5.

nec Jungermannia dubia Weber 1815. *Hist. Musc. Hep. Prodr.*: 79.

Type specimen: Norfolk Island, leg. F. Bauer.

Lejeunea alobifolia nom. nov.

Lejeunea aloba Steph. 1915. Spec. Hep. 5: 767.

nec Lejeunea aloba Sande Lac. 1856. Syn. Hepat. Jav.: 72.

Type specimen: New Caledonia, "Ile des Pins. Forêt de Kapan, Jun. 1909, Le Rat 53, hb G 14244.

Lejeunea gibbiloba (Steph.) comb. nov.

Eulejeunea gibbiloba Steph. 1923. Spec. Hep. 6: 418.

Lejeunea tutuilana (Pears.) comb. nov.

Eulejeunea tutuilana Pears. 1924. Carnegie Inst. Washington Dept. Marine Biol. Publ. 20: 149.

Lophocolea papulimarginata nom. nov.

Lophocolea papulosa Steph. 1922. Spec. Hep. 6: 286.

nec Lophocolea papulosa Steph. 1911. Kungl. Svenska Vet. Akad. Handl. 46: 50.

Type specimen: New Caledonia, Le Rat legit.

Phaeoceros parisii (Steph.) comb. nov.

Aspiromitus parisii Steph. 1916. Spec. Hep. 5: 968.

Riccardia angustissima (Steph.) comb. nov.

Aneura angustissima Steph. 1917. Spec. Hep. 6: 20.

Riccardia breviramosa (Steph.) comb. nov.

Aneura breviramosa Steph. 1911. Kungl. Svenska Vet. Akad. Handl. 46(9): 6.

Riccardia comata (Steph.) comb. nov.

Aneura comata Steph. 1917. Spec. Hep. 6: 22.

Riccardia comptonii (Pears.) comb. nov.

Aneura comptonii Pears. 1922. Journ. Linn. Soc. Bot. 46: 17.

Riccardia flagellaris (Gepp in Gibbs) comb. nov.

Aneura flagellaris Gepp in Gibbs. 1909. Journ. Linn. Soc. Bot. 39: 194.

Riccardia gunniana (Steph.) comb. nov.

Aneura gunniana Steph. 1914. Journ. Proc. Roy. Soc. New South Wales 48: 96.

Riccardia hebridensis (Steph.) comb. nov.

Aneura hebridensis Steph. 1914. Journ. Proc. Roy. Soc. New South Wales 48: 96.

Riccardia hyalina (Steph.) comb. nov.

Aneura hyalina Steph. 1917. Spec. Hep. 6: 31.

Riccardia intricata (Steph.) comb. nov.

Aneura intricata Steph. 1897. Engler's Bot. Jahrb. 23: 301.

Riccardia lichenoides (Steph.) comb. nov.

Aneura lichenoides Steph. 1897. Engler's Bot. Jahrb. 23: 301.

Riccardia macrantha (Pears.) comb. nov.

Aneura macrantha Pears. 1922. Journ. Linn. Soc. Bot. 46: 17.

Riccardia micropinna (Steph.) comb. nov.

Aneura micropinna Steph. 1893. Hedwigia 32: 94.

Riccardia palmatifida (Steph.) comb. nov.

Aneura palmatifida Steph. 1917. Spec. Hep. 6: 36.

Riccardia robusta (Steph.) comb. nov.

Aneura robusta Steph. 1917. Spec. Hep. 6: 40.

Riccardia tutuilana (Pears.) comb. nov.

Aneura tutuilana Pears. 1924. Carnegie Inst. Washington Dept. Marine Biol. Publ. 20: 132.



Miller, Harvey A. 1981. "Notulae hepaticarum polynesiae." *Phytologia* 47, 319–324. <https://doi.org/10.5962/bhl.part.4461>.

View This Item Online: <https://www.biodiversitylibrary.org/item/47388>

DOI: <https://doi.org/10.5962/bhl.part.4461>

Permalink: <https://www.biodiversitylibrary.org/partpdf/4461>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.