

1.1 Land Use

1.1.1 Data Collection


The Land Use data was obtained from the CSIR Land Cover database utilising a GIS desktop study and supplemented with visual observations on site.







1.1.2 Regional Description

Land-Use is dominated by maize and sunflower farming, coal mines and power stations. Land uses also occurring in the area include Commercial / Industrial, Conservation areas, Cultivated land, Forestry areas, Residential, Subsistence farming and Vacant or Unspecified land, however for the purposes of this report, land use of the region is grouped into urban, cultivation, grassland / plantations, mines and quarries and waterbodies / wetlands.

1.1.3 Site Description

The Land-Use on site is dominated by maize, grazed fields, quarries, residential and conservation. From the map below (Figure 2) it can be seen that the proposed by-pass line and buffer zone include all classes of land use. Water bodies are the only land use regarded as sensitive and as such certain mitigatory measures will be outlined. The study area is located in an area which is predominantly unimproved grassland and this type of land cover is associated with intensive grazing. It was noted that Corobrik make use of a quarry on site to manufacture bricks and that a small section of the ash dump from Duvha Power Station is located on site, the Golden Miles Estate is also located on site (Figure 1). With this in mind Alternative 1 is the most preferred alternative as it avoids the water bodies and also disturbs the shortest section of grazing land before entering the power station site.



	
<p>Corobrik quarry on site</p>	<p>Corobrik quarry on site</p>
	
<p>Golden Mile Residential Estate on site</p>	<p>Golden Mile Residential Estate on site</p>
	
<p>Residential properties surrounding the Witbank Dam</p>	<p>Residential properties surrounding the Witbank Dam</p>
<p>Figure 1: Land Uses encountered on site</p>	




Figure 2: Land Use Map

1.2 Vegetation

1.2.1 Data Collection

The floral study involved extensive fieldwork, a literature review and a desktop study utilizing GIS. The site was investigated during a one week site visit, conducted from the 10th-12th September 2008, in early spring. The area within the servitude was sampled using transects placed at 300m intervals. At random points along the transect an area of 20m x 20m was surveyed. All species within the 20m x 20m quadrant were identified, photographed and their occurrence noted. Sensitive features such as ridges or wetlands were sampled by walking randomly through the area concerned and identifying all species within the area.

The floral data below is taken from The Vegetation of South Africa, Lesotho and Swaziland (Mucina and Rutherford 2006). Also, while on site, the following field guides were used:

- Ü Guide to Grasses of Southern Africa (Frits van Oudtshoorn, 1999);
- Ü Field Guide to Trees of Southern Africa (Braam van Wyk and Piet van Wyk, 1997);
- Ü Field Guide to the Wild Flowers of the Highveld (Braam van Wyk and Sasa Malan, 1998);
- Ü Problem Plants of South Africa (Clive Bromilow, 2001);
- Ü Medicinal Plants of South Africa (Ben-Erik van Wyk, Bosch van Oudtshoorn and Nigel Gericke, 2002)

The occurrence of the species was described as either:

- Ü Very common (>50 % coverage);
- Ü Common (10 – 50 % coverage);
- Ü Sparse (5 – 10 % coverage); and
- Ü Individuals (< 5 % coverage).

1.2.2 Regional Description

According to the South African National Biodiversity Institute, the study area falls within the Grassland Biome, where most of the county's maize production occurs. The study area comprises of the Rand Highveld Grassland, Eastern Highveld Grassland and Eastern Temperate Freshwater Wetlands vegetation units as classified by Mucina and Rutherford^{Error! Bookmark not defined.}. Each of these vegetation units are described in more detail below.

Rand Highveld Grassland

Rand Highveld Grassland is found in the highly variable landscape with extensive sloping plains and ridges in the Gauteng, North-West, Free State and Mpumalanga Provinces. The vegetation type is found in areas between rocky ridges from Pretoria to Emahlaheni, extending onto ridges in the Stoffberg and Roossenekal regions as well as in the vicinity of Derby and Potchefstroom, extending southwards and north-eastwards from there. The vegetation is species rich, sour grassland alternating with low shrubland on rocky outcrops. The most common grasses on the plains belong to the genera *Themeda*, *Eragrostis*, *Heteropogon* and *Elionurus*. High numbers of herbs, especially *Asteraceae* are also found. In rocky areas shrubs and trees also prevail and are mostly *Protea caffra*, *Acacia caffra*, *Celtis africana* and *Rhus spp.*

This vegetation type is poorly conserved (approx 1 %) and has a target of 24 % of the vegetation type to be conserved. Due to the low conservation status this vegetation type is classified as endangered. Almost half of the vegetation type has been transformed by cultivation, plantations, urbanisation or dam-building. Scattered aliens (*Acacia mearnsii*) are present in the unit.

Eastern Highveld Grassland

The Eastern Highveld Grassland is found in the Mpumalanga and Gauteng Provinces on the plains between Belfast in the east and the eastern side of Johannesburg in the west and extending southwards to Bethal, Ermelo and west of Piet Retief. The landscape is dominated by undulating plains and low hills with short dense grassland dominating belong to the genera *Themeda*, *Aristida*, *Digitaria*, *Eragrostis*, *Tristachya* etc. Woody species are prevalent on the rocky outcrops. In terms of conservation and disturbance, 44 % of the vegetation type is already transformed by cultivation, plantations, mines, and urbanisation. No serious alien invasion, but *Acacia mearnsii* can dominate in certain areas

Eastern Temperate Freshwater Wetlands

Another vegetation type associated with the region is the Eastern Temperate Freshwater Wetlands, found around water bodies and embedded within the Grassland biome. Eastern Temperate Freshwater Wetlands are typically found in flat landscapes or shallow depressions filled with (temporary) water bodies supporting zoned systems of aquatic and hydrophilous (water loving) vegetation of temporarily flooded grasslands and ephemeral herblands. Important species include *Cyperus congestus*, *Phragmites australis*, *Marsilea farinose*, *Rorippa fluviatilis*, *Disa zuluensis*, *Crassula tuberella* and the carnivorous herb *Utricularia inflexa*. These wetlands are one of the most sensitive vegetation units found in the region and have been extensively modified by mining and industrial activities in the region.




Figure 3: Regional Vegetation

1.2.3 Site Description







Three main vegetation types were identified, namely Disturbed/Grazed Grassland, Undisturbed/Natural Grassland and Wetland and Riparian communities. Each of these vegetation types are described in more detail below and illustrated in Figure 7 below. The species list for the site is attached in Appendix 1. The species that could occur in the quarter degree grid was obtained from the Plants of Southern Africa (POSA) online database upheld by the South African National Botanical Institute (SANBI) and supplemented with field notes. The list provides species names, common names, as well as notes on which species were observed on site. In total 136 species could occur in the area with 43 confirmed species.

Disturbed/Grazed Grassland

Disturbed grassland or other disturbed areas such as road reserves or fallow fields, not cultivated for some years, are also usually *Hyparrhenia* dominated. However, while *Hyparrhenia* – is present in this vegetation unit, it is not dominate. This grassland is a result of high disturbance as a result of over-grazing, characterised by Bankrupt Bush (*Stoebe vulgaris*), an invader dwarf shrub which usually indicates grassland's degraded condition.

This grassland mostly has low species richness, with only a few other species able to establish or survive in the shade of the dense sward of tall grass. Most of these species are relict pioneers or early seral species. The most prominent species include the grasses *Cynodon dactylon*, *Eragrostis plana*, *E. racemosa*, *E. curvula* and *E. capensis*. Herbaceous species such as *Anthospermum rigidum*, *Conyza podocephala*, *Crabbea angustifolia* and *Helichrysum rugulosum* are present. Alien species such as *Acacia mearnsii* (Black Wattle) have also invaded this vegetation unit.


Figure 4 below provides an illustration of the Disturbed/Grazed Grassland unit.

	
<p>Disturbed/Grazed Grassland with the Duvha Power Station in the background.</p>	<p><i>Dovyalis zeyheri</i> (Wild Apricot) on a rocky outcrop that forms parts of the Disturbed/Grazed Grassland.</p>
	
<p><i>Stoebe vulgaris</i> (Bankrupt bush) characterising the Disturbed/Grazed Grassland.</p>	<p>Disturbed/Grazed Grassland with invasive <i>Acacia mearnsii</i> (Black wattle) in the background.</p>
	
<p>Disturbed/Grazed Grassland with the Corobrik quarry in the background.</p>	<p>Disturbed/Grazed Grassland with <i>Acacia mearnsii</i> (Black wattle).</p>
<p>Figure 4: Disturbed/Grazed Grassland</p>	

Undisturbed/Natural Grassland

This grassland comprises both the Eastern Highveld Grassland and the Rand Highveld Grassland and is dominated by the grasses of these vegetation types (Figure 5).

The vegetation is species rich located on a landscape is dominated by undulating plains and low hills with short dense, sour grassland alternating with low shrubland on rocky outcrops. The most common grasses on the plains belong to the genera *Themeda*, *Eragrostis*, *Heteropogon*, *Aristida*, *Digitaria*, *Tristachya* and *Elionurus*. High numbers of herbs, especially *Asteraceae* are also found. In rocky areas shrubs and trees also prevail and are mostly *Protea caffra*, *Acacia caffra*, *Celtis africana* and *Rhus spp.*




Wetland and Riparian communities

Wetland and Riparian communities are seasonally wet areas that occur in sandy areas where water seeps into lowlying drainage lines after rains. These areas are usually covered by hygrophytes such as sedges and reeds. The dominant sedge in the study area is *Juncus rigidus*. Sometimes bulrush (*Typha capensis*) and reeds (*Phragmites australis*) also occurs.

Wetlands are of a more permanent nature and occur in low-lying areas such as tributaries of streams and rivers. Wetlands are typically found in flat landscapes or shallow depressions filled with (temporary) water bodies supporting zoned systems of aquatic and hydrophilous (water loving) vegetation of temporarily flooded grasslands and ephemeral herblands. Typical plants are the

Orange River Lily (*Crinum bulbispermum*), bulrush (*Typha capensis*) and reeds (*Phragmites australis*), sedges of the *Cyperus*, *Fuirena* and *Scirpus* genera also occur (Figure 6).



1.2.4 Red data Flora Species

The Mpumalanga Parks Board provided information as to sensitive plant species occurring in the area and it was found that only one sensitive plant species, *Hypoxis hemerocallidea*, occurs in the quarter degree square 2529CD. *Hypoxis hemerocallidea* is rated as “Least Concern” in Mpumalanga and throughout the rest of South Africa, however populations are declining. It should be noted that during the site visit, *Hypoxis hemerocallidea* was not found to occur on site; however its presence can not be completely excluded.




Figure 7: Vegetation units found on site

1.3 Fauna

1.3.1 Data Collection

A literature review of the faunal species that could occur in the area was conducted. C-Plan data provided from the Mpumalanga provincial department was used to conduct a desktop study of the area. This data consists of terrestrial and aquatic components, ratings provide an indication as to the importance of the area with respect to biodiversity. Additionally, all fauna were noted during the site visit conducted on the 17th – 20th November 2008.

1.3.2 Regional Description

As a consequence of mining and farming in the area, it appears that only small animals are to be found at the site. Small mammals known to occur in the area include hedgehog, rabbits, polecat, meerkat and the ubiquitous rats and mice. Given the habitat, it is likely that korhaans, larks, longclaws, species of Euplectes (bishops and widows), weavers, starlings and sparrows occur in the grassland.

The area surrounding the proposed loop in lines does include areas of terrestrial and aquatic habitats. These areas should be treated as sensitive and should therefore be managed accordingly; if feasible they should be avoided.

1.3.3 Site Description

The scope of work indicated that an avifauna assessment was required. An avifaunal assessment will be undertaken and the report will be provided as part of the EIA. All herpetofauna and mammals observed on site were noted during the site visit.

Habitat

The habitat on site is described in the vegetation site description in Section 3.8.3 above. All of the vegetation types identified have been disturbed to a certain extent, as the main land use in the area is grazing of livestock. The largest portion of the site is comprised of Undisturbed Grassland, totalling approximately 50.1 % of the study site. The remainder of the site comprises Disturbed Grassland and Riparian and Wetland zones. All of these are suitable habitat to a number of protected species found in the region.

Species potentially occurring on site

A detailed list of the species potentially occurring on site is attached in Appendix 2.

Herpetofauna

Herpetofauna could potentially occur in all the habitat types. The Riparian and Wetland zones could potentially support amphibians representative of the region, specifically *Pyxicephalus adspersus* (African Bullfrog) which is a species rated as “near threatened” and is a protected species in South Africa.

The quarter degree square is known to contain *Geochelone pardalis* (Leopard tortoise), *Aparallactus capensis* (Cape Centipede Eater), *Atractaspis bibronii* (Southern or Bibron’s Burrowing Asp), *Causus rhombeatus* (Common Night Adder), *Crotaphopeltis hotamboeia* (Herald or Red-lipped Snake), *Dasypeltis scabra* (Common or Rhombic Egg Eater), *Hemachatus haemachatus* (Rinkhals), *Lycodonomorphus rufulus* (Common Brown Water Snake), *Naja annulifera annulifera* (Snouted Cobra), *Psammophylax tritaeniatus* (Striped Skaapsteker), *Agama atra* (Southern Rock Agama), *Bitens arietans* (Puff Adder), *Cordylus vittifer* (Transvaal Girdled Lizard), *Gerrhosaurus flavigularis* (Yellow Throated Plated Lizard), *Lygodactylus ocellatus* (Spotted Dwarf Gecko), *Pachydactylus affinis* (Transvaal Thick-toed Gecko), *Telescopus semiannulatus semiannulatus* (Eastern Tiger Snake), *Psammophis brevirostris brevirostris* (Leopard or Short-snouted Grass Snake) and *Varanus niloticus* (Water Monitor). *Hemachatus haemachatus* (Rinkhals), *Psammophis brevirostris brevirostris* (Leopard or Short-snouted Grass Snake) and *Cordylus vittifer* (Transvaal Girdled Lizard) are endemic to Southern Africa, while *Lygodactylus ocellatus* (Spotted Dwarf Gecko) and *Pachydactylus affinis* (Transvaal Thick-toed Gecko) are endemic to South Africa.

None of the above mentioned Herpetofauna were encountered on site during the site visit that took place from the 17th-20th November 2008.

Avifauna

Avifauna that could potentially occur on site is provided in Table 1 below. The avifaunal assessment (Appendix 3) focused on identifying a preferred alignment for the new Bravo-Vulcan 400kV line from a bird impact perspective, and the description of associated impacts on birds. Recommendations were also provided to mitigate for potential impacts.

Table 1: Avifauna Species List

Species	Common name
<i>Phalacrocorax africanus</i>	Reed Cormorant
<i>Ardea cinerea</i>	Grey Heron
<i>Ardea melanocephala</i>	Blackheaded Heron
<i>Bubulcus ibis</i>	Cattle Egret
<i>Bostrychia hagedash</i>	Hadedda Ibis
<i>Plegadis falcinellus</i>	Glossy Ibis
<i>Alopochen aegyptiacus</i>	Egyptian Goose
<i>Elanus caeruleus</i>	Blackshouldered Kite
<i>Francolinus swainsonii</i>	Swainson's Francolin
<i>Numida meleagris</i>	Helmeted Guineafowl
<i>Fulica cristata</i>	Redknobbed Coot
<i>Gallinula chloropus</i>	Moorhen
<i>Anthropoides paradisea</i>	Blue Crane
<i>Sagittarius serpentarius</i>	Secretary Bird
<i>Eupodotis cafra</i>	Whitebellied Korhaan
<i>Vanellus armatus</i>	Blacksmith Plover
<i>Vanellus coronatus</i>	Crowned Plover
<i>Streptopelia semitorquata</i>	Redeyed Dove
<i>Streptopelia senegalensis</i>	Laughing Dove
<i>Asio capensis</i>	Marsh Owl
<i>Colius striatus</i>	Speckled Mousebird
<i>Mirafra africana</i>	Rufousnaped Lark
<i>Corvus albus</i>	Pied Crow
<i>Saxicola torquata</i>	Stone Chat
<i>Phylloscopus trochilus</i>	Willow Warbler
<i>Cisticola fulvicapilla</i>	Neddicky
<i>Motacilla clara</i>	Cape Wagtail
<i>Anthus cinnamomeus</i>	Grassveld Pipit
<i>Passer domesticus</i>	House Sparrow
<i>Ploceus velatus</i>	Masked Weaver
<i>Euplectes orix</i>	Red Bishop
<i>Emberiza capensis</i>	Cape Bunting

The species that could potentially occur on site include waterfowl, grassland specialists and common generalists. This is attributed to the variety of habitats that occur on site, as well as the adequate supply of fresh water.

Sensitive species known to occur in the quarter degree square include *Oxyura maccoa* (Maccoa Duck) and *Geronticus calvus* (Bald Ibis) which is endemic to South Africa.

The following conclusions from the avifaunal impact assessment are put forward:

- Ü A number of power line sensitive, Red Data species could potentially occur along any of the alignments, although the occurrence of these species would be the exception rather than the rule.
- Ü The proposed power line, unless mitigated, will pose a limited collision risk to power line sensitive bird species in the study area. Another potential risk is the destruction of sensitive wetland habitat through the construction of access roads.
- Ü Of the Red Data species potentially present in the area, none are particularly at risk by the power line due to the very small densities at which the species occur. The latter is a result of the extensive habitat degradation that has taken place.
- Ü There is, however, a substantial risk of collisions for several non Red Data species which warrants the application of mitigation measures.

The following recommendations are put forward:

- Ü A sensitivity map indicating the areas where anti-collision devices need to be applied to the proposed line is shown in Figure 8 below.
- Ü The construction of access roads in sensitive wetland habitat should be avoided.



Figure 8: Avifaunal Sensitivity Map

From the analysis undertaken in the avifaunal impact assessment, Alternative 1 is the preferred alignment from a bird interaction perspective.

Mammals

Large mammals have to a large extent been removed from the area and the only indication of large mammal species that could have previously occurred in the area are re-introduced mammals found on a few game farms and lodges encountered during the site visit. These include Springbok (*Antidorcas marsupialis*), Blesbok (*Damaliscus dorcas phillipsi*), Blue Wildebeest (*Connochaetes taurinus*) and Burchell's Zebra (*Equus burchelli*). During the site visit, Yellow Mongoose (*Cynictis pencilata*) were spotted as well as signs of other small mammals such as droppings. Other small mammals known to occur in the area include Hedgehog (*Atelerix frontalis*), Striped Polecat (*Ictonyx striatus*), Suricate / Meerkat (*Suricata suricatta*), Aardvark / Antbear (*Orycteropus afer*) and the ubiquitous rats and mice. Sensitive mammal species that could occur in the quarter degree square 2529CD include *Genetta tigrina* (Large-spotted Genet), *Lepus saxatilis* (Scrub hare), *Hyaena brunnea* (Brown Hyaena), *Sylvicapra grimmia* (Common/Grey Duiker), *Tragelaphus scriptus* (Bushbuck), *Vulpes chama* (Cape Fox) None of these species were identified on site.

1.4 Wetland and Riparian Zone Delineation

1.4.1 Riparian Zones vs. Wetlands

Wetlands

The riparian zone and wetlands were delineated according to the Department of Water Affairs and Forestry (DWAF) guideline, 2003: A practical guideline procedure for the identification and delineation of wetlands and riparian zones. According to the DWAF guidelines a *wetland* is defined by the National Water Act as:

“land which is transitional between terrestrial and aquatic systems where the water table is usually at or near surface, or the land is periodically covered with shallow water, and which land in normal circumstances supports or would support vegetation typically adapted to life in saturated soil.”

In addition the guidelines indicate that wetlands must have one or more of the following attributes:

- Ü Wetland (hydromorphic) soils that display characteristics resulting from prolonged saturation;
- Ü The presence, at least occasionally, of water loving plants (hydrophytes); and

- Ü A high water table that results in saturation at or near surface, leading to anaerobic conditions developing in the top 50 centimetres of the soil.

During the site investigation the following indicators of potential wetlands were identified:

- Ü Terrain unit indicator;
- Ü Soil form Indicator;
- Ü Soil wetness indicator; and
- Ü Vegetation indicator.

Riparian Areas

According to the DWAF guidelines a *riparian area* is defined by the National Water Act as:

“Riparian habitat includes the physical structure and associated vegetation of the areas associated with a watercourse which are commonly characterised by alluvial soils, and which are inundated or flooded to an extent and with a frequency sufficient to support vegetation of species with a composition and physical structure distinct from those of adjacent land areas”

The difference between Riparian Areas and Wetlands

According to the DWAF guidelines the difference between a wetland and a riparian area is:

“Many riparian areas display wetland indicators and should be classified as wetlands. However, other riparian areas are not saturated long enough or often enough to develop wetland characteristics, but also perform a number of important functions, which need to be safeguarded... Riparian areas commonly reflect the high-energy conditions associated with the water flowing in a water channel, whereas wetlands display more diffuse flow and are lower energy environments.”

1.4.2 Delineation

The site was investigated for the occurrence of wetlands and riparian areas, using the methodology described above and described in more detail in the DWAF guidelines.

Terrain Unit Indicator

The terrain on site varies from 1600 mamsl to 1520 mamsl as illustrated in **Error! Reference source not found.** From **Error! Reference source not found.** it can be seen that the site is located in an area of undulating hills with the dominant terrain units on site being the midslope,

footslope and valley bottom units. According to the DWAF guidelines the valley bottom is the terrain unit where wetlands are most likely to occur, but they are not excluded from any of the other terrain units.

Soil Form Indicator

The site is located on a slope that drains towards the Witbank Dam. Water enters the soils profile and then flows through the profile down-slope. This action of water movement through the slope typifies the soils of the largest part of the site (eluvial and plinthic soils). Closer to the dam (within the valley bottom terrain unit) the soils gradually deepen due to the down-slope transport of soil (colluvium). In addition these soils have gradually higher percentages of clays that over time have been washed down-slope and accumulate at the valley bottom where the slope angle reduces. The detailed soil mapping exercise was limited to the footslope and valley bottom area in order to delineate the wetland / riparian zones.

During a three day site visit the soils on site were identified (Refer to Section 3.5). Of the soils identified on site the Katspruit soil form is indicative of the permanent wetland zone.

Soil Wetness Indicator

The soils on site were subjected to a soil wetness assessment. If soils showed signs of wetness within 50 cm of the soil surface, it was classified as a hydromorphic soil and divided into the following groups:

Temporary Zone

- Ü Minimal grey matrix (<10%);
- Ü Few high chroma mottles; and
- Ü Short periods of saturation.

Seasonal Zone

- Ü Grey matrix (>10%);
- Ü Many low chroma mottles present; and
- Ü Significant periods of wetness (>3 months / annum).

Permanent Zone

- Ü Prominent grey matrix;

- Ü Few to no high chroma mottles;
- Ü Wetness all year round; and
- Ü Sulphuric odour.


The Katspruit soil form had signs of wetness within the top 50 cm of the soil profile. The Katspruit soil form was classified as the permanent zone, the temporary and seasonal zone could not be delineated since the soil forms diagnostic of these zones probably occur at a depth greater than that of the soil auger used. Therefore, while the temporary and seasonal zones could not be delineated, they are still likely to occur. The soil forms are illustrated in **Error! Reference source not found.**

Vegetation Indicator

The vegetation units on site are described in Section 3.8.3 above and illustrated in Figure 7. The vegetation found in the Undisturbed/Natural Grassland and the Riparian and Wetland zone vegetation units both have species present to indicate the presence of wetlands.

1.4.3 Wetlands and Buffer Zones

According to the methodology that was followed for delineation of wetlands by DWAF, the permanent zone classifies as a wetland present on site. It should however be noted that several of the so-called wetlands could also be classified as riparian zones as they follow the drainage path of the non-perennial streams on site. All the areas identified above perform critical ecosystem functions and also provide habitat for sensitive species. It is suggested that a 50 m buffer be placed from the edge of the permanent zone in order to sufficiently protect the wetlands and riparian zones. Figure 9 above illustrates the various wetland zones as well as the buffer placed along the edge of the permanent zone. From the figure it is once again clear that Alternative 1 is the best alignment, as it avoids most of the sensitive wetlands as well as the buffer zones.



CLIENT CODE: ZIT001	PROJ CODE: ESC 228	REF NO: 01	DATE DRAWN: 2008/11/26 AUTHOR: G. Louwrens	PROJECTION: WGS 84 Hartebeesthoek	SCALE: 1:91,092	DATA SOURCES: Survey General's Office Eskom WR 90
-------------------------------	------------------------------	----------------------	---	--	---------------------------	---

Figure 9: Wetland and Riparian Zone Map

1.5 Biodiversity Rating

In order to quantify the sensitivity of the fauna, flora and wetlands, a biodiversity assessment is undertaken.

1.5.1 Biodiversity Assessment Methodology

Each vegetation unit and its associated fauna were subjected to a biodiversity assessment according to the following methodology. The biodiversity of an area is measured as a combination of the variety of species and habitats within the area, as well as the ecological processes and functional value of the site. This can be captured in two broader categories namely conservation status and functional status. The conservation status encompasses species diversity, habitat diversity and ecological processes. The functional status encompasses ecological services and human use services.

It is suggested, due to the number of variables to be considered, that the following scoring system is used to first determine the value of each of the components (conservation status and functional status) from which the overall biodiversity value is determined.

Conservation status

The conservation status of a particular habitat / vegetation unit is determined using the methodology described in Table 2 below. The conservation status encompasses species diversity, habitat diversity and ecological processes. Each of the habitats found on site are rated accordingly in Section 3.11.2 below.

Table 2: Conservation Status Determination

A. How much of the larger vegetation type or system of which the defined area is a representative example, still exists?	Rating
Only a small area still exists (< 500km ²)	5
A moderate area still exists (500 to 1000 km ²)	3
A large areas still exist (> 1000 km ²)	1
B. What is (based on a qualitative assessment) the species and habitat diversity of the defined area?	Rating
Noticeably high	5
Difficult to assess	3
Obviously low	1
C. What is the condition (qualitative assessment) of the defined area?	Rating
Pristine and largely undisturbed	5
Moderately disturbed	3
Highly disturbed	1

The possible results for the conservation status of the defined area are based on a combination of the attributes, as follows.

$$A \text{ (Size)} + B \text{ (Diversity)} + C \text{ (Condition)} = \text{Conservation Status}$$

Based on the combined score, the conservation status can range from very high to low, as described below in Table 3:

Table 3: Conservation Status Rating

Conservation Status	Rating
High conservation status, needs to be maintained and improved	11 – 15
Moderate conservation status, heavily disturbed and will require improvement	6 – 10
Low conservation status, heavily reduced and of limited value.	3 – 5

Functional status

The functional status encompasses ecological services and human use services. All these elements are rated according to the methodology described in Table 4 below. A detailed rating of each habitat is given in Section 3.11.2 below.

Table 4: Functional Status Determination

A. Are there currently any signs of obvious recreational use of the area, such as walking/hiking, bird watching, mountain biking, fishing, etc?	Rating
Obvious signs of regular use	5
Signs of periodic use	3
No noticeable signs of use	1
B. Does the area carry out any ecological service, such as water purification, flood attenuation, riverbank stabilisation, soil stabilisation, etc?	Rating
Has an obvious functional role	5
Difficult to determine its functional role	3
Clearly has no to very limited functional role	1
C. Does the area serve an aesthetic role?	Rating
Forms part of a larger landscape that is widely visible and has a high aesthetic appeal	5
Forms part of a landscape that has high aesthetic appeal but which is not widely visible	3
Forms part of a landscape that has low aesthetic appeal	1

The possible results for the functional status of the defined area are based on a combination of the attributes, as follows.

$$A \text{ (recreational use)} + B \text{ (ecological service)} + C \text{ (aesthetic value)} = \text{Functional Status}$$

Based on the combined score, the functional status can range from very high to low as illustrated in Table 5 below:

Table 5: Functional Status Rating

Functional Status	Rating
High service value	11 – 15
Moderate service value	6 – 10
Low service value	3 – 5

Biodiversity value

The perceived biodiversity value of an area to human development is not always easy to describe, but it includes the natural system and its variety of species, the ecological processes and the service or functional value that it provides. The combination of the conservation status and functional status scores provides a ranking of the overall biodiversity value for a defined area, as shown in the matrix in Table 6 below.

Table 6: Biodiversity Value Rating

Conservation status	Functional status		
	High service value	Moderate service value	Low service value
High	High	High	Moderate
Moderate	Moderate	Moderate	Low
Low	Moderate	Low	Low

1.5.2 Biodiversity Rating

The following vegetation units were identified on site:

- Ü Undisturbed/Natural grassland;
- Ü Disturbed/Grazed Grassland; and
- Ü Wetland and Riparian zones.

Each of the abovementioned vegetation units are rated for their biodiversity value below.

Undisturbed/Natural grassland

This vegetation unit has a **high** biodiversity rating as indicated in Table 7 below. The **high** conservation value is attributed to the high grassland species diversity in the unit and the large area of grassland not conserved remaining. The **high** functional rating is attributed to the obvious ecological services and the high aesthetic value of the grassland.

Table 7: Biodiversity Rating for the *Undisturbed/Natural grassland* unit

Conservation status	Size of vegetation unit	Species diversity	Condition
	3 – Moderate	5 - High	3 – Moderately Disturbed
Functional status	Use	Ecological service	Aesthetic value
	3 – Periodic	5 – Obvious	5 - High
Biodiversity Rating	Conservation status	Functional status	Biodiversity
	11 – High	13 - High	High

Disturbed/Grazed Grassland

This vegetation unit has a **moderate** biodiversity rating as indicated in Table 8 below. The **moderate** conservation value is attributed to the moderate grassland species diversity in the unit and the moderate area of moist grassland remaining. The **moderate** functional rating is attributed to the moderate ecological service and the moderate aesthetic value of this grassland.

Table 8: Biodiversity Rating for the *Disturbed/Grazed grassland* unit

Conservation status	Size of vegetation unit	Species diversity	Condition
	3 – Moderate	3 - Moderate	3 – Moderately Disturbed
Functional status	Use	Ecological service	Aesthetic value
	3 – Periodic	3 – Difficult to determine	3 - Moderate
Biodiversity Rating	Conservation status	Functional status	Biodiversity
	9 – Moderate	9 - Moderate	Moderate

Wetland and Riparian zones

This vegetation unit has a **high** biodiversity rating as indicated in Table 9 below. The **high** conservation value is attributed to the high grassland species diversity in the unit and the small area of wetlands remaining. The **high** functional rating is attributed to the obvious ecological services and the high aesthetic value of the wetlands and seepage areas.

Table 9: Biodiversity Rating for the seepage areas and wetlands

	Size of vegetation unit	Species diversity	Condition
Conservation status	5 – Small	5 – High	3 – Moderately Disturbed
	Use	Ecological service	Aesthetic value
Functional status	1 – none	5 – Obvious	5 - High
	Conservation status	Functional status	Biodiversity
Biodiversity Rating	13 – High	11 - High	High




Figure 10: Biodiversity Rating Map

2.0 VISUAL IMPACT ASSESSMENT

2.1 Introduction

The site and surrounding area may be characterised as agricultural land utilised mainly for the grazing of cattle. The topography of the region and study site is gently undulating to moderately undulating landscape of the Highveld plateau.

The proposed power lines are located in the area north-west of the Duvha Power Station with the power station and other existing power lines featuring prominently in the landscape. The Witbank Dam and its associated drainage features represent other significant features in the landscape.

2.2 Methodology

The methodology adopted for the visual assessment includes the following tasks:

- Ü Examine the baseline information (contours, building dimensions, vegetation, inter alia);
- Ü Determine the area from which any of the upgrade may be visible (viewshed);
- Ü Identify the locations from which views of the upgrade may be visible (observation sites), which include buildings and roads;
- Ü Analyse the observation sites to determine the potential level of visual impact that may result from the upgrade; and

Each component of the assessment process is explained in detail in the following sections of the Report.

2.2.1 The Viewshed

The viewshed represents the area from which the proposed site would potentially be visible. The extent of the viewshed is influenced primarily by the combination of topography and vegetation, which determine the extent to which the site would be visible from surrounding areas.

The viewshed was determined by Cymbian through the following steps and presumptions:

- Ü The likely viewshed was determined by desktop study (ArcGIS) using contour plans (20 m interval); and
- Ü An offset of 2 m (maximum) for the observer and an offset of 30 m (maximum) for the proposed power lines were utilized during the spatial analysis.

2.2.2 Visibility Assessment

Site visibility is an assessment of the extent to which the proposed upgrade would potentially be visible from surrounding areas. It takes account of the context of the view, the relative number of viewers, duration of view and view distance.

The underlying rationale for this assessment is that if the proposed upgrade (power lines) is not visible from surrounding areas then the development will not produce a visual impact. On the other hand if one or more power lines are highly visible to a large number of people in surrounding areas then the potential visual impact is likely to be high.

Based on a combination of all these factors an overall rating of visibility was applied to each observation point. For the purpose of this report, categories of visibility have been defined as high (H), moderate (M) or low (L).

2.2.3 Assessment Criteria

For the purpose of this report, the quantitative criteria listed in Table 10 have been determined and used in the Visibility Assessment. The criteria are defined in more detail in the subsection following.

Table 10: Visual Impact Assessment Criteria

CRITERIA	DEFINITIONS
Category of Viewer	
<i>Static</i>	<i>Farms, homesteads or industries</i>
<i>Dynamic</i>	<i>Travelling along road</i>
View Elevation	
<i>Above</i>	<i>Higher elevation than proposed upgrade.</i>
<i>Level</i>	<i>Level with upgrade view</i>
<i>Below</i>	<i>Lower elevation than upgrade viewed</i>
View Distance	
<i>Long</i>	<i>> 5 km</i>
<i>Medium</i>	<i>1 – 5 km</i>
<i>Short</i>	<i>200 m – 1000 m</i>
<i>Very Short</i>	<i>< 200 m</i>
Period of View	
<i>Long Term</i>	<i>> 120 minutes</i>
<i>Medium Time</i>	<i>1 – 120 minutes</i>
<i>Short Term</i>	<i>< 1 minute</i>

Category Viewer

The visibility of the upgrade will vary between static and dynamic view types. In the case of static views, such as views from a farmhouse or homestead, the visual relationship between an upgrade and the landscape will not change. The cone of vision is relatively wide and the viewer tends to scan back and forth across the landscape.

In contrast views from a moving vehicle are dynamic as the visual relationship between the upgrade / structures is constantly changing as well as the visual relationship between the upgrade and the landscape in which they are seen. The view cone for motorists, particularly drivers, is generally narrower than for static views.

View Elevation

The elevation of the viewer relative to the object observed, which in this case are the upgrade / structure, significantly influences the visibility of the object by changing the background and therefore the visual contrast. In situations where the viewer is at a higher elevation than the building/structure it will be seen against a background of landscape. The level of visual contrast between the upgrade and the background will determine the level of visibility. A white/bright coloured structure seen against a background of dark/pale coloured tree-covered slopes will be highly visible compared to a background of light coloured slopes covered by yellow/brown dry vegetation.

In situations where the viewer is located at a lower elevation than the proposed upgrade it will mostly be viewed against the sky. The degree of visual contrast between a white coloured structure will depend on the colour of the sky. Dark grey clouds will create a significantly greater level of contrast than for a background of white clouds.

View Distance

The influence of distance on visibility results from two factors:

- Ü With increasing distance the proportion of the view cone occupied by a visible structure will decline; and
- Ü Atmospheric effects due to dust and moisture in the air reduce the visual contrast between the structure and the background against which they are viewed.

Period of View

The visibility of structures will increase with the period over which they are seen. The longer the period of view the higher the level of visibility. However, it is presumed that over an extended period the level of visibility declines as people become accustomed to the new element in the landscape.

Long term views of the upgrade will generally be associated with rest camps located within the viewshed. Short term and moderate term views will generally relate to tourist moving through the viewshed mostly by vehicle.

Site Visibility

The procedure followed by Cymbian to assess Site Visibility involved:

- Ü Generate a viewshed analysis of the area utilizing ArcGIS 9.
- Ü Determine the various categories of observation points (e.g. Static, Dynamic).

2.2.4 Impact Assessment Methodology

Visual impact is defined as the significance and/or severity of changes to visual quality of the area resulting from a development or change in land use that may occur in the landscape.

Significance or severity is a measure of the response of viewers to the changes that occur. It represents the interaction between humans and the landscape changes that they observe. The response to visible changes in the landscape may vary significantly between individuals.

Perception results from the combination of the extent to which the proposed upgrade is visible (level of visibility) and the response of individuals to what they see. A major influence on the perception of people/tourist in relation to the proposed upgrade will be the visual character and quality of the landscape in which it would be located. Natural landscape areas such as national parks, mountain areas or undeveloped sections of coast are valued for their high visual quality. The introduction of buildings and associated infrastructure may be seen as a negative impact on these areas of high visual quality. In the case of rest camps many people perceive them in a positive manner because they represent tourism/conservation infrastructure usually elegantly designed, non-conspicuous and contributing the local and national economy.

The potential visual impact of the proposed upgrade will primarily result from changes to the visual character of the area within the viewshed. The nature of these changes will depend on the level of

the visual contrast between buildings/structures and the existing landscape within which they would be viewed.

The degree of contrast between the upgrade and the surrounding landscape will result from one or more of the following visual characteristics:

- Ü Colour;
- Ü Shape or form;
- Ü Scale;
- Ü Texture; and
- Ü Reflectivity.

2.3 Visual Character

2.3.1 Landscape Character

The site and the surrounding area can be described as an agricultural landscape with intermittent mining and power generation activities. Elevations along the slope range from 1520 mamsl and 1600 mamsl. All the power line alternatives are located on this slope with very little screening from topography or vegetation, however the presence of numerous existing power lines in the area would provide a screening effect for the proposed lines. Please refer to **Error! Reference source not found.** for the topography of the site.

The major drainage features in the area include the Witbank Dam and the Olifants River. Alternatives 2 and 3 cross large sections of the Witbank Dam while Alternative 1 crosses only a small section of the unnamed tributaries. For an illustration of the surface water features please refer to **Error! Reference source not found.**

The landscape surrounding the proposed power lines can be described as open grassland with numerous fields used for grazing. In addition a large section of the site is occupied by the Witbank Dam. The natural vegetation does not provide any screening of the power lines. There are several existing power lines on site, and in deed the intention of the project is to connect existing power lines and by-pass the Duvha Power Station.

Residential homes, mining infrastructure and the Duvha Power Station comprise infrastructure in the area. There numerous formal and informal roads in the area.

2.3.2 Viewshed

It should be noted that the viewshed for each of the alternatives, which is plotted on Figure 11, Figure 12 and Figure 13, is an approximation that may vary in some locations. Potential views to the proposed upgrade are likely to be blocked in some localised situations by buildings, vegetation or local landform features at specific locations within the viewshed. Similarly, glimpses of the proposed upgrade may be visible from some isolated high-elevation locations outside the plotted viewshed. The figures illustrate the visibility of each of the alternatives. The coloured areas indicate areas that are visible with the red areas having very high visibility and the green having lower visibility. It should be noted that Alternatives 2 and 3 are more visible than Alternative 1 due to the fact that they are located along the Witbank Dam while Alternative 1 is located within a less sensitive visually.

Notable features of the viewshed are summarised by the following points:

- Ü The viewshed for Alternative 1 is low to moderate while Alternatives 2 and 3 are moderate to high;
- Ü The area in the immediate vicinity of Alternative 2 has a high viewshed, this is compounded by it traversing the Witbank Dam; and
- Ü To the south of Alternative 3, the viewshed reaches a moderate to high;

2.4 **Impact Assessment**

The visual simulations prepared by Cymbian illustrate the extent to which the upgrade will be visible from key observation points (static and dynamic views).

The vertical form/dimensions of the buildings/structures would be hidden by their location among existing buildings and within a well vegetated area. The visual contrast is increased by the “shape” and scale of the buildings/structures, which generally will not be viewed along the skyline.

Static Views

The upgrade would potentially be visible from the surrounding farmland and the high-lying areas, as well as isolated areas in Witbank. Middelburg remains outside of the viewshed. The potential number of viewers from the surrounding farmland should be low as the farmlands are quite sparsely populated but the views would vary greatly depending on site specific conditions like the orientation of the homes as well as the location of other buildings, fences, vegetation and localized landforms. All these elements have the potential to block views from the surrounding areas to the proposed upgrade.

Dynamic Views

The power lines will be visible to a moderate number of viewers, mainly those travelling along the N4 highway and some travellers along the N12 highway. Views from the N4 extend approximately 9 km and represent a view period of approximately 36 seconds travelling at 120 km/h. The level of visibility would be moderate to high due to the view distance of less than 5 km. The effects are similar for the N12. Please refer to Table 11 for a summary of the dynamic impacts. This assessment is similar for all three route alternatives.

The power line upgrade would also be visible from several farm roads which are located around the proposed site. The viewing distance varies between 1 and 11 km for these roads and if the viewing distance is less than 2 km, the potential visual impact would be considered as moderate.

Table 11: Dynamic Impact Table

Road Name	Speed limit (km/h)	Length of Road (km)	Approximate Period of View (sec)	View Distance
Alternative 1 – N4	120	2.6	78	6.2 – 11 km
Alternative 2 – N4	120	1.2	36	8.2 – 11 km
Alternative 3 – N4	120	2.6	78	6.2 – 11 km
Alternative 1 – N12	120	2.2	79	5 – 6 km
Alternative 2 – N12	120	2.8	101	5 – 6 km
Alternative 3 – N12	120	2.2	79	5 – 6 km




Figure 11: Viewshed from the Alternative 1 alignment.




Figure 12: Viewshed from the Alternative 2 alignment




Figure 13: Viewshed from the Alternative 3 alignment

Conclusion

Table 12 lists the observation points together with the category of viewer, context of view, relative numbers of viewers and approximate distance of observation point to the proposed site. The location of these observation points are shown in Figure 11 and Figure 12.

Table 12: Visual Impact Matrix

<i>Potential Observation Point</i>	<i>Category of Potential Receptor</i>	<i>Context of View</i>	<i>Approximate View Distance</i>	<i>Period of View</i>	<i>Visibility Rating</i>
<i>Surrounding Farmland</i>	<i>Static</i>	<i>Level</i>	<i>0 – 11 km</i>	<i>Long Term</i>	<i>Medium</i>
<i>Emahlaheni</i>	<i>Static</i>	<i>Level Above</i>	<i>> 10 km</i>	<i>Long Term</i>	<i>Medium</i>
<i>Gravel Roads</i>	<i>Dynamic</i>	<i>Above & below</i>	<i>1 – 11 km</i>	<i>Medium</i>	<i>Low</i>
<i>Tar Roads</i>	<i>Dynamic</i>	<i>Level Above</i>	<i>5 – 11 km</i>	<i>Short</i>	<i>Low</i>

It should however be noted that there are a number of existing power lines in the study area as shown in the Figures above. Viewers in the viewshed have become accustomed to these power lines in the landscape and an additional 10 km of power line will not increase the impact significantly.

3.0 ALTERNATIVE SENSITIVITY ANALYSIS

This section provides a short sensitivity matrix, which compares the three different alternatives and their associated environmental sensitivities.

Table 13: Alternative Sensitivity Matrix

Sensitivity	Alternative 1	Alternative 2	Alternative 3
Geology	None	None	None
Climate	None	None	None
Topography	None	None	None
Land Use	Traverses short section of ash dump, surrounding land used as grazing for cattle	Traverses Witbank Dam and farmland	Traverses Witbank Dam and agricultural land
Surface Water	Traverses only a short section of the un-named tributaries on site	Traverses a large section of the Witbank Dam	Traverses the largest Section of the Witbank Dam
Soils & Land Capability	Mainly agricultural and non sensitive soils	Along sensitive wetland and clay soils	Along sensitive wetland and clay soils
Flora	None	Sensitive vegetation units and plants present	Sensitive vegetation units and plants present
Fauna	None	None	None
Wetlands	None	Traverses wetland	Traverses wetland
Visual	Low Visibility	Moderate visibility	Highly visibility
Total Sensitivities	1	4	4

On the basis of the matrix presented above, it is suggested that the Bravo 5 Alternative 1 be utilised as the preferred alternative for the proposed project, as it has the least sensitive features associated with the alignment.

4.0 ENVIRONMENTAL IMPACT ASSESSMENT METHODOLOGY

The impacts will be ranked according to the methodology described below. Where possible, mitigation measures will be provided to manage impacts. In order to ensure uniformity, a standard impact assessment methodology was utilised so that a wide range of impacts can be compared with each other. The impact assessment methodology makes provision for the assessment of impacts against the following criteria:

- Ü significance;
- Ü spatial scale;
- Ü temporal scale;
- Ü probability; and
- Ü degree of certainty.

A combined quantitative and qualitative methodology was used to describe impacts for each of the aforementioned assessment criteria. A summary of each of the qualitative descriptors along with the equivalent quantitative rating scale for each of the aforementioned criteria is given in Table 14.

Table 14: Quantitative rating and equivalent descriptors for the impact assessment criteria

Rating	Significance	Extent Scale	Temporal Scale
1	VERY LOW	<i>Isolated sites / proposed site</i>	<u>Incidental</u>
2	LOW	<i>Study area</i>	<u>Short-term</u>
3	MODERATE	<i>Local</i>	<u>Medium-term</u>
4	HIGH	<i>Regional / Provincial</i>	<u>Long-term</u>
5	VERY HIGH	<i>Global / National</i>	<u>Permanent</u>

A more detailed description of each of the assessment criteria is given in the following sections.

4.1 Significance Assessment

Significance rating (importance) of the associated impacts embraces the notion of extent and magnitude, but does not always clearly define these since their importance in the rating scale is very relative. For example, the magnitude (i.e. the size) of area affected by atmospheric pollution may be extremely large (1000km²) but the significance of this effect is dependent on the concentration or level of pollution. If the concentration is great, the significance of the impact would be HIGH or VERY HIGH, but if it is diluted it would be VERY LOW or LOW. Similarly, if 60 ha of a grassland type are destroyed the impact would be VERY HIGH if only 100 ha of that grassland type were known. The impact would be VERY LOW if the grassland type was common. A more detailed description of the impact significance rating scale is given in Table 15 below.

Table 15 : Description of the significance rating scale

Rating		Description
5	Very high	Of the highest order possible within the bounds of impacts which could occur. In the case of adverse impacts: there is no possible mitigation and/or remedial activity which could offset the impact. In the case of beneficial impacts, there is no real alternative to achieving this benefit.
4	High	Impact is of substantial order within the bounds of impacts, which could occur. In the case of adverse impacts: mitigation and/or remedial activity is feasible but difficult, expensive, time-consuming or some combination of these. In the case of beneficial impacts, other means of achieving this benefit are feasible but they are more difficult, expensive, time-consuming or some combination of these.
3	Moderate	Impact is real but not substantial in relation to other impacts, which might take effect within the bounds of those which could occur. In the case of adverse impacts: mitigation and/or remedial activity are both feasible and fairly easily possible. In the case of beneficial impacts: other means of achieving this benefit are about equal in time, cost, effort, etc.
2	Low	Impact is of a low order and therefore likely to have little real effect. In the case of adverse impacts: mitigation and/or remedial activity is either easily achieved or little will be required, or both. In the case of beneficial impacts, alternative means for achieving this benefit are likely to be easier, cheaper, more effective, less time consuming, or some combination of these.
1	Very low	Impact is negligible within the bounds of impacts which could occur. In the case of adverse impacts, almost no mitigation and/or remedial activity are needed, and any minor steps which might be needed are easy, cheap, and simple. In the case of beneficial impacts, alternative means are almost all likely to be better, in one or a number of ways, than this means of achieving the benefit. Three additional categories must also be used where relevant. They are in addition to the category represented on the scale, and if used, will replace the scale.
0	No impact	There is no impact at all - not even a very low impact on a party or system.

4.2 Spatial Scale

The spatial scale refers to the extent of the impact i.e. will the impact be felt at the local, regional, or global scale. The spatial assessment scale is described in more detail in Table 16.

Table 16 : Description of the significance rating scale

Rating	Description
5	Global/National
4	Regional/Provincial
3	Local
2	Study Area
1	Isolated Sites / proposed site

4.3 Duration Scale

In order to accurately describe the impact it is necessary to understand the duration and persistence of an impact in the environment. The temporal scale is rated according to criteria set out in Table 17.

Table 17: Description of the temporal rating scale

Rating		Description
1	Incidental	The impact will be limited to isolated incidences that are expected to occur very sporadically.
2	Short-term	The environmental impact identified will operate for the duration of the construction phase or a period of less than 5 years, whichever is the greater.
3	Medium term	The environmental impact identified will operate for the duration of life of plant.
4	Long term	The environmental impact identified will operate beyond the life of operation.
5	Permanent	The environmental impact will be permanent.

4.4 Degree of Probability

Probability or likelihood of an impact occurring will be described as shown in Table 18 below.

Table 18 : Description of the degree of probability of an impact occurring

Rating	Description
1	Practically impossible
2	Unlikely
3	Could happen
4	Very Likely
5	It's going to happen / has occurred

4.5 Degree of Certainty

As with all studies it is not possible to be 100% certain of all facts, and for this reason a standard “degree of certainty” scale is used as discussed in Table 19. The level of detail for specialist studies is determined according to the degree of certainty required for decision-making. The impacts are discussed in terms of affected parties or environmental components.

Table 19 : Description of the degree of certainty rating scale

Rating	Description
Definite	More than 90% sure of a particular fact.
Probable	Between 70 and 90% sure of a particular fact, or of the likelihood of that impact occurring.
Possible	Between 40 and 70% sure of a particular fact or of the likelihood of an impact occurring.
Unsure	Less than 40% sure of a particular fact or the likelihood of an impact occurring.
Can't know	The consultant believes an assessment is not possible even with additional research.
Don't know	The consultant cannot, or is unwilling, to make an assessment given available information.

4.6 Quantitative Description of Impacts

To allow for impacts to be described in a quantitative manner in addition to the qualitative description given above, a rating scale of between 1 and 5 was used for each of the assessment

criteria. Thus the total value of the impact is described as the function of significance, spatial and temporal scale as described below:

$\text{Impact Risk} = \left(\frac{\text{SIGNIFICANCE} + \text{Spatial} + \text{Temporal}}{3} \right) \times \frac{\text{Probability}}{5}$
--

An example of how this rating scale is applied is shown below:

Table 20 : Example of Rating Scale

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
	LOW	Local	Medium-term	Could Happen	
Impact to air	2	3	3	3	1.6

Note: The significance, spatial and temporal scales are added to give a total of 8, that is divided by 3 to give a criteria rating of 2,67. The probability (3) is divided by 5 to give a probability rating of 0,6. The criteria rating of 2,67 is then multiplied by the probability rating (0,6) to give the final rating of 1,6.

The impact risk is classified according to five classes as described in the table below.

Table 21 : Impact Risk Classes

Rating	Impact Class	Description
0.1 – 1.0	1	Very Low
1.1 – 2.0	2	Low
2.1 – 3.0	3	Moderate
3.1 – 4.0	4	High
4.1 – 5.0	5	Very High

Therefore with reference to the example used for air quality above, an impact rating of 1.6 will fall in the Impact Class 2, which will be considered to be a low impact.

4.7 Cumulative Impacts

It is a requirement that the impact assessments take cognisance of cumulative impacts. In fulfilment of this requirement the impact assessment will take cognisance of any existing impact sustained by the operations, any mitigation measures already in place, any additional impact to environment through continued and proposed future activities, and the residual impact after mitigation measures.

It is important to note that cumulative impacts at the national or provincial level will not be considered in this assessment, as the total quantification of external companies on resources is not possible at the project level due to the lack of information and research documenting the effects of existing activities. Such cumulative impacts that may occur across industry boundaries can also only be effectively addressed at Provincial and National Government levels.

Using the criteria as described above an example of how the cumulative impact assessment will be done is shown below:

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Initial / Existing Impact (I-IA)	2	2	2	<u>1</u>	0.4
Additional Impact (A-IA)	1	2	<u>0</u>	<u>1</u>	0.2
Cumulative Impact (C-IA)	3	4	<u>2</u>	<u>1</u>	0.6
Residual Impact after mitigation (R-IA)	2	1	<u>2</u>	<u>1</u>	0.3

As indicated in the example above the Additional Impact Assessment (A-IA) is the amount that the impact assessment for each criterion will increase. Thus if the initial impact will not increase, as shown for temporal scale in the example above the A-IA will be 0, however, where the impact will increase by two orders of magnitude from 2 to 4 as in the spatial scale the A-IA is 2. The Cumulative Impact Assessment (C-IA) is thus the sum of the Initial Impact Assessment (I-IA) and the A-IA for each of the assessment criteria.

In both cases the I-IA and A-IA are assessed without taking into account any form of mitigation measures. As such the C-IA is also a worst case scenario assessment where no mitigation measures have been implemented. Thus a Residual Impact Assessment (R-IA) is also made which takes into account the C-IA with mitigation measures. The latter is the most probable case scenario, and for the purpose of this report is considered to be the final state Impact Assessment.

4.8 Notation of Impacts

In order to make the report easier to read the following notation format is used to highlight the various components of the assessment:

Significance or magnitude- IN CAPITALS

Temporal Scale – in underline

Probability – *in italics and underlined*.

Degree of certainty - **in bold**

Spatial Extent Scale – *in italics*

5.0 ENVIRONMENTAL IMPACT ASSESSMENT

The Impact Assessment will highlight and describe the impact to the environment following the abovementioned methodology and will assess the following components:

- Ü Geology;
- Ü Climate;
- Ü Surface Water;
- Ü Topography;
- Ü Soils;
- Ü Land Capability
- Ü Land Use;
- Ü Flora;
- Ü Fauna; and
- Ü Visual Assessment.

The impact assessment was undertaken for the construction, operational and decommissioning phases of the project. The impact of each line/route alternative was also assessed separately, however, where the impact was not significantly different, only one impact assessment was undertaken. Also, at the time of writing this report, no technical data was available as to the type of tower to be used for the construction of the transmission lines. Therefore, it is assumed that the Self-supporting strain and suspension tower type would be used. Contained in this assumption is that the maximum distance between towers would be 300 m and that the tower would be erected on concrete footings with dimensions of 2 x 2 x 2 m (area = 4 m² and volume = 8 m³).

5.1 Construction Phase

During the construction phase, the 400 kV power line will be erected. A 400 kV Transmission line requires a servitude width of 55 m. Where there are physical constraints such as other power lines adjacent to the new servitude, a minimum of 35 m-separation distance from such lines is required. Without physical constraints, parallel lines will have at least 55 m-separation distance. The power line cables are strung between pylons / towers, which are steel structures erected on concrete footings fixed in the substrate (soil or rock) below the pylon.

The major impacts during construction are the construction activities associated with the erection of the power lines and include, amongst others, heavy vehicle movement, construction of an access road and any wastes generated.

5.1.1 Geology

Initial Impact

Impacts that could occur to geology are limited to the physical removal of geological strata, resulting in permanent damage to those strata. There are no present indications that any existing impacts to geology have occurred and therefore there is no initial impact rating.

Additional Impact

There is no additional impact resulting from the power line construction since there are no significant geological features on site. The impact would be limited to the construction of the pylon footings, and should be a maximum of three pylons and therefore 12 footings. The 12 footings will disturb a combined area of 96 m³ of geological strata. This VERY LOW impact **could** probably occur in *isolated sites* over the long term. This results in a final impact class of **Low** as rated in the table below.

Table 22: Geology Additional Impact Assessment

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Geology	VERY LOW	<i>Isolated sites</i>	<u>Long Term</u>	<u>Probably</u>	Low
	1	1	4	4	1.6

Cumulative Impact

Since there is no initial impact, the cumulative impact is the same as rated for the additional impact above.

Mitigation Measures

- Ü No blasting is undertaken on site without a suitable blast design, compiled in line with relevant SANS codes and approved by an appropriately qualified professional;

Residual Impact

Although mitigation measures will not reduce the significance of impact to geology they will ensure that the impacts are contained. Mitigation measures will ensure that the likelihood of secondary impacts occurring is significantly reduced. The residual impact to geology at the completion of the construction phase will be the same as for the additional impact assessment.

5.1.2 Topography

Initial Impact

There are no present indications that any existing impacts to topography have occurred and therefore there is no initial impact rating.

Additional Impact

The construction of the power lines should not impact on the topography and therefore there is no additional impact.

Cumulative Impact

Since there is no initial impact, the cumulative impact is the same as rated for the additional impact above.

Mitigation Measures

No mitigation measures are required as there is no impact to topography from the proposed development.

Residual Impact

There is no residual impact to topography from the proposed development.

5.1.3 Soils, Land Capability and Land Use

Initial Impact

The study site has predominantly been used for grazing of livestock and some agricultural uses. The section of soils that will be crossed by the power line alternatives are presently not impacted upon, but in the near future the construction of the new power line will impact the soils. Other existing impacts are the existing pylon footings and cultivation of soils.

The initial impact to soils and land capability is **definitely** a HIGH negative impact acting over the long term, and is presently occurring in the *study area*. As indicated in Table 23 below the impact rating class is a High Impact.

Table 23: Soil and Land Capability Initial Impact Assessment

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Soils	HIGH	<i>Study Site</i>	<u>Long Term</u>	<i>Is occurring</i>	High
	4	2	4	5	3.33

Additional Impact

The additional impact from the new power lines will mainly be as a result of the construction of the power line pylons and their footings. Alternatives 1, 2 and 3 are approximately 7.4 km, 10.5 and 9.5 in length respectively and each will have a double power line. Therefore if using the average pylon distance of 300 m it can be assumed that there would be 56 pylons constructed. At the time of writing this report, the proponent has not determined which of the various pylon designs will be utilised, and therefore the actual impact could vary. For this analysis it is assumed that pylons similar to the existing power lines will be utilised. This will result in 4 footings impacting on the soils per pylon.

In addition to the pylon footings the soils will also be disturbed by the establishment of a construction road as well as the movement of construction vehicles. The impact from each of the routes are summarised below.

Table 24: Soil Impact

Soil Type	Alternative 1 (km)	Alternative 2 (km)	Alternative 3 (km)
Katspruit	0.4 km	6.6 km	2.5 km
Mispah	2.75 km	0.25 km	1.5 km
Clovelly/Hutton	2.85 km	3.65 km	5.0 km

As indicated in Table 24 above, Alternatives 2 and 3 cross more sensitive soils than Alternative 1. That said, the impact rating class between the two alternatives differ and is therefore rated separately.

For Alternative 1 the additional impact to soils and land capability is **probably** a LOW negative impact acting over the long term, and will definitely occur at *isolated sites*. As indicated in Table 25 below the impact rating class is a Moderate Impact.

Table 25: Soil and Land Capability Additional Impact Assessment – Alternative 1

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
--------	--------------	---------------	----------------	-------------	--------

Impact to Soils	Low	<i>Isolated Site</i>	<u>Long Term</u>	<u>Will occur</u>	Moderate
	2	1	4	5	2.3

For Alternatives 2 and 3 the additional impact to soils and land capability is **probably** a MODERATE negative impact acting over the long term, and will definitely occur at *isolated sites*. As indicated in Table 26 below the impact rating class is a Moderate Impact.

Table 26: Soil and Land Capability Additional Impact Assessment – Alternatives 2 and 3

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Soils	Moderate	<i>Isolated Site</i>	<u>Long Term</u>	<u>Will occur</u>	Moderate
	3	1	4	5	2.67

Cumulative Impact

The cumulative impact remains as rated for the initial impact i.e. a High impact class.

Mitigation Measures

- Ü Avoid placement of pylon footings in the clay soils on site;
- Ü Spread absorbent sand on areas where oil spills are likely to occur, such as the refuelling area in the hard park;
- Ü Oil-contaminated soils are to be removed to a contained storage area and bio-remediated or disposed of at a licensed facility;
- Ü If soils are excavated for the footing placement, ensure that the soil is utilised elsewhere for rehabilitation/road building purposes; and
- Ü Ensure that soil is stockpiled in such a way as to prevent erosion from storm water.

Residual Impact

The residual impact remains a High Impact, as the mitigation measures will not reduce the overall impact.

5.1.4 Surface Water

Initial Impact

The surface water features on site constitute sensitive surface water features. The Witbank Dam and Olifants River constitute sensitive surface water features on site. The impact is assessed in Table 27 below.

Table 27: Surface Water Initial Impact Rating

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Surface water	VERY LOW	<i>Study Site</i>	<u>Medium Term</u>	<u>Could happen</u>	Low
	1	2	3	3	1.2

The initial impact to surface water is VERY LOW, occurs in *Isolated sites / proposed site* and will be Medium Term and It's going to happen / has occurred. This results in a rating of 1.2 or a Low impact class.

Additional Impact

During the construction phase there should be limited impacts to surface water features as all the wetlands and riparian zones have been declared no-go zones that should be avoided. It is anticipated that the placement of the pylons will be done in such a way as to avoid the surface water features on site. Note that the wetlands are assessed separately below.

Waste generated during the construction phase may enter the environment through surface water runoff i.e. litter or pollution such as hydrocarbons can be washed into aquatic systems affecting those systems negatively. Storm-water flowing over the site will also mobilise loose sediments, which may enter the surface water environment affecting water quality. Storm-water containing sediment can be discharged to grassland buffers to ensure sediments fall out prior to water entering surface water bodies. Care must be taken that storm-water containing hydrocarbons and other pollution sources are not discharged.

Impacts will be felt as wide as the *study area* when storm-water flows from the power line sites into the study area. The impact to the surface water will **probably** be of a VERY LOW negative significance, and will act in the short-term. This impact could happen. This results in a Very Low impact class as assessed in Table 28.

Table 28: Surface Water Additional Impact Rating

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Surface water	VERY LOW	<i>Study area</i>	<u>Short Term</u>	<u>Could happen</u>	Very Low
	1	2	2	3	1.0

Cumulative Impact

The cumulative impact of the current activities and the future activities will not increase the impact rating from a Low Impact as rated in the initial impact assessment.

Mitigation Measures

- Ü Demarcated areas where waste can be safely contained and stored on a temporary basis during the construction phase should be provided at the hard park;
- Ü When adequate volumes (not more than 1 month) have accumulated all waste is to be removed from site and disposed of at a licensed facility;
- Ü Waste is not to be buried on site;
- Ü Hydro-carbons should be stored in a bunded storage area;

- Ü All hazardous materials *inter alia* paints, turpentine and thinners must be stored appropriately to prevent these contaminants from entering the environment;
- Ü Spill-sorb or similar type product must be used to absorb hydrocarbon spills in the event that such spills should occur;
- Ü Care must be taken to ensure that in removing vegetation adequate erosion control measures are implemented;
- Ü No construction vehicles or activities will be allowed to work within 50 m of any of the streams or wetlands on site.
- Ü If possible utilise Alternative 1 as the preferred alternative.

Residual Impact

The mitigation measures proposed will reduce the risk of the additional impact occurring, but it will not reduce the residual impact class, which remains at a Low impact as rated in the initial impact assessment.

5.1.5 Flora

Initial Impact

The initial impacts to flora include extensive grazing, cultivation and alien invasive colonisation. The initial impact to flora is **definitely** a MODERATE negative impact acting over the long term, and is presently occurring in the *study area*. As indicated in Table 29 below the impact rating class is a Moderate Impact.

Table 29: Flora Initial Impact Assessment

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Flora	MODERATE	<i>Study Site</i>	<u>Long Term</u>	<u>Is occurring</u>	Moderate
	3	2	4	5	3.00

Additional Impact

The additional impact to flora during the construction phase will be as a result of vegetation clearance for access roads and the removal of vegetation in the areas of the pylon footings. Table 30 below illustrates the length that each route alternative will cross the vegetation types identified.

Table 30: Flora Impact

Soil Type	Alternative 1	Alternative 2	Alternative 3
Undisturbed/Natural Grassland	1.37 km	3.52 km	3.6 km
Disturbed/Grazed Grassland	4.2 km	0.2 km	2.76 km
Wetland and Riparian Zones*	0.45 km	6.45 km	1.7 km

* Indicates sensitive vegetation types

The additional impact from the Alternative 1 alignment to flora is **probably** a VERY LOW negative impact acting over the short term, and will occur in *isolated sites*. As indicated in Table 31 below the impact rating class is a Low Impact.

Table 31: Flora Additional Impact Assessment – Alternative 1

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Flora	Very Low	<i>Isolated Site</i>	<u>Short Term</u>	<u>Will occur</u>	Low
	1	1	2	5	1.33

Due to the alignment of Alternatives 2 and 3 in line with the sensitive vegetation types, the impact is higher and will be active for a longer period. As there is sensitive species along this alignment the additional impact from the Alternatives 2 and 3 to flora is **probably** a HIGH negative impact acting over the long term, and will occur in *isolated sites*. As indicated in Table 32 below the impact rating class is a Moderate Impact.

Table 32: Flora Additional Impact Assessment – Alternative 2

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Flora	High	<i>Isolated Site</i>	<u>Long Term</u>	<u>Will occur</u>	Moderate
	4	1	4	5	3

Cumulative Impact

The cumulative impact to flora will remain as assessed for the initial impact assessment with a Moderate impact class.

Mitigation Measures

- Ü All construction areas should be demarcated prior to construction to ensure that the footprint of the impacts are limited (including areas where vehicles may traverse);
- Ü The sensitive vegetation unit should be avoided and construction limited to 50 m from the edge of the wetlands and streams;
- Ü Alternative 1 should be considered as the preferred alternative;
- Ü All alien invasive species on site should be removed and follow up monitoring and removal programmes should be initiated once construction is complete;
- Ü Adhere to the ESKOM vegetation management guideline (Appendix 4).

Residual Impact

If the mitigation measures are implemented and Alternative 1 is constructed then the residual impact to flora is **probably** a MODERATE negative impact acting over the medium term, and will occur in the *study area*. As indicated in Table 33 below the impact rating class is a Moderate Impact.

Table 33: Flora Residual Impact Assessment

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Flora	MODERATE	<i>Study Site</i>	<u>Medium Term</u>	<u>Will happen</u>	Moderate
	3	2	3	5	2.33

5.1.6 Fauna

Initial Impact

As described in the habitat assessment in Section 3.9, the site is relatively disturbed with the disturbed/grazed grassland, the undisturbed/natural grassland and the wetland and riparian zones the main habitat still available for fauna. The site is 34.7 % disturbed and while this is not ideal habitat for fauna, it will still provide habitat for various fauna. The suitable areas did show high species diversity, indicating that the impact is limited to isolated sites throughout the study area.

The study area is criss crossed with existing high voltage power lines that could potentially impact on the faunal life, especially large avi-faunal species. While there appears to be no negative impacts associated with electro magnetic fields generated by the power lines, Eskom's document, *Transmission Bird Collision Prevention Guideline* (Ref. no.: TGL41-335)⁵, the major impact to birds or avi-fauna is in the form of collisions with power lines. According to the document, it was found

that the majority of birds affected are large flighted birds, which are also often endangered or threatened species.

These large flighted birds are also long lived, with low breeding rate and often mate for life. Therefore, a single mortality due to a collision with a power line should be viewed as a high impact. In addition some of the most sensitive species to power line collisions such as Blue Crane are found in the study site in addition to other sensitive species such as White-Bellied Korhaan and Secretary Birds.

The current impact on fauna on site is **probably** of a HIGH negative significance, affecting the *region*, and acting in the long-term. The impact can likely occur. The impact class is classified as a High impact.

Table 34: Fauna Initial Impact Assessment

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Fauna	HIGH	<i>Region</i>	<u>Long Term</u>	<u>Likely</u>	High
	4	4	4	4	3.2

Additional Impact

The impact to fauna during the construction phase of the power lines will mostly be in the form of disturbance from the construction workers and vehicle noise. Due to the fact that the area is habitat to sensitive species, the impact could be quite high. Once again Alternatives 2 and 3 are significantly closer to the habitat for the sensitive species and therefore the impacts are assessed separately.

The additional impact from the Alternative 1 alignment to fauna is **probably** a MODERATE negative impact acting over the short term, and will occur in *isolated sites*. As indicated in Table 35 below the impact rating class is a Low Impact.

Table 35: Fauna Additional Impact Assessment – Alternative 1

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Fauna	MODERATE	<i>Isolated Site</i>	<u>Short Term</u>	<u>Will occur</u>	Low
	3	1	2	5	2

The additional impact from the Alternative 2 and 3 alignments to fauna is **probably** a HIGH negative impact acting over the short term, and will occur in *isolated sites*. As indicated in Table 35 below the impact rating class is a Moderate Impact.

Table 36: Fauna Additional Impact Assessment – Alternative 1

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Fauna	High	<i>Isolated Site</i>	<u>Short Term</u>	<u>Will occur</u>	Moderate
	4	1	2	5	2.3

Cumulative Impact

The cumulative impact to fauna should remain as assessed for the initial impact assessment as the impacts are identical. Therefore the impact remains a High impact to Fauna.

Mitigation Measures

- Ü All construction areas should be demarcated prior to construction to ensure that the footprint of the impacts are limited (including areas where vehicles may traverse);
- Ü The sensitive habitat should be avoided and construction limited to 50 m from the edge of the wetlands and streams;
- Ü Alternative 1 should be considered as the preferred alternative;
- Ü All alien invasive species on site should be removed and follow up monitoring and removal programmes should be initiated once construction is complete;
- Ü Adhere to the ESKOM vegetation management guideline (Appendix 4); and
- Ü Install power lines according to the ESKOM bird collision prevention guideline.

Residual Impact

The mitigation measures proposed above will ensure that the construction of the proposed power line remains a Moderate impact but the Residual Impact remains High. If the mitigation measures were to be extended into the existing power lines and bird flappers be installed, the residual impact could be mitigated to a Moderate Impact Class.

5.1.7 Wetlands

The impact assessment for wetlands is the same as assessed for the surface water component in Section 6.1.4.

5.1.8 Visual Impact

Initial Impact

At present the viewers in the viewshed are seeing the Duvha Power Station and the various mining activities including the Corobrik works and the various coal collieries in the area. In addition to the Power Station there are numerous power lines already traversing the landscape. The initial impact to the visual environment is HIGH negative acting in the long term, and has already occurred. The impact has **definitely** impacted on the *local region*.

Table 37: Visual Impact Assessment – Initial Impact

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Visual	High	<i>Local</i>	<u>Long Term</u>	<u>Has occurred</u>	High
	4	3	4	5	3.6

As illustrated in Table 37 above the initial impact to the visual environment is rated as a High impact.

Additional Impact

The additional impact from the power lines as described in Section 4.4 indicated that the additional impact to the visual environment is **probably** a LOW negative impact acting in the short term and impacting on the *local region*. This impact will definitely occur.

Table 38: Visual Impact Assessment – Additional Impact

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Visual	Low	<i>Local</i>	<u>Short Term</u>	<u>Will occur</u>	Moderate
	2	3	2	5	2.3

From Table 38 above it is clear that the additional impact from the construction of the power lines will be a Moderate impact.

Cumulative Impact

There are a high number of existing visual impacts on site as well as a high number of power lines. The cumulative impact from the developments will remain as assessed for the initial impact above; therefore the impact remains a High negative impact.

Mitigation Measures

- Ü Only the footprint of the proposed power line should be exposed. In all other areas, the natural vegetation should be retained;
- Ü Dust suppression techniques should be in place at all times during the construction phase;
- Ü Access roads should be minimised to prevent unnecessary dust.

Residual Impact

The mitigation measures proposed above will ensure that the construction of the proposed power line remains a High impact to the visual environment.

5.2 Operational Phase

The main impacts during the operational phase are the electro magnetic field associated with the power lines and the occurrence of the physical structures in the landscape. See *Electric and Magnetic Fields – A summary of Technical and Biological Aspects* (2006)¹ for a detailed discussion regarding the impact of electro magnetic fields (Appendix 5).

5.2.1 Geology

The impact assessment does not change from that of the construction phase, refer to Section 6.1.1 above.

5.2.2 Topography

The impact assessment does not change from that of the construction phase, refer to Section 6.1.2 above.

5.2.3 Soils, Land Capability and Land Use

The impact assessment does not change from that of the construction phase, refer to Section 6.1.3 above.

¹ *Electric and Magnetic Fields – A summary of Technical and Biological Aspects*, Empetus cc, 2006.

5.2.4 Surface water

The impact assessment does not change from that of the construction phase, refer to section 6.1.4 above.

5.2.5 Vegetation

The impact assessment does not change from that of the construction phase, refer to section 6.1.5 above.

5.2.6 Fauna

Initial impact

The initial impact remains as assessed in Section 6.1.6, a High impact.

Additional impact

During the operational phase the proposed development will add approximately 10 km of high voltage power lines to the existing network of power lines in the area. Sensitive blue cranes occur in the area and a single death of one of these protected species would be seen as a high impact.. The additional impact to fauna will **probably** be a HIGH negative impact, acting in the long term, and affected the *local area* and this impact could occur. This calculates to a Moderate impact class as illustrated in Table 39 below.

Table 39: Fauna Additional Impact Rating – Operations

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Fauna	HIGH	<i>Local</i>	<u>Long Term</u>	<u>Could occur</u>	Moderate
	4	3	4	3	2.2

Cumulative impact

During the operational phase the proposed development will add approximately 10 km of high voltage power lines to the existing network of power lines in the area. The addition is relatively small in consideration of the approximately 180 km of existing high voltage powerlines in the area. The cumulative impact to fauna remains a High impact as assessed in the initial impact assessment.

Mitigation Measures

- Ü The sensitive habitat should be avoided and power lines limited to 50 m from the edge of the wetlands and streams;
- Ü Alternative 1 should be considered as the preferred alternative;
- Ü Adhere to the ESKOM vegetation management guideline (Appendix 4); and
- Ü Install power lines according to the ESKOM bird collision prevention guideline.

Residual impact

In order to prevent power line collisions from birds, anti-collision devices can be installed to the power lines. These include static, dynamic, reflective and illuminated devices. As mentioned in Appendix 3 these devices have resulted in a 60% reduction in bird collisions but they will not completely eliminate the impact risk to birds. In addition this reduction will only be effective if the anti-collision devices are installed on all the power lines in the region. If the anti collision devices are only installed for the proposed 10 km of new power line, the impact would remain a High impact. If the devices are to be installed on all the regional power lines the impact to fauna would **probably** be a HIGH negative impact, acting on the *regional scale* in the long term. The probability would however be reduced to unlikely.

Table 40: Fauna Residual Impact Rating

Impact	Significance	Spatial Scale	Temporal Scale	Probability	Rating
Impact to Fauna	HIGH	<i>Regional / Provincial</i>	<u>Long Term</u>	<u>Unlikely</u>	Low
	4	4	4	2	1.6

The residual impact to fauna as calculated in Table 40 above has a rating of 1.6 and a Low impact class.

5.2.7 Visual

The impact assessment does not change from that of the construction phase, refer to Section 6.1.7 above.

5.3 Decommissioning Phase

5.3.1 Geology

The impacts to geology during the decommissioning phase of the development remain as assessed in the construction phase in Section 6.1.1 above.

5.3.2 Topography

The impacts to topography during the decommissioning phase of the development remain as assessed in the construction phase in Section 6.2.2 above.

5.3.3 Soils, Land Capability and Land Use

The impacts to soils during the decommissioning phase of the development remain as assessed in the construction phase in Section 6.2.3 above.

5.3.4 Surface water

The impacts to surface water during the decommissioning phase of the development remain as assessed in the construction phase in Section 6.2.4 above.

5.3.5 Vegetation

The impacts to vegetation during the decommissioning phase of the development remain as assessed in the construction phase in Section 6.2.5 above.

5.3.6 Fauna

Even though the removal of the 10 km of proposed power lines will reduce the number of power lines in the area that could impact on fauna, the impact after decommissioning will remain as assessed in Section 6.2.6 above due to the remaining network of high voltage power lines.

5.3.7 Visual

Even though the removal of the 10 km of proposed power lines will reduce the number of power lines in the area that could impact on the visual environment, the impact after decommissioning will remain as assessed in Section 6.2.7 above due to the remaining network of high voltage power lines.

6.0 ENVIRONMENTAL MANAGEMENT

This section describes the suggested commitments that should be included in the Environmental Management Plan (EMP) to be compiled by the environmental consultant responsible for the EIA.

6.1 Geology and Soils

<i>Management Component</i>	<i>Geology and Soils</i>
Primary Objective	
<i>To ensure that the soils are stockpiled in the correct manner to prevent erosion and contamination of surface water runoff.</i>	
Core Criteria:	Monitoring Criteria
<i>No blasting is undertaken on site without a suitable blast design, compiled in line with relevant SANS codes and approved by an appropriately qualified professional.</i>	<i>Site Development Plan, EMP monitoring and Intermittent observation</i>
<i>Avoid placement of pylon footings in the clay soils on site</i>	
<i>Spread absorbent sand on areas where oil spills are likely to occur, such as the refuelling area in the hard park</i>	
<i>Oil-contaminated soils are to be removed to a contained storage area and bio-remediated or disposed of at a licensed facility</i>	
<i>If soils are excavated for the footing placement, ensure that the soil is utilised elsewhere for rehabilitation/road building purposes</i>	
<i>Ensure that soil is stockpiled in such a way as to prevent erosion from storm water.</i>	

6.2 Fauna

Management Component	Fauna – especially red data birds		
Primary Objective			
<i>To ensure that the development minimises the potential impact to endangered species and their habitat.</i>			
Core Criteria:		Monitoring Criteria	
<i>All construction areas should be demarcated prior to construction to ensure that the footprint of the impacts are limited (including areas where vehicles may traverse)</i>		Site Development Plan, EMP monitoring and Intermittent observation	
<i>No construction activity and disturbance will be permitted in the seasonal seepage zone where the red data birds were observed.</i>			
<i>Bird flappers are to be installed on all power lines in order to prevent bird collisions.</i>			
<i>Construction activities, people and vehicles will not be allowed outside of the area demarcated for construction.</i>			
<i>No hunting, snaring or collection of eggs will be allowed.</i>			
<i>If any Blue Crane nests or young are found, contact the Mpumalanga Parks Board for assistance. Also avoid the area at all cost (250m buffer)</i>			
<i>If adult birds are observed on site, avoid startling the birds, as they could fly into the already existing power lines.</i>			
<i>No animals/pets will be allowed in the construction site.</i>			
<i>Adhere to the ESKOM bird collision prevention guideline (Appendix 3)</i>			
<i>Poisoning of any sort is strictly forbidden.</i>			
<i>Remove all food wastes daily and discard at a licensed waste facility</i>			
<i>Provide vermin-proof bins for construction workers</i>			
<i>Designate eating areas and prevent food and waste build up</i>			
<i>No cooking fires will be permitted, the grassland is highly susceptible to veld fires and these destroy bird eggs</i>			

6.3 Vegetation

Management Component	Vegetation
Primary Objective	
<i>To ensure the control of alien invasive species and that the rehabilitation of indigenous vegetation to as close to the original state as possible.</i>	
Core Criteria:	Monitoring Criteria
<i>All construction areas should be demarcated prior to construction to ensure that the footprint of the impacts are limited (including areas where vehicles may traverse)</i>	Site Development Plan, EMP monitoring and Intermittent observation
<i>Take appropriate remedial action where vegetation establishment has not been successful or erosion is evident.</i>	
<i>Control of alien invasive species in line with the requirements of Conservation of Agricultural Resources Act will be undertaken.</i>	
<i>Alien invasive plant material will be preferentially removed in entirety through mechanical means (e.g. chainsaw, bulldozer, hand-pulling of smaller specimens). Chemical control is only required as a last resort.</i>	
<i>If during the establishment period, any noxious or excessive weed growth occurs, such vegetation will be removed.</i>	
<i>No construction activity and disturbance will be permitted in the seasonal seepage zone.</i>	
<i>It is the developer's responsibility to implement a monitoring programme that will be instituted to ensure that re-growth of alien invasive plants species does not occur, or that such re-growth is controlled.</i>	
<i>The sensitive vegetation unit should be avoided and construction limited to 50 m from the edge of the wetlands and streams</i>	
<i>Adhere to the ESKOM vegetation management guideline (Appendix 4)</i>	

6.4 Rivers, wetlands and Streams

Management Component	Rivers and streams
Primary Objective	
<i>To ensure that the rivers and streams are protected and incur minimal negative impact from the development as possible.</i>	
Core Criteria:	Monitoring Criteria
<i>The Contractor will minimise the extent of any damage to the flood plain that is necessary to complete the works, and will not pollute any river as a result of construction activities.</i>	Storm water Management Plan, Site Development Plan, EMP monitoring and Intermittent observations
<i>The Contractor will not cause any physical damage to any aspects of a watercourse, other than that necessary to complete the works as specified and in accordance with the accepted method statement.</i>	
<i>No construction vehicles or activities will be allowed to work within 50 m of any of the streams or wetlands on site</i>	
<i>Demarcated areas where waste can be safely contained and stored on a temporary basis during the construction phase should be provided at the hard park</i>	
<i>When adequate volumes (not more than 1 month) have accumulated all waste is to be removed from site and disposed of at a licensed facility</i>	
<i>Waste is not to be buried on site</i>	
<i>All hazardous materials inter alia paints, turpentine and thinners must be stored appropriately to prevent these contaminants from entering the environment</i>	
<i>Spill-sorb or similar type product must be used to absorb hydrocarbon spills in the event that such spills should occur</i>	
<i>Care must be taken to ensure that in removing vegetation adequate erosion control measures are implemented</i>	

7.0 CONCLUSION

In conclusion the proponent is proposing the construction and operation a 400 kV overhead power line, by-passing the existing Duvha Power Station, to form a new Bravo-Vulcan line near Emahlaheni, Mpumalanga.

Cymbian was appointed to investigate the biophysical aspects of the proposed site as well as the potential visual impact of the development. The aspects investigated include topography, soils, land use, land capability, wetland, fauna, flora and the visual environment.

It was found that the major areas of concern were the sensitive wetlands and riparian zones on site, along with the sensitive avifaunal and floral species that occur in these environments. In addition it was noted that the visual impact of the development could be high.

Upon review of the existing procedures and mitigation measures that Eskom have applied in the past and which are based on sound scientific research it was found that the impacts to fauna could be reduced.

The impacts to the wetland and riparian zones could be reduced by utilising the Alternative 1 route alignment, thereby avoiding most of the wetlands and seepage zones.

The visual impact was found to be relatively small, when considering the high number of existing power lines in the area as well as the imminent Duvha Power Station.

In conclusion the proposed development will impact on the environment, but these impacts can be managed and mitigated to the point where they are within acceptable norms. It is suggested that the Alternative 1 route alignment be utilised in order to decrease the risk of impacting in fauna and flora.

Appendix 1: Floral Species List

Family	Species	Common name	Present	Occurrence
AMARYLLIDACEAE	<i>Cyrthanthus breviflorus</i>	Yellow Fire lily	x	Individuals
ANACARDIACEAE	<i>Rhus magalismsontana</i> Sond. subsp. <i>magalismsontana</i>	Bergtaabos		
ANACARDIACEAE	<i>Sclerocarya birrea</i> (A.Rich.) Hochst. subsp. <i>caffra</i> (Sond.) Kokwaro	Marula		
APIACEAE	<i>Afroscidium</i> <i>magalismsontanum</i> (Sond.) P.J.D.Winter	Wild Parsley		
APIACEAE	<i>Heteromorpha arborescens</i> (Spreng.) Cham. & Schtdl. var. <i>abyssinica</i> (Hochst. ex A.Rich.) H.Wolff	Parsley Tree		
APOCYNACEAE	<i>Asclepias gibba</i> (E.Mey.) Schltr. var. <i>gibba</i>			
APOCYNACEAE	<i>Asclepias stellifera</i> Schltr.	Spring Stars		
APOCYNACEAE	<i>Brachystelma rubellum</i> (E.Mey.) Peckover			
APOCYNACEAE	<i>Pachycarpus schinzianus</i> (Schltr.) N.E.Br.	Bitterwortel		
APOCYNACEAE	<i>Parapodium costatum</i> E.Mey.			
APONOGETONACEAE	<i>Aponogeton natalensis</i> Oliv.	Wateruintjie	x	Individuals
AQUIFOLIACEAE	<i>Ilex mitis</i> (L.) Radlk. var. <i>mitis</i>	Cape Holly		
ASPHODELACEAE	<i>Kniphofia ensifolia</i> Baker subsp. <i>ensifolia</i>			
ASTERACEAE	<i>Bidens pilosa</i>	Blackjack	x	Common
ASTERACEAE	<i>Bidens formosa</i>	Cosmos	x	Individuals
ASTERACEAE	<i>Crassocephalum x</i> <i>picridifolium</i> (DC.) S.Moore			
ASTERACEAE	<i>Dicoma macrocephala</i> DC.			
ASTERACEAE	<i>Helichrysum aureonitens</i> Sch.Bip.		x	Individuals
ASTERACEAE	<i>Helichrysum nudifolium</i> (L.) Less. var. <i>nudifolium</i>	Hottentot's Tea		
ASTERACEAE	<i>Helichrysum setosum</i> Harv.	Yellow Everlasting		
ASTERACEAE	<i>Helichrysum splendidum</i> (Thunb.) Less.			
ASTERACEAE	<i>Nidorella hottentotica</i> DC.			
ASTERACEAE	<i>Stoebe vulgaris</i>	Bankrupt Bush	x	Sparse
ASTERACEAE	<i>Tagetes minuta</i>	Khaki weed	x	Common
ASTERACEAE	<i>Vernonia poskeana</i> Vatke & Hildebr. subsp. <i>botswanica</i> G.V.Pope			
BRYACEAE	<i>Bryum argenteum</i> Hedw.	Silver Moss		
CAPPARACEAE	<i>Maerua cafra</i> (DC.) Pax	Common bush-cherry, White- wood		
CARYOPHYLLACEAE	<i>Corrigiola litoralis</i> L. subsp. <i>litoralis</i> var. <i>perennans</i> Chaudhri			
CARYOPHYLLACEAE	<i>Dianthus mooiensis</i> F.N.Williams subsp. <i>mooiensis</i> var. <i>mooiensis</i>	Wild Pink		
CARYOPHYLLACEAE	<i>Dianthus transvaalensis</i> Burt Davy			

CONVOLVULACEAE	<i>Ipomoea crassipes</i> Hook. var. <i>crassipes</i>			
CONVOLVULACEAE	<i>Ipomoea magnusiana</i> Schinz			
CRASSULACEAE	<i>Crassula setulosa</i> Harv. var. <i>setulosa</i> forma <i>setulosa</i>			
CYPERACEAE	<i>Bulbostylis densa</i> (Wall.) Hand.-Mazz. subsp. <i>afromontana</i> (Lye) R.W.Haines			
CYPERACEAE	<i>Bulbostylis hispidula</i> (Vahl) R.W.Haines subsp. <i>pyriformis</i> (Lye) R.W.Haines			
CYPERACEAE	<i>Cyperus esculentus</i>	Yellow Nutsedge	x	Sparse
CYPERACEAE	<i>Lipocarpa nana</i> (A.Rich.) Chem.			
CYPERACEAE	<i>Pycnus pumilus</i> (L.) Domin			
DICRANACEAE	<i>Campylopus savannarum</i> (Müll.Hal.) Mitt.			
ERIOCAULACEAE	<i>Eriocaulon abyssinicum</i> Hochst.			
EUPHORBIACEAE	<i>Euphorbia inaequilatera</i> Sond. var. <i>inaequilatera</i>			
EXORMOTHECACEAE	<i>Exomotheca holstii</i> Steph.			
FABACEAE	<i>Eriosema psoraleoides</i> (Lam.) G.Don			
FABACEAE	<i>Indigofera arrecta</i> Hochst. ex A.Rich.			
FABACEAE	<i>Indigofera zeyheri</i> Spreng. ex Eckl. & Zeyh.			
FABACEAE	<i>Lotononis foliosa</i> Bolus		x	Individuals
FABACEAE	<i>Rhynchosia monophylla</i> Schltr.		x	Individuals
FABACEAE	<i>Rhynchosia nervosa</i> Benth. ex Harv. var. <i>nervosa</i>			
FABACEAE	<i>Virgilia divaricata</i> Adamson			
FABACEAE	<i>Zornia milneana</i> Mohlenbr.		x	Individuals
FOSSOMBRONIACEAE	<i>Fossombronia gemmifera</i> Perold			
HALORAGACEAE	<i>Myriophyllum aquaticum</i> (Vell.) Verdc.			
HALORAGACEAE	<i>Myriophyllum spicatum</i> L.			
HYACINTHACEAE	<i>Albuca setosa</i> Jacq.	Slymuintjie	x	Individuals
HYPOXIDACEAE	<i>Hypoxis acuminata</i>		x	Individuals
HYPOXIDACEAE	<i>Hypoxis filiformis</i> Baker		x	Individuals
HYPOXIDACEAE	<i>Hypoxis iridifolia</i>		x	Individuals
IRIDACEAE	<i>Gladiolus crassifolius</i> Baker		x	Individuals
IRIDACEAE	<i>Lapeirousia sandersonii</i> Baker			
ISOETACEAE	<i>Isoetes transvaalensis</i> Jemy & Schelpe			
LAMIACEAE	<i>Becium angustifolium</i> (Benth.) N.E.Br.			
LAMIACEAE	<i>Becium obovatum</i>		x	Individuals
LAMIACEAE	<i>Mentha aquatica</i> L.			
LAMIACEAE	<i>Pycnostachys reticulata</i> (E.Mey.) Benth.			
LENTIBULARIACEAE	<i>Utricularia arenaria</i>		x	Individuals
LENTIBULARIACEAE	<i>Utricularia stellaris</i> L.f.			

LILIACEAE	<i>Protaspargus setaceus</i>	Asparagus Fern	x	Individuals
LOBELIACEAE	<i>Monopsis decipiens</i>		x	Individuals
MALPIGHIACEAE	<i>Triaspis hypericoides</i> (DC.) <i>Burch. subsp. nelsonii</i> (Oliv.) Immelman			
MALVACEAE	<i>Pavonia transvaalensis</i> (Ulbr.) A.Meeuse	Klapperbossie		
MALVACEAE	<i>Triumfetta obtusicornis</i> Sprague & Hutch.	Maagbossie		
MENYANTHACEAE	<i>Nymphoides thunbergiana</i> (Griseb.) Kuntze			
MESEMBRYANTHEMACEAE	<i>Delosperma leendertziae</i> N.E.Br.			
MESEMBRYANTHEMACEAE	<i>Frithia humilis</i> Burgoyne			
MESEMBRYANTHEMACEAE	<i>Mossia intervallaris</i> (L.Bolus) N.E.Br.			
MOLLUGINACEAE	<i>Limeum viscosum</i> (J.Gay) <i>Fenzl subsp. viscosum var.</i> <i>glomeratum</i> (Eckl. & Zeyh.) Friedrich			
MORACEAE	<i>Ficus abutilifolia</i> (Miq.) Miq.			
MORACEAE	<i>Ficus salicifolia</i> Vahl			
MYRTACEAE	<i>Eucalyptus</i> spp	Blue Gum	x	Sparse
NYMPHAEACEAE	<i>Nymphaea nouchali</i> <i>Burm.f. var. caerulea</i> (Savigny) Verdc.			
OCHNACEAE	<i>Ochna gamostigmata</i> Du Toit			
ONAGRACEAE	<i>Epilobium hirsutum</i> L.			
ORCHIDACEAE	<i>Centrostigma occultans</i> (Welw. ex Rchb.f.) Schltr.			
ORCHIDACEAE	<i>Habenaria clavata</i> (Lindl.) Rchb.f.			
ORCHIDACEAE	<i>Satyrium hallackii</i> Bolus <i>subsp. ocellatum</i> (Bolus) A.V.Hall			
OROBANCHACEAE	<i>Striga gesnerioides</i> (Willd.) Vatke			
OXALIDACEAE	<i>Oxalis obliquifolia</i>	Sorrel	x	Individuals
PALLAVICINIACEAE	<i>Symphyogyna brasiliensis</i> Nees & Mont.			
PARMELIACEAE	<i>Canoparmelia</i> <i>pustulescens</i> (Kurok.) Elix			
PEDALIACEAE	<i>Dicerocaryum senecioides</i> (Klotzsch) Abels			
PHYLLANTHACEAE	<i>Phyllanthus</i> <i>maderaspatensis</i> L.	Kleurbossie	x	Individuals
POACEAE	<i>Andropogon eucomus</i> Nees	Old Man's Beard	x	Sparse
POACEAE	<i>Calamagrostis epigejos</i> (L.) Roth var. <i>capensis</i> Stapf			
POACEAE	<i>Cymbopogon excavatus</i>	Broad-leaved Turpentine Grass	x	Individuals
POACEAE	<i>Cynodon dactylon</i>	Coch Grass	x	Sparse
POACEAE	<i>Echinochloa jubata</i> Stapf			
POACEAE	<i>Elionurus muticus</i>	Wire Grass	x	Individuals
POACEAE	<i>Eragrostis capensis</i> (Thunb.) Trin.	Heart-seed Love Grass	x	Individuals
POACEAE	<i>Eragrostis chloromelas</i> Steud.	Narrow Curly leaf	x	Common
POACEAE	<i>Eragrostis hiemiana</i> Rendle			

POACEAE	<i>Eragrostis inamoena</i> K.Schum.			
POACEAE	<i>Eragrostis plana</i>	Tough Love Grass	x	Sparse
POACEAE	<i>Eragrostis racemosa</i> (Thunb.) Steud.	Narrow Heart Love Grass	x	Sparse
POACEAE	<i>Eragrostis tef</i> (Zuccagni) Trotter	Tef	x	Sparse
POACEAE	<i>Hyparrhenia hirta</i> (L.) Stapf	Common Thatching Grass	x	Common
POACEAE	<i>Hyparrhenia quarrei</i> Robyns			
POACEAE	<i>Hyparrhenia tamba</i> (Steud.) Stapf	Blue Thatching Grass		
POACEAE	<i>Hyperthelia dissoluta</i> (Nees ex Steud.) Clayton			
POACEAE	<i>Loudetia simplex</i>	Russet Grass	x	Sparse
POACEAE	<i>Miscanthus junceus</i> (Stapf) Pilg.	Wireleaf Daba Grass		
POACEAE	<i>Perotis patens</i> Gand.	Cat's Tail	x	Individuals
POACEAE	<i>Schizachyrium sanguineum</i>	Red Autumn Grass	x	Sparse
POACEAE	<i>Setaria nigrirostris</i> (Nees) T.Durand & Schinz			
POACEAE	<i>Setaria sphacelata</i> var. <i>sphacelata</i>	Common Bristle Grass	x	Individuals
POACEAE	<i>Sporobolus fimbriatus</i>	Dropseed Grass	x	Sparse
POACEAE	<i>Themeda triandra</i>	Red Grass	x	Sparse
POACEAE	<i>Urochloa brachyura</i> (Hack.) Stapf		x	Sparse
POLYGALACEAE	<i>Polygala ohlendorfiana</i> Eckl. & Zeyh.			
POLYGALACEAE	<i>Polygala transvaalensis</i> Chodat subsp. <i>transvaalensis</i>			
PORTULACACEAE	<i>Anacampseros subnuda</i> Poelln. subsp. <i>subnuda</i>			
PORTULACACEAE	<i>Portulaca hereroensis</i> Schinz			
PORTULACACEAE	<i>Portulaca quadrifida</i> L.			
POTAMOGETONACEAE	<i>Potamogeton schweinfurthii</i> A.Benn.			
PTERIDACEAE	<i>Cheilanthes involuta</i> (Sw.) Schelpe & N.C.Anthony var. <i>obscura</i> (N.C.Anthony) N.C.Anthony			
RANUNCULACEAE	<i>Ranunculus meyeri</i> Harv.		x	Individuals
RICCIACEAE	<i>Riccia atropurpurea</i> Sim			
RICCIACEAE	<i>Riccia okahandjana</i> S.W.Arnell			
RICCIACEAE	<i>Riccia volkii</i> S.W.Arnell			
RUBIACEAE	<i>Richardia scabra</i> L.			
SCROPHULARIACEAE	<i>Chaenostoma leve</i> (Hiern) Kornhall			
SELAGINELLACEAE	<i>Hebenstretia angolensis</i> Rolfe	Katstert	x	Individuals
SELAGINELLACEAE	<i>Selaginella dregei</i> (C.Presl) Hieron.			
SOLANACEAE	<i>Solanum sisymbriifolium</i>	Wild tomato	x	Individuals
THELYPTERIDACEAE	<i>Thelypteris confluens</i> (Thunb.) C.V.Morton			
THYMELAEACEAE	<i>Gnidia sericocephala</i> (Meisn.) Gilg ex Engl.			
XYRIDACEAE	<i>Xyris capensis</i> Thunb.			

Appendix 2: Animal Species List

Faunal Species List	
Species	Common name
Reptiles	
<i>Bitens arietans</i>	Puff Adder
<i>Varanus niloticus</i>	Water Monitor
Birds	
<i>Phalacrocorax africanus</i>	Reed Cormorant
<i>Ardea cinerea</i>	Grey Heron
<i>Ardea melanocephala</i>	Blackheaded Heron
<i>Bubulcus ibis</i>	Cattle Egret
<i>Bostrychia hagedash</i>	Hadedda Ibis
<i>Plegadis falcinellus</i>	Glossy Ibis
<i>Alopochen aegyptiacus</i>	Egyptian Goose
<i>Elanus caeruleus</i>	Blackshouldered Kite
<i>Francolinus swainsonii</i>	Swainson's Francolin
<i>Numida meleagris</i>	Helmeted Guineafowl
<i>Fulica cristata</i>	Redknobbed Coot
<i>Gallinula chloropus</i>	Moorhen
<i>Anthropoides paradisea</i>	Blue Crane
<i>Sagittarius serpentarius</i>	Secretary Bird
<i>Eupodotis cafra</i>	Whitebellied Korhaan
<i>Vanellus armatus</i>	Blacksmith Plover
<i>Vanellus coronatus</i>	Crowned Plover
<i>Streptopelia semitorquata</i>	Redeyed Dove
<i>Streptopelia senegalensis</i>	Laughing Dove
<i>Asio capensis</i>	Marsh Owl
<i>Colius striatus</i>	Speckled Mousebird
<i>Mirafra africana</i>	Rufousnaped Lark
<i>Corvus albus</i>	Pied Crow
<i>Saxicola torquata</i>	Stone Chat
<i>Phylloscopus trochilus</i>	Willow Warbler
<i>Cisticola fulvicapilla</i>	Neddicky
<i>Motacilla clara</i>	Cape Wagtail
<i>Anthus cinnamomeus</i>	Grassveld Pipit
<i>Passer domesticus</i>	House Sparrow
<i>Ploceus velatus</i>	Masked Weaver
<i>Euplectes orix</i>	Red Bishop
<i>Emberiza capensis</i>	Cape Bunting
Mammals	
<i>Antidorcas marsupialis</i>	Springbok
<i>Damaliscus dorcas phillipsi</i>	Blesbok
<i>Cynictis pencilata</i>	Yellow Mongoose
<i>Orycteropus afer</i>	Aardvark / Antbear

**Appendix 3: Bird Collision Prevention Guidelines and Bird Impact Assessment
Study**

Appendix 4: Vegetation Management Guideline

**Appendix 5: Electric and Magnetic Fields – A summary of Technical and Biological
Aspects**