

279-3462

4/16/2014

1/11



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Registration
Number:

279-3462

Date of Issuance:

APR 16 2014

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance: **Conditional**

Name of Pesticide Product:
F7214 3.2 RTS Herbicide

Name and Address of Registrant (include ZIP Code):

FMC Corporation
1735 Market Street
Philadelphia, PA 19103

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA §3(c)(7)(A) provided that you submit and/or cite all data required for registration review of your product when the Agency requires all registrants of similar products to submit data. Data requirements can be found at www.regulations.gov. Quinclorac data requirements are outlined in GDCI-128974-919 under docket ID EPA-HQ-OPP-2007-1135. Sulfentrazone data requirements are outlined in GDCI-129081-883 under docket ID EPA-HQ-OPP-2009-0624.

The basic and alternate #2 confidential statement of formula (CSF) dated August 30, 2013 and alternate CSF#1 dated September 12, 2013 are acceptable.

A stamped copy of the label is enclosed for your records. Submit one (1) copy of the final printed label before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions regarding this Notice, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov.

Signature of Approving Official:

Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)

Date:

APR 16 2014

2/11

F7214 3.2 RTS HERBICIDE
Note to Reviewer: Text in brackets is optional

4/16/2014

F7214 3.2 RTS Herbicide

EPA Reg. No. 279-3462

EPA Est. 279-XXX

Kills Crabgrass, Dandelion, Ground Ivy, Yellow & Purple Nutsedge, Kyllinga and over XXX other tough weeds

Selective Postemergence Control of Annual Grasses, Broadleaf weeds, and Sedges in Residential Lawns

Active Ingredients:	
Sulfentrazone.....	0.81%
Quinclorac.....	2.42%
Other Ingredients.....	96.77%
	100.00%

NET CONTENTS XX FL OZ / XXX mL

KEEP OUT OF REACH OF CHILDREN
CAUTION

See other panels for additional precautionary information.



FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia PA 19103

ACCEPTED
04/16/2014
Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 279-3462

Questions, Comments or Medical Information

[inside]FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-(800)-331-3148 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes, socks, and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

Keep children and pets out of the treated area until sprays have dried.

ENVIRONMENTAL HAZARDS

This product is toxic to marine/estuarine invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

This product may impact surface water quality due to runoff of rain water or spray drift. Under certain conditions this product may have a high potential for runoff into surface water for several months after application. This is especially true for poorly draining soils and soils with shallow ground water.

The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of labeled use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of F7214 3.2 RTS Herbicide not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

PRODUCT INFORMATION

F7214 3.2 RTS Herbicide selectively controls tough weeds like crabgrass, chickweed, dandelion, clover, ground ivy, purple and yellow nutsedge (nutgrass), Kyllinga, wild onion and garlic, broadleaf plantain, purslane, wild violet, spurge, and other listed annual, biennial, and broadleaf weeds in a variety of lawns, including:

4/11

Cool Season Grasses
Tall & Fine Fescue
Kentucky Bluegrass
Rough Bluegrass
Ryegrass (perennial)

Warm Season Grasses
Common & Hybrid Bermudagrass
Zoysiagrass
Centipedegrass
Buffalograss
Seashore Paspalum

This unique formula causes visual symptoms to develop quickly within 3 to 7 days after application. F7214 3.2 RTS Herbicide involves uptake by both weed roots and shoots. Complete kill should occur in 2-3 weeks. F7214 3.2 RTS Herbicide also controls yellow nutsedge tubers (nutlets) in the soil over time.

A postemergence application of F7214 3.2 RTS Herbicide is improved when adequate soil moisture is present at application. Best weed control results will be obtained when no rainfall or irrigation occurs within 24 hours after application. If no rainfall or irrigation occurs within 7 days after application of F7214 3.2 RTS Herbicide in the amount of 0.5 inches, then irrigation of at least 0.5 inches is recommended.

APPLICATION INSTRUCTIONS AND PRECAUTIONS

Where to Use	Lawns: Well established turfgrasses. Cool Season Grasses: Fine & Tall Fescue, Perennial Ryegrass, Kentucky & Rough Bluegrass Warm Season Grasses: Common & Hybrid Bermudagrass, Buffalograss, Centipedegrass, Seashore Paspalum, Zoysiagrass. Do not use on St. Augustinegrass.
Kills	Sedges, Grasses and Broadleaf Weeds: Yellow Nutsedge, Kyllinga, Crabgrass, Foxtail, Ground Ivy, Wild Violet, Plantain, Dandelion, Clover, Dollarweed, Spurge, Woodsorrel, Knotweed, Chickweed, Curly Dock, Wild Onion, Wild Garlic, and other listed weeds.
When to Use	As weeds are small and actively growing.
How Much to Use	4 fl oz per 1 gallon of water. Spot spray weed directly until thoroughly covered. [For RTS sprayer: Mixes automatically as you spray.] Some weeds may require repeat application. (See detailed instructions.)
Coverage	1 finished gallon covers 500 sq ft

[PowerFlex] SPRAYER: USE DIRECTIONS

The [AMDRO Spray Delivery System Name] is a patented innovative home lawn, landscape, and perimeter tank spray system. Spot treat weeds with this easy to use system.

1. Unscrew the tank cap and fill the tank with water up to the one gallon indicator. [The tank holds one gallon of water.]
2. Remove the [sprayer][wand][applicator] from the tank holster and unfold the [sprayer][wand][applicator].
3. Select the [cartridge][bottle] [herbicide] labeled for intended use. READ THE ENTIRE PRODUCT LABEL PRIOR TO USE.
4. Remove the cap from the [cartridge][bottle][herbicide]. [Each [cartridge][bottle] contains 4 fl. oz. of [product][liquid].] [Each 4 fl. oz. [cartridge][bottle] of concentrate makes one gallon of [liquid][ready to use] spray.]
5. Insert the [cartridge][bottle][herbicide] into the top of the sprayer and twist until tight.

5/11

6. Press the trigger to spot spray weeds as listed on the product label.
7. When finished spot treating remove the [cartridge][bottle][-herbicide]. Replace the cap on the [cartridge][bottle][-herbicide]. Follow label Storage and Disposal instructions for product [herbicide][cartridge][bottle].
8. After spot spraying rinse the [sprayer][wand][applicator] according to the RINSE: USE DIRECTIONS.

RINSE: USE DIRECTIONS

1. Remove the [cartridge][bottle][-herbicide]. Ensure that there is water in the one gallon tank.
2. Press and hold the [trigger][sprayer] for 10 seconds to flush the [sprayer][wand][applicator] with clean water from the tank.

CHANGE [CARTRIDGE][BOTTLE]: USE DIRECTIONS

1. Remove the [cartridge][bottle][-herbicide] currently loaded in the [AMDRO Spray Delivery System Name] .
2. Follow label Storage and Disposal instructions for product [herbicide][cartridge][bottle].
3. Press and hold the [trigger][sprayer] for 10 seconds to flush the [sprayer][wand][applicator] with clean water from the tank.
3. Select a new [cartridge][bottle][-herbicide] labeled for intended use. READ THE ENTIRE PRODUCT LABEL PRIOR TO USE.
4. Insert the [cartridge][bottle][-herbicide] into the top of the sprayer and twist until tight.
5. Press the trigger to spot spray weeds as listed on the product label.
6. When finished spot treating remove the [cartridge][bottle][-herbicide]. Replace the cap on the [cartridge][bottle][-herbicide]. Follow label Storage and Disposal instructions for [product] [herbicide][cartridge][bottle].
7. After spot spraying, rinse the [sprayer][wand][applicator] according to the RINSE: USE DIRECTIONS. Store the [AMDRO Spray Delivery System Name] OR RINSE and change [cartridge][bottle].

NOTE: IF OUTPUT SPRAY IS DIMINISHED OR SPUTTERING, DO THE FOLLOWING:

1. Check that there is water remaining in the one gallon tank. If not, refill the tank with water.
2. Check that there is liquid remaining in the [cartridge][bottle]. If not, change to a new [cartridge][bottle].

[Two [cartridges][bottles] can be stored on the back of the [AMDRO Spray Delivery System Name]] .

[Optional:]CHANGING BATTERIES:]

If sprayer does not operate or spray performance declines significantly, change the four AA alkaline batteries in the sprayer handle as follows:

1. Remove the [cartridge][bottle].
2. Slide the battery cover open in the underside of sprayer handle.
3. Allow battery cover to swing open.
4. Remove old batteries.
5. Replace with four AA alkaline batteries. Make sure the positive and negative ends match the direction indicated on the inside of the battery door.
6. Close battery door and slide the door until latch is closed.

Hose End Sprayer Applications

1. Shake Well.
2. Make sure control lever is in the **OFF** position. Attach to hose.
3. Turn on water.

Spray

1. Aim towards direction of application and away from you.

6/11

2. Turn control lever to **ON** to begin spraying.
3. Spray evenly and as directed for specified type of treatment area.

Stop

1. When finished turn control lever to **OFF**.
2. Turn faucet **OFF**. Discharge residual water pressure by turning control lever **ON** and **OFF** again.
3. Disconnect Sprayer and Store / Discard as instructed.

APPLICATION RESTRICTIONS AND PRECAUTIONS

For best results, apply when weeds are less than 3 inches high.

Try to avoid overlap spraying.

Do not spray over-the-top of flowers, ornamentals (including herbaceous) or garden vegetables.

Do not use on turfgrasses other than those listed on the label.

Do not use in greenhouses.

Do not graze or feed treated grass clippings.

Do not apply to newly planted, plugged, or sodded turfgrass.

Use only on well established non-stressed turfgrass. Applications to stressed plants (turfgrass or target weeds) may result in temporary discoloration or poor weed control. Allow turfgrass to develop good root system before applying.

If discoloration occurs, treated turfgrass will recover with new growth and discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turfgrass growth.

Wait at least 4 weeks after treatment before reseeding, winter overseeding or plugging lawns.

Avoid applying when windy to prevent product drifting to other plants.

Do not apply directly to or near water, storm drains, or drainage ditches.

Care should be used when spraying to avoid fish and reptile pets in/around ornamental ponds.

CONTROL OF SEDGES (INCLUDING NUTSEDGES) AND GRASSES

Spray directly on emerged weeds to control or suppress sedges and some grasses. F7214 3.2 RTS Herbicide is effective against newly emerged and established sedges and grasses. Good spray coverage is needed for optimum control.

Grass and sedge weed species controlled or suppressed by F7214 3.2 RTS Herbicide

Common Name	Scientific Name
Barnyardgrass	(<i>Echinochloa crus-galli</i>)
Broadleaf Signalgrass	(<i>Brachiaria platyphylla</i>)
Crabgrass, Large	(<i>Digitaria sanguinalis</i>)
Crabgrass, Smooth	(<i>Digitaria ischaemum</i>)
Deergrass	(<i>Panicum clandestinum</i>)
Goosegrass	(<i>Eleusine indica</i>)
Foxtail, giant	(<i>Setaria faberi</i>)
Foxtail, green	(<i>Setaria viridis</i>)
Foxtail, yellow	(<i>Setaria glauca</i>)
Kyllinga, cocks-comb	(<i>Kyllinga squamulata</i>)
Kyllinga, green	(<i>Kyllinga brevifolia</i>)
Kyllinga, false green	(<i>Kyllinga gracillima</i>)
Kyllinga, fragrant	(<i>Kyllinga sesquiflorus</i>)
Kyllinga, tufted	(<i>Kyllinga pumila</i>)
Japanese stiltgrass	(<i>Microstegium vimineum</i>)
Junglerice	(<i>Echinochloa colona</i>)
Nutsedge, purple	(<i>Cyperus rotundus</i>)
Nutsedge, yellow	(<i>Cyperus esculentus</i>)
Sedge, annual	(<i>Cyperus compressus</i>)
Sedge, globe	(<i>Cyperus globulosus</i>)
Sedge, cylindrical	(<i>Cyperus retrorsus</i>)

Sedge, Surinam	(<i>Cyperus surinamensis</i>)
Sedge, Texas	(<i>Cyperus polystachyos</i>)

For optimal control of Goosegrass, Junglerice, or Purple Nutsedge, repeat treatment every 3-4 weeks.

CONTROL OF ANNUAL, BIENNIAL & PERENNIAL BROADLEAF WEEDS

Spray directly on newly emerged weeds to control or suppress a wide range of broadleaf weeds.

Broadleaf weeds controlled or suppressed by F7214 3.2 RTS Herbicide

Common Name	Scientific Name
Bedstraw, catchweed	(<i>Galium aparine</i>)
Beggarticks	(<i>Bidens frondosa</i>)
Beggarweed, Florida	(<i>Desmodium tortuosum</i>)
Bindweed, field	(<i>Convolvulus arvensis</i>)
Bittercress	(<i>Cardamine spp.</i>)
Black medic	(<i>Medicago lupuline</i>)
Buttercups	(<i>Ranunculus spp.</i>)
Buttonweed, common [poorjoe]	(<i>Diodia teres</i>)
Carolina geranium	(<i>Geranium carolinianum</i>)
Carpetweed	(<i>Mollugo verticillata</i>)
Chickweed, common	(<i>Stellaria media</i>)
Chickweed, mouseear	(<i>Cerastium vulgatum</i>)
Cinquefoil	(<i>Potentilla spp.</i>)
Clover	(<i>Trifolium spp.</i>)
Clover, alyce	(<i>Alysicarpus ovalifolius</i>)
Clover, hop	(<i>Trifolium campestre</i>)
Clover, red	(<i>Trifolium pretense</i>)
Clover, white	(<i>Trifolium repens</i>)
Cudweed	(<i>Gnaphalium spp.</i>)
Daisy, English	(<i>Bellis perennis</i>)
Dandelion	(<i>Taraxacum officianale</i>)
Dandelion, false	(<i>Microseris cuspidate</i>)
Dayflower, common	(<i>Commelina diffusa</i>)
Dock, Curly	(<i>Rumex crispus</i>)
Dollarweed [pennywort]	(<i>Hydrocotyle spp.</i>)
Eclipta	(<i>Eclipta prostrata</i>)
Evening primrose	(<i>Oenothera biennis</i>)
Fiddleneck	(<i>Amsinckia spp.</i>)
Filaree	(<i>Erodium spp.</i>)
Filaree, redstem	(<i>Erodium cicutarium</i>)
Galinsoga	(<i>Galinsoga ciliate</i>)
Goldenrod	(<i>Solidago spp.</i>)
Ground ivy	(<i>Glechmea hederacea</i>)
Groundsel, common	(<i>Senecio vulgaris</i>)
Henbit	(<i>Lamium amplexicaule</i>)
Knawel	(<i>Scleranthus annuus</i>)
Knotweed, prostrate	(<i>Polygonum aviculare</i>)
Kochia	(<i>Kockia scoparia</i>)
Lambsquarters, common	(<i>Chenopodium album</i>)
Lawn burweed (Spurweed)	(<i>Soliva pterosperma</i>)
Lespedeza, common	(<i>Lespedeza striafa</i>)
London Rocket	(<i>Sisymbrium irio</i>)
Mallow, common	(<i>Malva neglecta</i>)
Morningglory	(<i>Ipomea spp.</i>)
Morningglory, entireleaf	(<i>Ipomoea hederacea var. integriuscula</i>)
Morningglory, ivyleaf	(<i>Ipomoea hederacea</i>)

8/11

Morningglory, palmleaf	(<i>Ipomoea wrightii</i>)
Morningglory, pitted	(<i>Ipomoea lacunose</i>)
Morningglory, tall	(<i>Ipomoea purpurea</i>)
Parsley piert	(<i>Alchemilla arvensis</i>)
Pigweed, Redroot	(<i>Amaranthus retroflexus</i>)
Pigweed, Smooth	(<i>Amaranthus hybridus</i>)
Pigweed, Tumble	(<i>Amaranthus albus</i>)
Pineapple weed	(<i>Matricaria matricariodes</i>)
Plantain, Broadleaf	(<i>Plantago comanalata</i>)
Plantain, Buckhorn	(<i>Plantago lanceolata</i>)
Plantain, Narrowleaf	(<i>Plantago narrowiforma</i>)
Puncturevine	(<i>Tribulus terrestris</i>)
Purslane, common	(<i>Portulaca oleracea</i>)
Pusley, Florida	(<i>Richardia scabra</i>)
Redweed	(<i>Melochia corchorifolia</i>)
Shepherd's-Purse	(<i>Capsella bursa-pastoris</i>)
Smartweed, Pennsylvania	(<i>Polygonum pennsylvanicum</i>)
Sorrel, Red	(<i>Rumex acetosella</i>)
Speedwell	(<i>Veronica spp.</i>)
Speedwell, slender	(<i>Veronica filiformis</i>)
Speedwell, thymeleaf	(<i>Veronica serpyllifolia</i>)
Spurge, (annuals)	(<i>Euphorbia spp.</i>)
Spurge, prostrate	(<i>Euphorbia humistrata</i>)
Spurge, spotted	(<i>Euphorbia maculate</i>)
Star-of-Bethlehem	(<i>Ornithogalum umbellatum</i>)
Velvetleaf	(<i>Abutilon theophrasti</i>)
Violet, wild	(<i>Viola pratensis</i>)
Violet, Johnny-Jump-Up	(<i>Viola rafeinesquii</i>)
Wild garlic	(<i>Allium vineale</i>)
Wild onion	(<i>Allium canadense</i>)
Woodsorrel, creeping	(<i>Oxalis corniculata</i>)
Woodsorrel, yellow	(<i>Oxalis stricta</i>)

Certain listed weeds may require more than one application to achieve season-long control. If listed weeds are not controlled by the initial application, apply a second application 4 to 6 weeks after the initial treatment.

STORAGE AND DISPOSAL

Pesticide Storage

Keep from freezing. Do not store below 32°F. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat.

Pesticide Disposal and Container Handling

Non-refillable container. Do not reuse or refill this container. If empty: Offer for recycling, if available, or dispose of in trash.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Notice: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.

[Spanish translation]

[on last page of booklet label attached to bottle, also attach the following]

[icon] Questions, Comments or Medical Information

Optional Marketing Claims:

- *Kills Crabgrass, Dandelion, Ground Ivy, Yellow & Purple Nutsedge, Kyllinga and over XXX other tough weeds*
- *Selective Postemergence Control of Annual Grasses, Broadleaf weeds, and Sedges in Residential Lawns*
- *Rainproof in 2 hours*
- *1 quart treats [up to] [covers] 4,000 sq. ft.*
- *100% Satisfaction Guaranteed* (or your money back)*
- *All-in-One Lawn Weed Killer*
- *All-in-One Weed Killer*
- *All-in-One Weed Killer for Lawns*
- *Also Kills Dandelion and Clover Plus XXX Other Listed Weeds*
- *Attach to [garden] hose and spray*
- *Consistently fast control on tough weeds like dandelion, spurge, clover, plantain, ground ivy (and various other listed weeds).*
- *Control lawn weeds plus crabgrass in your lawn with ease!*
- *Controls XXX weeds!*
- *Controls crabgrass, dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and many other listed broadleaf lawn weeds*
- *Controls crabgrass, foxtail and signalgrass*
- *Controls crabgrass, yellow nutsedge, dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and many other listed broadleaf lawn weeds.*
- *Controls even the tough weeds - To the Root!**
 - **Listed weeds*
- *Controls even the tough weeds ... roots and all!**
 - **Listed weeds*
- *[Double [Dual] [Two-Way] Action] - Kills Broadleaf and Crabgrass Weeds**
 - **Listed weeds*
- *Dual Action Weed Control - Kills the Plant and the Root*
- *Dual Action Killing – uptake by both leaves and roots*
- *Easy & Fast Coverage*
- *Easy to mix*
- *Easy to use*
- *Effective [Broadleaf] Weed Protection*
- *Effectively Kills Both Broadleaf and certain weed grasses*
- *Effectively Protects Lawns from Broadleaf Weeds*
- *Fast Coverage*
- *Fast Results*
- *For home lawn care*
- *Gets the Root*
- *Guaranteed* Not to Harm Lawns*
- *Guaranteed* Not to Harm Your Lawn*
- *Guaranteed* Results*
- *Ideal for Spot Treatment*
- *Kill crabgrass, yellow nutsedge & [any listed weed(s)] too!*
- *Kills [Controls] XXX broadleaf weeds.*
- *Kills [Dandelion], [Crabgrass], and [Clover], Plus [XXX] Other listed Weeds.*
- *Kills [Hard-to-Control] [Tough] Weeds*
- *Kills XXX Weeds (to the root)*
- *Kills All Major Broadleaf Weeds*
- *Kills Both Broadleaf and Crabgrass Weeds*
- *Kills Both Broadleaf and Grass Weeds*
- *Kills both Crabgrass & Broadleaf weeds*
- *Kills Crabgrass & Other Weeds in Your Lawn*
- *Kills Crabgrass plus Broadleaf Weeds*
- *Kills Even Tough Weeds - roots & all*
- *Kills even tough weeds - To the Root!*

10/11

- *Kills Hard-to-Control Weeds*
- *Kills Listed Lawn Weeds Plus Crabgrass*
- *Kills Listed Lawn Weeds plus Crabgrass in One (Easy) Step*
- *Kills listed lawn weeds plus crabgrass*
- *Kills Listed Weeds*
- *Kills Over XXX (Lawn) Weeds (Plus Crabgrass in One Easy Step)*
- *Kills Over XXX Weeds!*
- *Kills (to) the root*
- *Kills the Weed, Kills the Root*
- *Kills the weeds, not the lawn*
- *Kills Weeds Fast*
- *Mixes with Water - Concentrate*
- *No messy cleanup*
- *Save Time & Energy*
- *Simple to use*
- *Starts Working Immediately*
- *Starts Working on Contact*
- *Systemic [Herbicide] [Activity]*
- *Targets Weeds*
- *Two-in-One Product that Kills Crabgrass & Broadleaf Weeds*
- *Visible Results*
- *Visible results in [less than] 24 hours*
- *Visible results in [less than] a day*
- *Visible results overnight*
- *Visible results in [XX] hours*
- *Water based [formula]*
- *Won't Harm Lawns (when used as directed)*
- *Kill Crabgrass, Clover, Plantain, Chickweed and other listed weeds and sedges*
- *Controls Listed Annual Grasses, Broadleaf Weeds and Sedges*
- *Kills Tough Weeds [like] [Nutsedge] & [Crabgrass]*
- *Kills [Nutsedge] & [Crabgrass] in Warm and Cool Season Turf Grasses*
- *Kills [Nutsedge] & [Crabgrass] in Northern and Southern Turf Grasses*
- *Kills [Nutsedge] & [Crabgrass] in Northern Turf Grasses*
- *Kills [Nutsedge] & [Crabgrass] in Cool Season Turf Grasses*
- *Kills [Nutsedge] & [Crabgrass] in Southern Turf Grasses*
- *Kills [Nutsedge] & [Crabgrass] in Warm Season Turf Grasses*
- *[Nutsedge] & [Crabgrass] Killer*
- *[Nutsedge] & [Crabgrass] Killer in Warm and Cool Season Turf Grasses*
- *[Nutsedge] & [Crabgrass] Killer in Northern and Southern Turf Grasses*
- *[Nutsedge] & [Crabgrass] Killer in Cool Season Turf Grasses*
- *[Nutsedge] & [Crabgrass] Killer in Northern Turf Grasses*
- *[Nutsedge] & [Crabgrass] Killer in Warm Season Turf Grasses*
- *[Nutsedge] & [Crabgrass] Killer in Southern Turf Grasses*
- *[Nutsedge] & [Crabgrass] Killer for broadcast application*
- *For Northern and Southern Turf Grasses*
- *For Northern Turf Grasses*
- *For Warm and Cool Season Turf Grasses*
- *For Cool Season Turf Grasses*
- *Results are visible in days*
- *Complete kill occurs in 2-3 weeks*
- *Weed is yellowed in 1-2 days*
- *Connect and Spray Applicator – No mixing required*
- *Ready to Spray – no mixing*
- *Kills Roots, Shoots and nutlets*
- *Kills Tough Weeds, such as Purple and Yellow Nutsedge, Kyllinga, and other listed Sedges*
- *Also kills Plantain, Clover, Spurge, Woodsorrel, Knotweed, and other listed annual, biennial, and perennial broadleaf weeds*
- *Selective Herbicide for use in Northern Turf Grasses*

11/11

- *Selective Herbicide for use in Northern & Southern Turf Grasses*
- *Apply to growing and dormant grasses*
- *Uptake by both weed roots and shoots*
- *Dual action killing: Uptake by both leaves and roots*
- *Contains XXX Cartridges*
- *Makes up to XX gallons*
- *Kills Tough Weeds like [Nutsedge], [Crabgrass], [Dandelion], [Wild Onion], [Dollarweed]*
- *Kills purple and yellow Nutsedge*
- *[up to] 80% reduction in future weed populations**
 - **Listed weeds*
- *Activated by sunlight*
- *Great for [weed] resistance management{XXX}+ Weeds Controlled*
- *More than XXX weeds controlled*
- *Greater than XXX weeds controlled*
- *Over XXX weeds controlled*
- *Kills Nutsedge*
- *Nutsedge Killer*
- *[Also] Kills Dollarweed, Wild Onion, Sandbur, Annual Bluegrass*
- *Can be used treat listed weeds in dormant Bermudagrass*
- *Kills tough weeds in dormant Bermudagrass*
- *Kills summer and winter weeds*
- *For use on [Bermudagrass], [Zoysiagrass], [Tall Fescue], [Kentucky Bluegrass] and other listed grasses*
- *Kills [nutsedge], [chickweed], [white clover], [dandelion], [kyllinga] and other listed weeds*